



Additions to the Flora of China: seven new species of *Elatostema* (Urticaceae) from the karst landscapes of Guangxi and Yunnan

Y.G. WEI¹, A.K. MONRO² & W.T. WANG³

¹Guangxi Key Laboratory of Functional Phytochemicals Research and Utilization, Guangxi Institute of Botany, Chinese Academy of Sciences, Guilin 541006, China. E-mail: weiyigang@yahoo.com.cn

²Department of Botany, The Natural History Museum, London, SW7 5BD, United Kingdom. E-mail: alm@nhm.ac.uk (corresponding author)

³State Key Laboratory of Systematic and Evolutionary Botany, Institute of Botany, Chinese Academy of Sciences, Beijing 100093, China.

Abstract

Seven new species of *Elatostema* from southwestern China are described and illustrated. Affinities of the species are discussed and Global Species Conservation Assessments presented. The new species are: *E. celingense* (Critically Endangered) which most closely resembles *E. backeri*, *E. hezhouense* (Critically Endangered) which most closely resembles *E. ichangense*, *E. lui* (Vulnerable) which most closely resembles *E. platyceras*, *Elatostema multicaule* (Vulnerable) which most closely resembles *E. asterocephalum*, *E. nianbense* (Vulnerable) which most closely resembles *E. filipes*, *E. retrorstrigulosum* (Endangered) which most closely resembles *E. balansae* and *E. yachense* (Vulnerable) which most closely resembles *E. xinningense*. All of the new species are endemic to China.

Key words: cave flora, limestone karst, Global Conservation Assessments

Introduction

Elatostema Forster & Forster (1775: 53; Urticaceae) consists of several hundred species of understory herbs and subshrubs that occur in the deep shade of forests, stream sides, gorges and caves. *Elatostema* occurs throughout tropical and subtropical Africa, Asia, Australia, and Oceania but is notably absent from the Neotropics. In Southeast Asia the genus is most diverse on limestone karst (Yahara 1984, Lahav-Ginott & Cronk 1993, Wang & Chen 1995, Qi *et al.* 2003).

Elatostema forms part of a complex of genera, which includes *Elatostematoides*, *Pellionia* and *Procris* whose generic limits are difficult to define. Of these, *Elatostema* is the oldest name (Forster 1775). *Elatostematoides* has not been widely recognized or used since its description by Robinson (1911) and is treated as a synonym of *Elatostema* (Friis 1989). There is however little consensus as to the delimitation of *Elatostema*, *Pellionia* and *Procris* and disagreement has centred on the use and interpretation of two main suites of characters, inflorescence branching morphology and staminate flower perianth morphology. The recent incorporation of DNA sequence data from the plastid genome (Hadijah *et al.* 2003) is limited by small taxon sampling and has done little to resolve generic delimitation. Hadijah *et al.* (2003) recovered the complex as monophyletic but the genera within it paraphyletic. Monro *et al.* (in prep.) recovered *Elatostema* and a combined *Procris* and *Pellionia* as monophyletic. The taxa described here would be considered *Elatostema sensu stricto*. To date, 998 species names of the *Elatostema* complex of genera (*Elatostema*, *Elatostematoides*, *Pellionia* and *Procris*) have been published (IPNI 2008) and current published species estimates range from 250 (Chew 1989) to 400 species (Yahara 1984). In China 146 species are currently recognized (Qi *et al.* 2003). Assuming levels of synonymy of only 10% as for *Pilea* (Monro 2004), a genus with similar habit and

Acknowledgements

We like to acknowledge the support of the Innovation Fund of the Botany Department of The Natural History Museum, a Travel Grant from the Royal Society and the Science Research Foundation of Guangxi Institute of Botany (No. 10004) supporting the fieldwork. We also like to thank our driver Wei-Hua Li and local botanist Shi Nian Lu from Jingxi County, who showed us many interesting and rewarding localities. We thank Sun Ying-Bao and Ruth Harker for the illustrations and Norman Robson (BM) for help with the Latin diagnoses.

References

- Bourdy, A. & Walter, G. (1992) Maternity and medicinal plants in Vanuatu I. The cycle of reproduction. *Journal of Ethnopharmacology* 37: 179–196.
- Chew, W.-L. (1989) *Urticaceae*. In: *Flora of Australia*, vol. 3. Australian Government Publishing Service, Canberra. Pp. 68–93.
- Forster, J.R. & Forster, J.G.A. (1775) *Characteres Generum Plantarum*, second edition. London.
- Gagnepain, F. (1929) Quelques *Elatostema* et un *Procris* nouveaux d'Indo-Chine. *Bulletin de la Société Botanique de France* 76: 80–82.
- Hadijah, J.T., Quinn, C.J. & Conn, B.J. (2003) Phylogeny of *Elatostema* (Urticaceae) using chloroplast DNA data. *Telopea* 10: 235–246.
- He, S., Chen, L., Zhag, T., Fan, G., Pie, Q., Huang, Y. & Ran, M. (1991) New record of distribution of medicinal plants Guizhou Province China. *Journal of Chinese Materia Medica* 16: 643–701.
- IPNI [International Plant Names Index] (2008) <http://www.ipni.org> [accessed 13 March 2010]
- IUCN [International Union for Conservation of Nature and Natural Resources Species Survival Commission] (2001) *IUCN Red List Categories: Version 3.1*. International Union for Conservation of Nature and Natural Resources, Gland and Cambridge, pp. 70.
- Johannes, A. (1975) Medicinal plants of the nekematigi of the eastern highlands of New Guinea. *Economic Botany* 29: 268–277.
- Lahav-Ginott, S. & Cronk, Q.C.B. (1993) The mating system of *Elatostema* (Urticaceae) in relation to morphology: a comparative study. *Plant Systematics and Evolution* 186: 135–145.
- Monro, A.K. (2004) Three new species, and three new names in *Pilea* (Urticaceae) from New Guinea. *Kew Bulletin* 59: 573–579.
- Nandwani, D., Calvo, J.A., Tenorio, J., Calvo, F. & Manglona, L. (2008) Medicinal plants and traditional knowledge in the Northern Mariana Islands. *Journal of Applied Biosciences* 8: 323–330.
- Qi, L., Friis, I. & Wilmot-Dear, M.C. (2003) *Elatostema*. In: Wu, Z. & Raven, P.H. [eds.], *Flora of China*, vol. 5: 76–189. Science Press, Beijing and Missouri Botanical Garden Press, St. Louis, Missouri.
- Schroeter, H. & Winkler, H. (1936) Monographie der Gattung *Elatostema* s.l. *Repertorium Specierum Novarum Regni Vegetabilis* 83: 155–174.
- Schroeter, H. (1939) Nonnullae descriptiones specierum novarum generis *Elatostema* subgen *Eulatostema*. *Repertorium Specierum Novarum Regni Vegetabilis* 47: 220–221.
- Wang, W.T. & Chen, C.J. (1995) *Elatostema*. In: *Flora Reipublicae Popularis Sinicae* 23. Science Press, Beijing. Pp.187–317.
- Wang, W.T. (1980) Classificatio specierum Sinicarum *Elatostematis* (Urticaceae). *Bulletin of Botanical Laboratory of North-Eastern Forestry Institute* 7: 40–41, 49–50.
- Wang, W.T. (1985) Two new species of *Elatostema* (Urticaceae) from Hunan. *Guihaia* 5: 323–325.
- Wang, W.T. (1988) Notulae de *Elatostemate* Yunanensi. *Acta Botanica Yunnanica* 10: 343–348.
- Wang, W.T. & Wei, Y.G. (2007) Five new species of *Elatostema* J. R. et G. Forst. (Urticaceae) from Guangxi, China. *Guihaia* 27: 816.
- Wei, Y.G. & Wang, W.T. (2009) Notes on *Elatostema* J.R. et G. Forst. (Urticaceae) from Guangxi. *Guihaia* 29: 711–718.
- Yahara, T. (1984) *Pellionia* and *Elatostema* in Thailand. Taxonomic studies of Urticaceae II. *Journal of the Faculty of Science of the University of Tokyo III*, 13: 483–499.