





http://dx.doi.org/10.11646/phytotaxa.159.4.5

# Salacca acehensis (Arecaceae), A New Species from Sumatra, Indonesia

# ZUMAIDAR<sup>1,2</sup>, TATIK CHIKMAWATI<sup>3</sup>, ALEX HARTANA<sup>3,4</sup>, SOBIR<sup>5</sup>, JOHANIS P. MOGEA<sup>6</sup> & FINN BORCHSENIUS<sup>7</sup>

<sup>1</sup>Plant Biology Graduate Program, Department of Biology, Faculty of Mathematics and Natural Sciences, Bogor Agricultural University, Darmaga Campus, Bogor 16680, Indonesia.

<sup>2</sup>Department of Biology, Mathematics and Sciences Faculty, Syiah Kuala University, Jl. Syech Abdur Rauf, Darussalam Banda Aceh, Indonesia. Email: zumaidar@yahoo.com

<sup>3</sup>Department of Biology, Faculty of Mathematics and Natural Sciences, Bogor Agricultural University, Darmaga Campus, Bogor 16680, Indonesia. Email: tchikmawati@yahoo.com

<sup>5</sup>Department Agronomy and Horticulture, Bogor Agricultural University, Darmaga Campus, Bogor 16680, Indonesia Email: sobir@ipb.ac.id

<sup>6</sup>Herbarium Bogoriense, Botany Division, Research Center for Biology, Indonesian Institute of Sciences, Jl Raya Jakarta Bogor Km 47, Cibinong, Indonesia.

<sup>7</sup>Science Museums, Aarhus University, Building 1137 Universitetsparken, DK-8000 Aarhus C, Denmark. Email: finn.borchsenius@biology.au.dk.

#### Abstract

A new species of *Salacca* sect. *Salacca* from Sumatra is described and illustrated. It is characterized by dwarf habit, with leaves to 2.5 m long; erect and slender staminate inflorescences to 7 cm long; and small infructescences, to 10 cm long, with up to 3 branches The most similar species, *S. rupicola* from Borneo, is different in having the apical leaflets composed ca. 8 united segments versus 2 in the new species, and in having larger inflorescences, the staminate to 15 cm long and the pistillate to 40 cm long.

Key words: Aceh, dwarf salak, Gunung Leuser National Park, section Salacca

## Introduction

*Salacca* Reinwardt (1825: 3; Arecaceae, subfamily Calamoideae, tribe Calameae) comprises 22 species (Govaerts et al. 2013) distributed from Burma and Indochina south and eastwards to Borneo, Java and the southern most part of the Philippines (Dransfield et al. 2008) with the largest number of species and the greatest morphological diversity found in the Malay Peninsula and Borneo. Although there are widespread species, such as *S. affinis* Griffith (1845: 9) and *S. zalacca* Voss (1895: 1152), most of the species are restricted to very small areas (Dransfield 2009). The genus is neatly divisible into two sections (Beccari 1918; Dransfield et al. 2008): section *Salacca* with strigose ovary, hispid subspinous fruit, and pistillate rachillae bearing dyads of one fertile pistillate and one sterile staminate flower; and section *Leiosalacca* with ovaries and fruits covered in smooth scales, and pistillate rachillae bearing solitary pistillate flowers. Of the 22 species currently recognized, 20 belong to section *Salacca* and two to section *Leiosalacca* (Dransfield et al. 2008).

In this paper we describe a new species first collected by J. Dransfield and D. Saerudin at Gunung Kemiri, Southeast Aceh Regency, Aceh Province in 1971. The species was later recollected by one of the authors of this paper (JM) who annotated the specimens with the name *S. acehensis*. However, the name remained unpublished. In this paper we describe the new species and compare it to other small species of section *Salacca*.

<sup>&</sup>lt;sup>4</sup>*Email: ahartana@indo.net.id* 

#### Acknowledgements

We are grateful to John Dransfield, William J. Baker, and Lauren Gardiner at the Royal Botanic Gardens, Kew for their help and advice in connection with this study. The first author would like to thank Dikti (the Directorate General of Higher Education, Indonesia) for sponsoring her study and her Sandwich Program 2012 at Aarhus University, Denmark; Mien A. Rifai as her supervisors at Bogor Agricultural University for his wise guidance, advice, attention, patience and support during her study; Peter van Welzen, G. Thijsse, and P.A.J. Audiffred at Leiden Herbarium for making specimens available; Himmah Rustiami at Herbarium Bogoriense for kind help with her study. Thanks are also due to Anne Kusuma for making the drawing.

## References

Beccari, O. (1918) Asiatic palms—Lepidocaryeae. Part III. *Annals of the Royal Botanic Garden* 12(2). The Bengal Secretariat Book Depot, Calcutta. 231 pp.

Dransfield, J. (1980) A new species of *Salacca* Reinw. from the limestone of the Gunung Mulu National Park. *Botanical Journal of Linnean Society* 81: 36–37.

Dransfield, J. (2009). A new species of Salacca from Sarawak. Palms 53: 167-170.

Dransfield, J., Uhl, N.W., Asmussen-Lange, C.B., Baker, W.J., Harley, M.M. & Lewis, C.E. (2008) *Genera Palmarum. The Evolution and Clasification of Palms.* Royal Botanical Gardens, Kew, UK.732 pp.

Furtado, C. X., 1949. The Malayan species of Salacca. Gardens Bulletin Singapore 12: 378-403.

- Govaerts, R., Dransfield, J., Zona, S.F., Hodel, D.R. & Henderson, A. (2013) *World checklist of Arecaceae*. Facilitated by the Royal Botanical Gardens, Kew. Published on the internet; http://apps.kew.org/wcsp/ Retrieved 04 June 2013.
- Griffith, W. (1845) The palms of British East India. *Calcutta journal of natural history, and miscellany of the arts and sciences in India* 5: 1–103.

Hodel, D.R. (1997) New species of palms from Thailand. Palm Journal 134: 28-37.

- Mogea, J.P. (1980) The flabellate-leaved species of Salacca (Palmae). Reinwardtia 9: 461-479.
- Mogea, J.P. (1984) Three new species of Salacca (Palmae) from the Malay Peninsula. *Federation Museums Journal (Kuala Lumpur)* 29: 1–19.
- Reinwardt, C.G.C. (1825) In C. F. Hornschuch, Sylloge Plantarum Novarum Itemque Minus Cognitarum a Praestantissimis Botanicis adhuc Viventibus Collecta et a Societate Regia Botanica Ratisbonensi Edita. Vol. 2. Societate Regia Botanica, Regensburg.
- Voss, A. (1895) Vilmorin's Blumengärtnerei Beschreibung, Kultur und Verwendung des Gesamten Pflanzenmaterials fur Deutsche Garten. Dritte, neubearbeite Auflage. Part 1. Berlin. http://dx.doi.org/10.5962/bhl.title.67392