



Remarkable diversity of the lichen family *Graphidaceae* in the Amazon rain forest of Rondônia, Brazil

MARCELA E. S. CÁCERES¹, ANDRÉ APTROOT², SITTIPORN PARNMEN³ & ROBERT LÜCKING⁴

¹Departamento de Biociências, Universidade Federal de Sergipe, CEP: 49500-000, Itabaiana, Sergipe, Brazil;
email: mscaceres@hotmail.com

²ABL Herbarium, G.v.d. Veenstraat 107, NL-3762 XK Soest, The Netherlands; email: andreaptroot@gmail.com

³Toxicology and Biochemistry Section, Department of Medical Sciences, Ministry of Public Health, Nonthaburi 11000 Thailand;
email: sparnmen@gmail.com

⁴Science & Education, The Field Museum, 1400 South Lake Shore Drive, Chicago, IL 60605-2496, U.S.A.;
email: rlucking@fieldmuseum.org

Abstract

In a continuation of our investigation of lichenized fungi in Rondônia and adjacent areas, we present a preliminary treatment of the graphidoid and thelotremoid core Graphidaceae (subfamilies Fissurinoideae and Graphidoideae). A total of 122 identified species are reported here, almost all of which are new reports to Rondônia, and 37 of which are new to science. This includes three new, monospecific genera, viz. *Aggregatorygma triseptatum* M. Cáceres, Aptroot & Lücking, a new, phylogenetically distinct genus and species similar to *Diorygma* but with corticate thallus, aggregated and branched lirellae, very small, 3-septate ascospores and unknown secondary substances; *Byssotrema mirabile* M. Cáceres, Aptroot & Lücking, a new genus and species similar to *Glaucotrema* but with partially carbonized excipulum with pilose inner margin and cinchonarum unknown chemistry; and *Corticorygma stellatum* M. Cáceres, Feuerstein, Aptroot & Lücking, a new, phylogenetically distinct genus and species similar to *Diorygma* but with non-amyloid ascospores and corticate thallus. The following further species are described as new to science: *Cruentotrema amazonum* M. Cáceres, Aptroot & Lücking, differing from *Cruentotrema kurandense* in the 3-septate ascospores; *Fissurina amazonica* M. Cáceres, Aptroot & Lücking, differing from *F. dumastii* in the small, closed, much branched and dense lirellae and the apically smooth paraphyses; *F. amyloidea* M. Cáceres, Aptroot & Lücking, differing from *F. subnitidula* in the weakly carbonized lirellae and thick-walled, strongly amyloid ascospores; *F. chrysocarpa* M. Cáceres, Aptroot & Lücking, differing from *F. chrysocarpoides* in the short lirellae with distinct labia; *F. duplicans* M. Cáceres, Aptroot & Lücking, differing from *F. pseudostromatica* in the endoperidermal thallus and double margin of the lirellae; *F. macrospora* M. Cáceres, Aptroot & Lücking, differing from *F. undulata* in the much larger ascospores; *F. subfurfuracea* M. Cáceres, Aptroot & Lücking, differing from *F. furfuracea* in the thin margin of the lirellae (distinctly fissurinoid rather than hemithecioid); *Glaucotrema stegoboloides* M. Cáceres, Aptroot & Lücking, differing from *G. glaucophaenum* in the papillose thallus and complex columella; *Graphis amazonica* M. Cáceres, Aptroot & Lücking, differing from *G. pitmanii* in the interspersed hymenium and larger ascospores; *G. pustulosa* M. Cáceres, Aptroot & Lücking, differing from *G. hyphosa* in the pustulate thallus and larger ascospores; *G. rondoniana* M. Cáceres, Aptroot & Lücking, differing from *G. pinicola* in the pruinose labia and smaller ascospores; *Gyrotrema flavum* M. Cáceres, Aptroot & Lücking, differing from *G. sinuosum* in the yellow apothecial disc; *Myriotrema foliaceum* M. Cáceres, Aptroot & Lücking, differing from *M. rugiferum* in the gall-forming thallus; *M. inspersum* M. Cáceres, Aptroot & Lücking, differing from *M. foliicola* in the hyaline, smaller ascospores; *M. subclandestinum* M. Cáceres, Aptroot & Lücking, differing from *Myriotrema clandestinum* in the larger ascospores with more numerous septa; *Ocellularia brasiliensis* M. Cáceres, Aptroot & Lücking, differing from *O. africana* in the carbonized excipulum and columella and cinchonarum unknown chemistry; *O. diminuta* M. Cáceres, Aptroot & Lücking, differing from *O. papillata* in the smaller ascospores lacking a columella; *O. flavostroma* M. Cáceres, Aptroot & Lücking, differing from *O. fecunda* in the eolumellate ascospores; *O. halei* M. Cáceres, Aptroot & Lücking, differing from *O. protoinspersa* in the grey thallus, narrow columella, and shorter ascospores; *O. immersocarpa* M. Cáceres, Aptroot & Lücking, differing from *O. terebrata* in the immersed ascospores lacking carbonization; *O. lacerata* M. Cáceres, Aptroot & Lücking, differing from *O. margaritacea* in the irregularly chroodiscoid, weakly carbonized ascospores and the white medulla; *O. myriotrema* M. Cáceres, Aptroot & Lücking, differing from *M. inspersum* in the papillose



FIGURE 15. One of the sampled trees (*Ceiba samaumae*) in the Parque Municipal of Porto Velho.

References

- Adanson, M. (1763) *Familles des Plantes*. Vol. II. Paris.
<http://dx.doi.org/10.5962/bhl.title.271>
- Ahti, T. (2000) Cladoniaceae. *Flora Neotropica Monograph* 78: 1–362.

- Aptroot, A. (1997) Lichen biodiversity in Papua New Guinea, with the report of 173 species on one tree. *Bibliotheca Lichenologica* 68: 203–213.
- Aptroot, A. (1998) New lichens and lichen records from Papua New Guinea, with the description of *Crustospathula*, a new genus in the Bacidiaceae. *Tropical Bryology* 14: 25–34.
- Aptroot, A. & Cáceres, M.E.S. (2013) Pyrenocarpous lichens (except *Trypetheliaceae*) in Rondônia. *Lichenologist* 45: 763–785. <http://dx.doi.org/10.1017/s0024282913000534>
- Aptroot, A. & Cáceres, M.E.S. (2014a) New lichen species from termite nests in rainforest in Brazilian Rondônia and adjacent Amazonas. *Lichenologist* 46: 365–372. <http://dx.doi.org/10.1017/s0024282913000340>
- Aptroot, A. & Cáceres, M.E.S. (2014b) A key to the corticolous microfoliose, foliose and related crustose lichens from Rondônia, Brazil, with the description of four new species. *Lichenologist* 46(06): 763–799. <http://dx.doi.org/10.1017/S0024282914000358>
- Aptroot, A., Lücking, R., Sipman, H.J.M., Umaña, L. & Chaves, J.L. (2008) Pyrenocarpous lichens with bitunicate asci: a first assessment of the lichen biodiversity inventory in Costa Rica. *Bibliotheca Lichenologica* 97: 1–162.
- Aptroot, A., Sipman, H.J.M. & Cáceres, M.E.S. (2013) Twenty-one new species of *Pyrenula* from South America, with a note on over-mature ascospores. *Lichenologist* 45: 169–198. <http://dx.doi.org/10.1017/s0024282912000734>
- Archer, A.W. (2007) Key and checklist for the lichen family Graphidaceae (lichenised Ascomycota) in the Solomon Islands. *Systematics and Biodiversity* 5: 9–22. <http://dx.doi.org/10.1017/s1477200006002040>
- Archer, A. W. (2009). Graphidaceae. *Flora of Australia* 57: 84–194.
- Cáceres, M.E.S., Rivas Plata, E. & Lücking, R. (2012) *Malmographina*, a new genus for *Graphina malmei* (Ascomycota: Ostropales: Graphidaceae). *Lichenologist* 44: 115–120. <http://dx.doi.org/10.1017/s0024282911000697>
- Cáceres, M.E.S., Aptroot, A. & Ertz, D. (2014) Some new *Arthoniales* from the Amazon, Rondônia, Brazil. *Lichenologist* 46: 573–588. <http://dx.doi.org/10.1017/S0024282914000036>
- Clements, F.E. (1909) *The genera of fungi*. New York.
- Colinvaux, P.A., de Oliveira, P.E. & Bush, M.B. (2000) Amazonian and neotropical plant communities on glacial time-scales: the failure of the aridity and refuge hypotheses. *Quaternary Science Reviews* 19: 141–169. [http://dx.doi.org/10.1016/s0277-3791\(99\)00059-1](http://dx.doi.org/10.1016/s0277-3791(99)00059-1)
- Eschweiler, F.G. (1824) *Systema lichenum, genera exhibens rite distincta, pluribus novis adaucta*. Nürnberg.
- Fée, A.L.A. (1825) *Essai sur les cryptogames des écorces exotiques officinales*. Paris.
- Fée, A.L.A. (1874) Matériaux pour une flore lichénologique du Brésil, II. Les Graphidées. *Bulletin de la Société Botanique de France* 21: 21–32.
- Frisch, A. (2006) Contributions towards a new systematics of the lichen family Thelotremataceae I. The lichen family Thelotremataceae in Africa. *Bibliotheca Lichenologica* 92: 3–370.
- Frisch, A. & Kalb, K. (2006) Contributions towards a new systematics of the lichen family Thelotremataceae II. A monograph of Thelotremataceae with a complex structure of the columella. *Bibliotheca Lichenologica* 92: 371–516.
- Frisch, A. & Kalb, K. (2009) *Chapsa* species (Thelotremataceae) from Brazil. *Bibliotheca Lichenologica* 99: 133–142.
- Frisch, A., Kalb, K. & Grube, M. (2006) Contributions towards a new systematics of the lichen family Thelotremataceae. *Bibliotheca Lichenologica* 92: 1–556.
- Frisch, A. & Ohmura, Y. (2012) *Chaspa leprieurii*, *Ocellularia cavata* and *O. pyrenuloides* (Graphidaceae, lichenized Ascomycota) new to Japan. *Bulletin of the National Museum of Nature and Science Series B* 38: 87–92.
- Hale, M.E. (1973) Studies on the lichen family Thelotremataceae. 1. *Phytologia* 26: 413–420.
- Hale, M.E. (1974a) Morden-Smithsonian Expedition to Dominica: The lichens (Thelotremataceae). *Smithsonian Contributions to Botany* 16: 1–46. <http://dx.doi.org/10.5479/si.0081024x.16>
- Hale, M.E. (1974b) Studies on the lichen family Thelotremataceae. 2. *Phytologia* 27: 490–501.
- Hale, M.E. (1978) A revision of the lichen family Thelotremataceae in Panama. *Smithsonian Contributions to Botany* 38: 1–60. <http://dx.doi.org/10.5479/si.0081024x.38>
- Hale, M.E. (1980) Generic delimitation in the lichen family Thelotremataceae. *Mycotaxon* 11: 130–138.
- Hale, M.E. (1981) A revision of the lichen family Thelotremataceae in Sri Lanka. *Bulletin of the British Museum (Natural History), Botany Series* 8: 227–332.
- Hooghiemstra, H. & van der Hammen, T. (1998) Neogene and Quaternary development of the neotropical rain forest: the forest refugia hypothesis, and a literature overview. *Earth-Science Reviews* 44: 147–183. [http://dx.doi.org/10.1016/s0012-8252\(98\)00027-0](http://dx.doi.org/10.1016/s0012-8252(98)00027-0)
- Hue, A. (1891) Lichenes exoticos a professore W. Nylander descriptos vel recognitos et in herbario musei Parisiense pro maxima parte asservatos in ordine systematico deposuit. *Nouvelles Archives du Muséum d'Histoire Naturelle* 3, Sér. 3: 33–120. <http://dx.doi.org/10.5962/bhl.title.82290>

- Kalb, K. (2009) New taxa and new records of thelotremoid Graphidaceae. *Herzogia* 22: 17–42.
- Kalb, K., Staiger, B. & Elix, J.A. (2004) A monograph of the lichen genus *Diorygma* - a first attempt. *Symbolae Botanicae Upsalienses* 34(1): 133–181.
- Komposch, H. & Hafellner, J. (1999) List of lichenized fungi so far observed in the tropical lowland rain forest plot Surumoni (Venezuela, Estado Amazonas). *Fritschiana* 19: 1–10.
- Kraichak, E., Parmen, S., Lücking, R., Rivas Plata, E., Aptroot, A., Cáceres, M.E.S., Ertz, D., Mangold, A., Mercado-Díaz, J.A., Papong, K., Van den Broeck, D., Weerakoon, G. & Lumbsch, H.T. (2014) Revisiting the phylogeny of Ocellularieae, the second largest tribe within Graphidaceae (lichenized Ascomycota: Ostropales). *Phytotaxa* 189(1): 52–81.
<http://dx.doi.org/10.11646/phytotaxa.189.1.6>
- Lücking, R. (2008) Follicolous lichenized fungi. *Flora Neotropica Monograph* 103: 1–866.
- Lücking, R. (2009) The taxonomy of the genus *Graphis* sensu Staiger (Ascomycota: Ostropales: Graphidaceae). *Lichenologist* 41: 319–362.
<http://dx.doi.org/10.1017/s0024282909008524>
- Lücking, R. & Rivas Plata, E. (2008) Clave y guía ilustrada para géneros de Graphidaceae. *Glalia* 1: 1–41.
- Lücking, R., Archer, A.W. & Aptroot, A. (2009) A world-wide key to the genus *Graphis* (Ostropales: Graphidaceae). *Lichenologist* 41: 363–452.
<http://dx.doi.org/10.1017/s0024282909008305>
- Lücking, R., Seavey, F., Common, R.S., Beeching, S.Q., Breuss, O., Buck, W.R., Crane, L., Hodges, M., Hodgkinson, B.P., Lay, E., Lendemer, J.C., McMullin, R.T., Mercado-Díaz, J.A., Nelsen, M.P., Rivas Plata, E., Safranek, W., Sanders, W.B., Schaefer, H.P. Jr. & Seavey, J. (2011) The lichens of Fakahatchee Strand Preserve State Park, Florida: Proceedings from the 18th Tuckerman Workshop. *Bulletin of the Florida Museum of Natural History, Biological Sciences* 49: 127–186.
- Lumbsch, H.T., Ahti, T., Altermann, S., Amo De Paz, G., Aptroot, A., Arup, U., Bárcenas Peña, A., Bawingan, P.A., Benatti, M.N., Betancourt, L., Björk, C.R., Boonpragob, K., Brand, M., Bungartz, F., Cáceres, M.E.S., Candan, M., Chaves, J.L., Clerc, P., Common, R., Coppins, B.J., Crespo, A., Dal Forno, M., Divakar, P.K., Duya, M. V., Elix, J.A., Elvebakk, A., Fankhauser, J.D., Farkas, E., Ferraro, L.I., Fischer, E., Galloway, D.J., Gaya, E., Giralt, M., Goward, T., Grube, M., Hafellner, J., Hernández M., J.E., Herrera Campos, M.A., Kalb, K., Kärnefelt, I., Kantvilas, G., Killmann, D., Kirika, P., Knudsen, K., Komposch, H., Kondratyuk, S., Lawrey, J.D., Mangold, A., Marcelli, M.P., Mccune, B., Ines Messuti, M., Michlig, A., Miranda González, R., Moncada, B., Naikatini, A., Nelsen, M.P., Øvstedal, D.O., Palice, Z., Papong, K., Parmen, S., Pérez-Ortega, S., Printzen, C., Rico, V.J., Rivas Plata, E., Robayo, J., Rosabal, D., Ruprecht, U., Salazar Allen, N., Sancho, L., Santos De Jesus, L., Santos Vieira, T., Schultz, M., Seaward, M.R.D., Sérusiaux, E., Schmitt, I., Sipman, H.J.M., Sohrabi, M., Søchting, U., Zeuthen Søgaard, M., Sparrius, L.B., Spielmann, A., Spribille, T., Sutjaritturakan, J., Thammathaworn, A., Thell, A., Thor, G., Thüs, H., Timdal, E., Truong, C., Türk, R., Umaña Tenorio, L., Upreti, D.K., Van Den Boom, P., Vivas Reubelta, M., Wedin, M., Will-Wolf, S., Wirth, V., Wirtz, N., Yahr, R., Yeshitela, K., Ziemmeck, F., Wheeler, T. & Lücking, R. (2011) One hundred new species of lichenized fungi: a signature of undiscovered global diversity. *Phytotaxa* 18: 1–127.
- Lumbsch, H.T., Kraichak, E., Parmen, S., Rivas Plata, E., Aptroot, A., Cáceres, M.E.S., Ertz, D., Feuerstein, S.C., Mercado-Díaz, J.A., Staiger, B., Van den Broeck, D. & Lücking, R. (2014) New higher taxa in the lichen family Graphidaceae (lichenized Ascomycota: Ostropales) based on a three-gene skeleton phylogeny. *Phytotaxa* 189(1):39–51.
<http://dx.doi.org/10.11646/phytotaxa.189.1.5>
- Mangold, A., Eli, J.A. & Lumbsch, H.T. (2009) Thelotremataceae. *Flora of Australia* 57: 195–420.
- Massalongo, A.B. (1860) Esame comparativo di alcuni generi di licheni. *Atti Reale Istituto Veneto Scienza Letteri Arti ser. 3,5*: 247–276, 313–337.
- Mercado-Díaz, J.A., Lücking, R. & Parmen, S. (2014) Two new genera and twelve new species of Graphidaceae from Puerto Rico: a case for higher endemism of lichenized fungi in islands of the Caribbean? *Phytotaxa* 189(1): 186–203.
<http://dx.doi.org/10.11646/phytotaxa.189.1.14>
- Meyer, G.F.W. (1825) *Nebststunden meiner Beschaeftigungen im Gebiete der Pflanzenkunde. Erster Theil*. Göttingen.
- Montagne, C. (1842) Troisième centurie de plantes cellulaires exotiques nouvelles, Décades V-VIII. *Annales des Sciences Naturelles, Botanie* 18: 241–282.
- Montagne, C. (1845) Plantae cellulares quas in insulis Philippinensibus a cl. Cuming collectas recensuit observationibus nonnullis descriptionibusque illustravit. *London Journal of Botany* 4: 1–6.
- Müller, J. (1881) Lichenologische Beiträge 14. *Flora* 64: 505–511, 513–527.
- Müller, J. (1882) Lichenologische Beiträge 16. *Flora* 65: 483–490, 499–505, 515–519.
- Müller, J. (1887a) Lichenologische Beiträge 26. *Flora* 70: 268–273, 283–288, 316–322, 336–338, 396–402, 423–429.
- Müller, J. (1887b) Graphideae Fééanae inclus. trib. affinis nec non Graphidaea exoticae Acharii, El. Friesii et Zenkeri e novo studio speciminum originalium expositae et in novam dispositionem ordinatae. *Mémoires de la Société de Physique et Histoire naturelle de Genève* 29(8): 1–80.
- Müller, J. (1892) Lichenes exotici. *Hedwigia* 31: 276–288.
- Nakanishi, M., Kashiwadani, H. & Moon, K.H. (2003) Taxonomical notes on Japanese Graphidaceae (Ascomycotina), including some new combinations. *Bulletin of the National Science Museum (Tokyo), Series B (Botany)* 29: 83–90.
- Orange, A., James, P.W., White, F.J. (2010) *Microchemical methods for the identification of lichens*. 2nd Ed. British Lichen Society, London.
- Papong, K.B., Boonpragob, K., Mangold, A., Divakar, P.K. & Lumbsch, H.T. (2010) Thelotremoid lichen species recently

- described from Thailand: a re-evaluation. *Lichenologist* 42: 131–137.
<http://dx.doi.org/10.1017/s0024282909990405>
- Papong, K.B., Lücking, R., Kraichak, E., Parmen, S., Von Konrat, M. & Lumbsch, H.T. (2014) Preliminary insights into the diversity of the lichen family Graphidaceae in New Caledonia (Ostropales, Ascomycota), with the description of 23 new species. *Phytotaxa* 189(1): 204–231.
<http://dx.doi.org/10.11646/phytotaxa.189.1.15>
- Parmen, S., Lücking, R. & Lumbsch, H.T. (2012) Phylogenetic classification at generic level in the absence of distinct phylogenetic patterns of phenotypical variation: A case study in Graphidaceae (Ascomycota). *PLoS One* 7(12): e51392.
<http://dx.doi.org/10.1371/journal.pone.0051392>
- Parmen, S., Cáceres, M.E.S., Lücking, R. & Lumbsch, H.T. (2013) *Myriocephala* and *Nitidocephala*, two new genera in Graphidaceae (Ascomycota: Ostropales) for chroodiscoid species in the *Ocellularia* clade. *Bryologist* 116: 127–133.
<http://dx.doi.org/10.1639/0007-2745-116.2.127>
- Redinger, K. (1933) Neue und wenig bekannte Flechten aus Brasilien. *Hedwigia* 73: 54–67.
- Redinger, K. (1936) Thelotremataceae brasilienses imprimis e herbario Regnelliano cognitae praetereaue in herbariis Krempelhuberi, Mülleri Arg., Nylanderi, Wainionis et Zahlbruckneri asservatae. *Arkiv för Botanik* 28A(8): 1–122.
- Rivas Plata, E. & Lücking, R. (2013) High diversity of Graphidaceae (lichenized Ascomycota: Ostropales) in Amazonian Perú. *Fungal Diversity* 58: 13–52.
<http://dx.doi.org/10.1007/s13225-012-0172-y>
- Rivas Plata, E. & Lumbsch, H.T. (2011) Parallel evolution and phenotypic divergence in lichenized fungi: a case study in the lichen-forming fungal family Graphidaceae (Ascomycota: Lecanoromycetes: Ostropales). *Molecular Phylogenetics and Evolution* 61: 45–63.
<http://dx.doi.org/10.1016/j.ympev.2011.04.025>
- Rivas Plata, E., Lücking, R. & Lumbsch, H.T. (2008) When family matters: an analysis of Thelotremataceae (lichenized Ascomycota: Ostropales) as bioindicators of ecological continuity in tropical forests. *Biodiversity and Conservation* 17: 1319–1351.
<http://dx.doi.org/10.1007/s10531-007-9289-9>
- Rivas Plata, E., Lücking, R., Sipman, H.J.M., Mangold, A., Kalb, K. & Lumbsch, H.T. (2010). A world-wide key to the thelotremoid Graphidaceae, excluding the *Ocellularia-Myriotrema-Stegobolus* clade. *Lichenologist* 42: 187–189.
<http://dx.doi.org/10.1017/s0024282909990491>
- Rivas-Plata, E., Hernández, J.E.M., Lücking, R., Staiger, B., Kalb, K. & Cáceres, M.E.S. (2011) *Graphis* is two genera: A remarkable case of parallel evolution in lichenized Ascomycota. *Taxon* 60: 99–107.
- Rivas Plata, E., Lücking, R. & Lumbsch, H.T. (2012a) Molecular phylogeny and systematics of the *Ocellularia* clade (Ascomycota: Ostropales: Graphidaceae). *Taxon* 61: 1161–1179.
- Rivas Plata, E., Lücking, R. & Lumbsch, H.T. (2012b) A new classification for the family Graphidaceae (Ascomycota: Lecanoromycetes: Ostropales). *Fungal Diversity* 52: 107–121.
<http://dx.doi.org/10.1007/s13225-011-0135-8>
- Rivas Plata, E., Parmen, S., Staiger, B., Mangold, A., Frisch, A., Weerakoon, G., Hernández M., J.E., Cáceres, M.E.S., Kalb, K., Sipman, H.J.M., Common, R.S., Lücking, R. & Lumbsch, H.T. (2013) A molecular phylogeny of Graphidaceae (Ascomycota: Lecanoromycetes: Ostropales) including 437 species. *MycKeys* 5: 55–94.
<http://dx.doi.org/10.3897/mycokeys.6.3482>
- Schumm, F. & Aptroot, A. (2012) *A microscopical atlas of some lichens from SE-Asia (Thailand, Cambodia, Philippines, Vietnam)*. Books on Demand, Norderstedt. 2 Vols.
- Sipman, H.J.M. & Aptroot, A. (1992) Results of a botanical expedition to Mount Roraima, Guyana. II. Lichens. *Tropical Bryology* 5: 79–107.
- Sipman, H.J.M., Lücking, R., Aptroot, A., Kalb, K., Chaves, J.L. & Umaña, L. (2012) A first assessment of the Ticolichen biodiversity inventory in Costa Rica and adjacent areas: the thelotremoid Graphidaceae (Ascomycota: Ostropales). *Phytotaxa* 55: 1–214.
- Staiger, B. (2002) Die Flechtenfamilie Graphidaceae. Studien in Richtung einer natürlicheren Gliederung. *Bibliotheca Lichenologica* 85: 1–526.
- Staiger, B. & Kalb, K. (1999) *Acanthothecis* and other graphidioid lichens with warty periphysoids or paraphysis-tips. *Mycotaxon* 73: 69–134.
- Thell, A., Crespo, A., Divakar, P.K., Kärnefelt, I., Leavitt, S.D., Lumbsch, H.T. & Seaward, M.R.D. (2012) A review of the lichen family Parmeliaceae. *Nordic Journal of Botany* 30: 641–664.
<http://dx.doi.org/10.1111/j.1756-1051.2012.00008.x>
- Vainio, E.A. (1890) Étude sur la classification et la morphologie des lichens du Brésil I. *Acta Societatis pro Fauna et Flora Fennica* 7(2): 1–247.
<http://dx.doi.org/10.5962/bhl.title.2741>
- Vainio, E.A. (1921) Lichenes insularum Philippinarum III. *Annales Academia Scientia Fennica, ser. A* 15(6): 1–368.
- Zahlbruckner, A. (1923) *Catalogus Lichenum Universalis* 2. Borntraeger, Leipzig.
- Zahlbruckner, A. (1930) Lichenes (Übersicht über sämtliche bisher aus China bekannten Flechten). In: Handel-Mazetti, H., *Symbolae sinicae. Botanische Ergebnisse der Expedition der Akademie der Wissenschaften in Wien nach Südwest-China 1914-1918*. III. Springer, Wien.