



Phyllosticta species from banana (*Musa* sp.) in Chongqing and Guizhou Provinces, China

SHI-PING WU^{1,2,3}, YONG-XIANG LIU³, JIE YUAN², YONG WANG¹, KEVIN D. HYDE⁴ & ZUO-YI LIU^{3,5*}

¹ Agricultural College of Guizhou University, Guiyang, 550006, P.R. China

² Guizhou Institute of Plant Protections, Guiyang, 550006 P.R. China

³ Key Laboratory for Agricultural Biotechnology of Guizhou, Guiyang, 550006 P.R. China

⁴ Institute of Excellence in Fungal Research, and School of Science, Mae Fah Luang University, Chiang Rai 57100, Thailand

⁵ Guizhou Academy of Agricultural Science, 550006 P.R. China

*Corresponding author: Zuo-Yi Liu: gzliuzuoyi@163.com

Abstract

Six *Phyllosticta* strains were isolated from diseased leaves of *Musa* species in Chongqing and Guizhou provinces, China. Morphological and molecular analysis of LSU and combined ITS, ACT, TEF-1, and GPDH gene sequences, identified these strains as *P. capitalensis* (3 strains), *P. musarum* (1 strain) and two isolates were distinct from known *Phyllosticta* species. The latter is herein introduced as *Phyllosticta musaechinensis* sp. nov. A description and illustrations are provided, and the new species is compared with other species from *Musa* in this paper.

Key words: *Musa*, Phylogeny, New species, Taxonomy

Introduction

The genus *Phyllosticta* is an important causal agent of banana (*Musa* sp.) leaf and post-harvest diseases (Meredith 1968, Jones & Alcorn 1982, Wulandari *et al.* 2010). Seven species of *Phyllosticta* have been recorded from banana although their identification is confusing (Meredith 1968, Chuang 1981, Brown *et al.* 1998, Photita *et al.* 2001, van der Aa & Vanev 2002, Pu *et al.* 2008, Wong *et al.* 2013, Wulandari *et al.* 2010). Wulandari *et al.* (2010) investigated the *Guignardia/Phyllosticta* species associated with freckle disease on banana leaves, re-examined the holotype of each epithet, and reported that the agents of banana freckle are *Guignardia musae* Racib. and *G. musicola* Wulandari, L. Cai & K.D. Hyde. Wong *et al.* (2012) used the name *Phyllosticta* rather than *Guignardia*, found five species on *Musa* in Australia, designated the epitypes for *Phyllosticta maculata* M.H. Wong & Crous and *P. musarum* (Cooke) Aa, and described *P. cavendishii* M.H. Wong & Crous as a new species. The history of *Phyllosticta* on banana was also discussed by Wulandari *et al.* (2010) and Wong *et al.* (2012). *Phyllosticta capitalensis* Henn., *P. cocoicola* (Bat.) Sivan., *Phyllosticta musae* (as *Guignardia musae*), *Phyllosticta musicola* and *Guignardia sydowiana* Trotter have been also recorded as endophytes on banana (Brown *et al.* 1998, Photita *et al.* 2001, 2002).

Wikee *et al.* (2013) provided a multilocus backbone tree for *Phyllosticta* species based on combined ITS, TEF-1, ACT, LSU and GPDH region genes, however, they did not include pathogens from banana (except the ubiquitous endophyte, *P. capitalensis*). In the present study six, *Phyllosticta* strains were isolated from diseased leaves of *Musa* spp. from Chongqing and Guizhou provinces in China. Among them, one taxon differed from known *Phyllosticta* species from banana and other hosts. The aim of this paper is to describe the new species based on morphological and molecular data and investigate the relationship of *P. musaechinensis* with other species.

Materials and methods

Isolates

Symptomatic banana leaves with small to expanding lesions were selected for isolation. The leaves were cut into pieces

approximately 3 × 5 cm, surface sterilized in 70% ethanol for 1 min, and then air-dried. Isolations were performed using two different laboratory methods. Firstly, a fruiting body was transferred into sterile water, allowed to soak overnight, and then single spore isolates were prepared using the method described in Chomnunti *et al.* (2014). Secondly, single fruiting bodies, observed under a stereo microscope were removed with a scalpel and plated onto potato dextrose agar (PDA) containing streptomycin sulphate to inhibit bacterial growth (Wong *et al.* 2012). The ex-type strain has been deposited in Guizhou Academy of Agricultural Sciences Collection (GZAAS) and an ex-paratype strain has been sent to International Collection of Microorganisms from Plants (ICMP) and Mae Fah Luang University Culture Collection (MFLUCC), respectively.

DNA isolation, amplification and phylogeny

DNA was extracted from isolates growing on PDA at 28°C for 30 d following the protocol of Cubero *et al.* (1999). The primers used were LROR (Rehner & Samuels 1994) and LR5 (Vilgalys & Hester 1990) for LSU region, ITS1 and ITS4 (White *et al.* 1990) for ITS region, EF1-728F and EF1-986R (Carbone & Kohn 1999) for translation elongation factor 1- α gene (TEF-1), ACT-512F and ACT-783R (Carbone & Kohn 1999) for the actin gene (ACT) and GDF1 (Guerber *et al.* 2003) and Gpd2-LM (Myllys *et al.* 2002) or GDR1 (Guerber *et al.* 2003) for the glyceraldehyde 3-phosphate dehydrogenase gene (GPDH). Amplification conditions followed Arzanlou *et al.* (2008). DNA sequencing was performed at the SinoGenoMax Company (Beijing, China) using corresponding primer pairs in sequencing. Novel sequences have been deposited in the GenBank (Tab. 1)

Sequences of our isolates together with reference sequences obtained from GenBank (Table 1) were aligned with MAFFT (Kato *et al.* 2005). The alignments were checked visually and improved manually if necessary in BioEdit (v. 7.1.3.0). The index of substitution saturation was assessed using DAMBE v5.3.70 (Xia *et al.* 2003). The selection of conserved blocks was conducted using Gblocks v0.91b. The phylogenetic analysis of the aligned sequences was conducted using PAUP v.4.0b10 (Swofford 2003) for maximum-parsimonious analysis.

TABLE 1 Sources of isolates and GenBank accession number used in this study

Species	Strain no. ^a	Host	Locality	GenBank Accession number ^b			
				ITS	ACT	TEF-1	GPDH
<i>Botryosphaeria obtusa</i>	CMW8232	<i>Conifers</i>	South Africa	AY972105	AY972111	DQ280419	
<i>Guignardia alliacea</i>	MUCC0014*	<i>Allium fistulosum</i>	Japan	AB454263			
<i>G. mangiferae</i>	IMI260.576*	<i>Manifera indica</i>	India	JF261459	JF343641	JF261501	JF343748
<i>G. philoprina</i>	CBS447.68*	<i>Taxus baccata</i>	Netherlands	AF312014			
<i>G. rhodora</i>	CBS 901.69	<i>Rhododendron sp.</i>	Netherlands	KF206174	KF289256	KF289230	KF289166
<i>Phyllosticta abieticola</i>	CBS112067*	<i>Abies concolor</i>	Canada	KF170306	KF289238		
<i>P. aloicola</i>	CPC21020*	<i>Aloe ferox</i>	South Africa	KF154280	KF289311	KF289193	KF289124
<i>P. ampellicida</i>	ATCC200578*	<i>Vitis riparia</i>	USA	KC193586	KC193581	KC193584	
<i>P. ardisiicola</i>	NBRC102261*	<i>Ardisia crenata</i>	Japan	AB454274	AB704216		
<i>P. aspidisticola</i>	NBRC102244*	<i>Aspidistra elatior</i>	Japan	AB454260			
<i>P. beaumarisii</i>	IMI 298910 *	<i>Muehlenbekia adpressa</i>	Australia	AY042927	KF306232	KF289170	KF289074
<i>P. bifrenariae</i>	CBS128855*	<i>Bifrenaria harrisoniae</i>	Brazil	JF343565	JF343649	JF343586	JF343744
<i>P. brazilianiae</i>	CBS126270*	<i>Mangifera indica</i>	Brazil	JF343572	JF343656	JF343593	JF343758
<i>P. capitalensis</i>	CBS128856*	<i>Stanhopea sp.</i>	Brazil	JF261465	JF343647	JF261507	JF343776
<i>P. capitalensis</i>	CBS117118	<i>Musa acuminata</i>	Indonesia	FJ538339	FJ538455	FJ538397	KF289090
<i>P. cavendishii</i>	BRIP554196*	<i>Musa cv. Formosana</i>	Taiwan	JQ743562	KF014080	KF009743	
<i>P. cavendishii</i>	BRIP58008	<i>Musa sp.</i>	Australia	KC988365	KF014071	KF009742	
<i>P. citriasiana</i>	CBS 120486*	<i>Citrus maxima</i>	Thailand	FJ538360	FJ538476	FJ538418	JF343686
<i>P. citribraziliensis</i>	CBS100098*	<i>Citrus limon</i>	Brazil	FJ538352	FJ538468	FJ538410	JF343691
<i>P. citricarpa</i>	CBS127454*	<i>Citrus limon</i>	Australia	JF343583	JF343667	JF343604	JF343771

.....Continued on the next page

TABLE1 (Continued)

Species	Strain no. ^a	Host	Locality	GenBank Accession number ^b			
				ITS	ACT	TEF-1	GPDH
<i>P. cordylinophila</i>	CPC20261*	<i>Cordyline fruticosa</i>	Thailand	KF170287	KF289295	KF289172	KF289076
<i>P. cornicola</i>	CBS111639	<i>Cornus florida</i>	USA	KF170307	KF289234		
<i>P. cussoniae</i>	CBS136060*	<i>Cussonia</i> sp.	South Africa	JF343578	JF343662	JF343599	JF343764
<i>P. elongata</i>	CBS 126.22*	<i>Oxycoccus macrocarpos</i>	USA	FJ538353	FJ538469	FJ538411	KF289164
<i>P. ericarum</i>	CBS132534*	<i>Erica gracilis</i>	South Africa	KF206170	KF289291	KF289227	KF289162
<i>P. eugeniae</i>	CBS 445.82	<i>Eugenia aromatica</i>	Indonesia	AY042926	KF289246	KF289208	KF289139
<i>P. fallopiae</i>	MUCC0113*	<i>Fallopia japonica</i>	Japan	AB454307			
<i>P. foliorum</i>	CBS 447.68*	<i>Taxus baccata</i>	Netherlands	KF170309	KF289247	KF289201	KF289132
<i>P. gaultheriae</i>	CBS 447.70*	<i>Gaultheria humifusa</i>	USA	JN692543	KF289248	JN692531	JN692508
<i>P. hamamelidis</i>	MUCC149	<i>Hamamelis japonica</i>	Japan	KF170289	KF289309		
<i>P. hostae</i>	CGMCC3.14355*	<i>Hosta plantaginea</i>	China	JN692535	JN692511	JN692523	JN692503
<i>P. hubeiensis</i>	CGMCC3.14986*	<i>Viburnum odoratissimum</i>	China	JX025037	JX025032	JX025042	JX025027
<i>P. hymenocallidicola</i>	CBS 131309*	<i>Hymenocallis littoralis</i>	Australia	JQ044423	KF289242	KF289211	KF289142
<i>P. hypoglossi</i>	CBS 434.92*	<i>Ruscus aculeatus</i>	Italy	FJ538367	FJ538483	FJ538425	JF343695
<i>P. ilicis-aquifolii</i>	CGMCC3.14358*	<i>Ilex aquifolium</i>	China	JN692538	JN692514	JN692526	
<i>P. kerriae</i>	MAFF240047*	<i>Kerria japonica</i>	Japan	AB454266			
<i>P. leucothoicola</i>	MUCC0553*	<i>Leucothoe catesbaei</i>	Japan	AB454370	KF289310		
<i>P. ligustricola</i>	MUCC0024*	<i>Ligustrum obtusifolium</i>	Japan	AB454269	AB704212		
<i>P. maculata</i>	CPC18347*	<i>Musa cv. Goly-goly pot-pot</i>	Australia	JQ743570	KF014016	KF009700	
<i>P. maculata</i>	BRIP46622	<i>Musa</i> sp.	Australia: Northern Territory	JQ743567	KF014013	KF009692	
<i>P. mangifera-indica</i>	CPC20264*	<i>Mangifera indica</i>	Thailand	KF170305	KF289296	KF289190	KF289121
<i>P. minima</i>	CBS 585.84*	<i>Acer rubrum</i>	USA	KF206176	KF289249	KF289204	KF289135
<i>P. musarum</i>	BRIP55434*	<i>Hill banana</i>	India	JQ743584			
<i>P. musarum</i>	BRIP57803	<i>Musa</i> sp.	Australia	JX997138	KF014055	KF009737	
<i>P. musarum</i>	BRIP58028	<i>Musa</i> sp.	Australia	KC988377	KF014054	KF009738	
<i>P. musicola</i>	CBS123405*	<i>Musa acuminata</i>	Thailand	FJ538334	FJ538450	FJ538392	
<i>P. neopyrolae</i>	MUCC0125*	<i>Pyrola asarifolia</i>	Japan	AB454318	AB704233		
<i>P. owaniana</i>	CBS776.97*	<i>Brabejum stellatifolium</i>	South Africa	FJ538368	KF289254	FJ538426	JF343767
<i>P. pachysandricola</i>	MUCC0124*	<i>Pachysandra terminalis</i>	Japan	AB454317	AB704232		
<i>P. parthenocissi</i>	CBS111645*	<i>Parthenocissus quinquefolia</i>	USA	EU683672	JN692518	JN692530	
<i>P. paxistimae</i>	CBS112527*	<i>Paxistima mysinites</i>	USA	KF206172	KF289239	KF289209	KF289140
<i>P. philoprina</i>	CBS616.72	<i>Ilex aquifolium</i>	Germany	KF154279	KF289251	KF289205	KF289136
<i>P. podocarp</i>	CBS111647	<i>Podocarpus lanceolata</i>	South Africa	KF154276	KF289235	KF289232	KF289168
<i>P. podocarpicola</i>	CBS728.79*	<i>Podocarpus maki</i>	USA	KF206173	KF289252	KF289203	KF289134

.....Continued on the next page

TABLE 1 (Continued)

Species	Strain no. ^a	Host	Locality	GenBank Accession number ^b			
				ITS	ACT	TEF-1	GPDH
<i>P. spinarum</i>	CBS292.90	<i>Chamaecyparis pisifera</i>	France	JF343585	JF343669	JF343606	JF343773
<i>P. styracicola</i>	CGMCC3.14985*	<i>Styrax grandiflorus</i>	China	JX025040	JX025035	JX025045	JX025030
<i>P. telopeae</i>	CBS777.97*	<i>Telopea speciosissima</i>	Tasmania	KF206205	KF289255	KF289210	KF289141
<i>P. vaccinii</i>	ATCC46255*	<i>Vaccinium macrocarpon</i>	USA	KC193585	KC193580	KC193582	KC193583
<i>P. vacciniicola</i>	CPC18590*	<i>Vaccinium macrocarpum</i>	USA	KF170312	KF289287	KF289229	KF289165
<i>P. yuccae</i>	CBS117136	<i>Yucca elephantipes</i>	New Zealand	JN692541	JN692517	JN692529	JN692507
<i>P. capitalensis</i>	GZAAS6.1201	<i>Musa</i> sp.	China: Guizhou	KF955290	KM816623	KM816635	KM816629
<i>P. capitalensis</i>	GZAAS6.1202	<i>Musa</i> sp.	China: Guizhou	KF955291	KM816624	KM816636	KM816630
<i>P. capitalensis</i>	GZAAS6.1242	<i>Musa</i> sp.	China: Guizhou	KF955292	KM816625	KM816637	KM816631
<i>P. musarum</i>	GZAAS6.1228	<i>Musa</i> sp.	Thailand: Chiang Rai	KF955293	KM816626	KM816638	KM816632
<i>P. musaechinensis</i>	GZAAS6.1247	<i>Musa</i> sp.	China: Chongqing	KF955294	KM816627	KM816639	KM816633
<i>P. musaechinensis</i>	GZAAS6.1384	<i>Musa</i> sp.	China: Chongqing	KF955295	KM816628	KM816640	KM816634

Note:

a ATCC: American Type Culture Collection, Virginia, USA; **BRIP**: Plant Pathology Herbarium, Biosecurity Queensland, Dutton Park, Queensland, Australia; **CBS**: CBS KNAW Fungal Biodiversity Centre, Utrecht, the Netherlands; **CGMCC**: China General Microbial Culture Collection; **CPC**: Culture collection of P. W. Crous, housed at CBS; **IMI**: International Mycological Institute, Bioscience, Egham, Bakenham Lane, U. K.; **MAFF**: the Microbiological Genebank, National Institute of Agrobiological Sciences, Japan; **MUCC**: Culture Collection, Laboratory of Plant Pathology, Mie University, Tsu, Mie prefecture, Japan; **NBRC**: Biological Resource Center, the National Institute of 502 Technology and Evaluation, Japan; **ZJUCC**: Zhejiang University Culture Collection, China.

* indicates the ex-type cultures

b **ITS**: Internal transcribed spacers 1 and 2 together with 5.8S nrDNA; **TEF-1**: partial translation elongation factor 1- α gene; **ACT**: actin gene; **GAPDH**: glyceraldehyde-3-phosphate dehydrogenase gene.

The models of evolution were estimated with MrModeltest v2.3 (Nylander 2004). Bayesian analyses with the selected evolutionary model were performed with MrBayes 3.1.2 (Ronquist & Huelsenbeck 2003). Six simultaneous Markov chains were run for 1×10^6 generations and trees were sampled every 100th generation. The first 2000 trees, representing the burn-in phase of the analyses, were discarded and the remaining 8000 trees used for calculating posterior probabilities (PP) in the majority rule consensus tree (Liu *et al.* 2012).

The models of evolution were estimated with MrModeltest v2.3 (Nylander 2004). The alignments were converted by ALTER on <http://sing.ei.uvigo.es/ALTER> (Glez-Peña *et al.* 2010). A maximum likelihood analysis was performed (Silvestro *et al.* 2012). One thousand non parametric bootstrap iterations were run with the GTR model and a discrete gamma distribution. The resulting replicates were plotted on to the best scoring tree obtained previously. The representative sequences and phylogenetic analyses were performed according to Wong *et al.* (2012).

Morphological characters

Specimens were observed by a Nikon eclipse 80i compound microscope with DS-5Mc camera and an Olympus SZX2 stereomicroscope. Hand sections were made for microscopic examination. Measurements were made in water. The morphological characters of colony were assessed after 60 d growth on PDA, MEA (malt extract agar) and OA (oat agar).

Results

Phylogenetic analysis

The six isolates from banana obtained in this study were sequenced using five (LSU, ITS, ACT, TEF-1 and GPDH) genes.

The LSU alignment contained 22 sequences (including the outgroup *Botryosphaeria dothidea*) and consisted of 780 (including alignment gaps) total characters, of which 60 characters (7.7%) are parsimony informative. A heuristic search with random addition of taxa (1000 replicates) and treating gaps as missing characters generated 1 parsimonious tree (TL = 127, CI = 0.764, RI = 0.880, RC = 0.672, HI = 0.236) shown in Fig. 1. Bootstrap support values of MP (BS) (equal to or above 50%) and Bayesian posterior probabilities (PP) (equal to or above 0.90 based on 1,000,000 generations) are shown on the upper branches. The phylogenetic tree of the LSU region indicated that all *Phyllosticta* species and the six new isolates form a monophyletic lineage sister to *Botryosphaeria dothidea* (BS = 100). It confirms that the six isolates, GZAAS6.1201, GZAAS6.1202, GZAAS6.1228, GZAAS6.1242, GZAAS6.1247 and GZAAS6.1384 were members of *Phyllosticta sensu stricto*.

The combined dataset of ITS, ACT, TEF-1 and GPDH contained 75 combined 250 sequences from 64 taxa and comprised 1644 total characters including gaps, of which 887 characters were constant; 535 characters (32.5%) were parsimony informative; 211 variable characters are parsimony-uninformative. A heuristic search with random addition of taxa (1000 replicates) and treating gaps as missing characters generated 2496 equally parsimonious trees. All trees were similar in topology, first of which equally most parsimonious tree (TL = 2760, CI = 0.435, RI = 0.707, RC = 0.307, HI = 0.565) shown in Fig. 2. Bootstrap support values of most parsimonious (MPBS) and maximum likelihood (MLBS) (equal to or above 50%) are shown on the upper branches. The phylogenetic tree based on four gene loci analysis indicated that three isolates, GZAAS6.1201, GZAAS6.1202 and GZAAS6.1242, clustered with *P. capitalensis* and formed a branch with high support (MPBS = 83, MLBS = 87). GZAAS6.1228 was shown to be species of *P. musarum* (MPBS = 77, MLBS = 94). GZAAS6.1247 and GZAAS6.1384 formed a strong single lineage (MPBS = 100, MLBS = 100) relative to *P. musarum*, *P. maculata* and *P. cavendishii*.




FIGURE 1. The most parsimonious trees obtained from a heuristic search with 1000 random taxon additions of the LSU sequences using PAUP v. 4.0b10. The scale bar shows 5 changes. Bootstrap support values for maximum parsimony (MP) and Bayesian posterior probabilities above 0.90 are shown. A hyphen (-) indicates the value lower than 50% (BS) or 0.90 (PP). The tree is rooted to *Botryosphaeria dothidea*. Ex-type/ex-epitype isolates are marked by an asterisk *. Novel sequences are in boldface.




FIGURE 2. The first of 2496 equally most parsimonious trees obtained from a heuristic search with 1000 random taxon additions of the ITS, ACT, TEF-1 and GPDH sequences using PAUP v. 4.0b10. The scale bar shows 20 changes. Bootstrap support values for maximum parsimony (MPBS) and maximum likelihood (MLBS)(equal to or above 50 %) were shown. Thickened branches represent significant Bayesian posterior probability ($\geq 90\%$). A hyphen (-) indicates the value lower than 50%. The tree is rooted *Botryosphaeria dothidea*. An asterisk (*) indicates the ex-type strains. Novel sequences are printed in bold.




FIGURE 3. *Phyllosticta musaechinensis* (CQ12097, holotype). a. Symptom of disease. b. Pycnidia on infected leaf of *Musa* sp. c, d. Ostiole of pycnidium. e. Pycnidium and conidiogenous cell. f–h. Conidia. i. Conidia on colony. j. Conidiogenous cell in culture. k. Conidia from culture. l–n. Culture on PDA. Scale bars: e = 50 μ m, f–h, j, k = 20 μ m, n = 5 mm.

Taxonomy

Phyllosticta musaechinensis S.P. Wu, Z.Y. Liu & K.D. Hyde, *sp. nov.* (Fig 3) MycoBank MB 806057.

Phyllosticta musaechinensis is weakly pathogenic on leaves of *Musa* sp. It grows slowly on PDA. Sexual and spermatial states were not observed.

Type:—CHINA. Chongqing Municipality: Yubei District, Longtou Temple Park, on leaves of *Musa* sp., 21 September 2012, *Shiping Wu*, (CQ12097, holotype); ex-type living culture = GZAAS6.1247 = MFLUCC13-0907 = ICMP 20111).

Etymology:—From *Musa*, the host and *chinensis*, in reference to the first collection of the species on a banana host in China.

Weakly pathogenic on leaves of *Musa* sp., slightly discoloring leaves yellow, with black, shiny conidiomata forming on healthy green, or yellowing parts of leaves. Sexual state: Unknown. Asexual state: Pycnidia 45–145 μm (\bar{x} = 93 μm) diam. subcuticular to erumpent, solitary or clustered in small groups, black, shiny, globose or subglobose, with a rounded ostiole at the center. Conidiogenous cells cylindrical or conical. Conidia 14–18 \times 8–12 μm (\bar{x} = 17 \times 10 μm), hyaline, aseptate, coarsely guttulate, ellipsoidal or clavate, thin- and smooth-walled, surrounded by a mucilaginous sheath 0.5–3.5 μm thick, apex tapering, straight to curved, appendage 4.0–18.5 μm (\bar{x} = 12 μm) long. Spermatial state: unknown.

Colony on PDA bluish black to black, without aerial mycelium, irregular, raised to about 0.7 mm, reaching 14.2–12.5 mm diam after 60 d at 28°C. Pycnidiasolitary or aggregated in colony, black. Conidia 15.5–22.5 \times 8.5–13 μm (\bar{x} = 18 \times 11 μm), hyaline, aseptate, coarsely guttulate, ellipsoidal, clavate or irregular, thin- and smooth-walled, surrounded by a mucilaginous sheath or not, apex tapering, straight to curved 4–18 μm (\bar{x} = 11.6 μm). Spermatia not formed.

Known distribution:—China.

Host:—*Musa* sp.

Additional material examined:—CHINA. Chongqing Municipality: Yubei District, Longtou Temple Park, on leaves of *Musa* sp., 3 August 2013, *Shiping Wu* (CQ13018); living culture = GZAAS6.1384.

Discussion

After phylogenetic analysis based on four gene loci (Fig. 2), our taxon displayed a close relationship with *P. musarum*, *P. maculata* and *P. cavendishii* with strong statistical support, but formed an independent branch. All of these species are known from *Musa* sp. Additionally, four other species of *Phyllosticta*/*Guignardia* have also been reported from banana (van der Aa & Vanev 2002, Wulandari *et al.* 2010, Wong *et al.* 2012). Thus, we compare our taxon with these seven species. The asexual morphs of *G. stevensii* and *G. sydowiana* have never been reported (Wulandari *et al.* 2010). However, only the asexual states have been found for our new species. The new species shows greater variability than other related *Phyllosticta* spp. in pycnidia size, and the conidia are longer than those of *P. capitalensis*, *P. musicola*, and *P. cavendishii*. The conidial appendage of *P. musaechinensis* is shorter than in *P. maculata* and *P. musarum*. Thus, *P. musaechinensis* is distinguished from other *Phyllosticta* spp. from banana in pycnidia size, conidia size, mucilaginous sheath and appendage. The detailed information about morphological comparison is shown in Table 2.

Phyllosticta musaechinensis caused disease symptom similar to freckle disease, however, this weak pathogen mainly attacked older leaves which rapidly became yellow around the leaf spots. Additionally, leaves could not be infected by artificial inoculation.

Acknowledgements

We thank Dong-Qin Dai and Sajeewa S.N. Maharachchikumbura (cultures), Wen-Jing Li (photographic plates), and Jian-Kui Liu (phylogenetic analysis) for their invaluable assistance. This study was financially supported by the Innovation Ability Construction of Research Institutions in Guizhou Fund.

TABLE 2. Conidia morphology of *Phyllosticta* / *Guignardia* spp. described from *Musa*.

Taxa	Pycnidia size (µm)	Conidia size (µm)	Mucilaginous sheath thickness (µm)	Appendage length (µm)
<i>P. cavendishii</i>	78–137	12–17 × 8–10	1–3	8–20
<i>P. maculata</i>	84–137	15–21 × 9–13	2–6	12–37
<i>P. musarum</i>	69–118	12–20 × 7–11	1–3	14–20
<i>G. stevensii</i> (no asexual state)	–	–	–	–
<i>G. sydowiana</i> (no asexual state)	–	–	–	–
<i>G. musicola</i>	90–125	12–17 × 8–11	2–4	10–15
<i>P. capitalensis</i>	300	10–14 × 5–7	2–4	6–8
<i>P. musaechinensis</i>	45–145	15.5–22.5 × 8.5–13	0.5–3.5	4–18.5

References

- Brown, K.B., Hyde, K.D. & Guest, D.I. (1998) Preliminary studies on endophytic fungal communities of *Musa acuminata* species complex in Hong Kong and Australia. *Fungal Diversity* 1:27–51.
- Chomnunti, P., Hongsanan, S., Aguirre-Hudson, B., Tian, Q., Persoh, D., Dhimi, M.K., Alias, A.S., Xu, J., Liu, X.Z. & Stadler, M. (2014) The sooty moulds. *Fungal Diversity* 66: 1–36.
<http://dx.doi.org/10.1007/s13225-014-0278-5>
- Chuang, T.Y. (1981) Isolation of *Phyllosticta musarum*, causal organism of banana freckle. *Transactions of the British Mycological Society* 77(3):670–671.
[http://dx.doi.org/10.1016/s0007-1536\(81\)80127-1](http://dx.doi.org/10.1016/s0007-1536(81)80127-1)
- Cubero, O.F., Crespo, A., Fatehi, J. & Bridge, P.D. (1999) DNA extraction and PCR amplification method suitable for fresh, herbarium-stored, lichenized, and other fungi. *Plant Systematics and Evolution* 216(3–4):243–249.
<http://dx.doi.org/10.1007/bf01084401>
- Glez-Peña, D., Gómez-Blanco, D., Reboiro-Jato, M., Fdez-Riverola, F. & Posada, D. (2010) ALTER: program-oriented conversion of DNA and protein alignments. *Nucleic acids research* 38(suppl. 2):W14–W18.
<http://dx.doi.org/10.1093/nar/gkq321>
- Hoog, G.S. & Ende, A. (1998) Molecular diagnostics of clinical strains of filamentous Basidiomycetes. *Mycoses* 41(5–6):183–189.
<http://dx.doi.org/10.1111/j.1439-0507.1998.tb00321.x>
- Jones, D. & Alcorn, J. (1982) Freckle and black Sigatoka diseases of banana in far north Queensland. *Australasian Plant Pathology* 11(1):7–9.
- Katoh, K., Kuma, K.-I., Toh, H. & Miyata, T. (2005) MAFFT version 5: improvement in accuracy of multiple sequence alignment. *Nucleic Acids Research* 33(2):511–518.
<http://dx.doi.org/10.1093/nar/gki198>
- Liu, J.-K., Phookamsak, R., Doilom, M., Wikee, S., Li, Y.-M., Ariyawansa, H., Boonmee, S., Chomnunti, P., Dai, D.-Q. & Bhat, J.D. (2012) Towards a natural classification of Botryosphaerales. *Fungal Diversity* 57(1):149–210.
<http://dx.doi.org/10.1007/s13225-012-0207-4>
- Meredith, D. (1968) Freckle disease of banana in Hawaii caused by *Phyllostictina musarum* (Cke.) Petr. *Annals of Applied Biology* 62(2):329–340.
<http://dx.doi.org/10.1111/j.1744-7348.1968.tb02828.x>
- Nylander, J. (2004) *MrModeltest v2*. Program distributed by the author. Evolutionary Biology Centre, Uppsala University.
- Photita, W., Lumyong, S., Lumyong, P. & Hyde, K.D. (2001) Endophytic fungi of wild banana *Musa acuminata* at Doi Suthep Pui National Park, Thailand. *Mycological Research* 105(12):1508–1513.
<http://dx.doi.org/10.1017/S0953756201004968>
- Photita, W., Lumyong, S., Lumyong, P. & Hyde, K.D. McKenzie, E.H.C. (2002) Index of fungi described from the Musaceae. *Mycotaxon* 81(9): 491–503.
- Pu, J., Xie, Y., Zhang, X., Qi, Y., Zhang, C. & Liu, X. (2008) Preinfection behaviour of *Phyllosticta musarum* on banana leaves. *Australasian Plant Pathology* 37(1): 60–64.
<http://dx.doi.org/10.1071/ap07079>

- Rehner, S.A. & Samuels, G.J. (1994) Taxonomy and phylogeny of *Gliocladium* analysed from nuclear large subunit ribosomal DNA sequences. *Mycological Research* 98(6):625–634.
[http://dx.doi.org/10.1016/s0953-7562\(09\)80409-7](http://dx.doi.org/10.1016/s0953-7562(09)80409-7)
- Ronquist, F. & Huelsenbeck, J.P. (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19(12):1572–1574.
<http://dx.doi.org/10.1093/bioinformatics/btg180>
- Silvestro, D. & Michalak, I. (2012) raxmlGUI: a graphical front-end for RAxML. *Organisms Diversity & Evolution* 12(4):335–337.
<http://dx.doi.org/10.1007/s13127-011-0056-0>
- Swofford, D. (2003) *PAUP*: phylogenetic analysis using parsimony, version 4.0 b10*.
- van der Aa, H.A. & Vanev, S. (2002) A revision of the species described in *Phyllosticta*. In: Aptroot, A., Summerbell, R.C. & Verkley, G.J. (Eds.) *A revision of the species described in Phyllosticta*. CBS.
- Vilgalys, R. & Hester, M. (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172(8):4238–4246.
- White, T.J., Bruns, T., Lee, S. & Taylor, J. (Eds.) (1990) *PCR protocols: a guide to methods and applications*.
- Wikee, S., Lombard, L., Nakashima, C., Motohashi, K., Chukeatirote, E., Cheewangkoon, R., McKenzie, E.H.C., Hyde, K.D. & Crous, P.W. (2013) A phylogenetic re-evaluation of *Phyllosticta* (Botryosphaeraiales). *Studies in Mycology* 76: 1–29.
<http://dx.doi.org/10.3114/sim0019>
- Wong, M.-H., Crous, P.W., Henderson, J., Groenewald, J.Z. & Drenth, A. (2012) *Phyllosticta* species associated with freckle disease of banana. *Fungal Diversity* 56: 173–187.
<http://dx.doi.org/10.1007/s13225-012-0182-9>
- Wong, M.-H., Henderson, J., & Drenth, A. (2013) Identification and differentiation of *Phyllosticta* species causing freckle disease of banana using high resolution melting (HRM) analysis. *Plant Pathology* 62: 1285–1293.
<http://dx.doi.org/10.1111/ppa.12056>
- Wulandari, N.F., To-Anun, C., Lei, C., Abd-Elsalam, K.A. & Hyde, K.D. (2010) *Guignardia/Phyllosticta* species on banana. *Cryptogamie, Mycologie* 31(4):403–418.
- Xia, X., Xie, Z., Salemi, M., Chen, L. & Wang, Y. (2003) An index of substitution saturation and its application. *Molecular Phylogenetics and Evolution* 26(1):1–7.
[http://dx.doi.org/10.1016/S1055-7903\(02\)00326-3](http://dx.doi.org/10.1016/S1055-7903(02)00326-3)