



A new species of *Bulbophyllum* (Orchidaceae; Epidendroideae; Dendrobiinae) from Yunnan, China

QIANG LIU^{1,2}, JIAN-WU LI¹, XIAO-HUA JIN³ & JIANG-YUN GAO¹

¹Key Laboratory of Tropical Plant Resource and Sustainable Use, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Mengla, Yunnan 666303, China; E-mail: gjy@xtbg.org.cn

²University of Chinese Academy of Science, Beijing 100049, China

³State Key Laboratory of Systematic and Evolutionary Botany & Herbarium (PE), Institute of Botany, Chinese Academy of Sciences, Beijing 100093, China

Bulbophyllum Thouars (1822: 3) is one of the largest orchid genera, including more than 1900 species widely distributed from tropical America, Africa, Madagascar, and mainland Asia to Australasia (Lindley 1830, Pearce & Cribb 2002, Seidenfaden 1979, 1992, Chen *et al.* 2009, Pridgeon *et al.* 2014). The species possess a wide range of vegetative form and floral structure (Bose *et al.* 1980). There are about 103 species (33 endemic) in 18 sections in China (Chen *et al.* 2009).

Section *Stachysanthes* (Blume) Averynov (1994: 276) is distinguished by inconspicuous pseudobulbs and includes 72 species. There are five species (three endemic) of section *Stachysanthes* in China (Tsi 1999, Chen *et al.*, 2009). During our fieldwork in Mengla County of Xishuangbanna, southeastern Yunnan, a new species of *Bulbophyllum* of sect. *Stachysanthes* was found in the limestone forest and described below.

Taxonomy

Bulbophyllum mengyuanense Q.Liu, J.W.Li & X.H.Jin, *sp. nov.* (Figs. 1, 2)

Type:—China. Yunnan: Mengla County, Mengyuan Town, limestone forest, 1000 m, 7 Oct. 2010, Li 1099 (holotype, HITBC!).

Diagnosis: *Bulbophyllum mengyuanense* is similar to *B. drymoglossum* and *B. hainanense*, but can be distinguished from them by having yellow flowers with purple stripes, lateral sepals connate in their basal third, conspicuous column stelids and a labellum without a callus.

Epiphytic herbs. Rhizome creeping, slender, with internodes about 1 cm, each node bearing 1 leaf and 2–3 roots. Pseudobulb reduced to nearly absent. Leaf subsessile, elliptic, 13.0 × 8.5 mm, apex acuminate. Inflorescences arising from base of the reduced pseudobulb, erect, 31 mm, with 1 sheath, solitary flowered; floral bracts ovate, 1.3–3.0 mm, apex acute. Pedicel and ovary 8 mm. Flowers yellow, veins purple. Dorsal sepal ovate-oblong, 10 × 4 mm, apex acuminate, triveined; lateral sepals slightly larger, ovate-oblong, 12.0 × 4.5 mm, apex acuminate, connate in basal 1/3, 3-veined; petals oblong, 4.5 × 1.8 mm, entire, obtuse, univeined; lip elliptic, 5 × 4 mm, triveined, base attached to end of column foot by a mobile joint, apex obtuse; column stout, 2 × 1 mm, column-wing oblique triangular, 1.1 × 0.8 mm; column foot 4.5–5.0 mm, swollen in the middle, anther cap glabrous, pollinia 2.

Distribution:—Mengla, Yunnan, China (close to the border with Laos).

Ecology:—Epiphytic on trees in the limestone forest, which mainly is composed of *Pterospermum proteum* (Sterculiaceae), *Pistacia weinmannifolia* (Anacardiaceae), *Cleistanthus sumatranus* (Euphorbiaceae), *Quercus yiwuensis* (Fagaceae), *Tetrameles nudiflora* (Tetramelaceae) (Wang *et al.* 1997).

Phenology:—Flowering from October to November.

Conservation status:—At present, *Bulbophyllum mengyuanense* is known from two sites, and just two populations were discovered during four years of botanical surveys. However, it is expected that more populations may be found by thorough botanical investigation in the vicinity of the limestone forest. Moreover, it is possible that other populations occur in similar limestone habitats in the Laos and Myanmar. Therefore, it is premature to conduct a full conservation assessment based on the two known population. We regard the species as Data Deficient (DD: IUCN 2012).

Etymology:—Named for Mengyuan town in Xishuangbanna Autonomous Prefecture, Yunnan Province, where the type was collected.