

Order **Trombidiformes** Reuter, 1909^{1,2}

Suborder **Sphaerolichida** OConnor, 1984 (2 superfamilies)³

Superfamily **Lordalychioidea** Grandjean, 1939 (2 families)

Family **Lordalychidae** Grandjean, 1939 (1 genus, 16 species)

Superfamily **Sphaerolichoidea** Berlese, 1913

Family **Sphaerolichidae** Berlese, 1913 (1 genus, 5 species)

Suborder **Prostigmata** Kramer, 1877 (4 infraorders, 40 superfamilies)

Infraorder **Labidostommatina** Krantz, 1978 (1 superfamily)

Superfamily **Labidostommatoidea** Oudemans, 1904 (1 family)

Family **Labidostommatidae** Oudemans, 1904 (5 genera, 51 species; †0/2)⁴

Infraorder **Eupodina** Krantz, 1978 (5 superfamilies)

Superfamily **Bdelloidea** Dugès, 1834 (2 families)

Family **Bdellidae** Dugès, 1834 (15 genera, 256 species; †0/5)⁵

Family **Cunaxidae** Thor, 1902 (27 genera, 329 species)⁶

Superfamily **Eriophyoidea** Nalepa, 1898 (3 families)⁷

Family **Phytoptidae** Murray, 1877 (20 genera, 164 species)⁸

Family **Eriophyidae** Nalepa, 1898 (274 genera, 3,790 species)⁹

Family **Diptilomiopidae** Keifer, 1944 (63 genera, 450 species)

Superfamily **Eupodoidea** Koch, 1842 (9 families)¹⁰

Family **Cocceupodidae** Jesionowska, 2010 (3 genera, 23 species)¹¹

Family **Eupodidae** Koch, 1842 (10 genera, 69 species)¹²

Family **Dendrochaetidae** Olivier, 2008 (1 genus, 1 species)¹³

Family **Eriorhynchidae** Qin & Halliday, 1997 (1 genus, 5 species)

Family **Pentapalpidae** Olivier & Theron, 2000 (1 genus, 1 species)

Family **Penthaleidae** Oudemans, 1931 (5 genera, 16 species)¹⁴

Family **Penthalodidae** Thor, 1933 (6 genera, 35 species; ; †0/1)¹⁵

Family **Rhagidiidae** Oudemans, 1922 (28 genera, 157 species; ; †1/1)

Family **Strandtmanniidae** Zacharda, 1979 (1 genus, 2 species)¹⁶

1. **BY** Zhi-Qiang Zhang, Qing-Hai Fan, Vladimir Pesic, Harry Smit, Andre V. Bochkov, A. A. Khaustov, Anne Baker, Andreas Wohltmann, Tinghuan Wen, James W. Amrine, P. Beron, Jianzhen Lin, Grzegorz Gabrys & Robert Husband (for full author addresses, see **Contributor roles and addresses** after **References**). The title of this contribution should be cited as “Order Trombidiformes Reuter, 1909. *In*: Zhang, Z.-Q. (Ed.) *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*”. Recent Trombidiformes currently include 151 families, 2,235 genera, and 25,821 species, whereas fossil taxa include 10 genera and 24 species.
2. Higher classification and estimates of genera and species within families generally follow Walter *et al.* (2009), with revisions and updates as otherwise indicated. To be consistent with Beaulieu *et al.* (2011) and Schatz *et al.* (2011) in this volume, the rank cohort is not used and older names for higher taxa are used—also consistent with OConnor (1984) and Mironov & Bochkov (2009).
3. Based on Moraza (2008) and Walter *et al.* (2009), with updates.
4. Based on Dunlop & Bertrand (2011); throughout this paper, counts of genera and species are totals (Recent and fossil); †x/y refers x genera and y species, among the totals, are based on fossil.
5. Based on van der Schyff *et al.* (2003), Hernandez & Feres (2006), Hernandez *et al.* (2007), with updates.
6. Based on Den Heyer (2011).
7. Based on unpublished database of James Amrine.
8. One species of fossil (Dunlop *et al.* 2011).
9. Three species of fossil galls (Dunlop *et al.* 2011).
10. Species numbers do not include taxa that are evidently correctly placed in the Eupodoidea but are insufficiently defined for recognition. Many of these were described by C.L. Koch (see Thor & Willmann 1941). Jesionowska (2010) and Zacharda (1980) list species inquirendae for the Cocceupodidae and Rhagidiidae respectively.
11. Based on Jesionowska (2010).
12. Fain (1964) synonymized *Protereunetes* Berlese with *Ereynetes* Berlese. Despite Strandtmann (1970) reassigning eupodid species described in *Protereunetes* to *Eupodes* Koch, the generic name continued to be used by some authors. Fain’s synonymy is accepted and *Protereunetes* is not here recognized as a eupodid genus.
Jesionowska (2008) transferred *Hawaiieupodes* Strandtmann & Goff from the Eupodidae to the Penthalodidae. However, the unsettled diagnosis of the latter means that the new placement is also unsatisfactory. Consequently, the genus is here retained in the Eupodidae as originally assigned (Strandtmann & Goff 1978).
13. Abou-Awad (1984) erected *Egypteupodes* on the basis of the absence of internal vertical setae (*iv*). Olivier and Theron (1997) synonymized the genus with *Eupodes* because they observed that, although minute, *iv* were present. However, because it is not clear whether these authors examined type material, *Egypteupodes* is included in the generic complement.
14. Published in 2008, 2009, too late to be included by Walter *et al.* (2009).
15. See first footnote for the Penthalodidae.

- Superfamily **Halacaroidea** Murray, 1877 (2 families)
 Family **Halacaridae** Murray, 1877 (63 genera, 1,118 species)¹⁷
 Family **Pezidae** Harvey, 1990 (1 genus, 2 species)
 Superfamily **Tydeoidea** Kramer, 1877 (4 families)¹⁸
 Family **Triophtydeidae** André, 1980¹⁹ (3 genera, 40 species)
 Family **Ereynetidae** Oudemans, 1931 (29 genera, 180 species)
 Family **Tydeidae** Kramer, 1877 (30 genera, 340 species; †1/2)
 Family **Iolinidae** Pritchard, 1956 (36 genera, 125 species)
 Infraorder **Anystina** van der Hammen, 1972 (2 hyporders, 22 superfamilies)
 Hyporder **Anystae** Krantz, 1978 (5 superfamilies)
 Superfamily **Adamystoidea** Cunliffe, 1957 (1 family)
 Family **Adamystidae** Cunliffe, 1957 (4 genera, 20 species)²⁰
 Superfamily **Anystoidea** Oudemans, 1936 (3 families)
 Family **Anystidae** Oudemans, 1936 (18 genera, 107 species; †2/5)²¹
 Family **Pseudocheylidae** Oudemans, 1909 (3 genera, 15 species)²²
 Family **Teneriffidae** Thor, 1911 (5 genera, 20 species)²³
 Superfamily **Caeculoidea** Berlese, 1883 (1 family)
 Family **Caeculidae** Berlese, 1883 (7 genera, 73 species; †1/2)²⁴
 Superfamily **Pomerantzioidea** Baker, 1949 (1 family)
 Family **Pomerantziidae** Baker, 1949 (2 genera, 6 species)²⁵
 Superfamily **Paratydeoidea** Baker, 1949 (1 family)
 Family **Paratydeidae** Baker, 1949 (6 genera, 10 species)²⁶
 Family **Stigmocheylidae** Kethley, 1990 (1 genus, 4 species)²⁷
 Hyporder **Parasitengona** Oudemans, 1909 (17 superfamilies)²⁸
 Superfamily **Calyptostomatoidea** Oudemans, 1923 (1 family)²⁹
 Family **Calyptostomatidae** Oudemans, 1923 (1 genus, 6 species)
 Superfamily **Erythraeoidea** Robineau-Desvoidy, 1828 (2 families)³⁰
 Family **Erythraeidae** Robineau-Desvoidy, 1828 (55 genera, 771 species)³¹
 Family **Smarididae** Kramer, 1878 (10 genera, 53 species)³²

15. **Penthalodidae**: Qin (1997) considered that *Protopenthalodes* Jesionowska would be better accommodated in either the Eupodidae or Penthalidae, rather than the Penthalodidae as originally placed (Jesionowska 1989). However, he retained the genus in the latter until familial monophyly was resolved and a correct taxonomic position could be established.

Jesionowska (2008) reassessed the importance of the epistrosum as a diagnostic familial character state and concluded that *Callipenthalodes* Qin, *Turanopenthalodes* Barilo and *Stereotydeus* Berlese should be removed from the Penthalodidae. Because she neither reassigned them to existing families nor erected new ones for them, these genera have been treated here as penthalodids.

16. Strandmanniidae is considered a possible junior synonym of Rhagidiidae by Qin (1996).

17. Based on Bartsch (2009).

18. After Walter *et al.* (2009); we regret to note here that the data were not checked by a specialist.

19. H.M. André of Belgium (not M. André of France).

20. Based on Fuangarworn & Lekprayoon (2010).

21. Based on Meyer & Ueckermann (1987), Otto (1999, 2000), with updates.

22. Based on Ueckermann & Khanjani (2004).

23. Based on Judson (1994), Ueckermann & Khanjani (2002), Khanjani *et al.* (2011), with updates.

24. Based on Franz (1952), Coineau & Poinar (2001), with updates.

25. Based on Bochkov & Walter (2007).

26. Based on Seeman *et al.* (2000).

27. Based on Bochkov (2008a).

28. The classification above the family level is highly unsettled for this group (Zhang & Fan, 2007). One view is that water mites (Hydracarina) form a sister group to the rest of the terrestrial Parasitengona (Krantz 1978; Smith & Cook 1991; Witte 1991). A radically different view (Welbourn (1991), is that water mites and Trombidiiina are considered sister groups, and together they form a sister group to Erythraeina (Erythraeoidea + Calyptostomatoidea). Walter *et al.* (2009) proposed Stygothrombiae for Stygothrombidoidea, which was not analysed in Witte (1991) and Welbourn (1991). Future analyses sampling all superfamilies of both water mites and terrestrial Parasitengona are much needed before any reliable classification of the Parasitengona at the ordinal level can be reached. Therefore we provide a list of superfamilies without choosing any higher classification.

29. Based on unpublished list of Beron, who also provided data for Erythraeoidea.

30. †*Pararainbowia martilli* Dunlop, 2007 is unplaced within the superfamily.

31. Based on Southcott (1961), Beron (2008) with updates. Dunlop *et al.* (2011) listed 13 species described from Baltic amber in 1854; these require checking.

32. Based on Southcott (1961), Zhang (2003), Beron (2008), Wohltmann (2010) with updates.

Family †**Proterythraeidae** Vercammen-Grandjean, 1973 (1 genus, 1 species; †1/1)
 Superfamily **Amphotrombioidea** Zhang, 1998 (1 family)³³
 Family **Amphotrombiidae** Zhang, 1998 (1 genus, 1 species)
 Superfamily **Allotanaupodoidea** Zhang & Fan, 2007 (1 family)³⁴
 Family **Allotanaupodidae** Zhang & Fan, 2007 (3 genera, 6 species)
 Superfamily **Chyzerioidea** Womersley, 1954 (1 family)
 Family **Chyzeriidae** Womersley, 1954 (11 genera, 30 species)³⁵
 Superfamily **Tanaupodoidea** Thor, 1935 (1 family)
 Family **Tanaupodidae** Thor, 1935 (7 genera, 23 species; †1/1)³⁶
 Superfamily **Trombiculoidea** Ewing, 1929 (7 families)
 Family **Johnstoniidae** Thor, 1935 (10 genera, 53 species)³⁷
 Family **Neotrombidiidae** Feider, 1959 (5 genera, 24 species)³⁸
 Family **Trombellidae** Leach, 1815 (19 genera, 41 species)³⁹
 Family **Leeuwenhoekidae** Womersley, 1944 (33 genera, 230 species)
 Family **Trombiculidae** Ewing, 1929 (152 genera, 3,100 species)⁴⁰
 Family **Walchiidae** Ewing, 1946 (19 genera, 298 species)
 Family **Audyaniidae** Southcott, 1987 (1 genus, 1 species)⁴¹
 Superfamily **Trombidoidea** Leach, 1815 (4 families)
 Family **Achaemenothrombiidae** Saboori, Wohltmann & Hakimitabar, 2010 (1 genus, 2 species)⁴²
 Family **Neothrombiidae** Feider, 1959 (16 genera, 25 species)⁴³
 Family **Microtrombidiidae** Thor, 1935 (126 genera, 437 species)⁴⁴
 Family **Trombidiidae** Leach, 1815 (25 genera, 253 species)⁴⁵
 Superfamily **Yurebilloidea** Southcott, 1996 (1 family)⁴⁶
 Family **Yurebillidae** Southcott, 1996 (1 genus, 1 species)⁴⁷
 Superfamily **Hydryphantoidea** Piersig, 1896 (8 families)⁴⁸
 Family **Hydryphantidae** Piersig, 1896 (51 genera, 329 species)
 Family **Hydrodromidae** Viets, 1936 (2 genera, 28 species)
 Family **Rhynchohydracaridae** Lundblad, 1936 (5 genera, 13 species)
 Family **Teratothyadidae** Viets, 1929 (2 genera, 9 species)
 Family **Ctenothyadidae** Lundblad, 1936 (2 genera, 3 species)
 Family **Thermacaridae** Sokolow, 1927 (1 genera, 4 species)
 Family **Zelandothyadidae** Cook, 1983 (2 genera, 3 species)
 Family **Malgasacaridae** Tuzovskij, Gerecke & Goldschmidt 2007 (1 genus, 1 species)
 Superfamily **Eylaoidea** Leach, 1815 (4 families)
 Family **Eylaidae** Leach, 1815 (2 genera, 96 species)
 Family **Limnocharidae** Grube, 1859 (5 genera, 44 species)

-
33. Based on larval stage only with similarities to both early-derivative water mites and Trombidina (Zhang 1998).
 34. This superfamily was missed in the chapter on Trombidiformes in the *Manual of Acarology*, third edition (Walter *et al.* 2009). Zhang & Fan (2007) provided a key to superfamilies of terrestrial Parasitengona (based on post-larval stages).
 35. Based on Zhang & Saboori (1995); Haitlinger (1999), Saboori *et al.* (2005), with updates.
 36. Based on Newell (1957), Welbourn, W.C. (1991), Judson & Makol (2009) with updates.
 37. Based on Newell (1957, 1960), Wohltmann *et al.* (2004), with updates.
 38. Based on Lindquist & Vercammen-Grandjean (1971), Vercammen-Grandjean (1973), Feider (1977), Beron (1995), with updates.
 39. Based on Southcott (1987), Welbourn (1991), Harvey (1996), Saboori *et al.* (2005) with updates.
 40. An estimate to be varified.
 41. Also placed by some authors in Trombellidae (Southcott 1987).
 42. Known from larvae only; Saboori *et al.* (2010) provided a key to families of Trombidoidea based on larvae.
 43. Based on Zhang (1994), Zhang & Fan (2005), Haitlinger (2007), Saboori *et al.* (2008) and Małkol *et al.* (2009), with updates.
 44. Based on Southcott (1993, 1994), Gabrys (1999), Wohltmann (2006), Małkol & Laydanowicz (2010) with updates; Eutrombidiidae is merged into Microtrombidiidae based on phylogenetic work of Wohltmann (2006).
 45. Updated based on Małkol (2007) and Małkol & Sevsay (2011). Includes Podothrombiidae recognized in Małkol (2007) and some earlier papers cited by Małkol (2007). Dunlop *et al.* (2011) listed 5 species described from Baltic amber in 1854; these require checking.
 46. Updated from Southcott (1993) and Małkol & Gabrys (2005).
 47. Described by Southcott (1996) from larval stage only, initially placed in Trombidoidea *s.l.*, but was excluded from Trombidoidea *s.s.* by Saboori *et al.* (2010). Here it is placed in Yurebilloidea Southcott, 1996; superfamily rank is used here for the first time; decision by Z. Zhang. Based on larval characters, this superfamily is very closely related to Trombidoidea *sin* its strict sense.
 48. Davids *et al.* (2006) with updates.

- Family **Piersigiidae** Oudemans, 1902 (4 genera, 9 species)
- Family **Apheviderulicidae** Gerecke, Smith & Cook, 1999 (1 genus, 3 species)
- Superfamily **Hydrovolzioidea** Thor, 1905 (2 families)
- Family **Hydrovolziidae** Thor, 1905 (3 genera, 16 species)
- Family **Acherontacaridae** Cook, 1967 (2 genera, 19 species)
- Superfamily **Hydrachnoidea** Leach, 1815 (1 family)
- Family **Hydrachnidae** Leach, 1815 (1 genus, 131 species)
- Superfamily **Lebertioidea** Thor, 1900 (10 families)
- Family **Bandakiopsidae** Panesar, 2004 (3 genera, 4 species)
- Family **Stygotoniidae** Cook, 1992 (1 genus, 1 species)
- Family **Sperchontidae** Thor, 1900 (5 genera, 173 species)
- Family **Rutripalpidae** Sokolow, 1934 (1 genus, 2 species)
- Family **Teutonidae** Koenike, 1910 (2 genera, 6 species)
- Family **Anisitsiellidae** Koenike, 1910 (29 genera, 159 species)
- Family **Lebertiidae** Thor, 1900 (3 genera, 138 species)
- Family **Acucapitidae** Wiles, 1996 (1 genus, 2 species)
- Family **Oxidae** Viets, 1926 (2 genera, 126 species)
- Family **Torrenticolidae** Piersig, 1902 (6 genera, 397 species)
- Superfamily **Hygrobatoidea** Koch, 1842 (13 families)
- Family **Pontarachnidae** Koenike, 1910 (2 genera, 40 species)
- Family **Limnesiidae** Thor, 1900 (29 genera, 227 species)
- Family **Omartacaridae** Cook, 1963 (2 genera, 14 species)
- Family **Wettinidae** Cook, 1956 (4 genera, 9 species)
- Family **Frontipodopsidae** Viets, 1931 (1 genus, 9 species)
- Family **Ferradasiidae** Cook, 1980 (1 genus, 1 species)
- Family **Lethaxonidae** Cook, Smith & Harvey, 2000 (3 genera, 16 species)
- Family **Hygrobatidae** Koch, 1842 (82 genera, 846 species)
- Family **Aturidae** Thor, 1900 (81 genera, 800 species)
- Family **Feltriidae** Viets, 1926 (1 genus, 115 species)
- Family **Unionicolidae** Oudemans, 1909 (18 genera, 477 species)
- Family **Pionidae** Thor, 1900 (15 genera, 342 species)
- Family **Astacocrotonidae** Thor, 1927 (1 genus, 1 species)
- Superfamily **Arrenuroidea** Thor, 1900 (20 families)
- Family **Momoniidae** Viets, 1926 (13 genera, 55 species)
- Family **Nudomideopsidae** Smith, 1990 (3 genera, 31 species)
- Family **Mideopsidae** Koenike, 1910 (8 genera, 113 species)
- Family **Athienemanniidae** Viets, 1922 (16 genera, 43 species)
- Family **Chappuisididae** Motas & Tanasachi, 1946 (2 genera, 13 species)
- Family **Gretacaridae** Viets, 1978 (1 genus, 15 species)
- Family **Neocaridae** Motas & Tanasachi, 1947 (2 genera, 19 species)
- Family **Mideidae** Thor, 1911 (2 genera, 6 species)
- Family **Acalyptonotidae** Walter, 1911 (2 genera, 4 species)
- Family **Kantacaridae** Imamura, 1959 (1 genus, 1 species)
- Family **Nipponacaridae** Imamura, 1959 (1 genus, 3 species)
- Family **Harpagopalpidae** Viets, 1924 (1 genus, 7 species)
- Family **Arenohydracaridae** Cook, 1974 (1 genus, 3 species)
- Family **Amoenacaridae** Smith & Cook, 1997 (1 genus, 3 species)
- Family **Laversiidae** Cook, 1955 (1 genus, 1 species)
- Family **Krendowskiidae** Viets, 1926 (4 genera, 52 species)
- Family **Arrenuridae** Thor, 1900 (7 genera, 950 species)
- Family **Bogatiidae** Motas & Tanasachi, 1948 (2 genera, 3 species)
- Family **Hungarohydracaridae** Motas & Tanasachi, 1959 (5 genera, 17 species)
- Family **Uchidastygacaridae** Imamura, 1956 (3 genera, 15 species)
- Superfamily **Stygothrombidioidea** Thor, 1935 (1 family)

Family **Stygothrombiidae** Thor, 1935 (5 genera, 16 species)
 Infraorder **Eleutherengona** Oudemans, 1909 (12 superfamilies)
 Hyporder **Raphignathina** Kethley, 1982 (5 superfamilies)
 Superfamily **Cheyletoidea** Leach, 1815 (5 families)
 Family **Cheyletidae** Leach, 1815 (75 genera, 438 species)⁴⁹
 Family **Demodicidae** Nicolet, 1855 (7 genera, 108 species)⁵⁰
 Family **Harpirhynchidae** Dubinin, 1957 (14 genera, 93 species)⁵¹
 Family **Psorergatidae** Dubinin, 1955 (3 genera, 73 species)⁵²
 Family **Syringophilidae** Lavoipierre, 1953 (63 genera, 259 species)⁵³
 Superfamily **Myobioidea** Mégnin, 1877 (1 family)
 Family **Myobiidae** Mégnin, 1877 (54 genera, 503 species)⁵⁴
 Superfamily **Pterygosomatoidea** Oudemans, 1910 (1 family)
 Family **Pterygosomatidae** Oudemans, 1910 (10 genera, 166 species)⁵⁵
 Superfamily **Raphignathoidea** Kramer 1877 (11 families)⁵⁶
 Family **Barbutiidae** Robaux, 1975 (1 genus, 5 species)⁵⁷
 Family **Caligonellidae** Grandjean, 1944 (5 genera, 50 species)⁵⁸
 Family **Camerobiidae** Southcott, 1957 (7 genera, 146 species; †0/1)⁵⁹
 Family **Cryptognathidae** Oudemans, 1902 (3 genera, 52 species)⁶⁰
 Family **Dasythyreidae** Walter & Gerson, 1998 (2 genera, 5 species)⁶¹
 Family **Eupalopsellidae** Willmann, 1952 (5 genera, 40 species)⁶²
 Family **Homocaligidae** Wood, 1970 (2 genera, 8 species)⁶³
 Family **Mecognathidae** Gerson & Walter, 1998 (2 genera, 7 species)⁶⁴
 Family **Raphignathidae** Kramer, 1877 (3 genera, 57 species)⁶⁵
 Family **Stigmaeidae** Oudemans, 1931 (30 genera, 502 species)⁶⁶
 Family **Xenocaligonellididae** Gonzalez, 1978 (2 genera, 5 species)⁶⁷
 Superfamily **Tetranychoidae** Donnadieu 1875 (5 families)
 Family **Allochaetophoridae** Reck, 1959 (1 genus, 2 species)⁶⁸
 Family **Linotetranidae** Baker & Pritchard, 1953 (4 genera, 16 species)⁶⁹
 Family **Tenuipalpidae** Berlese, 1913 (34 genera, 895 species)⁷⁰
 Family **Tetranychidae** Donnadieu 1875 (95 genera, 1,270 species)⁷¹
 Family **Tuckerellidae** Baker & Pritchard, 1953 (1 genus, 28 species)⁷²
 Hyporder **Heterostigmata** Berlese, 1899 (7 superfamilies)

49. Based on Bochkov (2005) with additions.

50. Based on Bochkov (2009) with modifications.

51. Based on a unpublished database of A.V. Bochkov.

52. Based on Giesen (1990).

53. Based on Skoracki (2011) with additions.

54. Based on Bochkov (2009).

55. Based on Bochkov (2008b) with additions.

56. Based on a unpublished database of Q.-H. Fan.

57. Based on Fan *et al.* (2003) with additions.

58. Based on Fan (2005).

59. Based on Fan (2005) and Fan & Walter (2011) with additions.

60. Based on Doğan & Dönel (2010) and Fan (2005).

61. Based on Walter & Gerson (1998) with additions.

62. Based on Fan (2005) with additions.

63. Based on Fan (1997, 2005).

64. Based on Fan & Zhang (2005) and Fan (2005).

65. Based on Fan (2005) with additions.

66. Based on Fan & Zhang (2005), Meyer & Ueckermann (1989) and Fan (2005) with additions.

67. Based on Fan (2005).

68. Based on Meyer & Ueckermann (1997).

69. Based on Bagheri *et al.* (2008) and Khanjani *et al.* (2011), with updates.

70. Based on Mesa *et al.* (2009) and Beard & Gerson (2009), with updates by Q.-H. Fan.

71. Based on Spider Mites Web (<http://www.ensam.inra.fr/CBGP/spmweb/index.php>). Dunlop *et al.* (2011) listed 2 species described from Baltic amber in 1854; these require checking.

72. Based on Meyer & Ueckermann (1997) with additions.

- Superfamily **Dolichocyboidea** Mahunka, 1970 (1 family)
 Family **Dolichocybidae** Mahunka, 1970 (6 genera, 37 species)⁷³
 Family **Crotalomorphidae** Lindquist & Krantz, 2002 (1 genus, 1 species)
- Superfamily **Heterocheyleidea** Trägårdh, 1950 (1 family)
 Family **Heterocheylidae** Trägårdh, 1950 (1 genus, 25 species)
- Superfamily **Pyemotoidea** Oudemans, 1937 (4 families)
 Family **Acarophenacidae** Cross, 1965 (7 genera, 31 species; †1/1)
 Family **Caraboacaridae** Mahunka, 1970 (2 genera, 7 species)
 Family **Resinacaridae** Mahunka, 1976 (2 genera, 2 species; †1/1)
 Family **Pyemotidae** Oudemans, 1937 (1 genus, 24, species; †1/1)⁷⁴
- Superfamily **Pygmephoridea** Cross, 1965 (4 families)⁷⁵
 Family **Pygmephoridae** Cross, 1965 (27 genera, 310 species)
 Family **Scutacaridae** Oudemans, 1916 (24 genera, 802 species)
 Family **Microdispidae** Cross, 1965 (17 genera, 109 species)
 Family **Neopygmephoridae** Cross, 1965 (17 genera, 248 species)
- Superfamily **Tarsocheyleidea** Atyeo & Baker, 1964 (1 family)
 Family **Tarsocheylidae** Atyeo & Baker, 1964 (2 genera, 3 species)
- Superfamily **Tarsonemoidea** Kramer, 1877 (2 families)
 Family **Podapolipidae** Ewing, 1922 (30 genera, 239 species)
 Family **Tarsonemidae** Kramer, 1877 (41 genera, 566 species)⁷⁶
- Superfamily **Trochometridiidea** Mahunka 1970 (2 families)
 Family **Trochometridiidae** Mahunka 1970 (2 genera, 6 species)⁷⁷
 Family **Athyreacaridae** Lindquist, Kaliszewski & Rack, 1990 (1 genus, 1 species)
- Hyporder incertae sedis
 Superfamily **Cloacaroidea** Camin, Moss, Oliver & Singer, 1967 (2 families)⁷⁸
 Family **Cloacaridae** Camin, Moss, Oliver & Singer, 1967 (6 genera, 14 species)
 Family **Epimyodicidae** Fain, Lukoschus & Rosmalen, 1982 (1 genus, 4 species)

References

- Abou-Awad, B.A. (1984) The eupodid mites of Egypt (Acari: Eupodoidea: Eupodidae). *Acarologia*, 25(4), 329–335.
- Bagheri, M., Irani-Nejad, K.H., Kamali, K., Khanjani, M., Saboori, A. & Lotfollahi, P. (2008) A new species of *Linotetranus* (Acari: Prostigmata: Linotetranae) from Iran. *Zootaxa*, 1914, 65–68.
- Bartsch, I. (2009) Checklist of marine and freshwater halacarid mite genera and species (Halacaridae: Acari) with notes on synonyms, habitats, distribution and descriptions of the taxa. *Zootaxa*, 1998, 1–170.
- Beard, J.J. & Gerson, U. (2009) A new flat mite genus, *Acaricis* (Prostigmata: Tenuipalpidae), from Australian sedges (Cyperaceae). *Zootaxa*, 2073, 31–44.
- Beaulieu, F., Dowling, A.P.G., Klompen, H., Moraes, G.J. de & Walter, D.E. (2011) Superorder Parasitiformes Reuter, 1909. In: Zhang, Z.-Q. (Ed.), *Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*, 3148, 123–128.
- Beron, P. (1995) A new larval species of *Neotrombidium* (Acariformes, Actinedida: Neotrombidiidae) from Cuba. *Historia Naturalis Bulgarica*, 5, 13–18.
- Beron, P. (2008) *Acarorum catalogus. I. Acariformes: Calyptostomatoidea (Calyptostomatidae), Erythraeoidea (Smarididae, Erythraeidae)*. Pensoft & NMNH, Sofia, 271 pp.
- Bochkov, A. (2005) Synopsis of the described Actinedida of the world. Superfamily Cheyletoidea. In: Hallan, J. (Ed.), *Synopsis of the Described Arachnida of the World*. Available from: <http://insects.tamu.edu/research/collection/hallan/acari/Family/Actinedida1.htm> (accessed 23 November 2011)
- Bochkov, A.V. (2008a) A review of the mite family Stigmocheylidae Berlese (Acari: Prostigmata). *Annales Zoologici*, 58, 311–325.
- Bochkov, A.V. (2008b) Origin and evolution of parasitism in mites of the infraorder Eleutherengona (Acari: Prostigmata). Report I. Lower Raphignathae. *Parazitologiya*, 42, 337–359. [In Russian with English Summary]

73. Based on Hajiyanbar & Khaustov (2010) and Rahiminejad *et al.* (2011).

74. Based on Yu *et al.* (2010) and a recent fossil species in Khaustov & Perkovsky (2010).

75. The system of Pygmephoridea follows Khaustov (2008).

76. Based on Lin & Zhang (2002) with updates.

77. Based on Mortazavi *et al.* (2011).

78. Based on Bochkov & OConnor (2008), see this work for discussion about position of this superfamily.

- Bochkov, A.V. (2009) A review of mites of the parvorder Eleutherengona (Acariformes: Prostigmata) – permanent parasites of mammals. *Acarina*, Supplement 1, 1–149.
- Bochkov, A.V. & OConnor, B.M. (2008) A new mite superfamily Cloacaroida and its position within the Prostigmata (Acariformes). *The Journal of Parasitology*, 94, 335–344.
- Bochkov, A.V. & Walter, D.E. (2007) The life-cycle of *Pomerantzia philippina* sp.n. (Prostigmata: Pomerantziidae) described from the Philippines. *Acarina*, 15, 159–170.
- Coineau, Y. & Poinar, G. (2001) A Caeculidae from Dominican amber. *Acarologia*, 41(1–2), 141–144.
- Davids, C., Di Sabatino, A., Gerecke, R., Gledhill, T. & Smit, H. (2006) Acari, Hydrachnidia I. In: Gerecke, R. (ed.): *Süßwasserfauna von Mitteleuropa*, Vol. 7, 2-1 (2007). Spektrum Elsevier, pp. 241–388.
- Den Heyer, J. (2011) Some statistics on the taxonomy of the family Cunaxidae (Acari: Prostigmata). In: Moraes, G.J. de & Proctor, H. (Eds), *Acarology XIII: Proceedings of the International Congress. Zoosymposia*, 6, 34–38.
- Doğan, S. & Dönel, G. (2010) *Cryptofavognathus*, a new genus of the family Cryptognathidae Oudemans (Acari: Raphignathoidea), with the description of a new species from Turkey. *Zootaxa*, 2533, 36–42.
- Dunlop, J.A. (2007) A large parasitengonid mite (Acari, Erythraeoidea) from the Early Cretaceous Crato Formation of Brazil. *Fossil Record*, 10, 91–98.
- Dunlop, J.A. & Bertrand, M. (2011) Fossil labidostomatid mites (Prostigmata: Labidostomatidae) from Baltic amber. *Acarologia*, 51(2), 191–198.
- Dunlop, J.A., Penney, D. & Jekel, D. (2011) A summary list of fossil spiders and their relatives. In: Platnick, N. I. (ed.) The world spider catalog, version 12.0. American Museum of Natural History, online at <http://research.amnh.org/entomology/spiders/catalog/index.html> DOI: 10.5531/db.iz.0001.
- Fain, A. (1964) Les Ereyetidae de la collection Berlese à Florence designation d'une espèce type pour le genre *Ereynetes* Berlese. *Redia*, 49, 87–111.
- Fan, Q.-H. (1997) The Homocaligidae (Acari: Raphignathoidea) from China, with description of two new species. *Entomologia Sinica*, 4, 337–342.
- Fan, Q.-H. (2004) A catalogue of the genus *Eupalopsellus* Sellnick (Acari: Prostigmata, Eupalopsellidae) with the description of a new species from China. *Biologia, Bratislava*, 59(5), 533–545.
- Fan, Q.-H. (2005) Superfamily Raphignathoidea. In: J. Hallan, (Ed.), *Synopsis of the described Actinedida of the World*. Available from: <http://insects.tamu.edu/research/collection/hallan/acari/Family/Actinedida1.htm> (accessed 5 Dec. 2011).
- Fan, Q.-H. & Walter, D.E. (2011) *Acamerobia inflatus* gen. n. & sp. n. from Australia (Acari: Prostigmata: Raphignathoidea: Camerobiidae) with notes on the idiosomal chaetotaxy. *Zootaxa*, 3045, 45–56.
- Fan, Q.-H., Walter, D.E. & Proctor, H.C. (2003) A Review of the family Barbutiidae (Acari: Raphignathoidea), with the description of two new species from Australia. *Systematic and Applied Acarology*, 8, 107–130.
- Fan, Q.-H. & Zhang, Z.-Q. (2005) *Raphignathoidea (Acari: Prostigmata)*. *Fauna of New Zealand* 52. Manaaki Whenua Press. 400 pp.
- Feider, Z. (1977) Quelques acariens cavernicoles de Cuba de la super-famille Trombiculoidea. In: T. Orghidan, A. Núñez Jimenez et al. (Eds.), *Résultats des expéditions biospéologiques cubano-roumaines à Cuba*. 2. Institut de Spéologie "Emil Racovitza" Romania, pp. 201–208.
- Franz, H. (1952) Revision der Caeculidae Berlese 1883 (Acari). *Bonner zoologische Beiträge*, 3(1–2), 91–124.
- Fuangularwon, M. & Lekprayoon, C. (2010) *Adamystis thailandensis* sp. nov. (Acari: Prostigmata: Adamystidae), a new species of soil mites from Thailand with a key to world species of Adamystidae. *Zootaxa*, 2649, 61–68.
- Gabryś, G. (1999) The world genera of Microtrombidiidae (Acari, Actinedida, Trombidoidea). *Monographs of the Upper Silesian Museum*, 2, 1–361.
- Giesen, K.M.T. (1990) A review of the parasitic mite family Psorergatidae (Cheyletoidea: Prostigmata: Acari) with hypotheses on the phylogenetic relationships of species and species groups. *Zoologische Verhandlungen*, 259, 1–69.
- Haitlinger, R. (1999) Three new species of larval Chyzeriidae associated with Orthoptera (Insecta) from Cyprus, Crete and Peru, with description of the new subfamily Perumaroptinae and three new genera *Napassenia*, *Cretessenia* and *Perumaropta* (Acari: Prostigmata). *Bollettino della Società Entomologica Italiana*, 131, 3–13.
- Haitlinger, R. (2007) A new larval neothrombiid, *Giftitrombium skalaensis* gen. n., sp. n. (Acari: Prostigmata: Neothrombiidae) from Greece. *Zeszyty Naukowe Uniwersytetu Przyrodniczego we Wrocławiu, Biologia Hodowla Zwierząt LV*, (559), 17–22.
- Hajjiganbar, H. & Khaustov, A. (2010) A new species group and five new species of the genus *Pavania* (Acari: Dolichocybidae) associated with insects, with notes on leg chaetotaxy and the distribution of genera. *European Journal of Entomology*, 107(3), 441–453.
- Harvey, M.S. (1996) A revised placement for *Austrotrombella* Southcott (Acarina: Hydryphantidae). *Transactions of the Royal Society of South Australia*, 120, 37–40.
- Hernandes, F.A., Daud, R.D. & Feres, R.J.F. (2007) A new species of *Hexabdella* (Acari: Bdellidae) from Brazil. *Zootaxa*, 1501, 57–63.
- Hernandes, F.A. & Feres, R.J.F. (2006) *Tetrabdella neotropica* (Acari: Bdellidae), a new genus and species from Brazil. *Zootaxa*, 1135, 57–68.
- Jesionowska, K. (1989) New genus and new species of mite of the family Penthalodidae (Actinotrichida, Actinedida, Eupodoidea) from Poland. *Acta Zoologica Cracoviensia*, 32(3), 57–67.
- Jesionowska, K. (2008) Redescription of *Hawaiieupodes thermophilus* Strandtmann et Goff, 1978 (Acari: Prostigmata: Eupodoidea: Penthalodidae) from Hawaii, with a discussion of the systematic status of the taxon. *Annales Zoologici*, 58(2), 337–346.
- Jesionowska, K. (2010) Coceupodidae, a new family of eupodoid mites, with description of a new genus and two new species from Poland. Part I. (Acari: Prostigmata: Eupodoidea). *Genus*, 21(4), 637–658.
- Judson, M. (1994) Studies on the morphology and systematics of the Teneriffiidae (Acari, Prostigmata). I: a new species of *Neotener-*

- iffiola* from Namibia. *Acarologia*, 35(2), 115–134.
- Judson, M.L.I. & Mąkol, J. (2009) A mite of the family Tanaupodidae (Arachnida, Acari, Parasitengona) from the Lower Cretaceous of France. *Geodiversitas*, 31(1), 41–47.
- Judson, M. & Wunderlich, J. (2003) Rhagidiidae (Acari, Eupodoidea) from Baltic amber. *Acta Zoologica Cracoviensia*, 46(supplement), 146–152.
- Khanjani, M., Fayaz, B.A. & Ueckermann, E.A. (2011) A new species of the genus *Austroteneriffia* (Acari: Anystina: Teneriffiidae) from western Iran. *International Journal of Acarology*, 37(6), 550–555.
- Khanjani, M., Fayaz, B.A. & Khanjani, M. (2011) Two new species of *Linotetranus* (Parasitiformes: Tetranychoida: Linotetranae) from Iran. *Zootaxa*, 2834, 47–56.
- Khaustov, A.A. (2008) *Mites of the family Scutacaridae of Eastern Palaearctic*. Akademperiodyka, Kiev, 291 pp.
- Khaustov, A.A. & Perkovsky, E.E. (2010) The first fossil record of mites of the family Pyemotidae (Acari: Heterostigmata), with description of a new species of the genus *Pyemotes* from Rovno amber. *Paleontologicheskii Zhurnal*, 4, 56–58.
- Krantz, G.W. (1978) *A Manual of Acarology (second edition)*. Oregon State University Book Stores Inc., Corvallis. 509 pp.
- Lin, J.-Z. & Zhang, Z.-Q. (2002) *Tarsonemidae of the world (Acari: Prostigmata): key to genera, geographical distribution, systematic catalogue & annotated bibliography*. Systematic & Applied Acarology Society, London, 440 pp.
- Lindquist, E.E. & Vercammen-Grandjean, P.H. (1971) Revision of the chigger-like larvae of the genera *Neotrombidium* Leonardi and Monunguis Wharton, with a redefinition of the subfamily Neotrombidiinae Feider in the Trombidiidae (Acarina: Prostigmata). *Canadian Entomologist*, 103, 1557–1590.
- Mąkol, J. (2007) Generic level review and phylogeny of Trombidiidae and Podothrombiidae (Acari: Actinotrichida: Trombidoidea) of the World. *Annales Zoologici*, 57(1), 1–194.
- Mąkol, J. & Gabryś, G. (2005) *Caecothrombium deharvengi* sp. nov. (Acari: Actinotrichida: Eutrombidiidae) from Vietnam, with a proposal of Caecothrombiinae subfam. nov. *Zoologischer Anzeiger*, 243, 227–237.
- Mąkol, J. & Łaydanowicz, J. (2010) A new species of *Valgothrombium* Willmann, 1940, with additional taxonomic data for Valgothrombiinae genera known as larvae (Acari: Prostigmata: Microtrombidiidae). *Zootaxa*, 2647, 16–34.
- Mąkol, J., Łaydanowicz, J. & Kłosińska, A. (2009) A redescription of *Neothrombium neglectum* (Bruyant, 1909) with revision of *Neothrombium* (Acari: Neothrombidiidae) and notes on the biology of species. *International Journal of Acarology*, 35, 275–301.
- Mąkol, J. & Sevsay, S. (2011) Notes on the genus *Dolicho-thrombium* (Acari: Prostigmata: Trombidiidae) with description of a new species. *Zootaxa*, 2971, 1–16.
- Mesa, N.C., Ochoa, R., Welbourn, W.C., Evans, G. & Moraes, G.J. de (2009) A catalog of the Tenuipalpidae (Acari) of the World with a key to genera. *Zootaxa*, 2098, 1–185.
- Meyer, M.K.P. (Smith) & Ueckermann, E.A. (1987) A taxonomic study of some Anystidae (Acari: Prostigmata). *Entomology Memoir 68. Department of Agriculture and Fisheries, Republic of South Africa*, 1–37.
- Meyer, M.K.P. (Smith) & Ueckermann, E.A. (1989) African Raphignathoidea (Acari: Prostigmata). *Entomology Memoir 74. Department of Agriculture and Water Supply, Republic of South Africa*, 1–58.
- Meyer, M.K.P. (Smith) & Ueckermann, E.A. (1997) A review of some species of the families Allochaetophoridae, Linotetranae and Tuckerellidae (Acari: Tetranychoida). *International Journal of Acarology*, 23, 67–92.
- Moraza, M. (2008) First records of Endeostigmata and Sphaerolichina mites (Acari: Sarcoptiformes and Trombidiformes) from the Iberian Peninsula and the Canary Islands. *Boletín de la Asociación Española de Entomología*, 32 (3–4), 293–304.
- Mortazavi, A., Hajiqaanbar, H. & Saboori, A. (2011) A new species of the family Trochometridiidae (Acari: Heterostigmatina) associated with *Paulusiella* sp. (Coleoptera: Elateridae) from Iran. *Zootaxa*, 2746, 57–68.
- Newell, I.M. (1957) Studies on the Johnstonianidae (Acari, Parasitengona). *Pacific Science*, 11 (4), 396–466.
- Newell, I.M. (1960) Charadracarus new genus, Charadracarinae new subfamily (Acari: Johnstonianidae) and the status of *Typhlothrombium* Berlese, 1910. *Pacific Science*, 14, 156–172.
- Olivier, P.A.S. (2008) Dendrochaetidae, a new family of mites (Acari: Prostigmata), with descriptions of a new genus and species from South Africa. *African Zoology*, 43(1), 16–24.
[http://dx.doi.org/10.3377/1562-7020\(2008\)43\[16:DANFOM\]2.0.CO;2](http://dx.doi.org/10.3377/1562-7020(2008)43[16:DANFOM]2.0.CO;2)
- Olivier, P.A.S. (2009) A replacement name for *Dendrodus* Olivier, 2008, and validation of Dendrochaetidae Olivier, 2008 nomen nudum (Acari: Prostigmata). *Zootaxa*, 2293, 68.
- Olivier, P.A.S. & Theron, P.D. (1997) The genus *Eupodes* Koch, 1835 (Acari: Prostigmata: Eupodidae) from southern African soil and vegetation. Part 1. Characterisation of the genus, designation of the type species and descriptions of three new species. *Koedoe*, 40(2), 57–73.
- Otto, J.C. (1999) The taxonomy of *Tarsotomus* Berlese and *Paratarsotomus* Kuznetsov (Acarina: Anystidae: Erythracarinae) with observations on the natural history of *Tarsotomus*. *Invertebrate Taxonomy*, 13, 749–803.
- Otto, J.C. (2000) A cladistic analysis of Erythracarinae (Acarina: Prostigmata: Anystidae), with the description of a new genus. *Systematic Entomology*, 25, 447–484.
- Hajiqaanbar, H. & Khaustov, A. (2010) A new species group and five new species of the genus *Pavania* (Acari: Dolichocybidae) associated with insects, with notes on leg chaetotaxy and the distribution of genera. *European Journal of Entomology*, 107, 441–453.
<http://dx.doi.org/10.1603/AN11006>.
- Qin, T.-K. (1996) A review and cladistic analysis of the Eupodoidea (Acari: Acariformes). *Systematic & Applied Acarology*, 1, 77–105.
- Qin, T.-K. (1997) A reconsideration of the taxonomic position of *Protopenthalodes* (Prostigmata: Eupodoidea). *Systematic & Applied Acarology*, 2, 253–255.
- Rahiminejad, V., Hajiqaanbar, H. & Fathipour, Y. (2011) Redefinition of the genus *Dolichocybe* (Acari: Dolichocybidae), with description of two new species associated with insects. *Annals of the Entomological Society of America*, 104(4), 627–635.

<http://dx.doi.org/10.3161/000345411X603391>

- Saboori, A., Ueckermann, E.A. & Harten, A. van (2008) A new genus of Neothrombiidae (Acari: Trombidioidea) from Yemen. *Zootaxa*, 1925, 23–30.
- Saboori, A., Zhang, Z.-Q. & Nemati, A. (2005) A new genus and species of larval Chyzeriidae (Acari: Prostigmata: Chyzerioidea) from Iran. *International Journal of Acarology*, 31, 51–56.
- Saboori, A., Nemati, A. & Mossahebi, G. (2005). A new genus and species of Trombellidae (Acari: Trombidioidea) from Iran. *International Journal of Acarology*, 31 (1), 45–50.
- Saboori, A., Wohltmann, A. & Hakimitabar, M. (2010) A new family of trombidoid mites (Acari: Prostigmata) from Iran. *Zootaxa*, 2611, 16–30.
- Sayer, R.M., Smiley, R.L. & Walter, D.E. (1992) Report of a teneriffid mite (Acari) in Baltic amber and notes on recent discoveries, *International Journal of Acarology*, 18, 303–305.
- Schatz, H., Behan-Pelletier, V.M., OConnor, B.M. & Norton, R.A. (2011) Suborder Oribatida van der Hammen, 1968. In: Zhang, Z.-Q. (Ed.), *Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*, 3148, 141–148.
- Seeman, O. & Walter, D.E. (2000) A review of the Paratydeidae (Acari: Prostigmata) with descriptions of the first Australian representatives, *Tanytydeus lamington* sp. nov. and *T. kakadu* sp. nov. *Acarologia*, 40, 393–400.
- Skoracki, M. (2011) Quill mites (Acari: Syringophilidae) of the Palaearctic region. *Zootaxa*, 2840, 1–414.
- Smith, I.M. & Cook, D.R. (1991) Water mites. In: Thorp, J.H. & Covich, A.P. (Eds.), *Ecology and classification of North American Freshwater Invertebrates*, Academic Press, New York, pp. 523–592.
- Southcott, R.V. (1961). Studies on the systematics and biology of the Erythraeoidea (Acarina) with a critical revision of the genera and subgenera. *Australian Journal of Zoology*, 9(3), 367–610.
- Southcott, R.V. (1987) The classification of the mite families Trombellidae and Johnstonianidae and related groups, with the description of a new larva (Acarina: Trombellidae: Nothotrombidium) from North America. *Transactions of the Royal Society of South Australia*, 111, 25–42.
- Southcott, R.V. (1993) Revision of the taxonomy of the larvae of the subfamily Eutrombidiinae (Acarina: Microtrombidiidae). *Invertebrate Taxonomy*, 7, 885–959.
- Southcott, R.V. (1994) Revision of the larvae of the Microtrombidiinae (Acarina: Microtrombidiidae) with notes on life histories. *Zoologica*, 144, 1–155.
- Southcott, R.V. (1996) Description of a new Australian mite (Acarina: Trombidioidea), with comments on superfamily classification. *Records of The South Australian Museum*, 29, 55–62.
- Strandtmann, R.W. (1970) Acarina: Eupodiform Prostigmata of South Georgia. *Pacific Insects Monograph*, 23, 89–106.
- Strandtmann, R.W. & Goff, M.L. (1978) The Eupodoidea of Hawaii (Acarina: Prostigmata). *Pacific Insects*, 19(3–4), 121–143.
- Thor, S. & Willmann, C. (1941) Acarina. 71a Eupodidae, Penthaleidae, Penthaleidae, Rhagidiidae, Pachygnathidae, Cunaxidae. *Das Tierreich*, 71, 1–186.
- Ueckermann, E.A. & Khanjani, M. (2002) A new species of the genus *Austroteneriffia* (Acari: Teneriffiidae) from Iran. *Systematic & Applied Acarology*, 7, 167–172.
- Ueckermann, E.A. & Khanjani, M. (2004) A revision of the genus *Anoplocheylus* Berlese (Acari: Pseudocheylidae), with the description of two new and re-description of four known species. *Systematic & Applied Acarology*, 9, 53–68.
- van der Hammen, L. (1972) A revised classification of the mites (Arachnida, Acarida) with diagnoses, a key, and notes on phylogeny. *Zoologische Mededelingen*, 47(22), 273–292.
- van der Schyff, J., Theron, P.D. & Ueckermann, E.A. (2003) Polytrichinae, a new subfamily of Bdellidae (Acari: Prostigmata) from the Afrotropical region. *African Plant Protection*, 9, 19–22.
- Vercammen-Grandjean P.H. (1973) Sur les statuts de la famille des Trombidiidae Leach, 1815 (Acarina: Prostigmata). *Acarologia*, 15(1), 102–114.
- Walter, D.E. & Gerson, U. (1998) Dasythyreidae, new family, and *Xanthodasythyreus* n. g. (Acari: Prostigmata: Raphignathoidea) from Australia. *International Journal of Acarology*, 24(3), 189–197.
- Walter, D.E., Lindquist, E.E., Smith, I.M., Cook, D.R. & Krantz, G.W. (2009) Chapter thirteen Order Trombidiformes. In: Krantz, G.W. & Walter, D.E. (Eds.), *A Manual of Acarology Third edition*. Texas Tech University Press, Lubbock, pp. 233–420.
- Welbourn, W.C. (1991) Phylogenetic studies of terrestrial Parasitengona. In: Dusbabek, F. & Bukva, V. (Eds.), *Modern Acarology Vol. 2*, The Hague: SPB Academic Publishing bv/Prague: Academia, pp. 163–170.
- Witte, H. (1991) The phylogenetic relationships within the Parasitengonae. In: Dusbabek, F. & Bukva, V. (eds.), *Modern Acarology Vol. 2* Academia, Prague and SFB Academic publishing bv, The Hague, pp.171–182.
- Wohltmann, A. (2006) The phylogenetic relationships of and within the Microtrombidiidae (Acari: Prostigmata: Parasitengona). In: Gabryś, G. & Ignatowicz, S. (Eds), *Advances in Polish Acarology*. Wydawnictwo SGGW, Warszawa, pp. 436–457.
- Wohltmann, A. (2010) Notes on the taxonomy and biology of Smarididae (Acari: Prostigmata: Parasitengona). *Annales Zoologici* (Warsawa), 60(3), 355–381.
- Wohltmann, A., Makol, J. & Gabryś, G. (2004) A revision of European Johnstonianinae (Acari: Prostigmata: Parasitengona: Trombidioidea). *Annales Zoologici* (Warsawa), 54 (3), 595–630.
- Yu, L., Zhang, Z.-Q. & He, L. (2010) Two new species of *Pyemotes* closely related to *P. tritici* (Acari: Pyemotidae). *Zootaxa*, 2723, 1–40.
- Zacharda, M. (1980) Soil mites of the family Rhagidiidae (Actinedida: Eupodoidea). Morphology, systematics, ecology. *Acta Universitatis Carolinae Biologica*, 1978(5–6), 489–785.
- Zhang, Z.-Q. (1994) Neothrombiidae (Acari: Trombidioidea) of the world: systematic review with a phylogenetic analysis and descriptions of two new genera. *Oriental Insects*, 28, 205–237.
- Zhang, Z.-Q. (1998) An unusual early-derivative larva of Parasitengona (Acari: Prostigmata) and proposal of a new superfamily. *Sys-*

tematic & Applied Acarology, 3, 159–170.

- Zhang, Z.-Q. (2003) Taxonomic status of *Veithia* Oudemans, 1941 and redescriptions of two species described by Oudemans (Acari: Smarididae). *Zootaxa*, 145, 1–8.
- Zhang, Z.-Q. & Fan, Q.-H. (2005) A new genus of Neothrombiidae (Acari: Trombidioidea) from New Zealand. *Systematic & Applied Acarology*, 10, 155–162.
- Zhang, Z.-Q. & Fan, Q.-H. (2007) Allotanaupodidae, a new family of early derivative Parasitengona (Acari: Prostigmata). *Zootaxa*, 1517, 1–52.
- Zhang, Z.-Q. & Saboori, A. (1995) A new record of the little known genus *Ralphaudyna* (Acari: Chyzeriidae) from Iran, with description of a new species. *International Journal of Acarology*, 21, 11–15.

Contributor roles and addresses (order of authors based on the amount of data contributed)

- Zhi-Qiang Zhang (coordination, higher taxonomy and superfamilies/families not covered by others) New Zealand Arthropod Collection, Landcare Research, Private Bag 92170, Auckland, New Zealand; zhangz@landcareresearch.co.nz
- Qing-Hai Fan (Raphignathoidea & Tetranychoidae) Plant Health & Environment Laboratory, MAF Biosecurity New Zealand, 231 Morrin Road, St Johns, PO Box 2095, Auckland 1072, New Zealand; qinghai.fan@maf.govt.nz
- Vladimir Pesic (fresh water mite families except Limnesiidae, Unionicolidae, Arrenuridae, Pionidae) Department of Biology, University of Montenegro, Cetinjski put b.b., 81000 Podgorica, Montenegro; vladopesic@gmail.com
- Harry Smit (fresh water mite families specially Limnesiidae, Unionicolidae, Arrenuridae, Pionidae) Netherlands Centre for Biodiversity Naturalis, P.O. Box 9517, 2300 RA Leiden, The Netherlands; Harry.Smit@ncbnaturalis.nl
- Andre V. Bochkov (Cheyletoidea, Myobioidea, Pomerantizioidea, Pterygosomatoidea & Stygmochelidae) Zoological Institute of the Russian Academy of Sciences, Universitetskaya Embankment 1, 199034 St. Petersburg, Russia; andrevbochkov@gmail.com
- A. A. Khaustov (Pygmephoroidae, Acarophenacidae, Caraboacaridae & Trochometriviidae) Nikita Botanical Gardens, National Scientific Center, Yalta, 98648 Ukraine; alkhaustov@mail.ru
- Anne Baker (Eupodoidea) Department of Entomology, The Natural History Museum, London SW7 5BD, UK
- Andreas Wohltmann (Tanaupodidae, Chyzeriidae, Johnstonianidae, Trombellidae, Microtrombidiidae) Findorffstr. 11, D-27721 Ritterhude, Germany; wohltman@uni-bremen.de
- Tinghuan Wen (Trombiculidae, Leeuwenhoekidae & Walchiidae) Shanghai Medical School, Fudan University, Shanghai, China
- James W. Amrine (Eriophyoidea) Division of Plant and Soil Sciences, P.O. Box 6108, West Virginia University, Morgantown, WV 26506-6018, USA
- P. Beron (Calyptostomatidae, Erythraeidae & Smarididae) National Museum of Natural History, Sofia, Hungary; beron@mail.bg
- Jianzhen Lin (Tarsonemidae & Bdellidae) Research Institute of Plant Protection, Fujian Academy of Agricultural Sciences, Fuzhou 350013, China
- Grzegorz Gabrys (Microtrombidiidae) Department of Zoology, University of Zielona Góra, Szafrana 1, 65-516 Zielona Góra, Poland; g.gabrys@wnb.uz.zgora.pl
- Robert Husband (Podapolipidae) Biology Department, Adrian College, Adrian, Michigan 49221, U.S.A.; husbandadrian@aol.com