

Table 1. Diagnostic characters for genera of Haplomunnidae

Diagnostic characters	<i>Haplomunna</i>	<i>Thylakogaster</i>	<i>Abyssamea</i>	<i>Munella</i>	New species
Head lateral depth < or = depth of pereonite 1	<	=	=	=	=
Body and pleotelson surfaces with spines	x	x	x		
Body and pleotelson surfaces with setae only				x	x
Pereonites 1-4 with dorsal transverse sculpture	x				
Pleotelson:					
• strongly inflated, balloonlike, reflected ca. 90° over pereonites, dorsal surface overreaching pereonite 2		x			x
• moderately inflated, reflected ca. 90° over pereonites, dorsal surface barely reaching pereonite 4	x				
• dorsoproximal part inflated, covering pereonite 6, not recurved			x		
• not inflated, not recurved				x	
Antennula and antenna inserting on joint long projection					x
Antennula and antenna inserting directly on head surface, at most on low hump	x	x	x	x	
Mandible with (+) or without (-) palp	+	-	+	+	+
Mandible molar apex:					
• truncate, bi-ridged	x				
• truncate with fringe of setae			x		
• wedge shaped with thin spines		x		x	
• wedge shaped with row of short, curved setae					x
Pereopod I ischium shorter (<) or longer (>) than basis	<	<	<	<	>
Pereopod I carpus broad, ovoid, length ca. twice width		x	x	x*	
Pereopod I carpus slender, elongate, length 3-5 times width	x				x
Pereopod I propodus with posterior row of sharp robust setae (+) or not (-)	+	-	-	-	+
Pereopod VII present (+) or absent (-)	+	+	-	-	+
Uropods:					
• single article, simple	x		x		
• single article, apex bilobed		x			
• two article, apex simple				x	x

* According to description in Wilson and Hessler (1974).