



## A new species of *Coelogyne* (Orchidaceae) from western Yunnan, China

RONG LI<sup>1</sup> & ZHI-LING DAO<sup>1</sup>

<sup>1</sup>Key Laboratory for Plant Diversity and Biogeography of East Asia, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, Yunnan, China; email: daozhl@mail.kib.ac.cn

### Abstract

A new species of Orchidaceae, *Coelogyne pianmaensis*, is described and illustrated from western Yunnan, China. Morphologically, it is similar to *Coelogyne corymbosa*, but differs from the latter by having dark brown eyelike blotches surrounded by brownish-yellow on lip, 1-veined obovate-oblong leaves, 7-veined oblong-lanceolate sepals, shorter pedicel and ovary, and 7-veined petals.

**Key words:** Arethuseae, Epidendroideae, Gaoligong Mountains, taxonomy

### Introduction

The genus *Coelogyne* Lindley (1821: tab. 33) includes about 200 species distributed from India and Sri Lanka across southern Asia and the Malay archipelago to the Philippines, New Guinea, and the south west Pacific Islands (Gravendeel *et al.* 2005, Wu *et al.* 2006, Mabberley 2008). Thirty-one species are distributed in China, six of which are endemic (Chen & Clayton 2009).

During the past fifteen years, a series of expeditions to the western Yunnan for the project “Biodiversity Survey of Gaoligong Mountains” have been carried out jointed by the Kunming Institute of Botany, Chinese Academy of Sciences and Department of Botany, California Academy of Sciences. Among the *Coelogyne* specimens collected, we found one species with morphological features that differed from the other species of *Coelogyne* known. After a morphological study, literature search (Chen & Tsi 1998, Chen 1999, Chen *et al.* 1999, Li & Bartholomew 1999, Clayton 2002, Pearce & Cribb 2002, Li & Li 2003, Chen & Clayton 2009, George & George 2011, Subedi 2011), and examination of many other specimens in the KUN and PE herbaria, we concluded it represents a new species of *Coelogyne*, described and illustrated below.

### *Coelogyne pianmaensis* R. Li & Z. L. Dao, sp. nov. (Fig. 1)

*Species Coelogynis corymbosae affinis, sed labelli maculis oculis atrobrunneis a marginatis vaccinis, folia obovato-oblonga uninervia, sepala oblongo-lanceolata 7-nerviis, pedicello cum ovario breviori circiter 1.5 cm longo, petala 7-nerviis differt.*

**Type:**—CHINA. Yunnan: Nujiang Lisu people autonomous Prefecture, Lushui County, Pianma township, Yaojiaping, E side of Gaoligong Mountains, primary evergreen broad-leaved forest, on *Tsuga* trees, 2,800 m, 25°58'14.1"N, 98°41'57.1"E, 22 April 2003, Liu 846 (holotype KUN!).

Plant epiphytic, 8–14.5 cm tall. Rhizome stout, 3–4 mm in diameter, densely covered by brown, scaly sheaths. Pseudobulbs clustered, developing serially, less than 1 cm apart from each other, oblong-ovoid or rhombic-oblong, brightly yellow and strongly wrinkled when dried, 20–35 × 6–10 mm, 2 leaves at apex, sheathed at base; sheaths ovate, papery, dark brown, 1.2–2.2 cm long. Leaf blade obovate-oblong, subcoriaceous, 5–10 × 1–2 cm, 1-veined, apex acute; petiole grooved, 5–8 mm long. Inflorescence proteranthous to synanthous, erect, 6–11 cm long, covered by overlapping sheaths in lower part; raceme 2- to 3-flowered (opening together); floral bracts oblong-lanceolate, 9–11 × 1–2 mm, apex acute, caducous. Flowers fragrant, fleshy, white with 4 dark brown eyelike

## **Identification key to the species of *Coelogyne* sect. *Ocellatae***

1. Inflorescence hysteranthous .....	2
- Inflorescence proteranthous or synanthous .....	3
2. Lowermost bract of rachis fertile .....	<i>C. hysterantha</i>
- Lowermost bract of rachis sterile .....	<i>C. punctulata</i>
3. Pseudobulb ovoid, born distantly, 1.3–5.0 cm apart .....	<i>C. occultata</i>
- Pseudobulb ellipsoid, born clustered, less than 1.3 cm apart .....	4
4. Lamellae on lip plate-like shaped .....	<i>C. platylamellata</i>
- Lamellae on lip clavate or rod-like shaped .....	5
5. Lamellae extending ca. 3 mm away from lip base .....	<i>C. gongshanensis</i>
- Lamellae extending from lip base .....	6
6. Lateral lobes length 3–3.2 cm when flattened .....	<i>C. wardii</i>
- Lateral lobes length 1–2.5 cm when flattened .....	7
7. Margin of lateral lobes crispate .....	<i>C. taronensis</i>
- Margin of lateral lobes crenulate .....	8
8. Lateral lamellae with entire margins .....	9
- Lateral lamellae with crenulate margins .....	10
9. Flowers 4–8, mid-lobe acute at apex .....	<i>C. nitida</i>
- Flowers 2–3, mid-lobe cuspidate at apex .....	<i>C. tyyuii</i>
10. Yellow eyelike blotches surrounded by reddish orange on lip .....	<i>C. corymbosa</i>
- Dark brown eyelike blotches surrounded by brownish-yellow on lip .....	<i>C. pianmaensis</i>

**Additional specimens examined (paratypes):**—CHINA. Yunnan: Nujiang Lisu people autonomous Prefecture, Lushui County, Pianma township, Yaojiaping, E side of Gaoligong Mountains, primary evergreen broad-leaved forest, on *Tsuga* trees, 2952 m, 25°58'46.2"N, 98°41'52.1"E, 28 April 2003, *Liu* 953 (KUN!); Pianma township, Pianma pass, E side of Gaoligong Mountains, coniferous-broad leaved mixed forest, on *Tsuga* trees, 3130 m, 25°58'21.4"N, 98°41'01.4"E, 14 April 2005, *Li & Dao* 127 (KUN!); Luzhang township, Gangfang, E side of Gaoligong Mountains, primary evergreen broad-leaved forest, on *Tsuga* trees, 2600 m, 25°57'49.9"N, 98°41'22.3"E, 20 April 2006, *Li & Dao* 378 (KUN!); Luzhang township, Tingming lake, E side of Gaoligong Mountains, coniferous-broad leaved mixed forest, on *Tsuga* trees, 3010 m, 25°59'01.8"N, 98°41'26.1"E, 28 April 2007, *Li & Dao* 509 (KUN!).

## **Acknowledgements**

The authors kindly thank Mr. Yitao Liu from Kunming Institute of Botany, Chinese Academy of Sciences for the illustration. The study was supported by the National Natural Science Foundation of China (grant no. 31370243), the open fund of Key Laboratory of Biodiversity and Biogeography, Kunming Institute of Botany, Chinese Academy of Sciences (grant no. KLBB201204), and the international cooperation program of Bureau of International Cooperation of Chinese Academy of Sciences (grant no. GJHZ1140).

## **References**

- Chen, X.-Q. & Clayton, D. (2009) *Coelogyne* Lindley. In: Wu, Z.-Y. & Raven, P.H. (eds.) *Flora of China*, 25. Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, pp. 315–325.
- Chen, X.-Q. & Tsai, Z.-H. (1998) *The orchids of China*. China Forestry Publishing House, Beijing, 282 pp.
- Chen, X.-Q. (1999) *Coelogyne* Lindley. In: Wu, Z.-Y. (ed.) *Flora Republicae Popularis Sinicae*, 18. Science Press, Beijing, pp. 338–364.
- Chen, X.-Q., Tsai, Z.-H. & Luo, Y.-B. (1999) *Native orchids of China in colour*. Science Press, Beijing, 416 pp.
- Clayton, D. (2002) *The genus Coelogyne: a synopsis*. Natural History Publications, Kota Kinabalu, 316 pp.
- George, E. & George, J.C. (2011). *Les Coelogynes*. Belin, Paris, 607 pp.
- Li, H. & Bartholomew, B. (1999) The diversity of Orchidaceae in Gaoligong Mountains, Yunnan. *Acta Botanica Yunnanica* (Suppl.) 11: 65–78.
- Li, R. & Li, H. (2003) *Coelogyne* Lindley. In: Li, H. (ed.) *Flora Yunnanica*, 14. Science Press, Beijing, pp. 511–530.
- Lindley, J. (1821) *Collectanea Botanica*. Richard & Arthur Taylor, Shoe-Lane, London, 41 pp.

- Lindley, J. (1854) *Coelogynie*. In: Lindley, J. (ed.) *Folia orchidacea: an enumeration of the known species of orchids*. Published for the author by Matthews J., London, pp. 1–18.  
<http://dx.doi.org/10.5962/bhl.title.51709>
- Mabberley, D.J. (2008) *Mabberley's plant-book, a portable dictionary of plants, their classification and uses*. Cambridge University Press, New York, 1021 pp.
- Pearce, N.R. & Cribb, P.J. (2002) *The orchids of Bhutan*. Royal Botanic Garden Edinburgh, Edinburgh, 643 pp.
- Pfitzer, E. & Kraenzlin, F. (1907). *Coelogynie*. In: Engler, H.G.A. (ed.) *Das Pflanzenreich, Heft 32*. Akademie Verlag, Berlin, pp. 20–82.
- Gravendeel, B., de Vogel, E.F. & Schuiteman, A. (2005). *Coelogynie*. In: Pridgeon, A.M., Cribb, P.J., Chase, M.W. & Rasmussen, F.N. (eds.) *Genera Orchidacearum*, 4. Oxford University Press, New York, pp. 44–51.
- Subedi, A. (2011) *New species, pollinator interactions and pharmaceutical potential of Himalayan orchids*. Dissertation, Leiden University, 58 pp.
- Wu, Z.-Y., Zhou, Z.-K., Sun, H., Li, D.-Z. & Peng, H. (2006) *The areal-types of seed plants and their origin and differentiation*. Yunnan Science and Technology Press, Kunming, 566 pp.