



<http://dx.doi.org/10.11646/phytotaxa.162.4.7>

## Notes on Early Land Plants Today. 48. *Drepanolejeunea longifolia*, a new synonym of *Drepanolejeunea angustifolia* (Marchantiophyta, Lejeuneaceae)

QIONG HE<sup>1,2</sup> & RUI-LIANG ZHU<sup>1,3,4</sup>

<sup>1</sup>Department of Biology, School of Life Sciences, East China Normal University, 3663 Zhong Shan North Road, Shanghai 200062, China; rlzhu@bio.ecnu.edu.cn

<sup>2</sup>Ningbo Binhai International Cooperative School, Ningbo Binhai Education Group Ltd., 11 Mingyue Road, Binhai New City, Beilun, Ningbo, Zhejiang 315830, China

<sup>3</sup>Shanghai Key Lab for Urban Ecological Processes and Eco-Restoration, East China Normal University, 500 Dongchuan Road, Shanghai 200241, China

<sup>4</sup>Author for correspondence

*Drepanolejeunea* (Spruce 1884: 186) Stephani (1891: 209), with about 110 currently accepted species, is the fourth largest genus of Lejeuneaceae (following *Cololejeunea* (Spruce 1884: 291) Stephani (1891: 208), *Lejeunea* Libert (1820: 373), and *Cheilolejeunea* (Spruce 1884: 251) Stephani (1890: 284) (He *et al.* 2012). Although *Drepanolejeunea* is a diverse pantropical genus, it is characterized and easily distinguished from other related genera by the 1) erect to widely divergent lobes of underleaves without a basal disc, 2) transverse section of stem with 7 cortical cells and 3 medullary cells, 3) presence of ocelli at least at base of leaf lobe, 4) proximal hyaline papilla, 5) pycnolejeuneoid leaf sequence of gynoecial innovation (if present), 6) inflated perianths usually with various projections near apex, and 7) asexual reproduction by means of cladia or caducous leaves (Zhu & So 2001; He *et al.* 2012; Pócs *et al.*, 2013). The genus has been classified in a separate subtribe, *Drepanolejeuneinae* Gradstein (2013: 14), together with *Vitalianthus* Schuster & Giancotti (1993: 447).

During our recent studies on Asian *Drepanolejeunea*, we found that *Drepanolejeunea longifolia* Singh & Nath (2007: 248), previously known only from India is conspecific with *Drepanolejeunea angustifolia* (Mitten 1861: 116) Grolle (1965: 206), a common epiphytic and epiphyllous species in southeastern Asia (e.g., Udar & Awasthi 1982; Zhu & So 2001).

### Formal treatment

The format of this note follows Söderström *et al.* (2012).

***Drepanolejeunea angustifolia* (Mitt.) Grolle, J. Jap. Bot. 40: 206. 1965 (Grolle 1965).**

≡ *Lejeunea angustifolia* Mitt., J. Proc. Linn. Soc., Bot. 5: 116. 1861 “1860” (Mitten 1861). Type:—INDIA. “Sikkim Himalaya”: Tonglo, 10000 ft, *J. D. Hooker* 1498 (holotype NY!; isotype BM!).

= *Drepanolejeunea longifolia* Ajit P.Singh et V.Nath, Hepat. Khasi Jaintia Hills E. Himal.: 248, 2007 (Singh & Nath 2007); Bryologist 111: 477, 2008 (Singh *et al.* 2008). Type:—INDIA. Meghalaya: East Khasi Hills, Elephant Falls, on bark of tree, ca. 1500 m, 8 June 1965, *S. Chandra* 201213-B (holotype LWG!), *syn. nov.*

For further synonyms, see Mizutani (1961), Grolle (1965), Zhu & So (2001).

### Acknowledgements

This research was sponsored by the National Natural Science Foundation of China (nos. 31370238, 31170190), the Special Program for the National Basic Work of the Ministry of Science and Technology, China (No. 2012FY110600), and 211 Project for the East China Normal University. We also thank the curators of BM, LWG and NY, Virendra Nath (LWG ) and A.K. Asthana (LWG) for making specimens available for study through loans or visits.

## References

- Gradstein, S.R. (2013) A classification of Lejeuneaceae (Marchantiophyta) based on molecular and morphological evidence. *Phytotaxa* 100: 6–20.  
<http://dx.doi.org/10.11646/phytotaxa.100.1.2>
- Grolle, R. (1965) Miscellanea hepaticologica. 41–50. *Journal of the Hattori Botanical Laboratory* 40: 206–219.
- He, Q., Zhu, R.-L., Chantanaorapint, S., Kornochalert, S. & Printarakul, N. (2012) *Drepanolejeunea laciniata* (Lejeuneaceae), a new species from northern Thailand. *Cryptogamie Bryologie* 33: 291–298.  
<http://dx.doi.org/10.7872/cryb.v33.iss3.2012.291>
- Libert, M.-L. (1820) Sur un genre nouveau d'hépatiques, Lejeunia. *Annales générales des sciences physiques* 6: 372–374.
- Mitten, W. (1861) Hepaticae Indiae Orientalis: An enumeration of the Hepaticae of the East Indies. *Journal of the Proceedings of the Linnean Society. Botany* 5: 89–128.  
<http://dx.doi.org/10.1111/j.1095-8312.1861.tb01343.x>
- Mizutani, M (1961) A revision of Japanese Lejeuneaceae. *Journal of the Hattori Botanical Laboratory* 24: 115–302.
- Pócs, T., Luong, T.-T. & Ho, B.-C. (2013) New or little known epiphyllous liverworts, XVIII. Records from the Bidoup-Núi Bà National Park, Vietnam, with the description of *Drepanolejeunea bidoupensis*, sp. nov. *Cryptogamie Bryologie* 34: 287–298.  
<http://dx.doi.org/10.7872/cryb.v34.iss3.2013.287>
- Schuster, R.M. & Giancotti, C. (1993). On *Vitalianthus* Schust. & Giancotti, a new genus of Lejeuneaceae. *Nova Hedwigia* 57: 445–56.
- Singh, A.P. & Nath, V. (2007) Hepaticae of Khasi and Jaintia Hills: Eastern Himalayas. Bishen Singh Madendra Pal Singh, Dehra Dun, 1–382.
- Singh, A.P., Nath, V. & Asthana, A.K. (2008) Studies on two noteworthy species of *Drepanolejeunea* (Lejeuneaceae) from Meghalaya (India). *The Bryologist* 111: 476–481.  
[http://dx.doi.org/10.1639/0007-2745\(2008\)111\[476:sotnso\]2.0.co;2](http://dx.doi.org/10.1639/0007-2745(2008)111[476:sotnso]2.0.co;2)
- Söderström, L., Hagborg, A. & von Konrat, M. (2012) Notes on Early Land Plants Today. *Phytotaxa* 65: 41–42.  
<http://dx.doi.org/10.11646/phytotaxa.112.1.3>
- Spruce, R. (1884) Hepaticae of the Amazon and the Andes of Peru and Ecuador. *Transactions and Proceedings of the Botanical Society of Edinburgh* 15: 1–308.
- Stephani, F. (1890) Hepaticae africanae novae in insulis Bourbon, Maurice et Madagascar lectae. *Botanical Gazette* 15: 281–292.  
<http://dx.doi.org/10.1086/326585>
- Stephani, F. (1891) Hepaticae africanae. *Hedwigia* 30: 201–217.
- Udar, R. & Awasthi, U.S. (1982) The genus *Drepanolejeunea* in India. *Journal of the Hattori Botanical Laboratory* 53: 419–437.
- Zhu, R.-L. & So, M.L. (2001) Epiphyllous liverworts of China. *Beihefte zur Nova Hedwigia* 121: 1–418.