



## Notes on Early Land Plants Today. 60. Circumscription of Gymnomitriaceae (Marchantiophyta)

JIRÍ VÁŇA<sup>1</sup>, LARS SÖDERSTRÖM<sup>2</sup>, ANDERS HAGBORG<sup>3</sup> & MATT VON KONRAT<sup>3</sup>

<sup>1</sup>Department of Botany, Charles University, Benátská 2, CZ-12801 Praha 2, Czech Republic; email: vana@natur.cuni.cz

<sup>2</sup>Department of Biology, Norwegian University of Science and Technology, N-7491 Trondheim, Norway; lars.soderstrom@bio.ntnu.no

<sup>3</sup>Department of Science and education, The Field Museum, 1400 South Lake Shore Drive, Chicago, IL 60605–2496, USA; hagborg@pobox.com, mvonkonrat@fieldmuseum.org

Gymnomitriaceae Klinggräff (1858: 16) was thoroughly treated by Váňa *et al.* (2010). Since the publication, a series of papers have shown that the circumscription of the family and genera within the family must be revised.

*Herzogobryum* Grolle (1963: 160) and *Nothogymnomitrium* Schuster (1996: 43) was removed from the family by Váňa *et al.* (2013). Already de Roo *et al.* (2007) showed that *Apomarsupella revoluta* (Nees 1836: 419) Schuster (1996: 82) was nested in *Gymnomitrium* Corda (1829: 651) and Cailliau *et al.* (2013) showed that *Apomarsupella rubida* (Mitten 1861: 90) Schuster (1996: 85) also belongs there. Thus the genus *Apomarsupella* Schuster (1996: 79) should be synonymized and the necessary combinations made. The necessary combinations are made elsewhere.

Two invalidly published names in *Marsupella* Dumortier (1822: 114) were used in Váňa *et al.* (2010). The first of these, *Marsupella emarginata* (Ehrhart 1784: 141) Dumortier (1835: 24) var. *tubulosa*, was assumed to have been established as an autonym under subsp. *tubulosa* (Stephani 1897: 99) Kitagawa (1960: 76) via the publication of var. *patens* (Stephani 1901: 162) Kitagawa (1960: 77) or of var. *apertifolia* (Stephani 1901: 162) Kitagawa (1963:89). However, autonoms are only established directly under genera and species (cf. ICN Art. 22.1 & 26.1; McNeill *et al.* 2012) and thus var. *tubulosa* is invalid, which Váňa *et al.* (2010) overlooked. The second name, *Marsupella sparsifolia* (Lindberg 1868: 280) Dumortier (1874: 128) subsp. *childii* Schuster (1996: 61), was invalidly published by Schuster (1996) as he did not indicate the herbarium in which the type specimen was lodged (as required under ICN Art. 40.7; McNeill *et al.* 2012). Váňa *et al.* (2010) explicitly did not validate the subspecies as they had not located the type material. However, with the incorporation of Schuster's herbarium in F, the type material has now been found. Those two names are thus validated here.

Vilnet *et al.* (2010, 2011) showed that *Nardia* Gray (1821: 694) is related to Gymnomitriaceae and not to Solenostomataceae Crandall-Stotler *et al.* (2009: 190), a position also indicated by other studies. As *Nardia* forms a separate lineage sister to the rest of the Gymnomitriaceae, and differs morphologically in several important characters (see bold characters in the description below), it seems appropriate to separate out two subfamilies: Gymnomitrioideae Schuster (1996: 80) and Nardioideae, subfam. nov.

### Formal treatment

The format of this note follows that which is outlined in Söderström *et al.* (2012).

### Nardioideae Váňa, subfam. nov.

Description:—Leaves succubous, never transverse, undivided or bilobed, not interlocking dorsally (insertions not extending across the stem midline). Underleaves always present, mostly small, lanceolate. Gametangia on leading axes. Sporophytes enclosed by a shoot calyptra, perianth and perigynium (erect, of *Isotachis*-type or short pendent, of *Nardia geoscyphus*-type) always present. Capsules subsphaeroidal to shortly ellipsoidal, innermost wall cells with semiannular thickenings. Gemmae absent.

Type:—*Nardia* Gray.

*Marsupella emarginata* var. *tubulosa* (Steph.) N.Kitag. ex Váňa et L.Söderstr., comb. nov., Basionym:—*Marsupella tubulosa* Steph., *Bull. Herb. Boissier* 5: 99, 1897 (Stephani 1897). ≡ *Marsupella emarginata* var. *tubulosa* N.Kitag., *Mem. Coll. Sci. Kyoto Imp. Univ., Ser. B, Biol.* 27: 77, 1960 (Kitagawa 1960) *nom. inval.* (ICN Art. 41.5; basionym not cited).

*Marsupella sparsifolia* subsp. *childii* R.M.Schust. ex Váňa et L.Söderstr., subsp. nov., Based on:—*Marsupella sparsifolia* subsp. *childii* R.M.Schust., *J. Hattori Bot. Lab.* 80: 61, 1996 (Schuster 1996) *nom. inval.* (ICN Art. 40.7; herbarium not indicated). Type:—NEW ZEALAND. Fiordland Nat. Park: Margin of tarn, snow tussock zone, Mt. Burns track, *Schuster 84-142* (F, holotype). Note:—The reference to the description in Schuster (1996: 61) together with the specification of the holotype here validates the name.

## Acknowledgements

The Early Land Plants Today project (ELPT) has been generously supported in part by the Global Biological Information Facility (GBIF) Seed Money Award No.2007-41, activities facilitated in part by funding from the Biodiversity Synthesis Center of the Encyclopedia of Life (BioSynC), partial funding from the National Science Foundation (Award No's 0749762, 1115002), the Warwick Foundation, and the Negaunee Foundation.

## References

- Cailliau, A., Long, D.G., Price, M.J. & Perret, M. (2013) Phylogeny and systematic position of *Mesoptychia* (Lindb.) A. Evans. *Plant Systematics and Evolution* 299: 1243–1251.  
<http://dx.doi.org/10.1007/s00606-013-0792-z>
- Corda, A.J.C. (1829) Genera Hepaticarum. Die Gattungen der Lebermoose. *In*: Opiz, P.M. (Ed.) *Beiträge zur Naturgeschichte als Fortsetzung des Naturalientausches No. 12*. C.W. Enders, Praha, pp. 643–655.
- Crandall-Stotler, B.J., Stotler, R.E. & Long, D.G. (2009) Phylogeny and classification of the Marchantiophyta. *Edinburgh Journal of Botany* 66: 155–198.  
<http://dx.doi.org/10.1017/s0960428609005393>
- De Roo, R.T., Hedderson, T.A. & Söderström, L. (2007) Molecular insights into the phylogeny of the leafy liverwort family Lophoziaceae Cavers. *Taxon* 56: 301–314.
- Dumortier, B.C. (1822) *Commentationes botanicae*. Ch. Casterman-Dien, Tournay, 118 pp.  
<http://dx.doi.org/10.5962/bhl.title.10534>
- Dumortier, B.C. (1835) *Recueil d'Observations sur les Jungermanniacées*. J.-A. Blanquart, Tournay, 27 pp.
- Dumortier, B.C. (1874) Jungermannideae Europae post semiseculum recensitae, adjunctis hepaticis. *Bulletin de la Société Royale de Botanique de Belgique* 13: 5–203.  
<http://dx.doi.org/10.5962/bhl.title.26517>
- Ehrhardt, J.F. (1784) Botanische Bemerkungen. *Hannoversches Magazin* 22: 113–144.
- Gray, S.F. (1821) *Natural arrangement of British Plants, according to their relation to each other*. Baldwin, Craddock, and Joy, London, 824 pp.
- Grolle, R. (1963 [1964]) Notulae hepaticologicae VII–IX. *Revue Bryologique et Lichénologique* 32: 157–165.
- Kitagawa, N. (1960) New or interesting species of *Marsupella* in Japan. *Memoirs of the College of Science, Kyoto Imperial University. Series B. Biology* 27: 75–82.
- Kitagawa, N. (1963) A revision of the family Marsupellaceae of Japan. *Journal of the Hattori Botanical Laboratory* 26: 76–118.
- Klinggräff, H.V. (1858) *Die höheren Cryptogamen Preussens*. Vilhelm Koch, Königsberg, 220 pp.
- Lindberg, S.O. (1868) Musci Novi Scandinavici. *Notiser ur Sällskapet pro Fauna et Flora Fennica Förhandlingar* 9: 253–299.
- McNeill, J., Barrie, F.R., Buck, W.R., Demoulin, V., Greuter, W., Hawksworth, D.L., Herendeen, P.S., Knapp, S., Marhold, K., Prado, J., Prud'homme van Reine, W.F., Smith, G.F., Wiersema, J.H. & Turland, N.J. (2012) International Code of Nomenclature for algae, fungi and plants (Melbourne Code) adopted by the Eighteenth International Botanical Congress Melbourne, Australia, July 2011. *Regnum Vegetabile* 154: 1–240.
- Mitten, W. (1861 [1860]) Hepaticae Indiae Orientalis: An enumeration of the Hepaticae of the East Indies. *Journal of the Proceedings of the Linnean Society. Botany* 5: 89–128.  
<http://dx.doi.org/10.1111/j.1095-8312.1861.tb01343.x>
- Nees von Esenbeck, C.G. (1836) *Naturgeschichte der Europäischen Lebermoose mit besonderer Beziehung auf Schlesien und die*

*Oertlichkeiten des Riesengebirgs, vol. 2.* August Rucker, Berlin, 499 pp.

- Schuster, R.M. (1996) Studies on Antipodal Hepaticae. XII. Gymnomitriaceae. *Journal of the Hattori Botanical Laboratory* 80: 1–147.
- Söderström, L., Hagborg, A. & von Konrat, M. (2012) Notes on Early Land Plants Today. *Phytotaxa* 65: 41–42.
- Stephani, F. (1897) Hepaticae Japonicae. *Bulletin de l'Herbier Boissier* 5: 76–108.
- Stephani, F. (1901) Species Hepaticarum 2. *Bulletin de l'Herbier Boissier (sér. 2)* 1: 140–177.
- Vilnet, A.A., Konstantinova, N.A. & Troitsky, A.V. (2010 [2011]) Molecular insight on phylogeny and systematics of the Lophoziaceae, Scapaniaceae, Gymnomitriaceae and Jungermanniaceae. *Arctoa* 19: 31–50.
- Vilnet, A.A., Konstantinova, N.A. & Troitsky, A.V. (2011) Taxonomical rearrangements of Solenostomataceae (Marchantiophyta) with description of a new family Endogemmataceae based on trnL-F cpDNA analysis. *Folia Cryptogamica Estonica* 48: 125–133.
- Váňa, J., Söderström, L., Hagborg, A., von Konrat, J. & Engel, J.J. (2010) Early Land Plants Today: Taxonomy, systematics and nomenclature of Gymnomitriaceae. *Phytotaxa* 11: 1–80.  
<http://dx.doi.org/10.11646/phytotaxa.11.1.1>
- Váňa, J., Söderström, L., Hagborg, A. & von Konrat, M.J. (2013) Notes on Early Land Plants Today. 40. Notes on Cephaloziellaceae (Marchantiophyta). *Phytotaxa* 112: 1–6.  
<http://dx.doi.org/10.11646/phytotaxa.112.1.1>