



<http://dx.doi.org/10.11646/phytotaxa.201.4.2>

A new species of *Kali* (Salsoloideae, Chenopodiaceae) from Sicily, supported by molecular analysis

CRISTIAN BRULLO¹, SALVATORE BRULLO^{1*}, JOHN F. GASKIN², GIANPIETRO GIUSSO DEL GALDO¹, G. FREDERIC HRUSA³ & CRISTINA SALMERI⁴

¹ Dipartimento di Scienze Biologiche, Geologiche e Ambientali, Università di Catania, 95125 Catania, Italy; salvo.brullo@gmail.com

² USDA Agricultural Research Service, Northern Plains Agricultural Research Laboratory, MT 59270 Sidney, U.S.A.; john.gaskin@ars.usda.gov

³ California Department of Food and Agriculture, Plant Pest Diagnostics Branch, 95832.1448 Sacramento, U.S.A.; fhrusa@cdfa.ca.gov

⁴ Dipartimento di Biologia Ambientale e Biodiversità, Università di Palermo, 90123 Palermo, Italy; cristinasalmeri@gmail.com

* Corresponding author

Abstract

Nomenclatural and taxonomical considerations on *Kali*, a controversial genus recently segregated from the polyphyletic *Salsola* s. lat. (*Chenopodiaceae*), are provided. The *Kali* group includes annual plants with leaves ending in a spine and lacking hypodermis, having also a cortex alternate to longitudinal chlorenchymatous striae. The species belonging to this genus mainly have a paleotemperate distribution (Europe, Asia and North Africa), occurring as aliens in North America, Australia and South Africa. A new species collected on Mt. Etna (Sicily), and closely related to *K. australis*, is described and illustrated as *K. basalticum*. Its morphological and molecular features, karyology (2n=54), ecology, distribution, phylogeny and conservation status are examined. In addition, a list of the currently known species of *Kali* is provided, with some new combinations: *Kali nepalense* (Grubov) comb. nov., *Kali pellucidum* (Litvinov) comb. nov., *Kali sinkiangense* (A.J. Li) comb. nov., *Kali gobicolum* (Iljin) comb. nov., and *Kali ryanii* (G.F. Hrusa & Gaskin) comb. nov.

Key words: Chenopodiaceae, haplotype, ITS, *Kali*, karyology, new species, phylogeny, Sicily, taxonomy

Introduction

Salsola Linnaeus (1753: 222) is a rather controversial genus within the tribe *Salsoleae* Moq. (*Chenopodiaceae* Vent.) including succulent plants (at least succulent in the leaves and young branches), which are usually characterized by the fruiting perianth with scarious wings that assist in wind dispersion, and spiral embryos (Moquin-Tandon 1840, Bunge 1862, Freitag 1997, Zhu *et al.* 2003, Kadereit *et al.* 2005).

Several morphological and molecular investigations showed that there are many taxonomic and phylogenetic questions, some of which are still unresolved (Meyer 1829, Moquin-Tandon 1840, Bunge 1862, Volkens 1893, Ulbrich 1934, Iljin 1936, Botschantzev 1969a, 1969b, 1970, 1974a, 1974b, 1981, Aellen & Townsend 1972, Brullo 1984, Tzvelev 1993, Hedge *et al.* 1997, Grubov 2000, Pyankov *et al.* 2001, Kadereit *et al.* 2003, 2005, Kapralov *et al.* 2006, Akhani *et al.* 2007, Wen *et al.* 2010, Kadereit & Freitag 2011, Fëderova 2011, Akopian 2011, Hrusa 2012, Brullo *et al.* 2013). In particular, some sections, subsections or unranked groups formerly recognized within *Salsola* s.l. were segregated at generic rank. As a consequence, many new genera were described, as well as some misappreciated or neglected ones, such as *Climacoptera* Botschantzev (1956: 111) (see Botschantzev 1969a, and Pratov 1986), *Halo-thamnus* Jaubert & Spach (1846: 50) (see Botschantzev 1981, and Kothe-Heinrich 1993), *Traganum* Delile (1813: 204), *Darniella* Maire & Weiller (1939: 301) (see Brullo 1984), *Fadenia* Aellen & Townsend (1972: 501), *Xylosalsola* Tzvelev (1993: 80), *Nitrosalsola* Tzvelen (1993: 80), *Kali* Miller (1754: without pagination) (see Akhani *et al.* 2007, and Wen *et al.* 2010), *Turania* Akhani & Roalson in Akhani *et al.* (2007: 946), *Kaviria* Akhani & Roalson in Akhani *et al.* (2007: 248), *Pyankovia* Akhani & Roalson in Akhani *et al.* (2007: 249), *Caroxylon* Thunberg (1782: 37) (see Botschantzev 1974a, Tzvelev 1993, Akhani *et al.* 2007, Brullo *et al.* 2013), *Halimocnemis* C. A. Meyer (1829: 381) (see Ghobadnejah *et al.* 2004, and Akhani *et al.* 2007), and *Nanophyton* Lessing (1834: 197) (see Pratov 1982, 1985). In some cases, the taxonomic treatment based on morphologic features is not well supported by the molecular studies,

≡ *Salsola kali* L. var. *tragus* (Linnaeus) Moquin-Tandon in Candolle (1849: 187).
 ≡ *Salsola kali* L. subsp. *tragus* (Linnaeus) Čelakovsky (1871: 155).
 Type (lectotype designated by Degen 1937):—EUROPE. *Herb. Linneaus* 315.3 (LINN). Type (epitype designated by Rilke 1999):—FRANCE. *Kali spinosum cochleatum, Monspelii, Dania, Beleg, Herb Burser XVI(2)-24* (UPS).
 = *Salsola caroliniana* Walter (1788: 111).
 Type (lectotype designated by Rilke 1999):—U.S.A. America borealis, Carolina, *s.coll.* (BM).
 = *Salsola tragus* L. subsp. *iberica* Sennen & Pau in Sennen (1908:476).
 ≡ *Salsola iberica* (Sennen & Pau) Botschantzev ex Czerepanov (1973: 192).
 ≡ *Salsola kali* L. subsp. *iberica* (Sennen & Pau) Rilke in Rech. fil, (1997: 183).
 Type (lectotype designated by Rilke 1999):—SPAIN. Miranda de Ebro, 6 September 1907, *Elias s.n.* (BSB).
 = *Salsola ruthenica* Iljin (1934: 137).
 ≡ *Salsola kali* Linnaeus subsp. *ruthenica* Soó in Soó & Jávorka (1951: 786).
 Type (lectotype designated by Crompton & Bassett 1985):—EUROPE. In pratis vere inundatis, solo arenoso, 12 September 1899, *I. Schirajewsky s.n.* (K).

18) *Kali turgidum* (Dumontier) Gutermann (2011: 98)

Bas.:—*Salsola turgida* Dumortier (1827: 23).
 Type (lectotype designated by Rilke 1999):—NEDERLAND. In arenosis marit. Holl. près Harlem a Zandvoort (BR).
 ≡ *Salsola kali* Linnaeus (1753: 222).
 Type (lectotype designated by Jonsell & Jarvis 1994):—*Herb. Burser* (UPS XVI(2)-24).
 Type (epitype designated by Rilke 1999):—ORIGIN UNKNOWN. *Herb. Clifford*: 86, *Salsola 1* (BM).
 = *Kali sodium* Moench (1794: 331), non Scopoli (1772: 175), nom. illeg. (art. 53.1, McNeill *et al.* 2012).
 = *Salsola decumbens* Lamarck (1779: 241), nom illeg. (Art.52.1, McNeill *et al.* 2012).
 = *Salsola acicularis* Salisbury (1796: 153), nom. illeg. (Art.52.1, McNeill *et al.* 2012).

19) *Kali zaimadium* (Iljin) Akhani & Roalson in Akhani *et al.* (2007: 946).

Bas.:—*Salsola zaimadica* Iljin (1955:122).
 Type:—IRAN. Zaidam, August 1879, *Przewalski* 838 (holotype LE).

Collections information:

K. australe(a): L. Smith S-036, Fresno, CA, beside RR tracks, USA.

K. australe(b): F. Ryan A1, Avondale, Arizona, USA.

K. basalticum: S. Brullo & G. Giusso, Sicily, C.da Ruvolita (Bronte), ITALY.

References

- Aellen, P. & Townsend, C.C. (1972) *Fadenia* - a new genus of Chenopodiaceae from Tropical Africa. *Kew Bulletin* 27: 501–503.
<http://dx.doi.org/10.2307/4114362>
- Akhani, H., Edwards, G. & Roalson, E.H. (2007) Diversification of the old world Salsoleae s.l. (Chenopodiaceae): Molecular phylogenetic analysis of nuclear and chloroplast data set and a revised classification. *International Journal of Plant Sciences* 168: 931–956.
<http://dx.doi.org/10.1086/518263>
- Akhani, H., Greuter, W. & Roalson, E. (2014) Notes on the typification and nomenclature of *Salsola* and *Kali* (Chenopodiaceae). *Taxon* 63(3): 647–650.
<http://dx.doi.org/10.12705/633.1>
- Akopian, J.A. (2011) Genus *Salsola* L. sensu lato (Chenopodiaceae) in South Transcaucasia. *Takhtajania* 1: 124–132.
- Ayers, D., Ryan, F.J., Grotkoo, E., Bailey, J. & Gaskin, J. (2009) Tumbleweed (*Salsola* section *Kali*) species and speciation in California. *Biological Invasions* 1: 1175–1187.

- http://dx.doi.org/10.1007/s10530-008-9380-5
- Bailey, J.P. (1992) Contribution to a cytological catalogue of the British and Irish flora, 2. *Watsonia* 19: 134–137.
- Béguinot, A. (1913) Il polimorfismo nel ciclo di *Salsola kali* L. ed i suoi fattori. [ser. 3] *Atti Accademia delle Scienze Veneto-Trentino-Istriana* 4: 90–126.
- Bòlos, O. de & Vigo, J. (1974) Notes sobre taxonomia i nomenclatura de plantes, I. [Sec. Bot.] *Bulletí Institució Catalana d'Història Natural* 38(1): 61–89.
- Borger, C.P.D., Yan, G., Scott, J.R., Walsh, M. & Powles, S.B. (2008) *Salsola tragus* or *S. australis* (Chenopodiaceae) in Australia – Untangling the taxonomic confusion through random amplified microsatellite polymorphism (RAMP) and cytological analysis. *Australian Journal of Botany* 56: 600–608.
http://dx.doi.org/10.1071/BT08043
- Botschantzev, V.P. (1956) Dva novych rody iz semeistva Marevych. *Sbornik Rabot Po Geobotanike Lesovedenio Paleogeografii i Floristike*: 108–118.
- Botschantzev, V.P. (1969a) The genus *Salsola*: a concise history of its development and dispersal. *Botanicheskii Zhurnal (Moscow & Leningrad)* 54(7): 989–1001.
- Botschantzev, V.P. (1969b) *Malpighipila* Botsch.: generis *Salsola* L. secto nova. *Novosti Sistematički Vysshikh Rastenii* 6: 45–52.
- Botschantzev, V.P. (1970) Sectionis Caroxylon (Thunbg.) Fenzl generis *Salsola* L. species annuae. *Novosti Sistematički Vysshikh Rastenii* 7: 142–145.
- Botschantzev, V.P. (1974a) Species subsectionis Caroxylon sectionis Caroxylon (Thunbg.) Fenzl generis *Salsola*. *Novosti Sistematički Vysshikh Rastenii* 11: 110–174.
- Botschantzev, V.P. (1974b) A synopsis of *Salsola* (Chenopodiaceae) from South and South-West Africa. *Kew Bulletin* 29: 597–614.
http://dx.doi.org/10.2307/4108004
- Botschantzev, V.P. (1981) Revisio generis *Halothamnus* Jaub. et Spach (Chenopodiaceae). *Novosti Sistematički Vysshikh Rastenii* 18: 146–176.
- Brown, R. (1810) *Prodromus Florae Novae Hollandie et Insulae van Diemen* 1. Richard Taylor & Soch, London, 590 pp.
- Brullo, S. (1984) Taxonomic considerations on the genus *Darniella* (Chenopodiaceae). *Webbia* 38: 301–328.
http://dx.doi.org/10.1080/00837792.1984.10670309
- Brullo, S. & Pavone, P. (1987) *Cremnophyton lanfrancoi*: a new genus and species of Chenopodiaceae from Malta. *Candollea* 42: 621–625.
- Brullo, S. & Siracusa, G. (1994) *Crassula basaltica* (Crassulaceae), a new species from Mt. Etna (Sicily). *Flora Mediterranea* 4: 175–178.
- Brullo, C., Brullo, S., Giusso del Galdo, G., Guarino, R. & Iamonico, D. (2013) Il genere *Salsola* s.l. (Chenopodiaceae) in Italia. In: Peccenini, S. & Domina, G. (Eds.) *Contributi alla ricerca floristica in Italia*, Euroservice Punto Grafica, Palermo, pp. 35–38.
- Bunge, A. (1862) Anabasearum revisio. [sér. 6] *Mémoires présentés à l'Académie Impériale des Sciences de St.-Pétersbourg* 4(11): 11–102.
- Bunge, A. (1879) Enumeratio Salsolacearum omnium in Mongolia hucusque collectaru. *Bulletin de l'Academie Imperiale des Sciences de Saint-Pétersbourg* 25: 350–366.
- Čelakovský, L. (1871) *Prodromus der Flora von Böhmen* 2. Selbstverlag, Prag, 275 pp.
- Chinnock, R.J. (2010) Some observations on *Salsola* L. (Chenopodiaceae) in Australia. *Journal of the Adelaide Botanic Gardens* 24: 75–79.
- Crompton, C.V. & Bassett, I.J. (1985) The biology of Canadian weeds. 65. *Salsola pestifer* A. Nels. *Canadian Journal of Plant Science* 65: 379–388.
http://dx.doi.org/10.4141/cjps85-053
- Czerepanov, S.H. (1973) *Svod dopolnenii i izmenenii k Flore SSSR*. Nauka, Leningrad, 667 pp.
- Degen A. von (1937) *Flora Velebitica*.2. Ungarischen Akademie der Wissenschaften. Budapest, 667 pp.
- Delile, A..R. (1813) *Flore d'Egypte*. Imprimerie Impériale, Paris, 355 pp.
- Dumortier, B.C. (1827) *Florula Belgica, Operi Majoris Prodromus. Staminacia*. J. Caterman, Tornaci Nerviorum, 172 pp.
- Fèderova, T.A. (2011) Phylogenetic relations of the South African species of Caroxylon sect. Caroxylon and Tetragonae (Chenopodiaceae) based on the morphology and nrITS sequences. *Turczaninowia* 14(3) : 69–76.
- Freitag, H. (1997) *Salsola* L. In: Rechinger, K.H. (Ed.) *Flora Iranica* 172. Akademische Druck und Verlagsanstalt, Graz, pp. 154–255.
- Freitag, H. & Kani, K. (1987) *Salsola griffithii* (Chenopodiaceae) comb. nova –a remarkable species from the sand deserts of SW. Asia. *Plant Systematic and Evolution* 155 (1–4): 49–54.
http://dx.doi.org/10.1007/BF00936286
- Gandoger, M. (1913) L'herbier africain de Sonder. *Bulletin de la Société Botanique de France* 60: 414–422.
http://dx.doi.org/10.1080/00378941.1913.10836655

- Garcke, A. (1869) *Flora von Nord- und Mittel-Deutschland: Zum Gebrauche auf Excursionen, in Schulen und beim Selbstunterricht*. Ed. 9. Wiegandt und Hempel, Berlin, 520 pp.
- Gaskin, J., Ryan, F., Hrusa, G.F. & Londo, J.P. (2006) Genotype diversity of *Salsola tragus* and potential origins of a previously unidentified invasive *Salsola* from California and Arizona. *Madroño* 53: 244–251.
[http://dx.doi.org/10.3120/0024-9637\(2006\)53\[244:GDOSTA\]2.0.CO;2](http://dx.doi.org/10.3120/0024-9637(2006)53[244:GDOSTA]2.0.CO;2)
- Grubov, V.I. (1961) Duae species novae e familia Chenopodiaceae. *Botanicheskie Materialy Gerbariya Botanicheskogo Instituta Imeni V. L. Komarova Akademii Nauk SSSR, Leningrad* 21: 125–129.
- Grubov, V.I. (2000) *Plants of Central Asia – Plant Collection from China and Mongolia. Chenopodiaceae 2*. Science Publisher, Enfield, 198 pp.
- Gussone, G. (1825) *Index Seminum anni 1825 quae ab Horto Regio in Boccadifalco pro mutua commutatione exhibentur*. Ex Regia Typografia, Napoli, 12 pp.
- Gutermann, W. (2011) Notulae nomenclaturales 41–45. (New names in *Cruciata*, *Kali*, and some small corrections). *Phyton (Horn)* 51(1): 95–102.
- Hedge, I.C., Akhani, H., Freitag, H., Kothe-Heinrich, G., Podlech, D., Rilke, S. & Uotila, P. (1997) Chenopodiaceae. In: Rechinger, K.H. (Ed.) *Flora Iranica* 172. Akademische Druck und Verlagsanstalt, Graz, pp. 1–370.
- Hrusa, G.F. (2012) *Salsola*. In: Baldwin, B.G., Goldman, D.H., Keil, D.J., Patterson, R., Rosatti, T.J. & Wilken, D.H. (Eds.) *The Jepson Manual: Vascular Plants of California* 2nd edition. University California Press, Berkeley. Available from <http://ucjeps.berkeley.edu/IJM.html> (accessed 29 January 2014).
- Hrusa, G.F. & Gaskin, J.F. (2008) The *Salsola tragus* complex in California (Chenopodiaceae): Characterization and Status of *Salsola australis* and the autochthonous allotetraploid *Salsola ryanii* sp. nov. *Madroño* 55(2): 113–131.
[http://dx.doi.org/10.3120/0024-9637\(2008\)55\[113:TSTCIC\]2.0.CO;2](http://dx.doi.org/10.3120/0024-9637(2008)55[113:TSTCIC]2.0.CO;2)
- Iljin, M.M. (1932) Novye vidy roda *Salsola* L.. *Izvestiya Botanicheskogo Sada, Akademii Nauk SSSR* 30: 747–748.
- Iljin, M.M. (1933) Novye vidy roda *Salsola* L. kak predpoagaemyi mezhdurodovoj gibrid. *Botanicheskii Zhurnal (Moscow & Leningrad)* 18(4): 274–277,
- Iljin, M.M. (1934) Chenopodiaceae. In: Keller, B.A. (Ed.) *Sornye Rasteniya SSSR* 2. Tbilisi State University, Leningrad, pp. 90–146.
- Iljin, M.M. (1936) Chenopodiaceae. In: Komarov, V.L. (Ed.) *Flora of the URSS*, 6. Izdatel'stvo Akademii Nauk S.S.R., Leningrad, pp. 2–354.
- Iljin, M.M. (1955) Novye vidy sem. Marevych iz Central'noj Azii. *Botanicheskie Materialy Gerbariya Botanicheskogo Instituta imeni V. L. Komarova, Akademii Nauk SSSR* 17: 122–140.
- IUCN (2014) *Guidelines for Using the IUCN Red List Categories and Criteria*. Version 11. Prepared by the Standards and Petitions Subcommittee, Cambridge U.K. Available from: <http://jr.iucnredlist.org/documents/RedListGuidelines.pdf> (accessed 10 October 2014).
- Jaubert, C. & Spach, E. (1846) *Illustrationes Plantarum Orientalium* 2. Roret Bibliopolam, Paris, 50 pp.
- Jonsell, B. & Jarvis, C.F. (1994) Lectotypifications of Linnean names for Flora Nordica Vol. 1 (Lycopodiaceae – Papaveraceae). *Nordic Journal of Botany* 14: 143–164.
<http://dx.doi.org/10.1111/j.1756-1051.1994.tb00581.x>
- Kadereit, G. & Freitag, H. (2011) Molecular phylogeny of Camphorosmeae (Camphorosmoideae, Chenopodiaceae): Implications fro biogeography, evolution of C4-photosynthesis and taxonomy. *Taxon* 60(1): 51–78.
- Kadereit, G., Borsch, T., Weising, K. & Freitag, H. (2003) Phylogeny of Amaranthaceae and Chenopodiaceae and the evolution of C4 photosynthesis. *International Journal of Plant Sciences* 164: 959–986.
<http://dx.doi.org/10.1086/378649>
- Kadereit, G., Gotzek, D., Jakobs, S. & Freitag, H. (2005) Origin and age of Australian Chenopodiaceae. *Organisms Diversity & Evolution* 5: 59–80.
<http://dx.doi.org/10.1016/j.ode.2004.07.002>
- Kapralov, M.V., Akhani, H., Voznesenskaya, E.V., Edwards, G., Franceschi, V. & Roalson E.H. (2006) Phylogenetic relationships in the Salicornioideae/Suaeoideae/Salsoloideae s.l. (Chenopodiaceae) clade and a clarification of the phylogenetic position of *Bienertia* and *Alexandra* using multiple DNA sequence datasets. *Systematic Botany* 31: 571–585.
<http://dx.doi.org/10.1600/036364406778388674>
- Khatoon, S. & Ali, S.I. (1993) *Chromosome Atlas of the Angiosperms of Pakistan*. Department of Botany, University of Karachi. Karachi. BCC & T. Press, Karachi, 232 pp.
- Kockx van Roon, M. & Wieffering, J.H. (1982) In IOPB chromosome number reports LXXV. *Taxon* 31: 367.
- Kothe-Heinrich, G. (1993) Revision der Gattung *Halothamnus* (Chenopodiaceae). *Bibliotheca Botanica* 143: 1–180.
- Kung, H.-W., Chu, G.-L., Tsien, C.-P., Li, A.-J. & Ma, C.-G. (1978) The Chenopodiaceae in China. *Acta Phytotaxonomica Sinica* 16(1): 99–123.

- Lago Canzobre, E. (1989) Numeros cromosomaticos de plantas occidentales, 521–526. *Anales del Jardin Botanico de Madrid* 45: 501–504.
- Lamarck de, M. (1779) *Flore Françoise* 3. L’Imprimerie Royale, Paris, pp. 16–54
- Lessing, G.F. (1834) Beitrag zur Flora des südlichen Urals und der Steppen. *Linnaea* 9: 145–213.
- Linnaeus, C. (1753) *Species Plantarum* 1. Impensis Laurentii Salvii, Holmiae, 560 pp.
- Linnaeus, C. (1756) *Centuria II. Plantarum*. L.M. Hojer Reg. Acad. Typogr., Upsaliae, 38 pp.
- Litvinov, D. (1902) *Schedae ad Herbarium Florae Rossicae* 4. St. Petersburg, without pagination.
- Litvinov, D. (1905) Rasteniya beregov Aral’skago morya i Syr-Dar’inskoi oblasti, sobrannyya L.S.Bergom. *Izvestiya Turkestanskogo Otdela Imperatorskago Geograficheskago Obshchestva* (St. Petersburg) 4(5): 1–40.
- Litvinov, D. (1922) *Schedae ad Herbarium Florae Rossicae*, 8. St. Petersburg, without pagination.
- Lomonosova, M.N. (2005) Chromosome numbers of Chenopodiaceae species from Russia and Kazakhstan. *Botanicheskii Zhurnal (Moscow & Leningrad)* 90(2): 1132–1134.
- Lomonosova, M.N. (2012) *Konspekt Flory Aziatskoi Rossii: sosudistye rastenilia*. Sibirskoe oldelenie RAN, Novosibirsk, 630 pp.
- Lomonosova, M.N. & Krasnikov, A.A. (1993) Chromosome numbers in some members of the Chenopodiaceae. *Botanicheskii Zhurnal (Moscow & Leningrad)* 78(3): 158–159.
- Lomonosova, M.N., Krasnikov, A.A. & Krasnikova, S.A. (2001) Chromosome numbers of Chenopodiaceae from Siberia. *Botanicheskii Zhurnal (Moscow & Leningrad)* 86(9): 145–146.
- Lomonosova, M.N., Krasnikova, S.A., Krasnikov, A.A., Ebel A.L. & Rudaja, N.A. (2003) Chromosome numbers of the family members Chenopodiaceae representatives from Mongolia and Kazakhstan. *Botanicheskii Zhurnal (Moscow & Leningrad)* 88(8): 113–115.
- Lövkvist, B. & Hultgård, U.M. (1999) Chromosome numbers of South Swedish vascular plants. *Opera Botanica* 137: 1–42.
- Maire, R. & Weiller, M. (1939) Contributions à l’étude de la Flore de l’Afrique du Nord. Fascicule 27. Contributions à l’étude de la Flore de la Libye. *Bulletin de la Société d’Histoire Naturelle de l’Afrique du Nord* 30: 255–314.
- McNeill, J., Barrie, F.R., Buck, W.R., Demoulin, V., Greuter, D.L., Hawksworth, D.L., Herendeen, P.S., Knapp, S., Marhold, K., Prado, J., Proud’Homme van Reine, W.F., Smith, J.F. & Wiersema, J.H. (Eds.) (2012) *International Code of Nomenclature for algae, fungi and plants (Melbourne Code): Adopted by the Eighteenth International Botanical Congress, Melbourne, Australia, July 2011*. [Regnum Vegetabile 154.] Ruggell.
- Meyer, C.A. (1829) *Salsola*. In: Ledebour, C.F. (Ed.) *Flora Altaica* 1. G. Reimer, Berolini, pp. 387–397.
- Miller, P. (1754) *The Gardeners Dictionary*. Abridged, ed. 4, 2. Revigton, London, without pagination.
- Moench, C. (1794) *Methodus Plantas Horti Botanici et Agri Marburgensis, a staminum situ describendi*. Officina nova libraria academiae, Marburgi Cattorum, 780 pp.
<http://dx.doi.org/10.5962/bhl.title.304>
- Moquin-Tandon, A. (1840) *Chenopodearum Monographica Enumeratio*. P.J. Loss, Paris, 182 pp.
<http://dx.doi.org/10.5962/bhl.title.15484>
- Moquin-Tandon, A. (1949) Salsolaceae. In: Candolle, A.P. de (Ed.) *Prodromus Systematis Naturalis Regni Vegetabilis* 13(2). Subtibus Victoris Masson, Parisiis, pp. 43–219.
- Mosyakin, S.L. (1996) A taxonomic synopsis of the genus *Salsola* (Chenopodiaceae) in North America. *Annals of the Missouri Botanical Garden* 83: 387–395.
<http://dx.doi.org/10.2307/2399867>
- Mosyakin, S.L. (2003) *Salsola*. In: Morin, N. (Ed.) *Flora of North America* 4. Oxford Univerty Press, Oxford & New York, pp. 398–403.
- Nishikawa, I. (1985) Chromosome counts of flowering plants of Hokkaido (8). [sect. 2B] *Journal of Hokkaido University of Education* 35: 97–111.
- Pallas, P.S. (1803) *Illustrationes Plantarum*. Subtybus Godofredi Matini, Lipsiae, 129 pp.
- Pignatti, S. (1973) Note critiche sulla flora d’Italia. 1. *Giornale Botanico Italiano* 107(5): 207–221.
<http://dx.doi.org/10.1080/11263507309426323>
- Pratov UP (1985) Obzor roda *Nanophyton* Less. (Chenopodiaceae). *Novosti Sistematički Vysshikh Rastenii* 22: 81–88.
- Pratov, U.P. (1986) *Rod Climacoptera Botsch.: sistematika, geografiya, filogeniya i voprosy okhrany*. Izdatel’stvo Fan Uzbekskoi, Tashkent, 65 pp.
- Probatova, N.S. (2005) Chromosome numbers of some dicotyledons of the flora of the Amur Region. *Botanicheskii Zhurnal (Moscow & Leningrad)* 90(5): 779–792.
- Probatova, N.S., Seledets, V.P., Rudyka, E.G., Gnutikov, A.A., Kuzhevnikova, Z.V. & Barkalov, V.Y. (2009) IAPT/IOPB chromosome data 8. *Taxon* 58(4): 1284–1288.
- Pyankov, V.I., Ziegler, H., Kuz’mín, A. & Edwards, G.E. (2001) Origin and evolution of C4 photosynthesis in the tribe Salsoleae (Chenopodiaceae) based on anatomical and biochemical types in leaves and cotyledons. *Plant Systematic and Evolution* 230: 43–

<http://dx.doi.org/10.1007/s006060170004>

- Rilke, S. (1997) *Salsola* sect. *Salsola*. In: Rechinger, K.H. (Ed.) *Flora Iranica* 173. Akademische Druck und Verlagsanstalt, Graz, pp. 178–191
- Rilke, S. (1999) Revision der Section *Salsola* s.l. del Gattung *Salsola* (Chenopodiaceae). *Bibliotheca Botanica* 149: 1–190.
- Salisbury, R.A. (1796) *Prodromus Stirpium in Horto ad Chapel Allerton Vigentium*. Londini, 442 pp.
<http://dx.doi.org/10.5962/bhl.title.427>
- Schrader, H.A. (1834) *Index Seminum Horti Academicus Gottingensis* 5. Publisher, Gottingae.
- Scopoli, I.A. (1772) *Flora Carniolica*, ed. 2, 1. Ioannis Pauli Krauss, Viennae, 448 pp.
- Sennen, F. (1908) Plantes d' Espagne: Années 1906 & 1907. *Bulletin de l'Académie Internationale de Géographie Botanique* 18: 449–480.
- Soó, R. & Jávorka, S. (1951) *A magyar növényvilág kézikönyve*. Akadémiai Kiadó, Budapest, 1120 pp.
- Strobl, G. (1881) Flora des Etna. *Oesterreichische Botanische Zeitschrift* 31(12): 396–400.
<http://dx.doi.org/10.1007/BF01653457>
- Strobl, G. (1885) Flora des Etna. *Oesterreichische Botanische Zeitschrift* 35(10): 360–364.
<http://dx.doi.org/10.1007/BF01666946>
- Sukhorukov, A.P. (2011) On some new and critical taxa of the Family Chenopodiaceae in the flora of the Caucasus. *Novosti Sistematičeskoy Výsshikh Rastenii (Moscow & Leningrad)* 42: 106–110.
- Swofford, D.L. (2000) PAUP* *Phylogenetic analysis using parsimony (* and other methods)*. Version 4. Sunderland, Sinauer Associates.
- Tamura, K., Steche, G., Peterson, D., Filipski, A., Kumar, S. (2013) MEGA6: Molecular Evolutionary Genetics Analysis Version 6.0. *Molecular Biology and Evolution* 30(12): 2725–2729.
<http://dx.doi.org/10.1093/molbev/mst197>
- Thunberg, C. P. (1782) *Nova genera Plantarum, quorum partem secundam*. Johan Edman, Upsaliae, 11 pp.
- Toderich, K.N., Li, V.V., Black, C.C., Yunusov, T.R., Shuiskay, E.V., Mardonova, G.R. & Gismatullina, L.G. (2006) Linkage studies of structure, isoenzymatic diversity and some biotechnological procedure for *Salsola* species under desert saline environments. In: Öztürk, M., Waisel, Y., Khan, M.A. & Görk, G. (Eds.) *Biosaline agriculture and salinity tolerance in plants*. Birkhäuser, Basel, pp. 73–82.
http://dx.doi.org/10.1007/3-7643-7610-4_8
- Tornabene, F. (1855) Sopra un nuovo albero indigeno dell'Etna del genere *Celtis*. [ser. 2] *Atti Accademia Gioenia Scienze Naturali, Catania* 11: 187–200.
- Tournefort, J.P. (1719) *Institutiones rei Herbariae* 1. Typographia Regia, Parisiis, 697 pp.
- Tzvelev, N.N. (1993) Zametki o marevykh (Chenopodiaceae) Vostochnoy Evropy. *Ukrayins'kyi Botanicnyi Zhurnal, Kiev* 50(1): 78–85.
- Ulbrich, E. (1934) Chenopodiaceae. In: Engler, A. & Harms, A. (Eds.) *Die natürlichen Pflanzenfamilien*, 2., 16c. Engelmann, Leipzig, Berlin, pp. 379–584.
- Volkens, G. (1893) Chenopodiaceae. In: Engler, A & Prantl, K. (Eds.) *Die natürlichen Pflanzenfamilien* 3. Ia. Engelmann, Leipzig, Berlin, pp. 36–91.
- Walter, T. (1788) *Flora Caroliniana*. J. Fraser, Londini, 263 pp.
- Wen, Z.-B., Zhang, M.-L., Zhu, G.-L. & Sanderson, S.C. (2010) Phylogeny of Salsoleae s.l. (Chenopodiaceae) based on DNA sequence data from ITS, psbB-psbH, and rbcL, with emphasis on taxa of northwestern China. *Plant Systematics and Evolution* 288: 25–42.
<http://dx.doi.org/10.1007/s00606-010-0310-5>
- Wilson, B. (2009) *Salsola* of the Pacific Coast States. Available from <http://www.carexworkinggroup.com/pages/salsolakey.pdf> (accessed 10 November 2014).
- Zhu, G.-L., Mosyankin, S.L. & Clemants, S.E. (2003) Chenopodiaceae. In: Wu, Z.Y. & Raven, P.H. (Eds.) *Flora of China* 5. Science Press, Beijing, pp. 354–414.

Appendix. Specimens examined.

Kali australis. U.S.A. California: San Joaquin Valley ½ mi. NW of Holt townsite. Sandy low field, 8 November 2013, Hrusa & Rhode 18529 (CAT!, CDA!); California: San Diego Co., Sacramento, Mission Valley, November 1999, Ryan s.n. (CAT!; CDA!); California: Modesto Co., Santa Nella, November 1999, Ryan s.n. (CAT!; CDA!).