



<http://dx.doi.org/10.11646/phytotaxa.204.3.1>

Revision of *Navicula striolata* (Grunow) Lange-Bertalot and *N. rumanensis* Hustedt with the description of *N. friedelhinziae* sp. nov.

AGATA Z. WOJTAL^{1,*}, NADJA OGNJANOVA-RUMENOVA², KRISZTINA BUCZKÓ³, JANUSZ SIWEK⁴ & BART VAN DE VIJVER^{5,6}

¹ Institute of Nature Conservation, Polish Academy of Sciences, A. Mickiewicza 33, 31-120 Kraków, Poland

² Department of Paleontology, Stratigraphy and Sedimentology, Geological Institute, Bulgarian Academy of Sciences, Acad. G. Bonchev-Street Build. 24, 1113 Sofia, Bulgaria

³ Department of Botany, Hungarian Natural History Museum, Könyves Kálmán krt. 40, 1476 Budapest, Hungary

⁴ Jagiellonian University, Institute of Geography and Spatial Management, Gronostajowa 7, 30-387 Kraków, Poland

⁵ Botanic Garden, Meise, Department of Bryophyta & Thallophyta, Nieuwelaan 38, B-1860 Meise, Belgium

⁶ University of Antwerp, Department of Biology, ECOBE, Universiteitsplein 1, B-2610 Wilrijk, Belgium

* Corresponding author (e-mail: wojtal@iop.krakow.pl)

Abstract

Navicula striolata was originally described as *N. digitoradiata* var. *striolata* from modern material collected in Sweden. After examination of a sample collected from Belgium, the variety was transferred to *N. reinhardtii* as *N. reinhardtii* var. *gracilior*. From this time a large mix up of these and related taxa was observed in the literature. A similar species, *Navicula rumanensis* had also been established in 1934 from Neogene Romanian materials but there has been much confusion regarding the status of these taxa, leading to a poor understanding of their distribution. In this study, type material of *Navicula digitoradiata* var. *striolata*, *N. reinhardtii* var. *gracilior* and *N. rumanensis* are revised using light and scanning electron microscopy in order to clarify their identity and to investigate possible conspecificity. The results indicate that these species are not synonyms. Conspecificity of the modern *N. digitoradiata* var. *striolata* and *N. reinhardtii* var. *gracilior* was confirmed and lectotypes of both varieties have been designated whereas *N. rumanensis* proved to be a separate species. In addition, the study of Neogene material from Bulgaria revealed the presence of a new *Navicula* taxon—*N. friedelhinziae*. The morphology of these and similar taxa is discussed.

Key words: Bacillariophyta, collections, diatoms, *Navicula*, taxonomy, type material

Introduction

Navicula digitoradiata var. *striolata* Cleve in Cleve & Grunow (1880: 32) was originally described from modern material collected by Cleve in 19th century from Degeberga (province of Skåne, southern Sweden). Grunow (1880) indicated slide Cleve & Möller n°25 as being the type material (although a formal holotype was never indicated by him). Slides of this collection were dispatched through all major diatom collections worldwide (Cleve & Möller 1877). Five years later, Grunow synonymized *N. digitoradiata* var. *striolata* with a variety of *N. reinhardtii* (*reinhardtii*) (Grunow 1860: 566) Grunow (1880: 32), namely *N. reinhardtii* (*reinhardtii*) var. *gracilior* Grunow in Van Heurck (1885: 87). Grunow probably based his synonymization on the observation of this taxon in a sample collected by Delogne from Rouge-Cloître (Belgium) and used in the Types du Synopsis de Belgique (Van Heurck 1880, 1885; Type n° 80). In both samples (Degeberga and Rouge-Cloître), *N. digitoradiata* var. *striolata* /*N. reinhardtii* var. *gracilior* co-occurred with the abundantly present *N. reinhardtii* and likely their coexistence and some faint morphological similarity influenced Grunow's decision to transfer *N. digitoradiata* var. *striolata* to *N. reinhardtii*. Krammer & Lange-Bertalot (1986, plate 40) erroneously indicated Type du Synopsis n° 79 as material for the var. *gracilior*. In fact, the latter contains only the nominate variety *N. reinhardtii*.

Later observations of *N. reinhardtii* var. *gracilior* are, however, scarce and usually confused with *N. reinhardtii* var. *reinhardtii* (Krammer & Lange-Bertalot 1986). In 1985, the status of the taxon was once again altered and the variety *gracilior* was elevated to the species level (with a name change) as *Navicula striolata* [Grunow in Cleve

Conclusions

Our investigation of the type materials of *N. digitoradiata* var. *striolata* and *N. reinhardti* var. *gracilior* confirms their conspecific character to *N. striolata*. Our data suggests that fossil taxon—*N. rumaniensis* should remain a separate species. In addition, it is clear that the morphology of *N. friedelhinziae* allows for description of a new species.

Acknowledgements

We thank Mrs. Friedel Hinz (Hustedt Collection, AWI, Bremerhaven) for providing LM and SEM pictures the type slide/material of *Navicula rumaniensis* and her valuable comments. Mrs. Myriam de Haan (Botanic Garden Meise) is thanked for her help with the SEM. We are grateful for valuable comments made by the anonymous Reviewers and the Editor—Dr. Saúl Blanco Lanza. This research was supported by the Synthesys projects HU-TAF-4229 and HU-TAF-3690, Project FP 6 (Structuring the European Research Area Programme, <http://www.synthesys.info>), and the Natural History Museum, Budapest, Hungary; National Science Centre in Poland, project NN3050236406: Environmental and human impacts on springs from the Krakowsko-Wieluńska and Miechowska Uplands (Southern Poland) and the role of the springs in the natural and cultural landscape, with the help of bilateral agreements between the Hungarian and Bulgarian, and Polish and Hungarian Academies of Sciences. Support of the Hungarian Scientific Fund (OTKA 83999) is acknowledged.

References

- Bory de Saint-Vincent, J.B.G.M. (1822) Bacillarées. *Dictionnaire Classique d'Histoire naturelle* 2: 127–129.
<http://dx.doi.org/10.5962/bhl.title.33901>
- Buczkó, K., Ognjanova-Rumenova, N., Magyari, E.K., Morales, E.A., Wojtal, A.Z., Magyar, I. & Ector, L. (2014) Structure, biostratigraphy and distribution of *Navicula haueri* Grunow, a Miocene diatom (Bacillariophyta) from the Carpathian Realm. *Nova Hedwigia, Beiheft* 143: 369–379.
<http://dx.doi.org/10.1127/1438-9134/2014/018>
- Cantonati, M. (Ed.) (1998) 2. La microflora. In: Cantonati, M. (Ed.) *Le sorgenti del Parco Adamello–Brenta. Parco Adamello Brenta Documenti*, 1. Parco Adamello-Brenta, Strembo, Trento, pp. 55–101.
- Cleve, P.T. & Grunow, A. (1880) Beiträge zur Kenntniss der Arctischen Diatomeen. *Kongliga Svenska-Vetenskaps Akademiens Handlingar* 17 (2): 1–121.
- Cleve, P.T. & Möller, J.D. (1877) *Diatoms. Part I.* Esatas Edquists Boktryckeri, Uppsala, pp. 1–48.
- Grunow, A. (1860) Ueber neue oder ungenügend bekannte Algen. Erste Folge, Diatomeen, Familie Naviculaceen. *Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien* 10: 503–582.
- Grunow, A. (1880) Bemerkungen zu den Diatomeen von Finnmark, dem Karischen Meere und vom Jenissej nebst Vorarbeiten für Monographie der Gattungen *Nitzschia*, *Achnanthes*, *Pleurosigma*, *Amphiprora*, *Plagiotropis*, *Hyalodiscus*, *Podosira* und einiger *Navicula*-Gruppen. *Kongliga Svenska Vetenskaps-Akademiens Handligar, Series 4* 17 (2): 16–121.
- Grunow, A. (1882) Beiträge zur Kenntnis der fossilen Diatomeen Österreich-Ungarns. In: Mojsisovics, E. & Neumayer, N. (Eds.) *Beiträge zur Paläontologie Österreich-Ungarns und des Orients* 2/4: 136–159, pl. XXIX–XXX.
- Hasle, G.R. & Fryxell, G.A. (1970) Cleaning and Mounting for Light and Electron Microscopy. *Transactions of the American Microscopical Society* 89 (4): 469–474.
<http://dx.doi.org/10.2307/3224555>
- Hustedt, F. (1945) Diatomeen aus Seen und Quellgebieten der Balkan-Halbinsel. *Archiv für Hydrobiologie* 40 (4): 867–973.
- Krammer, K. & Lange-Bertalot, H. (1985) Naviculaceae Neue und wenig bekannte Taxa, neue Kombinationen und Synonyme sowie Bemerkungen zu einigen Gattungen. *Bibliotheca Diatomologica* 9: 5–230.
- Krammer, K. & Lange-Bertalot, H. (1986): Bacillariophyceae. 1. Teil: Naviculaceae. In: Ettl, H., Gärtner, G., Gerloff, J., Heyning, H. & Mollenhauer, D. (Eds.) *Süßwasserflora von Mitteleuropa* 2/1. Gustav Fischer Verlag, Stuttgart, Jena, pp. 1–876.
- Kulikovskiy, M.S., Lange-Bertalot, H., Metzeltin, D. & Witkowski, A. (2012) Lake Baikal: Hotspot of endemic diatoms I. *Iconographia Diatomologica* 23: 7–607.
- Lange-Bertalot, H. (2001) Diatoms of the European inland waters and comparable habitats. *Navicula* sensu stricto, 10 genera separated from *Navicula* sensu lato. *Frustulia. Diatoms of Europe* 2: 1–526.
- Levkov, Z., Krstic, S., Metzeltin, D. & Nakov, T. (2007) Diatoms of Lakes Prespa and Ohrid. *Iconographia Diatomologica* 16: 1–611.

- Levkov, Z., Edlund, M.B. & Nakov, T. (2008) Identity and typification of *Navicula hasta* (Bacillariophyceae). *Phycological Research* 56: 46–57.
<http://dx.doi.org/10.1111/j.1440-1835.2008.00484.x>
- Magyar, I., Geary, D.H. & Müller, P. (1999) Paleogeographic evolution of the Late Miocene Lake Pannon in Central Europe. *Palaeogeography, Palaeoclimatology, Palaeoecology* 147: 151–167.
[http://dx.doi.org/10.1016/s0031-0182\(98\)00155-2](http://dx.doi.org/10.1016/s0031-0182(98)00155-2)
- Ognjanova-Rumenova, N.G. (1991) *Neogene diatoms from sediments of Sofia Valley and their stratigraphic significance*. Ph.D. Thesis, Geological Institute, Bulgarian Academy of Sciences, 330 pp. [in Bulgarian]
- Pantocsek, J. (1892) *Beiträge zur Kenntnis der Fossilen Bacillarien Ungarns. III. Süßwasser Bacillarien Anhang. Analysen 15 neuer Depots von Bulgarien, Japan, Mahern, Russland und Ungarn (mit 42 Tafeln)*. Nagy-Tapolcsány, 584 figs. [without text]
<http://dx.doi.org/10.5962/bhl.title.14916>
- Pantocsek, J. (1905) *Beiträge zur Kenntnis der fossilen Bacillarien Ungarns III. Beschreibung der auf Tafeln abgebildeten Arten*. Buchdruckerei C.F. Wigand, Bratislava, 118 pp.
<http://dx.doi.org/10.5962/bhl.title.14916>
- Rakowska, B. (1996a) Diatom communities occurring in Niebieskie źródła near Tomaszów Mazowiecki, Central Poland (1963–1990). *Fragmenta Floristica et Geobotanica* 41 (2): 639–655.
- Rakowska, B. (1996b) Uwagi o taksonomii i ekologii kilkunastu gatunków okrzemek. *Acta Universitatis Lodziensis, Folia Botanica* 11: 127–146.
- Round, F.E., Crawford, R.M. & Mann, D.G. (1990) *The diatoms: biology and morphology of the genera*. Cambridge University Press, Cambridge, 747 pp.
- Ross, R., Cox, E.J., Karayeva, N.I., Mann, D.G., Paddock, T.B.B., Simonsen, R. & Sims, P.A. (1979) An amended terminology for the siliceous components of the diatom cell. *Nova Hedwigia Beiheft* 64: 513–533.
- Schmidt, A., Schmidt, M., Fricke, F., Heiden, H., Müller, O. & Hustedt, F. (1934) *Atlas der Diatomeen-Kunde. Tafel 395*. Leipzig.
- Simonsen, R. (1987) *Atlas and catalogue of the diatom types of Friedrich Hustedt*. Verlag J. Cramer, Berlin & Stuttgart, 521 pp.
- Schrader, H.-J. (1973) Proposal for a Standardized Method of Cleaning Diatom-bearing Deep sea and Land-exposed Marine Sediments. *Nova Hedwigia, Beiheft* 45: 403–409.
- Stefanova, M., Ivanov, D., Yaneva, N., Marinov, S., Grasset, L. & Ambles, A. (2008) Palaeoenvironment assessment of Pliocene Lom lignite (Bulgaria) from bitumen analysis and preparative off line thermochemolysis. *Organic Geochemistry* 39: 1589–1605.
<http://dx.doi.org/10.1016/j.orggeochem.2008.07.008>
- Van Dam, H., Mertens, A. & Sinkeldam, J. (1994) A Coded Checklist and Ecological Indicator Values of Freshwater Diatoms from Netherlands. *Netherlands Journal of Aquatic Ecology* 28: 117–184.
<http://dx.doi.org/10.1007/bf02334251>
- Van der Werff, A. (1955) A new method of concentrating and cleaning diatoms and other organisms. *Verhandlungen Internationalen Vereinigung für Theoretische und Angewandte Limnologie* 2: 276–277.
- Van Heurck, H.F. (1880) *Synopsis des Diatomées de Belgique. Atlas*. Ducaju & Cie, Antwerp.
- Van Heurck, H.F. (1885) *Synopsis des Diatomées de Belgique. Texte*. Antwerp, Martin Brouwers & Co., 235 pp.
<http://dx.doi.org/10.5962/bhl.title.1990>
- Wojtal, A.Z. (2013) Species composition and distribution of diatom assemblages in spring waters from various geological formations in southern Poland. *Bibliotheca Diatomologica* 59: 1–436.
- Yaneva, M., Ognjanova-Rumenova, N., Shanov, S., Nikolov, G. & Nikolov, N. (2010) Characteristic and distribution of the coal bearing sediments in Lom Depression. In: Nakov, R. (Ed.) *Proceedings “Geosciences 2010”, December 2010*. Bulgarian Geological Society (BGD), Sofia, pp. 85–86.
- Żelazna-Wieczorek, J. (2011) Diatom flora in Springs of Łódź Hills (Central Poland). Biodiversity, taxonomy, and temporal changes of epipsammic diatom assemblages in springs affected by human impact. *Diatom Monographs* 13: 1–420.
- Żelazna-Wieczorek, J. & Bik, A. (2009) Dynamika zbiorowisk okrzemek w źródłach o różnym typie hydrobiologicznym. *Fragmenta Floristica et Geobotanica Polonica* 16 (1): 155–167.