



## ***Acartia* (*Odontacartia*) *ohtsukai* Ueda and Bucklin, 2006 (Copepoda, Calanoida, Acartiidae): First Record of its Occurrence in Korean Waters and Habitat Segregation from its Sibling Species *A. pacifica* Steuer, 1915**

SEONG YONG MOON<sup>1</sup>, SUSUMU OHTSUKA<sup>2</sup>, HIROSHI UEDA<sup>3</sup> & HO YOUNG SOH<sup>1\*</sup>

<sup>1</sup>Division of Marine Technology, Chonnam National University, Yeosu 550-749, Korea

<sup>2</sup>Takehara Marine Science Station, Setouchi Field Science Center, Graduate School of Biosphere Science, Hiroshima University, 5-8-1 Minato-machi, Takehara 725-0024, Japan

<sup>3</sup>Utsunomiya Marine Biological Institute, Kochi University, 194 Inoshiri, Usa, Tosa, Kochi 781-1164, Japan

\* Corresponding author: Ho Young Soh (hysoh@chonnam.ac.kr)

Species of the genus *Acartia* Dana, 1846 (Copepoda, Calanoida, Acartiidae) are predominant and widespread in estuarine and coastal waters worldwide, but the spatio-temporal distribution of each species is very restricted (Bradford 1976). *Acartia pacifica* Steuer, 1915 has been recognized as a common species in euryhaline waters on the Korean coasts from summer to fall (Suh *et al.* 1991; Soh & Suh 1993; Park 2005). Recently, Ueda and Bucklin (2006) described *A. ohtsukai* from low saline waters in the Ariake Sea, Western Japan, which has been confused with *A. pacifica* s. s. on the basis of their morphological and genetic differences (0-3% in mtCOI sequences). According to Ueda and Bucklin (2006), *A. ohtsukai* has a restricted distribution in continental brackish waters of East Asia with a relict population in the Ariake Sea, while *A. pacifica* is broadly distributed in coastal waters in the tropical/subtropical Indo-West Pacific regions. During our investigation of planktonic copepods *A. ohtsukai* was recorded in Korean waters for the first time. This species may have been confused in the past with its sibling species *A. pacifica* by Korean researchers. Therefore, we compare the spatio-temporal distributions of *A. pacifica* and *A. ohtsukai* in Gwangyang Bay, southern Korea.

Zooplankton samples were collected from 40 stations within Gwangyang Bay located in the southern central region of Korea. A Norpac net (mouth diameter 45cm; mesh size 0.2 mm) was towed vertically from the bottom to the surface layer during November 2005, and February, May and August, 2006. The volume of water filtered was estimated from a flowmeter (General Oceanic Co.) attached to the net mouth. Copepods were immediately fixed in buffered formaldehyde/seawater solution (final concentration 5%). Also, water temperature and salinity were continuously measured from surface to bottom *in situ* with a CTD (Alec Electronics Co. Model ACL1150-DK). Specimens identified as *A. ohtsukai* and *A. pacifica* were sorted from the samples. Patterns of their occurrence were plotted on a Temperature-Salinity diagram.

Morphological characteristics of *A. ohtsukai* from Korea waters are similar to those described by Ueda and Bucklin (2006) from Japan, except for a slightly larger length/width ratio of the female caudal ramus, shorter spines on the third urosomite of the female, a square medial projection of the second exopodal segment of the male right fifth leg and the shorter first exopodal segment of the male left fifth leg (Fig. 1A-D). DNA sequences for mtCOI differed by 2-5 % between Korean and Japanese (Ariake Sea) populations, while these did by less than 5% among Korean populations and by less than 3% among Japanese individuals (Soh *et al.* unpublished data). However, these differences between *Acartia* species ranged from 21 to 42%: GeneBank Accession no. EU274431 for *A. hudsonica* Pinhey, 1926; AY145426 for *A. omorii* Bradford, 1976; DQ071177 for *A. pacifica* Steuer, 1915; DQ665253 and DQ665254 for *A. spinicauda* Giesbrecht, 1889; EU274464 for *A. tonsa* Dana, 1849. In addition to their morphological similarity, these facts suggest that Korean and Japanese populations belong to same species (Soh *et al.* unpublished data).

The habitats of specimens identified as *A. ohtsukai* and *A. pacifica* in Gwangyang Bay, Korea were pronouncedly segregated (Fig. 2). *A. ohtsukai* was found in Gwangyang Bay and in the adjacent coastal zone after the rainy season in late June to July when a large volume of freshwater flows into Gwangyang Bay from the Seomjin River. This inflow decreases salinity to less than 30.0 psu. In contrast, its sibling species *A. pacifica* occurred offshore in more than 32.0 psu (KORDI 2005). As the salinity of the regions again increased above 32.0 psu after the rainy season, *A. ohtsukai* was