Copyright © 2008 · Magnolia Press



## A new genus of Neothrombiidae (Acari: Trombidioidea) from Yemen

ALIREZA SABOORI<sup>1</sup>, EDWARD A. UECKERMANN<sup>2</sup> & ANTONIUS VAN HARTEN<sup>3</sup>

<sup>1</sup>Department of Plant Protection, College of Agriculture, University of Tehran, Karaj, Iran. E-mail: saboori@ut.ac.ir <sup>2</sup>ARC-Plant Protection Research Institute Private Bag X134 Queenswood, Pretoria 0121 South Africa. E-mail: UeckermannE@arc.agric.za <sup>3</sup>Cha de Vaca, São Jorge dos Orgãos, Santiago, Cape Verde. E-mail: vanharten@zmail.pt

## Abstract

*Solistrus mitrae* gen. et sp. nov. is described and illustrated from an unknown host collected with a light trap at Al Kowd, Yemen. This is the first record of the family Neothrombiidae in Yemen. The new taxon is based on the larva; deutonymphs and adults remain unknown.

Key words: Neothrombiidae, new genus, larva, key, Yemen

## Introduction

The taxonomy of the family Neothrombiidae Feider is in a state of flux. Zhang (1994) revised this family and recognized nine genera, seven of which were known only as larvae and presented a key to world genera. He described two new genera and transferred *Acritrombium* Southcott, 1986, *Paputrombidium* Fain, 1992, *Won-declia* Southcott, 1987 and *Falsivagothrombium* Zhang & Xin, 1989 (known for postlarval instars only) from the Trombidiidae to the Neothrombiidae.

Saboori (2002) described a new genus, *Southcottella* (larva), from Iran and presented an updated key to world genera.

The family Neothrombiidae was last revised by Zhang & Fan (2005), who described a new genus, *Opiliotrombium* (larva), parasitic on Opiliones from New Zealand and transferred *Silphitrombium* Fain, 1992, *Neosilphitrombium* Fain, 1992, *Machilitrombium* Fain, 1994 and *Thaitrombidium* Fain, 1998 from the Trombidiidae to this family.

This is the first record of this family from Yemen. In this paper, we describe a new genus and species from Yemen.

## Material and methods

Specimens were preserved in 75% ethanol, cleared in lactophenol solution and mounted in Hoyer's medium. Figures were drawn using a BX-51 Olympus microscope equipped with a drawing tube and magnification changer. Measurements were calculated using a Wild microscope. Terminology and abbreviations follow Robaux (1974). Measurements are made in micrometers (m).