

## New Neotropical genera and species of Dryobiini and Rhopalophorini (Coleoptera: Cerambycidae: Cerambycinae)

DILMA SOLANGE NAPP<sup>1,3</sup> & MIGUEL A. MONNÉ<sup>2,3</sup>

<sup>1</sup>Departamento de Zoologia, Universidade Federal do Paraná. Caixa Postal 19020, 81531-980 Curitiba, Paraná, Brazil.  
E-mail: napp@ufpr.br

<sup>2</sup>Museu Nacional, Universidade Federal do Rio de Janeiro. Quinta da Boa Vista, São Cristóvão, 20940-040 Rio de Janeiro, Rio de Janeiro, Brazil. E-mail: monne@uol.com.br

<sup>3</sup>Fellow of the Conselho Nacional de Desenvolvimento Científico e Tecnológico - CNPq

### Abstract

*Anisotyma* Napp & Monné, new genus, *A. soteri* Napp & Monné, new species (type species) from El Salvador (La Libertad), are described in Dryobiini, and *Potiapua* Napp & Monné, new genus, *P. antonia* Napp & Monné, new species (type species) from Brazil (Espírito Santo), in Rhopalophorini.

**Key words:** *Anisotyma*, Cerambycinae, *Potiapua*, Neotropical, new taxa

### Resumo

*Anisotyma* Napp & Monné, gen. nov., *A. soteri* Napp & Monné, sp. nov. (especie-tipo) de El Salvador (La Libertad), são descritos em Dryobiini, e *Potiapua* Napp & Monné, gen. nov., *P. antonia* Napp & Monné, sp. nov. (espécie-tipo) do Brasil (Espírito Santo), em Rhopalophorini.

**Palavras-chave:** *Anisotyma*, Cerambycinae, novos táxons, *Potiapua*, Região Neotropical

### Introduction

The tribe Dryobiini Linsley, 1964 includes two monotypic genera: *Dryobius* LeConte, 1850 recorded only from The United States of America, and *Ornithia* Thomson, 1864, with the distribution ranging from Central America to southeastern Texas (Linsley 1964; Monné 2005; Monné & Bezark 2008).

Rhopalophorini Blanchard, 1845 includes, at present, 23 American genera predominantly South American. Only one genus, *Rhopalophorella* Linsley, 1942 was described from the United States of America, and one, *Muxbalia* Giesbert & Chemsak, 1993, from Central America. The remaining 21 genera were described from South America, some of them extending their range northwards into Central and North America (Giesbert & Chemsak 1993; Monné 2005; Monné & Bezark 2008).

Herein, two Neotropical genera, both monotypic, are added to these tribes, one from Central America (El Salvador) assigned to Dryobiini, and one from South America (Brazil), to Rhopalophorini.

The types are deposited in the Departamento de Zoologia, Universidade Federal do Paraná, Curitiba, Brazil (DZUP) and National Museum of Natural History, Washington, D.C., United States of America (USNM).