

<http://dx.doi.org/10.11646/zootaxa.3709.2.5>
<http://zoobank.org/urn:lsid:zoobank.org:pub:BE65D2AB-00C7-456C-ACAC-530DA96386E0>

Collembola Poduromorpha from Guelma Province (Northeastern Algeria), with description of a new *Superodontella* species (Collembola: Odontellidae)

JAVIER I. ARBEA^{1,3}, HAYETTE BRAHIM-BOUNAB² & SALAH HAMRA KROUA²

¹C/ Ría de Soria 3, Ch. 39, 39610 El Astillero, España. E-mail: jarbeapo@gmail.com

²Laboratoire de Biosystématique et Écologie des Arthropodes, Faculté des Sciences de la Nature et de la Vie, Université Constantine 1, Algérie. E-mail: salahamra@yahoo.fr

³Corresponding author

Abstract

Ten Poduromorpha species have been found in the Taya Region, three of which are new for Algeria: *Hypogastrura vernalis* (Carl, 1901), *Xenylla brevisimilis mediterranea* Gama, 1964 and *Superodontella tayaensis* sp. nov. The new species is morphologically closest to *S. subalpina* Arbea, 1990, with three subequal dental chaetae. They differ in the number of tibiotarsal chaetae and the sensilar chaetotaxy of Ant IV.

Key words: *Superodontella tayaensis*, new species, species list, taxonomy, Taya Region

Introduction

Our study is based on material collected by the last two authors during a series of trips in the region of Taya (Guelma Province, Algeria). We present also a description of a new *Superodontella* species.

Abbreviations used: In the descriptions we used the morphological nomenclature as follows: antennal sensilla (S-chaetae) are numbered after D'Haese (2003); the nomenclature of tergite chaetotaxy follows the classical a,m,p pattern, after Yosii (1956) and Arbea and Jordana (1997); for the perilabial chaetotaxy, after Gama (1969) and Yosii (1971); for the labial chaetotaxy, after Massoud (1967); for the structure of maxilla, after Deharveng (1981) and Bedos and Deharveng (1990); for the chaetae classification of anal valves, after Hüther (1962).

Abd tr—abdominal tergum, Abd st—abdominal sternum, Ant—antennal segment, EPP—postero-external processus of fulcrum, IPP—postero-internal processus of fulcrum, ms—microsensillum, PAO—postantennal organ, Th tr—thoracic tergum, Th st—thoracic sternum.

Material deposit. LBEA—Laboratoire de Biosystématique et Écologie des Arthropodes, Faculté des Sciences de la Nature et de la Vie, Université Constantine 1 (Algérie); MZNA—Museo de Zoología, University of Navarra, Pamplona (Spain); JIAP—Private collection of J.I. Arbea (Spain).

Results

Poduromorpha Collembola from Taya (Guelma Province, Algeria)—species list

Ten species of Poduromorpha, including the new species described in this paper, are recorded from the Taya Region.

Family Hypogastruridae

Hypogastrura vernalis (Carl, 1901), was found in moss on soil.

Distribution: Palaearctic, new to the Algerian fauna.