

<http://dx.doi.org/10.111646/zootaxa.3745.1.6>
<http://zoobank.org/urn:lsid:zoobank.org:pub:6C7641F1-2F41-421C-948B-0DF410AECB17>

A new *Philautus* (Anura: Rhacophoridae) from northern Laos allied to *P. abditus* Inger, Orlov & Darevsky, 1999

BRYAN L. STUART^{1,4}, SOMPHOUTHONE PHIMMACHAK², SENGVILAY SEATEUN³ & JENNIFER A. SHERIDAN¹

¹*North Carolina Museum of Natural Sciences, 11 West Jones Street, Raleigh NC 27601, USA*

²*Kasetsart University, Faculty of Science, Department of Zoology, Ngam Wong Wan Road, Chatuchak, Bangkok, 10900, Thailand*

³*National University of Laos, Faculty of Science, Department of Biology, P.O. Box 2273, Dong Dok Campus, Vientiane, Laos*

⁴*Corresponding author. E-mail: bryan.stuart@naturalsciences.org*

Abstract

The small rhacophorid frog *Philautus abditus* is geographically restricted to central Vietnam and adjacent Cambodia. Our fieldwork in northern Laos resulted in the discovery of a *Philautus* species that very closely resembles *P. abditus*, but is at least 330 km from the nearest known locality of that species. The Laos population differs from *P. abditus* in mitochondrial DNA and coloration, and is described here as a new species. *Philautus nianae sp. nov.* is distinguished from its congeners by having the combination of a hidden tympanum; no nuptial pads; smooth skin; large black spots on the hidden surfaces of the hind limbs; light venter with dark spotting; and a bronze iris. A second species of *Philautus* from northern Laos, *P. petilus*, is transferred on the basis of morphology to the genus *Theloderma*.

Key words: Laos; new species; *Philautus abditus*; *Philautus petilus*; Rhacophoridae

Introduction

The rhacophorid frog genus *Philautus* Gistel, 1848 has a turbulent taxonomic history (Bossuyt & Dubois 2001; Hertwig *et al.* 2012; Frost 2013). Eighty-four species of *Philautus* were recognized in a recent nomenclatural review (Bossuyt & Dubois 2001), but many of these species have since been transferred to other genera of rhacophorids, primarily on the basis of evidence derived from molecular phylogenetic analyses (e.g., Frost *et al.* 2006; Li *et al.* 2008; Li *et al.* 2009). No morphological synapomorphy is currently known for *Philautus*. The genus has been defined in the past as rhacophorids lacking vomerine teeth (Liem, 1970), or lacking a free-swimming aquatic larva (Bossuyt & Dubois 2001), but these definitions have not been supported by molecular phylogenetic analyses (Hertwig *et al.* 2012). As currently recognized, the genus *Philautus* contains approximately 50 species (Frost 2013), most of which are relatively small and arboreal. The genus is distributed from India to the Philippines, with the greatest species diversity occurring in Borneo and the Philippines (Brown & Alcala 1994; Hertwig *et al.* 2012; Frost 2013).

Our recent fieldwork in northern Laos revealed a *Philautus* that bears remarkable similarity to *P. abditus* Inger, Orlov & Darevsky 1999, a species that is restricted to central Vietnam and adjacent northeastern Cambodia (Nguyen *et al.* 2009; Stuart *et al.* 2010) at least 330 km southeast of the Laos locality (the nearest known record of *P. abditus* is in Quang Nam Province, Vietnam; Figure 1). Despite the similarity, the Laos specimens differ from *P. abditus* in coloration and mitochondrial DNA (the call of *P. abditus* is unknown), and are herein described as a new species.

References

- Bain, R.H. & Nguyen, T.Q. (2004) Herpetofaunal diversity of Ha Giang Province in northeastern Vietnam, with descriptions of two new species. *American Museum Novitates*, 3453, 1–42.
[http://dx.doi.org/10.1206/0003-0082\(2004\)453<0001:hdohgp>2.0.co;2](http://dx.doi.org/10.1206/0003-0082(2004)453<0001:hdohgp>2.0.co;2)
- Bioacoustics Research Program (2011) Raven Pro: Interactive Sound Analysis Software (Version 1.4) The Cornell Lab of Ornithology, Ithaca, New York. Available from: <http://www.birds.cornell.edu/raven> (accessed 12 November 2013)
- Bossuyt, F. & Dubois, A. (2001) A review of the frog genus *Philautus* Gistel, 1848 (Amphibia, Anura, Ranidae, Rhacophorinae). *Zeylanica*, 6, 1–112.
- Brown W.C. & Alcala, A.C. (1994) Philippine frogs of the family Rhacophoridae. *Proceedings of the California Academy of Sciences*, 48, 185–220.
- Bourret, R. (1937) Notes herpétologiques sur l'Indochinese française. XIV. Les batraciens de la collection du Laboratoire des Sciences naturelles de l'Université. Descriptions de quinze espèces ou variétéés nouvelles. *Annexe au Bulletin Général de l'Instruction Publique*, 4, 5–56.
- Das, I. (2005) A new species of *Polypedates* (Anura: Rhacophoridae) from Gunung Murud, Sarawak (northwestern Borneo). *The Raffles Bulletin of Zoology*, 53, 265–270.
- Frost, D.R., Grant, T., Faivovich, J., Bain, R.H., Haas, A., Haddad, C.F.B., de Sá, R.O., Channing, A., Wilkinson, M., Donnellan, S.C., Raxworthy, C.J., Campbell, J.A., Blotto, B.L., Moler, P.E., Drewes, R.C., Nussbaum, R.A., Lynch, J.D., Green, D.M. & Wheeler, W.C. (2006) The amphibian tree of life. *Bulletin of the American Museum of Natural History*, 297, 1–370.
[http://dx.doi.org/10.1206/0003-0090\(2006\)297\[0001:tatol\]2.0.co;2](http://dx.doi.org/10.1206/0003-0090(2006)297[0001:tatol]2.0.co;2)
- Frost, D.R. (2013) Amphibian Species of the World: an Online Reference. Version 5.6 (9 January 2013). American Museum of Natural History, New York. Available from: <http://research.amnh.org/herpetology/amphibia/index.html> (accessed 1 June 2013)
- Gistel, J.v.N.F.X. (1848) *Naturgeschichte des Thierreichs. Für höhere Schulen*. Hoffmann, Stuttgart, 216 pp.
- Glaw, F. & Vences, M. (1997) Anuran eye colouration: definitions, variation, taxonomic implications and possible functions. In: Böhme, W., Bischoff, W. & Ziegler, T. (Eds.), *Herpetologia Bonnensis. Proceedings of the 8th Ordinary General Meeting of the Societas Europaea Herpetologica, 23–27 August 1995*. Societas Europaea Herpetologica, Deutsche Gesellschaft für Herpetologie und Terrarienkunde and Zoologisches Forschungsinstitut und Museum Alexander Koenig, Bonn, pp. 125–138.
- Hertwig, S.T., Das, I., Schweizer, M., Brown, R. & Haas, A. (2012) Phylogenetic relationships of the *Rhacophorus everetti*-group and implications for the evolution of reproductive modes in *Philautus* (Amphibia: Anura: Rhacophoridae). *Zoologica Scripta*, 41, 29–46.
<http://dx.doi.org/10.1111/j.1463-6409.2011.00499.x>
- Inger, R.F., Orlov, N. & Darevsky, I. (1999) Frogs of Vietnam: a report on new collections. *Fieldiana, New Series*, 92, 1–46.
<http://dx.doi.org/10.5962/bhl.title.3478>
- Li, J., Che, J., Bain, R.H., Zhao, E. & Zhang, Y. (2008) Molecular phylogeny of Rhacophoridae (Anura): a framework of taxonomic reassignment of species within the genera *Aquixalus*, *Chiromantis*, *Rhacophorus*, and *Philautus*. *Molecular Phylogenetics and Evolution*, 48, 302–312.
<http://dx.doi.org/10.1016/j.ympev.2008.03.023>
- Li, J., Che, J., Murphy, R.W., Zhao, H., Zhao, E., Rao, D. & Zhang, Y. (2009) New insights to the molecular phylogenetics and generic assessment in the Rhacophoridae (Amphibia: Anura) based on five nuclear genes and three mitochondrial genes, with comments on the evolution of reproduction. *Molecular Phylogenetics and Evolution*, 53, 509–522.
<http://dx.doi.org/10.1016/j.ympev.2009.06.023>
- Liem, S.S. (1970) The morphology, systematics, and evolution of the Old World treefrogs (Rhacophoridae and Hyperoliidae). *Fieldiana Zoology*, 57, 1–145.
- Liu, C.-C. & Hu, S-Q. (1962) A herpetological report of Kwangsi. *Acta Zoologica Sinica*, 14 (Suppl.), 73–104.
- Nguyen, V.S., Ho, C.T. & Nguyen, T.Q. (2009) *Herpetofauna of Vietnam*. Edition Chimaira, Frankfurt am Main, 768 pp.
- Ohler, A., Swan, S.R. & Daltry, J.C. (2002) A recent survey of the amphibian fauna of the Cardamom Mountains, Southwest Cambodia with descriptions of three new species. *The Raffles Bulletin of Zoology*, 50, 465–481.
- Orlov, N.L. & Ho, C.T. (2005) A new species of *Philautus* from Vietnam (Anura: Rhacophoridae). *Russian Journal of Herpetology*, 12, 135–142.
<http://dx.doi.org/10.5358/hsj.22.51>
- Orlov, N.L., Ho, C.T. & Nguyen, T.Q. (2004) A new species of the genus *Philautus* from Central Vietnam (Anura: Rhacophoridae). *Russian Journal of Herpetology*, 11, 51–64.
<http://dx.doi.org/10.5358/hsj.22.51>
- Palumbi, S.R. (1996) Nucleic acids II: the polymerase chain reaction. In: Hillis, D.M., Moritz, C. & Mable, B.K. (Eds.), *Molecular Systematics*. Second edition. Sinauer Associates, Inc., Sunderland, Massachusetts, pp. 205–247.
- Pyron, A. & Wiens, J.J. (2011) A large-scale phylogeny of Amphibia including over 2800 species, and a revised classification of extant frogs, salamanders, and caecilians. *Molecular Phylogenetics and Evolution*, 61, 543–583.
<http://dx.doi.org/10.1016/j.ympev.2011.06.012>

- Rowley, J.J.L., Le, D.T.T., Hoang, H.D., Dau, V.Q. & Cao, T.T. (2011) Two new species of *Theloderma* (Anura: Rhacophoridae) from Vietnam. *Zootaxa*, 3098, 1–20.
- Schlegel, H. (1837) *Abbildungen neuer oder unvollständig bekannter Amphibien, nach der Natur oder dem Leben entworfen, herausgegeben und mit einem erläuternden Texte begleitet. Part 1.* Arnz & Co., Düsseldorf, 32 pp.
- Sterling, E.J., Hurley, M.M. & Le, M.D. (2006) *Vietnam: a natural history*. Yale University Press, New Haven and London, 423 pp.
- Stuart, B.L. & Heatwole, H.F. (2004) A new *Philautus* (Amphibia: Rhacophoridae) from northern Laos. *Asiatic Herpetological Research*, 10, 17–21.
- Stuart, B.L., Inger, R.F. & Voris, H.K. (2006) High level of cryptic species diversity revealed by sympatric lineages of Southeast Asian frogs. *Biology Letters*, 2, 470–474.
<http://dx.doi.org/10.1098/rsbl.2006.0505>
- Stuart, B.L., Rowley, J.J.L., Neang, T., Emmett, D.A. & Som, S. (2010) Significant new records of amphibians and reptiles from Virachey National Park, northeastern Cambodia. *Cambodian Journal of Natural History*, 2010, 38–47.
- Stuart, B.L., Rowley, J.J.L., Tran, D.T.A., Le, D.T.T. & Hoang, H.D. (2011) The *Leptobrachium* (Anura: Megophryidae) of the Langbian Plateau, southern Vietnam, with description of a new species. *Zootaxa*, 2804, 25–40.
- Stuebing, R.B. & Wong, A. (2000) A new species of frog, *Philautus erythrophthalmus* (Rhacophoridae) from southwestern Sabah, Malaysia. *The Raffles Bulletin of Zoology*, 48, 293–296.
- Swofford, D.L. (2002) *PAUP*: Phylogenetic Analysis Using Parsimony *(and other methods)*. Version 4.0b10. Sinauer Associates, Sunderland, Massachusetts.
- Tschudi, J.J.v. (1838) *Classification der Batrachier, mit Berücksichtigung der fossilen Thiere dieser Abtheilung der Reptilien*. Petitpierre, Neuchâtel, 99 pp.
- Vieites, D.R., Wollenberg, K.C., Andreone, F., Köhler, J., Glaw, F. & Vences, M. (2009) Vast underestimation of Madagascar's biodiversity evidenced by an integrative amphibian inventory. *Proceedings of the National Academy of Sciences*, 106, 8267–8272.
<http://dx.doi.org/10.1073/pnas.0810821106>
- Yu, G., Rao, D., Yang, J. & Zhang, M. (2008) Phylogenetic relationships among Rhacophorinae (Rhacophoridae, Anura, Amphibia), with an emphasis on the Chinese species. *Zoological Journal of the Linnean Society*, 153, 733–749.
<http://dx.doi.org/10.1111/j.1096-3642.2008.00404.x>