



<http://dx.doi.org/10.11646/zootaxa.3745.2.8>

<http://zoobank.org/urn:lsid:zoobank.org:pub:0DE6D26C-A64A-49E1-9A43-D933CF8A7F44>

**RAFAL GOSIK & PETER SPRICK (2013) Morphology and identification of the pupae of several species of soil-dwelling broad-nosed weevils from Central Europe (Coleoptera, Curculionidae, Entiminae). *Zootaxa*, 3731(4): 445–472.**

1) Figure 1–10 on page 448 to be replaced by the following:



**FIGURES 1–10.** Weevils (adults, dorsal view): **1**—*Barypeithes pellucidus*: immature; **2**—*Brachyderes incanus*; **3**—*Liophloeus tessulatus*; **4**—*Otiorhynchus dieckmanni*; **5**—*O. tenebricosus* form *lugdunensis*; **6**—*Peritelus sphaeroides*; **7**—*Phyllobius intrusus*; **8**—*P. pyri* s. str.; **9**—*Strophosoma sus*; **10**—*Tanymecus palliatus* (photos: P. Sprick).

2) On page 445, 447, 449, 462, 463 and 467 to change c.f. to cf.

3) New Doi for references: De Fluiter, H.J. & Blijdorp, P.A. (1935) De grauwe dennensnuitkever *Brachyderes incanus* L. *Tijdschrift over Plantenziekten*, 41, 141–211.

<http://dx.doi.org/10.1007/BF02807371>

4) Delete Doi under reference: Scherf, H. (1964) Die Entwicklungsstadien der mitteleuropäischen Curculioniden (Morphologie, Bionomie, Ökologie). *Abh. Senckenberg. Naturf. Ges.*, 506, 1–335.

<http://dx.doi.org/10.1002/mmnd.4820240510>