



<http://dx.doi.org/10.11646/zootaxa.3895.3.2>

<http://zoobank.org/urn:lsid:zoobank.org:pub:44081058-1ACC-4DFE-AF20-096062256345>

## Two new species of the genus *Wellsopsyllus* (Copepoda; Harpacticoida; Paramesochridae) from the Yellow Sea

JINWOOK BACK<sup>1</sup> & WONCHOEL LEE<sup>2,3</sup>

<sup>1</sup>Marine Biological Resources Division, National Marine Biodiversity Institute of Korea, Seocheon, Korea

<sup>2</sup>Department of Life Science, College of Natural Sciences, Hanyang University, Seoul 133-791, Korea

<sup>3</sup>Corresponding author. E-mail: [wlee@hanyang.ac.kr](mailto:wlee@hanyang.ac.kr)

### Abstract

A study of benthic harpacticoids from the intertidal zone of Gosapo Beach, on the west (Yellow Sea) coast of Korea, resulted in the discovery of two species of *Wellsopsyllus* Kunz, 1981. The first new species, *Wellsopsyllus* (*Wellsopsyllus*) *egregius* sp. nov., was allocated to the nominotypical subgenus by a combination of three characters: 1) a 3-segmented P4 exopod, 2) 1-segmented P2–P3 endopods, and 3) a P1 endopod-2 with one apical seta. However, it can be clearly distinguished from its congeners by the following characters: 1) a caudal rami with two blunt setae and the seta VII located subdistally, 2) a P2 and P3 endopod represented by a small blunt process, and 3) the semioval shape of P5. The second new species was placed in the subgenus *Scottopsyllus* Apostolov & Marinov, 1988 according to the 3-segmented exopod and 2-segmented endopod of P4. *Wellsopsyllus* (*S.*) *koreanus* sp. nov. exhibits a unique combination of characters: 1) a dimpled distal margin of the female P5 endopodal lobe armed with two subequal setae, 2) a caudal seta V consisting of a stout spine-like base with a long cylindrical seta-like portion arising subdistally, and 3) a distinct bundle of five long inner setules on the P1 basis. We present a summary of the characters of all species in the genus *Wellsopsyllus* and propose an updated key for their identification.

**Key words:** *Wellsopsyllus*, *Scottopsyllus*, interstitial, meiofauna, biodiversity, Korea

### Introduction

More than 222,000 accepted eukaryotic marine species have been described worldwide (Appeltans et al. 2012; Costello 2013). According to a current review of marine biodiversity, Australia and Japan are the most speciose countries with respect to marine life (Costello et al. 2010). Interestingly, the most speciose region per unit area is South Korea (Costello et al. 2010). The National Marine Biodiversity Institute of Korea (MABIK) has invested significant effort in studying the marine biodiversity of Korean waters and runs an inventory project on marine organisms that includes registration of specimens in a national database. As a part of this project's ongoing effort to survey marine biodiversity, seven species belonging to the family Paramesochridae Lang, 1944 have been described in Korea. Six of these species have been found on sandy beaches while one was identified in brackish water (Back and Lee 2010; 2012; 2013; Back et al. 2011; Song et al. 2012). In the present study, two new species and six potential new species belonging to the family Paramesochridae were collected from coastal zones (Figure 1).

The family Paramesochridae harbors 13 genera (Wells 2007; Huys 2009). Huys (1987) mentioned that Paramesochridae consists of the subfamily Diarthrodellinae Huys, 1987 (*Diarthrodella* Klie, 1949, *Tisbisoma* Božić, 1964, and *Rossopsyllus* Soyer, 1975) and Paramesochrinae Lang, 1944 (*Wellsopsyllus*, *Paramesochra* groups and *Remanea* Klie, 1929). In Huys' (1987) phylogenetic tree, the *Wellsopsyllus* group was distinguished from the *Paramesochra* group by two apomorphies: 1) a 1-segmented P2–P3 endopod and 2) a reduction in setation of the distal segment in P2–P4.

Following the suggestion by Kunz (1962) to form the new genus *Scottopsyllus*, it was further divided into three subgenera (Kunz 1981) as follows: *Scottopsyllus* Kunz, 1981, *Intermediopsyllus* Kunz, 1962, and *Wellsopsyllus*

## References

- Appeltans, W., Ahyong, S.T., Anderson, G., Angel, M.V., Artois, T., Bailly, N., Bamber, R., Barber, A., Bartsch, I., Berta, A., Błażewicz-Paszkowycz, M., Bock, P., Boxshall, G., Boyko, C.B., Brandão, S.N., Bray, R.A., Bruce, N.L., Cairns, S.D., Chan, T.-Y., Cheng, L., Collins, A.G., Cribb, T., Curini-Galletti, M., Dahdouh-Guebas, F., Davie, P.J.F., Dawson, M.N., De Clerck, O., Decock, W., De Grave, S., de Voogd, N.J., Domning, D.P., Emig, C.C., Erséus, C., Eschmeyer, W., Fauchald, K., Fautin, D.G., Feist, S.W., Fransen, C.H.J.M., Furuya, H., Garcia-Alvarez, O., Gerken, S., Gibson, D., Gittenberger, A., Gofas, S., Gómez-Daglio, L., Gordon, D.P., Guiry, M.D., Hernandez, F., Hoeksema, B.W., Hopcroft, R.R., Jaume, D., Kirk, P., Koedam, N., Koenemann, S., Kolb, J.B., Kristensen, R.M., Kroh, A., Lambert, G., Lazarus, D. B., Lemaitre, R., Longshaw, M., Lowry, J., Macpherson, E., Madin, L.P., Mah, C., Mapstone, G., McLaughlin, P.A., Mees, J., Meland, K., Messing, C.G., Mills, C.E., Molodtsova, T.N., Mooi, R., Neuhaus, B., Ng, P.K.L., Nielsen, C., Norenburg, J., Opresko, D.M., Osawa, M., Paulay, G., Perrin, W., Pilger, J.F., Poore, G.C.B., Pugh, P., Read, G.B., Reimer, J.D., Rius, M., Rocha, R.M., Saiz-Salinas, J.I., Scarabino, V., Schierwater, B., Schmidt-Rhaesa, A., Schnabel, K., Schotte, M., Schuchert, P., Schwabe, E., Segers, H., Self-Sullivan, C., Shenkar, N., Siegel, V., Sterrer, W., Stöhr, S., Swalla, B., Tasker, M.L., Thuesen, E.V., Timm, T., Todaro, M.A., Turon, X., Tyler, S., Uetz, P., van der Land, J., Vanhooorne, B., van Ofwegen, L.P., van Soest, R.W.M., Vanaverbeke, J., Walker-Smith, G., Walter, T.C., Warren, A., Williams, G.C., Wilson S.P. & Costello, M.J. (2012) The magnitude of global marine species diversity. *Current Biology*, 22, 2189–2202.  
<http://dx.doi.org/10.1016/j.cub.2012.09.036>
- Back, J. & Lee, W. (2010) A new species of the genus *Paramesochra* (Copepoda: Harpacticoida) from Korean waters. *Proceedings of the Biological Society of Washington*, 123, 47–61.  
<http://dx.doi.org/10.2988/09-15.1>
- Back, J. & Lee, W. (2012) Two new species of *Apodopsyllus* (Copepoda, Harpacticoida, Paramesochridae) from Jeju Island, Korea. *Zootaxa*, 3368, 128–145.
- Back, J. & Lee, W. (2013) Three new species of the genus *Paramesochra* T. Scott, 1892 (Copepoda; Harpacticoida; Paramesochridae) from Yellow Sea, Korea with a redescription of *Paramesochra similis* Kunz, 1936. *Journal of Natural History*, 47, 769–703.  
<http://dx.doi.org/10.1080/00222933.2012.742585>
- Back, J., Lee, W. & Huys, R. (2011) A new species of the genus *Remanea* (Copepoda; Harpacticoida; Paramesochridae) with a redescription of *R. arenicola* Klie. *Journal of Natural History*, 45, 2939–2964.  
<http://dx.doi.org/10.1080/00222933.2011.622057>
- Becker, K.H., Noodt, W. & Schriever, G. (1979) Eidonomie und Taxonomie abyssaler Harpacticoida (Crustacea, Copepoda). Teil II. Paramesochridae, Cyliandropsyllidae und Cletodidae. *Meteor Forsch-Ergebnisse, Reihe D-Biologie*, 31, 1–37.
- Costello, M.J., May, R.M. & Stork, N.E. (2013) Can we name Earth's species before they go extinct? *Science*, 339, 413–416.  
<http://dx.doi.org/10.1126/science.1230318>
- Costello, M.J., Coll, M., Danovaro, R., Halpin, P., Ojaveer, H. & Miloslavich, P. (2010) A census of marine biodiversity knowledge, resources, and future challenges. *PLoS ONE*, 5, e12110.  
<http://dx.doi.org/10.1371/journal.pone.0012110>
- Huys, R. (1987) *Paramesochra* Scott, T. 1892 (Copepoda, Harpacticoida) – a revised key, including a new species from the SW Dutch coast and some remarks on the phylogeny of the Paramesochridae. *Hydrobiologia*, 144, 193–210.  
<http://dx.doi.org/10.1007/BF00005554>
- Huys, R. (2009) Unresolved cases of type fixation, synonymy and homonymy in harpacticoid copepod nomenclature (Crustacea: Copepoda). *Zootaxa*, 2183, 1–99.
- Huys, R., Gee, J.M., Moore, C.G. & Hamond, R. (1996) *Synopses of the British Fauna (New Series) No. 51. Marine and brackish water harpacticoid copepods, Part 1*. Field Studies Council, Shrewsbury, 352 pp.
- International Commission on Zoological Nomenclature (1999) *International Code of Zoological Nomenclature, Fourth Edition*. The International Trust for Zoological Nomenclature, London, xxx + 306 pp.
- Kornev, P.N. & Chertoprud, E.C. (2008) *Harpacticoid Copepods of the White Sea: Morphology, Systematics, Ecology*. KMK Scientific Press Ltd., Moscow, 379 pp. [in Russian]
- Kottmann, J., Kihara, T.C., Glatzel, T. & Veit-Köhler, G. (2012) A new species of *Wellsopsyllus* (Copepoda, Harpacticoida, Paramesochridae) from the deep Southern Ocean and remarks on its biogeography. *Helgoland Marine Research*, 67, 33–48.  
<http://dx.doi.org/10.1007/s10152-012-0302-7>
- Kunz, H. (1937) Zur Kenntnis der Harpacticoiden des Küstengrundwassers der Kieler Förde. *Kieler Meeresforschungen*, 2, 95–115.
- Kunz, H. (1962) Revision der Paramesochridae (Crust. Copepoda). *Kieler Meeresforschungen*, 18, 245–257.
- Kunz, H. (1981) Beitrag zur Systematik der Paramesochridae (Copepoda, Harpacticoida) mit Beschreibung einiger neuer Arten. *Mitteilungen aus dem Zoologischen Museum der Universität Kiel*, 1, 1–33.
- Kunz, H. (1992) Beitrag zur Kenntnis mariner Copepoda Harpacticoida (Fam. Paramesochridae Lang) mit Beschreibung zweier neuer Arten und einer neuen Unterart. *Crustaceana*, 62, 85–97.  
<http://dx.doi.org/10.1163/156854092X00073>
- Lang, K. (1965) Copepoda Harpacticoida from the Californian Pacific coast. *Kungl. svenska Vetenskapsakademiens*

*Handlingar*, 10, 1–560.

- Letova, V.N. (1982) Marine invertebrates of coastal biocenoses of the arctic ocean and the pacific ocean. [Harpacticoida (Crustacea, Copepoda) from the mud-sandy littoral of the East Murman]. *Issledovaniya Fauny Morei*, 29, 46–75. [in Russian]
- Mielke, W. (1975) Systematik der Copepoda eines Sandstrandes der Nordseeinsel Sylt. *Mikrofauna des Meeresboden*, 52, 1–134.
- Mielke, W. (1984) Interstitielle Fauna von Galapagos. XXXVI. Paramesochridae (Harpacticoida). *Microfauna Marina*, 5, 95–147.
- Nicholls, A.G. (1939) Some new sand-dwelling copepods. *Journal of the Marine Biological Association of the United Kingdom*, 23, 237–341.  
<http://dx.doi.org/10.1017/S0025315400013928>
- Plum, C. & George, K.H. (2009) The paramesochrid fauna of the Great Meteor Seamount (Northeast Atlantic) including the description of a new species of *Scottopsyllus* (*Intermedopsyllus*) Kunz (Copepoda: Harpacticoida: Paramesochridae). *Marine Biodiversity*, 39, 265–289.  
<http://dx.doi.org/10.1007/s12526-009-0022-7>
- Scott, T. & Scott, A. (1895a) On some new and rare Crustacea from Scotland. *The Annals and magazine of natural history, Series 6*, 15, 50–59.
- Scott, T. & Scott, A. (1895b) On new and rare species of Copepoda from Scotland. *The Annals of Scottish Natural History*, 13, 28–35.
- Song, S.J., Park, J., Kwon, B.-O., Ryu, J. & Khim, J.S. (2012) Ecological checklist of the marine brackish-water harpacticoid copepod fauna in Korean waters. *Zoological Studies*, 51, 1397–1410.
- Soyer, J. (1975) Harpacticoïdes (Crustacés Copépodes) de l'archipel de Kerguelen. I. Quelques forms mésopsammiques. *Bulletin du Muséum national d'histoire naturelle. 3e série, Zoologie*, 168, 1169–1223.
- Thompson, I.C. & Scott, A. (1900) Some recent additions to the Copepoda of Liverpool Bay. *Proceedings and transactions of the Liverpool Biological Society*, 14, 139–144.
- Veit-Köhler, G. (2000) Habitat preference and sexual dimorphism in species of *Scottopsyllus* (Copepoda, Harpacticoida) with the description of *Scottopsyllus* (*S.*) *praecipuus* sp n. from the Antarctic. *Vie Milieu*, 50, 1–17.
- Wells, J.B.J. (1965) Copepoda (Crustacea) from the meiobenthos of some Scottish marine sub-littoral muds. *Proceedings of the Royal Society of Edinburgh, Section B, Biological Sciences*, 69, 1–33.  
<http://dx.doi.org/10.1017/S0080455X00010110>
- Wells, J.B.J. (2007) An annotated checklist and keys to the species of Copepoda Harpacticoida (Crustacea). *Zootaxa*, 1568, 1–872.