

<http://dx.doi.org/10.11164/zootaxa.3918.3.9>

<http://zoobank.org/urn:lsid:zoobank.org:pub:47A64C4E-D6C6-42B9-8A8B-BAB5AD99D04D>

## Two new species of the genus *Arboridia* Zachvatkin (Hemiptera: Auchenorrhyncha: Cicadellidae: Typhlocybinae) from Korea

SUMIN OH, KWANG-RYUL CHOE & SUNGHOON JUNG<sup>1</sup>

Laboratory of Systematic Entomology, Department of Applied Biology, College of Agriculture and Life Sciences, Chungnam National University, Daejeon 305-764, Korea

<sup>1</sup>Corresponding author. E-mail: jung@cnu.ac.kr

### Abstract

Two new species of the genus *Arboridia* Zachvatkin, *A. koreana* sp. nov., *A. septempunctata* sp. nov., are described from Korea. Photographs of dorsal habitus, illustrations of male genitalia, abdomens and forewings, and biological information such as host plants and distributional records and a key to the Korean *Arboridia* species are provided.

**Key words:** Hemiptera, Cicadellidae, Typhlocybinae, *Arboridia*, new species

### Introduction

The leafhopper genus *Arboridia* Zachvatkin, 1946 belongs to the tribe Erythroneurini of subfamily Typhlocybinae (Hemiptera: Auchenorrhyncha: Cicadellidae). Several members of the genus *Arboridia* are known as grape pests in various Eurasian countries including Korea, and some of them are also known as pests on cherry, hawthorn, pear, peach, apple, mulberry, maple, woodbine, dogwood, and several other plants (Song & Li, 2013). A total of 69 valid species have been described in the Palaearctic and Oriental region (Song & Li, 2013). So far, seven species of this genus have been recorded in the Korean Peninsula (Kwon & Huh, 2001).

In this study, two new species of *Arboridia* are described and illustrated from Korea, and a key to the Korean *Arboridia* species is also provided.

The materials used in this study are stored in the Laboratory of Systematic Entomology, Chungnam National University (CNU), Daejeon, Korea. Morphological terminology follows Dietrich (2005). The genitalia segment was detached from abdomen and macerated in 10% KOH for about 60min. in 70° for dissection to examine male genitalia.

### Genus *Arboridia* Zachvatkin, 1946

*Arboridia* Zachvatkin, 1946: 153 (Type species: *Typhlocyba parvula* Boheman, 1845)  
*Khoduma* Dworakowska, 1972: 403 (Type species: *Khoduma jacobii* Dworakowska, 1972)

**Diagnosis.** Vertex with one or two pair of dark spots.. Pronotum without conspicuous pits. Abdominal apodemes small and pygofer lobes without macrosetae. Subgenital plates with group of macrosetae at middle of outer margin. Style apex with 3 points.

### Key to the *Arboridia* species of the Korean Peninsula

- |   |   |
|---|---|
| 1. Forewings with white round markings. Pronotum anterior margin with one black anterior spot ..... | 2 |
| - Forewings without white round markings. Pronotum anterior margin without black spot .....         | 5 |

**Type materials.** Holotype: [CNU] 1♂, Korea, Suwon, 23.x.1974, coll. T.Okada. Paratypes: 1♂, 2♀♀, same data as holotype.; [CNU] 2♀♀, Korea, Suwon, 15.v.1979. coll. K.R.Choe.

**Host plants.** *Vitis* sp. (Vitales: Vitaceae)

**Distributions.** Korea (Suwon)

**Remarks.** This species is similar to *Arboridia maculifrons* (Vilbaste, 1968), but differs in forewing pattern and the black spot above the antennal pit.

**Etymology.** The specific name is derived from the characteristic 7 prominent spots on the head, pronotum and mesonotum.

## Acknowledgements

We thank Prof. Kamitani and Dr. Ohara (Kyushu University) for kindly helping us to examine leafhopper specimens deposited at insect collections of Kyushu University. We also thank Dr. C. Dietrich, the editor of *Zootaxa* (Membracoidea), and an anonymous reviewer for their suggestions and corrections, which certainly improved this paper. This work was supported by a grant from the National Institute of Biological Resources (NIBR), funded by the Ministry of Environment (MOE) of the Republic of Korea (NIBR 201401203).

## References

- Anufriev, G.A. (1969) New and little known leaf-hoppers of the subfamily Typhlocybinae from the Soviet Maritime Territory (Homopt., Auchenorrhyncha). *Acta Faunistica Entomologica Musei Nationalis Pragae*, 13 (153), 163–190.
- Boheman, C.H. (1845) Nya Svenska Homoptera. *Ofversigt af Kongliga Svenska Vetenskaps-Akademiens Förfärlingar*, 1, 154–164.
- Dietrich, C.H. (2005) Keys to the families of Cicadomorpha and subfamilies and tribes of Cicadellidae (Hemiptera: Auchenorrhyncha). *Florida Entomologist*, 88 (4), 502–517.  
[http://dx.doi.org/10.1653/0015-4040\(2005\)88\[502:KTTFOC\]2.0.CO;2](http://dx.doi.org/10.1653/0015-4040(2005)88[502:KTTFOC]2.0.CO;2)
- Dworakowska, I. (1972) On some oriental Erythroneurini (Auchenorrhyncha, Cicadellidae, Typhlocybinae). *Bulletin de l' Académie Polonaise des Sciences Série des Sciences Biologiques*, 20 (6), 395–405.
- Kwon, Y.J. & Huh, E.Y. (2001) HOMOPTERA (Suborder Auchenorrhyncha). National Institute of Agricultural Science and Technology, Suwon, 460pp. [Korea]
- Matsumura, S. (1916) *Oyo konchugaku*. Keiseisha Shoten, Taisha, Tokyo, 731 pp.
- Matsumura, S. (1932) A revision of the Palaearctic and Oriental Typhlocybid-genera with descriptions of new species and new genera. *Insecta Matsumura*, 6 (3), 93–120
- Nawa, U. (1913) *Zigina apicalis* Mats. (Erythroneura apicalis Nawa). *Insect World*, 17 (12), 480–486
- Song, Y.H. & Li, Z.Z. (2013) Some new species and new record of the genus *Arboridia* Zachvatkin (Hemiptera: Cicadellidae: Typhlocybinae) from six provinces of China. *Zootaxa*, 3613 (3), 229–244.  
<http://dx.doi.org/10.11646/zootaxa.3613.3.2>
- Vilbaste, J. (1968) *Über die Zikadenfauna des Primorje Gebietes*. Valgus, Tallinn, 195 pp.
- Zachvatkin, A.A. (1946) Studies on the Homoptera of Turkey. I–VII. *Transactions of the Royal Entomological Society of London*, 97 (6), 149–176.  
<http://dx.doi.org/10.1111/j.1365-2311.1946.tb00278.x>