

<http://dx.doi.org/10.11646/zootaxa.3949.3.3>

<http://zoobank.org/urn:lsid:zoobank.org:pub:CBEF804D-5FBF-48D2-8CCD-036B70FE6ECD>

***Onuphis* and *Aponuphis* (Annelida: Onuphidae) from southwestern Europe, with the description of a new species**

ANDRÉS ARIAS^{1,4} & HANNELORE PAXTON^{2,3}

¹Departamento de Biología de Organismos y Sistemas (Zoología), Universidad de Oviedo, Oviedo 33071, Spain

Department of Biological Sciences, Macquarie University, Sydney, NSW 2109, Australia

³Australian Museum Research Institute, Australian Museum, 6 College Street, Sydney, NSW 2010, Australia

⁴Corresponding author. E-mail: ariasandres.uo@uniovi.es

Abstract

This study recognises five species of *Onuphis* and four species of *Aponuphis* from southwestern Europe, represented in the Bay of Biscay, the Atlantic coasts of Iberia and the western Mediterranean Sea. One species, *Onuphis anadonae* sp. nov., is newly described and another species, *Aponuphis willsiei* Cantone & Bellan, 1996 is a new record for the Iberian Peninsula waters. *Aponuphis ornata* (Fauvel, 1928) and *A. willsiei* are shown to be tube brooders and the development of the juveniles is described. The usefulness of some *Onuphis* characters, as well as the size-dependent distribution of brachiae and chaetae, complicating *Aponuphis* species identification, are discussed. We are presenting descriptions and photographs of the distinctive *Aponuphis* colour pattern which we recommend as an identificatory tool instead. A dichotomous key to all species is included.

Key words: polychaetes, Eunicida, biodiversity, direct development, ontogenetic changes, chaetal replacement, Iberian Peninsula, Bay of Biscay, western Mediterranean

Introduction

Onuphis Audouin & Milne Edwards, 1833 is the most speciose genus of the polychaete family Onuphidae Kinberg, 1865 after *Diopatra* Audouin & Milne Edwards, 1833 with about 40 described valid species (Maekawa & Hayashi 1999; Gil & Machado 2014; Arias & Paxton 2014). However, the genus has only been the subject of partial revisions (Fauchald 1982; Maekawa & Hayashi 1999; Arias & Paxton 2014) and it is quite possible that more undescribed species may be discovered in the near future. The genus *Onuphis* is a cosmopolitan taxon and most of its species are characteristic for the intertidal and relatively shallow waters of warm-temperate regions, nevertheless individual species of *Onuphis* seem to have limited restricted geographical distributions (Paxton 1986).

The genus *Aponuphis* Kucheruk, 1978 was erected for a group of slender onuphids without peristomial cirri that had previously been regarded as members of the genus *Hyalinoecia* Malmgren, 1867. Paxton (1986) recognised seven species of *Aponuphis* and underscored the morphological differences between the two genera by placing *Aponuphis* into the subfamily Onuphinae and *Hyalinoecia* into the subfamily Hyalinoeciinae.

Southwestern Europe, represented by the Bay of Biscay, the Atlantic coasts of Iberia and the western Mediterranean Sea, harbours five species of *Onuphis*, of which one is new to science, and four species of *Aponuphis*. Here, we present an updated revision of the *Onuphis* and *Aponuphis* fauna of this region with the description and illustration of the new species of *Onuphis*, newly report *A. willsiei* from the Iberian Peninsula and present notes on the ecology and reproduction together with a taxonomic key to the species of *Onuphis* and *Aponuphis* of southwestern Europe.

2.	Colour pattern and branchiae absent	<i>A. willsie</i>
-	Colour pattern and branchiae present	3
3.	Branchiae from chaetiger 1–2, colour pattern of two large dorsal spots per segment, by about chaetiger 10 becoming thin band (Fig. 7C)	<i>A. brementi</i>
-	Branchiae from chaetiger 3–5, colour pattern otherwise	4
4.	Pseudocompound hooks tridentate; branchiae from chaetiger 3–4; colour pattern of horizontal bars and bands (Fig. 7D–F)	<i>A. ornata</i>
-	Pseudocompound hooks bi- and tridentate; branchiae from chaetiger 4–5; colour pattern otherwise	5
5.	Two dorsal longitudinal pigment stripes and two segmental spots from chaetiger 1 to posterior region (Fig. 7A)	<i>A. bilineata</i> morph 1 (typical)
-	Peristomium and chaetigers 3–4 almost solidly light brownish, from chaetiger 4–5 two lateral longitudinal stripes close to parapodia (Fig. 7B)	<i>A. bilineata</i> morph 2
6.	Branchiae from chaetiger 1	7
-	Branchiae from chaetiger 5, all branchiae simple and strap-like	<i>O. farensis</i>
7.	All branchiae simple and strap-like, colour pattern practically absent, first four pairs of parapodia with pseudocompound hooks	8
-	Branchiae pectinate from about the 20 th chaetiger, colour pattern consisting of distinctive bars and spots on dorsum, first three pairs of parapodia with pseudocompound hooks	9
8.	Bi-, tri- and quadridentate pseudocompound hooks with falcate hoods (Fig. 3F–I), SAH from chaetiger 9, subulate ventral cirri in first four chaetigers	<i>O. anadonae</i> sp. nov.
-	Only tridentate pseudocompound hooks with moderately long, triangular and pointed hoods (Fig. 6A–D), SAH from chaetiger 14, subulate ventral cirri in first five to six chaetigers	<i>O. rullieriana</i>
9.	Bi- and tridentate (or even multidentate) pseudocompound hooks, eyespots on prostomium absent, maximum of eight branchial filaments, robust species	<i>O. pancerii</i>
-	Only tridentate pseudocompound hooks, eyespots on prostomium present, maximum of five branchial filaments, slender species	<i>O. eremita</i>

Acknowledgements

We are grateful to Javier Sánchez Almazán and Francisco Yagüe, Museo Nacional de Ciencias Naturales of Madrid, for the generous loan of most of the material on which this work is based, including the material from the Fauna Ibérica Project, to Helmut Sattmann and Stefan Szeiler, Naturhistorisches Museum Wien, for photographs of syntypes on slides, and to the curator and librarian of the Musée Océanographique de Monaco for help with locating specimens and literature. Our thanks go to Jorge Núñez and an anonymous reviewer for their careful reviews and helpful remarks.

We also thank Nuria Anadón for her constant support, Alfredo J. Quitana, Scientific-Technical Services of the University of Oviedo, for assistance with SEM microscopy and Sue Lindsay, Microscopy Unit of the Australian Museum, Sydney, for assistance with photography. The first author (AA) also wants to thank the Department of Biological Sciences, Macquarie University, Sydney, for hosting him for his overseas study period in 2013. This work was supported by the project RES-14-CI-058, University of Oviedo, and the FICYT foundation (Principado de Asturias).

References

- Aguirrezabalaga, F., Ceberio, A. & Paxton, H. (2002) Onuphidae (Polychaeta) from the Capbreton Canyon (Bay of Biscay, NE Atlantic) with the description of *Paradiopatra capbretonensis* sp. nov. *Steenstrupia*, 27 (1), 19–28.
- Amoureaux, L. (1971) Problèmes biologiques suggérés par certaines répartitions d'Annélides Polychètes dans la zone meuble intertidale. Divers problèmes sont évoqués à propos de la répartition de *Travisia forbesi* et *d'Ophelia bicornis*. *Vie et Milieu*, Supplément 22 (2), 701–705.
- Amoureaux, L. (1972) Annélides polychètes recueillies sur les pentes du talus continental, au large de la Galice (Espagne). Campagnes 1967 et 1968 de la "Thalassa". *Cahiers de Biologie Marine*, 13, 63–89.
- Amoureaux, L. (1973) Annélides polychètes recueillies sur les pentes du talus continental au Nord de la côte espagnole. Campagne 1970 de la "Thalassa". *Cahiers de Biologie Marine*, 19, 429–452.
- Amoureaux, L. (1974) Annélides polychètes recueillies sur les pentes du talus continental au nord-ouest de l'Espagne et du Portugal (Campagne 1972 de la "Thalassa"). *Cuadernos de Ciencias Biológicas*, 3, 121–54.
- Amoureaux, L. (1976) Annélides polychètes récoltées par J. Stirn en 1969, sur les côtes marocaines du Détrroit de Gibraltar. *Cuadernos de Ciencias Biológicas de la Universidad de Granada*, 5, 5–33.

- Amoureux, L. (1977) Annélides polychètes errantes recueillies sur les pentes du talus continental à l'entrée de la Manche, avec la description de deux espèces nouvelles. Campagne 1973 de la "Thalassa". *Cahiers de Biologie Marine*, 18 (4), 391–411.
- Arias, A., Anadón, N. & Paxton, H. (2010) New records of *Diopatra marocensis* (Annelida: Onuphidae) from northern Spain. *Zootaxa*, 2691, 67–68.
- Arias, A., Paxton, H. & Anadón, N. (2013) *Mooreonuphis vespa*, a new brooding species of Onuphidae (Annelida) from northern Spain (Bay of Biscay). *Zootaxa*, 3741 (4), 583–592.
<http://dx.doi.org/10.11646/zootaxa.3741.4.8>
- Arias, A. & Paxton, H. (2014) Hidden diversity within the polychaete *Onuphis eremita sensu lato* (Annelida: Onuphidae)—redescription of *O. eremita* Audouin & Milne-Edwards, 1833 and reinstatement of *Onuphis pancerii* Claparède, 1868. *Zootaxa*, 3861 (2), 145–169.
- Audouin, J.V. & Milne Edwards, H. (1833) Classification des Annélides et description de celles qui habitent les côtes de la France. *Annales des sciences naturelles*, 29, 195–269.
- Baird, W. (1870) Remarks on several genera of annelids belonging to the group Eunicea with a notice of such species as are contained in the collection of the British Museum and a description of some others hitherto undescribed. *Journal of the Linnean Society of London*, 10, 341–361.
<http://dx.doi.org/10.1111/j.1096-3642.1869.tb00665.x>
- Bellan, G. (1960) Annélides polychètes. Résultats Scientifiques de la Campagne du N.R.P. "Faial" dans les eaux côtières du Portugal (1957). *Lisboa*, 2, 1–31.
- Bellan, G. (1964) *Contribution à l'étude systématique, bionomique et écologique des annélides polychètes de la Méditerranée*. Thèse, Faculté des Sciences de l'Université d'Aix-Marseille, 371 pp.
- Bellan, G. (1969) Annélides Polychètes recueillis dans l'Archipel de Madère au cours de la campagne scientifique du navire océanographique "Jean Charcot". *Cahiers de Biologie Marine*, 10 (1), 35–57.
- Blake, J.A. (1974) The larval development of Polychaeta from the Northern California coast. II. *Nothria elegans* (Family Onuphidae). *Ophelia*, 13, 43–61.
<http://dx.doi.org/10.1080/00785326.1974.10430591>
- Borja, A., Franco, J. & Pérez, V. (2000) A Marine Biotic Index to establish the ecological quality of soft-bottom benthos within European estuarine and coastal environments. *Marine Pollution Bulletin*, 40 (12), 1100–1114.
[http://dx.doi.org/10.1016/S0025-326X\(00\)00061-8](http://dx.doi.org/10.1016/S0025-326X(00)00061-8)
- Campoy, A. (1982) Fauna de España. Fauna de anélidos poliquetos de la Península Ibérica. *EUNSA Publicaciones de Biología de la Universidad de Navarra*, Serie Zoológica, 7, 1–781. [Pamplona]
- Cantone, G. & Bellan, G. (1996) *Aponuphis willsieei*, una nuova specie di Onuphidae (Annelida, Polychaeta) delle coste marsigliesi. *Animalia*, 21 (1/3), 27–30.
- Castelli, A. (1982) *Onuphis falesia*, a new species of Onuphinae (Polychaeta, Eunicidae). *Bollettino di Zoologia*, 49, 45–49.
<http://dx.doi.org/10.1080/11250008209439371>
- Çinar, M.E. (2009) Alien polychaete species (Annelida: Polychaeta) on the southern coast of Turkey (Levantine Sea, eastern Mediterranean), with 13 new records for the Mediterranean Sea. *Journal of Natural History*, 43, 2283–2328.
<http://dx.doi.org/10.1080/00222930903094654>
- Claparède, E. (1868) Les Annélides Chétopodes du Golfe de Naples. *Mémoires de la Société de physique et d'histoire naturelle de Genève*, 19, 313–584.
- Day, J.H. (1967) *A monograph on the Polychaeta of Southern Africa I*. British Museum (Natural History) Publications, London, 458 pp.
- Dexter, D. (1992) Soft bottom invertebrates of the Portuguese benthos. *Boletim do Instituto Nacional de Investigação das Pescas (Lisboa)*, 17, 61–88.
- Ehrenberg, C.G. (1830) Neue Beobachtungen über blutartige Erscheinungen in Aegypten, Arabien und Sibirien, nebst einer Uebersicht und Kritik der früher bekannten. *Annual Review of Physical Chemistry*, 94, 4, 477–514.
<http://dx.doi.org/10.1002/andp.18300940402>
- Fauchald, K. (1982) Revision of *Onuphis*, *Nothria*, and *Paradiopatra* (Polychaeta: Onuphidae) based upon type material. *Smithsonian Contributions to Zoology*, 356, 1–109.
<http://dx.doi.org/10.5479/si.00810282.356>
- Fauvel, P. (1916) Deux polychètes nouvelles (*Disoma watsoni* n. sp., et *Hyalinoecia brementi* n. sp.). *Bulletin d'Institut Oceanographique* (Monaco), 316, 1–10.
- Fauvel, P. (1923) *Faune de France: Polychètes errantes*. Paul Lechevalier, Paris, 488 pp.
- Fauvel, P. (1928) Annélides Polychètes nouvelles du Maroc. *Bulletin de la Société Zoologique de France*, 53, 9–13.
- Fauvel, P. (1936) Contribution à la faune des Annélides Polychètes du Maroc. *Mémoires de la Société des Sciences Naturelles du Maroc*, 43, 1–143.
- Fauvel, P. (1938) Annelida Polychaeta della Laguna di Venezia. *R. Comitato Talassografico Italiano*, 246, 1–27.
- Gil, J. & Machado, M. (2014) A new species of *Onuphis* (Polychaeta: Onuphidae) from Southern Portugal, with comments on the validity of *O. pancerii* Claparède, 1868. *Zootaxa*, 3860 (4), 343–360.
- Glémarec M. (1969) *Les peuplements benthiques du plateau continental Nord-Gascogne*. Thèse d'Etat, Faculté des Sciences, Paris, 167 pp.
- Guille, A. (1971) Bionomie benthique de plateau continental de la côte catalane française. VI. Données autoécologiques

- (macrofaune). *Vie et Milieu*, 22, 469–527.
- Hartman, O. (1951) The littoral marine annelids of the Gulf of Mexico. *Publications of the Institute of Marine Sciences*, 2, 7–124.
- Hartmann-Schröder, G. (1977) Polychaeten aus dem Sublitoral und Bathyal vor der portugiesischen und marokkanischen Küste. Auswertung der Fahrt 8 (1967) von F.S. 'Meteor'. *METEOR Forsch.-Ergebnisse*, Series D, 26, 65–99.
- Hilbig, B. (1995) Family Onuphidae Kinberg, 1865. In: Blake, J.A., Hilbig, B. & Scott, P.H. (Eds.), *Taxonomic Atlas of the Benthic Fauna of the Santa Maria Basin and Western Santa Barbara Channel. 5 - The Annelida Part 2. Polychaeta: Phyllodocida (Syllidae and scale-bearing families), Amphipomida, and Eunicida*. Santa Barbara Museum of Natural History, Santa Barbara, pp. 341–364.
- Hobson, K.D. (1971) Some polychaetes of the superfamily Eunicea from the North Pacific and North Atlantic Oceans. *Proceedings of the Biological Society of Washington*, 83, 527–544.
- Ibáñez, M. (1972) Notas sobre algunas especies de Anélidos Poliquetos nuevas para las costas de España, con especial referencia a *Nereis (Neanthes) oxyopoda* Marenzeller y *Onuphis (Nothria) geophiliformis* Moore. *Boletín de la Real Sociedad Española de Historia Natural (Sección Biológica)*, 70, 23–31.
- Ibáñez, M. (1973) Catálogo de los anélidos poliquetos citados en las costas españolas. *Cuadernos de Ciencias Biológicas, Universidad de Granada*, 2, 121–140.
- Imajima, M. (1960) Description of a new polychaete, *Nothria shirikishinaiensis* n. sp. of the family Eunicidae. *Publications of the Seto Marine Biological Laboratory*, 8 (1), 55–58.
- Imajima, M. (1986) Eight species of Onuphidae (Polychaeta) in and Offshore of Otsuchi Bay, Northeastern Japan. *Bulletin of the National Science Museum of Tokyo*, 12, 93–116.
- Kinberg, J.C.H. (1865) *Annulata nova. Öfversigt af Königlich Vetenskapsakademiens förhandlingar (Stockholm)*, 21, 559–574.
- Kucheruk, N.V. (1978) Deep-water Onuphidae (Polychaeta) from the collections of the 16th cruise of the R/V Dmitry Mendeleev (to the generic classification of the family Onuphidae). *Trudy Instituti Okeanologii Akademii Nauk SSSR*, 113, 88–106.
- Langerhans, P. (1880) Wurmsfauna von Madeira. *Zeitschrift für wissenschaftliche Zoologie*, 33, 267–316.
- Laubier, L. & Paris, J. (1962) Faune marine des Pyrénées Orientales 4: Annélides Polychètes. *Vie et Milieu*, 13 (Supplement), 1–80.
- López, E. & San Martín, G. (1992) Familias de poliquetos errantes (Polychaeta), excepto Syllidae, recolectadas en las Islas de Cabo Verde por la "I Expedición Ibérica". *Revista de Biología Tropical*, 40 (2), 161–169.
- Louzao, M., Anadón, N., Arrontes, J., Álvarez-Claudio, C., Fuente, D.M., Ocharan, F., Anadón, A. & Acuña, J.L. (2010) Historical macrobenthic community assemblages in the Avilés Canyon, N Iberian Shelf: baseline biodiversity information for a marine protected area. *Journal of Marine Systems*, 80, 47–56.
<http://dx.doi.org/10.1016/j.jmarsys.2009.09.006>
- Maekawa, N. & Hayashi, I. (1989) Onuphid polychaetes from Wakasa Bay, Sea of Japan. *Memoirs of the College of Agriculture, Kyoto University*, 134, 61–93.
- Maekawa, N. & Hayashi, I. (1999) Taxonomic study on the genus *Onuphis* (Polychaeta, Onuphidae) from Japan and adjacent seas, with descriptions of six new species. *Bulletin of the National Science Museum, Series A (Zoology)*, 25 (3), 163–214.
- Malmgren, A.J. (1866) *Annulata Polychaeta Spetsbergiae, Groenlandiae, Islandiae et Scandinaviae hactenus cognita*. Ex Officina Frenckelliana, Helsingfors, 127 pp.
- Marenzeller, E. von (1879) Südjapanische Anneliden I. *Dankschrift der Mathematisch-Naturwissenschaftlichen Classe der Kaiserlichen Akademie der Wissenschaften*, 41 (2), 109–152. [Wien]
- Martínez, J. & Adarraga, I. (2006) Programa de vigilancia y control de la introducción de especies invasoras en los ecosistemas litorales de la costa vasca. 2. Costa de Bizkaia. Departamento de Medio Ambiente y Ordenación del Territorio del Gobierno Vasco /Sociedad Cultural de Investigación Submarina, Donostia-San Sebastián, 267 pp.
- Martínez, J., Adarraga, I. & Ruiz, J.M. (2010) Tipificación de poblaciones bentónicas de los fondos blandos de la plataforma continental de Guipúzcoa (sureste del golfo de Vizcaya). *Boletín del Instituto Español de Oceanografía*, 23 (1–4), 85–110.
- Moore, J.P. (1903) Polychaeta from the coastal slope of Japan and from Kamchatka and Bering Sea. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 55, 401–490.
- Mucha, A.P. & Costa M.H. (1999) Macrozoobenthic community structure in two Portuguese estuaries: relationship with organic enrichment and nutrient gradients. *Acta Oecologica*, 20 (4), 363–376.
[http://dx.doi.org/10.1016/S1146-609X\(99\)00130-7](http://dx.doi.org/10.1016/S1146-609X(99)00130-7)
- Orensanz, J.M. (1990) The eunicemorph polychaete annelids from Antarctic and Subantarctic Seas. With addenda to the Eunicemorpha of Argentina, Chile, New Zealand, Australia, and the Southern Indian Ocean. *Biology of the Antarctic Seas. XXI. Antarctic Research Series*, 52, 1–183.
<http://dx.doi.org/10.1029/AR052p0001>
- Paxton, H. (1986) Generic revision and relationships of the family Onuphidae (Annelida: Polychaeta). *Records of the Australian Museum*, 38, 1–74.
<http://dx.doi.org/10.3853/j.0067-1975.38.1986.175>
- Paxton, H. (1993) *Diopatra* Audouin and Milne Edwards (Polychaeta: Onuphidae) from Australia, with a discussion of developmental patterns in the genus. *The Beagle, Records of the Northern Territory Museum of Arts and Sciences*, 10, 115–154.

- Paxton, H., Fadlaoui, S. & Lechapt, J.P. (1995) *Diopatra marocensis*, a new brooding species of Onuphidae (Annelida: Polychaeta). *Journal of Marine Biological Association of the United Kingdom*, 75, 949–955.
<http://dx.doi.org/10.1017/S0025315400038273>
- Paxton, H. (1996) *Hirsutonuphis* (Polychaeta: Onuphidae) from Australia, with a discussion of setal progression in juveniles. *Invertebrate Taxonomy*, 10, 77–96.
<http://dx.doi.org/10.1071/IT9960077>
- Paxton, H. (1998) The *Diopatra chiliensis* confusion - redescription of *D. chiliensis* (Polychaeta, Onuphidae) and implicated species. *Zoologica Scripta*, 27, 31–48.
<http://dx.doi.org/10.1111/j.1463-6409.1998.tb00427.x>
- Redondo, M.S. & San-Martín, G. (1997) Anélidos Poliquetos de la costa comprendida entre el cabo de San Antonio y el Puerto de Valencia. *Publicaciones Especiales del Instituto Español de Oceanografía*, 23, 225–233.
- Rioja, E. (1918) Datos para el conocimiento de la fauna de anélidos poliquetos del Cantábrico (2^a Parte). *Trabajos del Museo Nacional de Ciencias Naturales, Madrid, Serie Zoológica*, 37, 1–99.
- Rivain, V. (1983) *Contribution à l'étude dynamique et fonctionnelle des peuplements de sables fins du Golfe normano-breton*. Thèse 3^{ème} cycle, Océanographie Biologique, Université de Paris VI, Paris, 166 pp.