

Table. 1. Selected meristic values for species of *Caragobius*. For *C. urolepis*, condition in lectotype is underscored. Meristic values, except for pectoral-fin rays, are based on radiographs, and eight cleared and stained specimens of *C. urolepis* - two removed from USNM 241794; three from USNM 243403; and three from USNM 243404.

<b>Dorsal-fin rays (total elements)</b>																	
	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52
<i>C. rubristriatus</i>								1			1		3	3	2	1	1
<i>C. urolepis</i>	1	2	4	<u>8</u>	3	2	1	1									
<b>Anal-fin rays (total elements)</b>																	
	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45		
<i>C. rubristriatus</i>					1				1	2	1	2	4		1		
<i>C. urolepis</i>	2	5	<u>8</u>	3	3	1											
<b>Caudal Vertebrae</b>																	
	18	19	20	21	22	23	24	25	26	27							
<i>C. rubristriatus</i>								2	5	5							
<i>C. urolepis</i>	1	2	11	<u>7</u>	1												
<b>Pectoral-fin rays</b>					<b>Anal-fin Pterygiophores (AP)</b>												
	16	17	18	19	20	3	4	5	6	7							
<i>C. rubristriatus</i>	1	8	7	3	1	2	9	1									
<i>C. urolepis</i>		4	<u>11</u>	12	3		1	<u>11</u>	9	1							