

Krogmann, L. & Burks, R.A. (2009) *Doddifoenus wallacei*, a new giant parasitoid wasp of the subfamily Leptofoeninae (Chalcidoidea: Pteromalidae), with a description of its mesosomal skeletal anatomy and a molecular characterization. *Zootaxa*, 2194, 21–36.

The locality data of the four specimens collected in Thailand (1 holotype, 3 paratypes) was incorrectly cited from the labels. The term “Thale Ban National Park” was erroneously added, but Thale Ban National Park is in Satun province, and some 60 km from the type locality in Songkhla province. The locality information of the fifth specimen (paratype from Laos) is correct.

page	section	error	corrected
24	Specimens examined.	Holotype ♀: THAILAND. <i>Songkhla</i> : Thale Ban National Park, Nam Tok Ton Pliu, 100m, 7°00'02''N 100°14'07''E, 17.ii.2005, D.Yanega. (UCRC, accession no. 104069). 2 paratype ♀♀: same data as holotype (UCRC, accession nos. 104070, 142549). 1 paratype ♂: same data as before, but collected by G. Ballmer (UCRC, accession no. 142548).	Holotype ♀: THAILAND. <i>Songkhla</i> : Nam Tok Ton Pliu, 100m, 7°00'02''N 100°14'07''E, 17.ii.2005, D.Yanega. (UCRC ENT 104069). 2 paratype ♀♀: same data as holotype (UCRC ENT 104070, 142549). 1 paratype ♂: same data as before, but collected by G. Ballmer (UCRC ENT 142548).