

***Andraegoidus* Aurivillius: new species, synonymies and key to the species (Insecta: Coleoptera: Cerambycidae)**

JUAN PABLO BOTERO R.¹ & MARCELA L. MONNÉ^{1,2}

¹Departamento de Entomologia, Museu Nacional, Universidade Federal do Rio de Janeiro, UFRJ, Quinta da Boa Vista, São Cristóvão, CEP 20940-040, Rio de Janeiro, RJ, Brazil. E-mail: jp_bot@yahoo.com

²Fellow of the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq)

Abstract

Andraegoidus translucidus sp. nov. is described from Bolivia (Santa Cruz). The following synonyms are proposed: *Andraegoidus homoplatus hassenteufeli* Fuchs, 1958 = *Andraegoidus homoplatus* (Dupont, 1838), and *Andraegoidus lacordairei punctipennis* Hüdepohl, 1985 = *Andraegoidus lacordairei* (Dupont, 1838). A key to the species of this genus is provided. The geographical distribution of *Andraegoidus lacordairei* is now widened to Trinidad and Tobago (Talparo).

Key words: Cerambycinae, Trachyderini, new species, synonymies

Resumo

Andraegoidus translucidus sp. nov. é descrita para a Bolívia (Santa Cruz). Os seguintes sinônimos são propostos: *Andraegoidus homoplatus hassenteufeli* Fuchs, 1958 = *Andraegoidus homoplatus* (Dupont, 1838) e *Andraegoidus lacordairei punctipennis* Hüdepohl, 1985 = *Andraegoidus lacordairei* (Dupont, 1838). Chave para as espécies é fornecida. A distribuição geográfica de *Andraegoidus lacordairei* é ampliada para Trinidad e Tobago (Talparo).

Palavras-chave: Cerambycinae, Trachyderini, nova espécie, sinonímias

Introduction

The subfamily Cerambycinae includes approximately 3.000 species and 771 genera in the Neotropical region. The tribe Trachyderini Dupont, 1836 is allocated in Cerambycinae, and includes 139 genera and almost 400 species that are widely distributed in the Americas (Monné, 2005).

Andraegoidus Aurivillius, 1920 currently includes seven species and six subspecies, which occur only in South America. The genus was described by Aurivillius (1920) for a single species, *Andraegoidus humeralis* from Argentina. The species was considered a synonym of *Trachyderes cruentatus* (Dupont, 1838) by Bosq & Rufinelli (1951), but was later resurrected and returned to *Andraegoidus* by Hüdepohl (1985).

Tippmann (1953) described *A. laticollis* from Colombia, and Fuchs (1958) described *A. hassenteufeli* from Paraguay. Hüdepohl (1985) described *A. distinguendus* from Brazil and *A. lacordairei punctipennis* from Venezuela, and transferred *Trachyderes homoplatus* Dupont, 1838, *T. lacordairei* Dupont, 1838, *T. rufipes* (Fabricius, 1787), and *T. variegatus* Perty, 1832 to *Andraegoidus*. *Trachyderes fabricii* Dupont, 1838, *T. fulvipennis* Dupont, 1838, *T. richteri* Bruch, 1908, and *T. zonatus* Dalman, 1823 were transferred to *Andraegoidus* and considered subspecies of *A. rufipes*, and the species *A. hassenteufeli* was considered a subspecies of *A. homoplatus*.