



## A new species of *Attheyella* (*Canthosella*) from Colombia and redescription of *Attheyella* (*Delachauxiella*) *freyi* (Copepoda: Harpacticoida: Canthocamptidae)

SANTIAGO GAVIRIA<sup>1</sup> & DANIELLE DEFAYE<sup>2</sup>

<sup>1</sup>University of Vienna, Dept. of Limnology & Technisches Büro für Biologie, Fred-Raymondgasse 19/2/4, A-1220 Vienna, Austria  
E-mail: santiago.gaviria@gmx.at

<sup>2</sup>Muséum national d'Histoire Naturelle, Département Milieux et Peuplements aquatiques, UMR 7208, CP 53, 61 rue de Buffon, F-75005 Paris, France. E-mail: ddefaye@mnhn.fr

### Table of content

Abstract	1
Introduction	1
Material and methods	2
Taxonomic descriptions	3
Family Canthocamptidae Brady, 1880	3
Genus <i>Attheyella</i> Brady, 1880	3
Subgenus <i>Canthosella</i> Chappuis, 1931	3
<i>Attheyella</i> ( <i>Canthosella</i> ) <i>chocoensis</i> n. sp.	3
<i>Attheyella</i> ( <i>Delachauxiella</i> ) <i>freyi</i> Löffler, 1963	21
Identification key for Colombian canthocamptids	35
Conclusions	36
Acknowledgements	36
References	37

### Abstract

Occasional collections of organisms from water bodies of the Andean Region and the Pacific coast of Colombia yielded two species of canthocamptid harpacticoids belonging to the genus *Attheyella* Brady, 1880. A new species, *Attheyella* (*Canthosella*) *chocoensis* n. sp., inhabiting phytotelmata of the forest at the Pacific Chocó State is described. *Attheyella* (*Delachauxiella*) *freyi* Löffler, 1963, described from Ecuador, was found in a high mountain pond in the southern Colombian Andes, and redescribed here. Both species are illustrated with line drawings and SEM photographs. A special morphological structure was discovered on the antenna of male and female *Attheyella* (*D.*) *freyi*: it is probably present in other species of the subgenus *Delachauxiella* Brehm, 1926, as well. An identification key to Colombian species of the family Canthocamptidae is provided.

**Key words:** taxonomy, Colombia, Pacific forest, phytotelmata, “páramo” region, Andes

### Introduction

Harpacticoid copepods from Colombian fresh waters have been little studied compared to calanoid and cyclopoid copepods (Gaviria 1994; Gaviria & Aranguren 2007). To date, 14 species of harpacticoids belonging to 4 families have been reported. *Nitocra lacustris colombianus* Reid, 1988 was reported within the Ameiridae, *Parastenocaris colombianus* Noodt, 1972, *Parastenocaris kubitzkii* Noodt, 1972 and *Parastenocaris roettgeri* Noodt, 1972 within the Parastenocarididae, *Phyllognathopus viguieri* (Maupas, 1892) within the Phyllognathopodidae, and 9 species within the Canthocamptidae. The family Canthocamptidae is represented by two species of *Attheyella* (*Chappuisiella*) Brehm, 1926, namely *Attheyella* (*Chappuisiella*) *fuhrmanni* (Thiébaud, 1912) (syn. *Attheyella derelicta*