

<http://dx.doi.org/10.11646/zootaxa.3895.3.4>
<http://zoobank.org/urn:lsid:zoobank.org:pub:B317ADE2-F627-4892-A3E6-141C2908B8BC>

A new species of *Meharia* Chrétien, 1915 (Lepidoptera, Cossidae) from the United Arab Emirates, with a world catalogue of the genus

ROMAN V. YAKOVLEV

Altai State University, South Siberian Botanical Garden, Lenina 61, RF-656049, Barnaul, Russia.

National Research Tomsk State University, Laboratory of Biodiversity and Ecology, 36, Lenin Prospekt, 634050 Tomsk, Russia

E-mail: cossus_cossus@mail.ru

Abstract

Meharia breithaupti sp. nov. is described from the United Arab Emirates.

Key words: *Meharia breithaupti*, new species, Arabian Peninsula

Introduction

The genus *Meharia* Chrétien, 1915 constitutes a monotypic subfamily, the Mehariinae, which is highly distinctive within the Cossidae (Yakovlev 2011). It was established for *Meharia incurvariella* Chrétien, 1915. Despite the descriptions of several additional species in the 20th Century, morphological features of the genus were described in details only recently (Komarov & Zolotuhin 2005; Saldaitis & Ivinskis 2010; Borth *et al.* 2011; Yakovlev 2011). Currently, twelve species and one subspecies are included in the genus. Half of them were described very recently (Saldaitis & Ivinskis 2010; Borth *et al.* 2011; Ivinskis *et al.* 2011; Yakovlev 2011; Yakovlev *et al.* 2013). They occur from the southern Volga River area through the Near East and the Sahara, south to Kenya and Malawi (Figs 1–13). Everywhere representatives of *Meharia* inhabit extremely arid biotopes (Yakovlev & Dubatolov 2013). During a survey of the Lepidoptera of Abu Dhabi, UAE, by Roland Breithaupt, a new species of *Meharia* was discovered, and its description is given below.

List of abbreviations

BMNH	Natural History Museum (London, G.B.);
MNHN	Muséum National d'Histoire Naturelle (Paris, France);
MNHW	Naturhistorisches Museum (Wien, Austria);
MWM	Museum of Thomas Witt (Munich, Germany);
ZSSM	Zoologische Staatssammlung der Bayerischen Staaten (Munich, Germany).

Description of a new species

Meharia breithaupti Yakovlev, sp. nov.

(Figs 3, 18–19, 43–49)

Type. Holotype: male, UAE, Ras Al Khaimah, Esfai garbage dump, 25° 06' 19" N / 56°07' 52 E, 488 m, 1.12.2013, et light, R. Breithaupt leg. Deposited in the Zoological Institute, St. Petersburg, Russia. **Paratype:** female, UAE, Ras Al Khaimah, Wadi Sifuni, 25° 10' 09" N/ 56° 04' 32" E, 290 m, 6.12.2012, et light, R. Breithaupt leg. Deposited in private collection of Thomas Sobczyk (Hoyerswerda, Germany).

***Meharia semilactea* (Warren et Rothschild, 1905) (Figs 10, 35–37)**

Alavona semilactea Warren et Rothschild, 1905: 32.

Type locality: Nakheila, R. Atbara [NW Sudan].

Type material (holotype) in BMNH, examined.

Distribution: Israel, Jordan, Saudi Arabia, Oman, Yemen, United Arabian Emirates, Egypt (Sinai peninsula), Morocco, N. Sudan, Mauritania (Rungs 1979; Wiltshire 1982, 1990; Legrain & Wiltshire 1998; Hacker 1999; Hacker *et al.* 1999, 2001; Yakovlev & Saldaitis 2008; Yakovlev & Breithaupt 2013).

***Meharia tancredi* Sutton, 1963 (Figs 11, 38)**

Meharia tancredi Sutton, 1963: 365.

Type locality: Meyan Kaleh peninsula, N. Iran

Type material (holotype) in BMNH, examined.

Distribution: N. Iran.

***Meharia tanganyikae* Bradley, 1952 (Figs 12, 39–40)**

Meharia tanganyikae Bradley, 1952: 242.

Type locality: Tanganyika, Ngaruka.

Type material (holotype) in BMNH, examined.

Distribution: Tanzania.

***Meharia yakovlevi* Saldaitis et Ivinskis, 2010 (Figs 13, 41–42)**

Meharia yakovlevi Saldaitis et Ivinskis, 2010: 379.

Type locality: Socotra Island, N hills near Hadibu.

Type material (holotype) in MWM, examined.

Distribution: Yemen (Socotra Isl.).

Acknowledgements

The author is grateful to R. Breithaupt (Abu Dhabi, UAE) for making available the material used in this study, and Th. Sobczyk (Hoyerswerda, Germany) for the photos of the paratype (habitus and genitalia). I am indebted to the BMNH Council of Trustees for kindly granting me the permission to publish images of type specimens preserved in the Natural History Museum.

References

- Borth, R., Ivinskis, P., Saldaitis, A. & Yakovlev, R. (2011) Cossidae of the Socotra Archipelago (Yemen). *ZooKeys*, 122, 45–69.
<http://dx.doi.org/10.3897/zookeys.122.1213>
- Bradley, J.D. (1952) Two new species of *Meharia* (Lep., Cossidae). *The Entomologist*, 85, 241–244.
- Chrétien, P. (1915) Contribution à la connaissance des Lépidoptères du Nord de l'Afrique. *Annales de la Société entomologique de France*, 84 (3), 289–374.
- Daniel, F. (1965) Österreichische Entomologische Iran-Afghanistan-Expeditionen Beiträge zur Lepidopterenfauna, Teil 4. Weitere Beiträge zur Bombyces et Sphinges Fauna. *Zeitschrift der Wiener Entomologischen Gesellschaft*, 50, 121–145.
- Hacker, H. (1999) Systematic List of the Lepidoptera of the Arabian Peninsula with a survey of the spread with special reference to the fauna of Yemen. *Esperiana*, 7, 15–237.
- Hacker, H.H., Schreier, H.-P. & Bischof, A. (1999) Lepidoptera of Yemen Arab Republic, collected by A. Bischof, H. Hacker and H.-P. Schreier in autumn 1996 and B. Müller in summer 1987. *Esperiana*, 7, 267–282.
- Hacker, H.H., Aulombard, F., Bischof, A., Bittermann, J., Fibiger, M. & Schreier, H.-P. (2001) Lepidoptera of Yemen Arab Republic, collected in 1996, 1998 and 2000. *Esperiana*, 8, 597–600.
- Komarov, D. & Zolotuhin, V. (2005) A new species of *Meharia* Chrétien, 1915 (Cossidae) from the Lower Volga Region. *Nota lepidopterologica*, 28 (1), 49–54.
- Legrain, A. & E.P. Wiltshire (1998) Provisional Checklist of the Macro-heterocera of UAE. *Tribulus*, 8 (2), 5–8.
- Rungs, Ch. (1942) III. Notes de lépidoptérologie Marocaine (XI). Additions à la faune du Maroc: Lepidoptères des régions

- Sahariennes. *Bulletin de la Societe des Sciences Naturelles du Maroc*, 22, 172–177.
- Rungs, Ch.E.E. (1979) Catalogue raisonné des Lépidoptères du Maroc. Inventaire Faunistique et observations écologiques 1. *Travaux de l'Institute Scientifique, Sér. Zoologie, Rabat*, 39, 1–244.
- Saldaitis, A. & Ivinskis, P. (2010) *Meharia yakovlevi*, a new species (Lepidoptera, Cossidae) from Yemen. *Esperiana*, 15, 379–381.
- Sutton, S.L. (1963) South Caspian Insect fauna 1961. *Annals and Magazine of Natural History*, 6 (13 Ser.), 23–370.
- Warren W. & Rothschild, Ch. (1905) Lepidoptera from the Sudan. *Novitates Zoologicae*, 12, 21–28.
- Wiltshire, E.P. (1946) Middle East Lepidoptera. V. A new genus, a new species, and two new races from Iran, with taxonomic notes on other Persian Heterocera. *Proceedings of the Royal Entomological Society of London Series B*, 15, 118–128.
- Wiltshire, E.P. (1982) Insects of Saudi Arabia. Lepidoptera: Fam. Cossidae, Zygaenidae, Sesiidae, Lasiocampidae, Bombycidae, Sphingidae, Thaumetopoeidae, Thyretidae, Notodontidae, Geometridae, Lymantriidae, Noctuidae, Ctenuchidae (Part 2). *Fauna of Saudi Arabia*, 4, 271–331.
- Wiltshire, E.P. (1990) An Illustrated Annotated Catalogue of the Macro-Heterocera of Saudi Arabia. *Fauna of Saudi Arabia*, 11, 91–250.
- Yakovlev, R.V. (2005) New records of distribution and systematic of Cossidae (Lepidoptera) of Europe and adjacent territories. *Eversmannia*, 3/4, 18–27. [in Russian]
- Yakovlev, R.V. (2011) Catalogue of the Family Cossidae of the Old World. *Neue Entomologische Nachrichten*, 66, 1–129.
- Yakovlev, R.V. & Breithaupt, R. (2013) The first data on the Cossidae fauna (Lepidoptera) from the Eastern Emirates of the UAE. *Amurian zoological journal*, 5 (2), 1–176. [in Russian]
- Yakovlev, R.V. & Dubatolov, V.V. (2013) Distribution of Carpenter-Moths (Lepidoptera, Cossidae) in the Palaearctic Deserts. *Entomological Review*, 93 (8), 991–1004.
<http://dx.doi.org/10.1134/S0013873813080071>
- Yakovlev, R., Ivinskis, P., Rimsaite, J. & Saldaitis, A. (2013) Description of two new species of *Meharia* Chrétien, 1915 (Lepidoptera: Cossidae) from East Africa. *Zootaxa*, 3635 (5), 587–590.
<http://dx.doi.org/10.11646/zootaxa.3635.5.9>
- Yakovlev, R.V. & Saldaitis, A. (2008) New species of Palaearctic and Oriental Cossidae (Lepidoptera). III. New species of genus *Meharia* Chretien, 1915 from Morocco. *Eversmannia*, 15–16, 49–52. [in Russian]