

## A review of the genus *Lamellipalpodes* Maulik (Coleoptera: Lampyridae)

MILADA BOCAKOVA<sup>1</sup>, LADISLAV BOCAK<sup>2</sup>, MATTHEW L. GIMMEL<sup>1</sup> & TEREZA FRIEDLOVA<sup>1</sup>

<sup>1</sup>Department of Biology, Faculty of Education, Palacky University, Purkrabska 2, CZ-77140 Olomouc, Czech Republic.  
E-mails: milada.bocakova@upol.cz, phalacrid@gmail.com, and tera.fri@seznam.cz

<sup>2</sup>Department of Zoology, Faculty of Science, Palacky University, 17. listopadu 50, CZ-77146 Olomouc, Czech Republic.  
E-mail: ladislav.bocak@upol.cz

### Abstract

The Ototretinae genus *Lamellipalpodes* Maulik, 1921 is reviewed. Four species are proposed as new to science, *L. bajhangensis* Bocakova sp. nov., *L. holzschuhi* Bocakova sp. nov. and *L. godawarensis* Bocakova sp. nov., all from Nepal, and *L. yunnanensis* Bocakova sp. nov. from China (Yunnan) and northern Laos. Six species are redescribed. Illustrations of diagnostic characters are included, with a checklist and a key to *Lamellipalpodes* species. The genus is newly recorded from China and Laos.

**Key words:** taxonomy, Ototretinae, checklist, key, new species, Oriental Region

### Introduction

*Lamellipalpodes* Maulik, 1921 is a small genus of Ototretinae fireflies distributed from southern India to the foothills of the Himalayas. The genus was originally monotypic (Maulik 1921). Later, two new species from Nepal were described by Wittmer (1995) and another two species from Thailand and Myanmar (Kawashima 2007). Brancucci and Geiser (2009) recognized another species, *L. lineatus* (Pic, 1921), originally placed in the genus *Eugeusis* Westwood, 1853.

*Lamellipalpodes* is one of the more attractive ototretine representatives. Conspicuously enlarged terminal palpomeres of maxillary and labial palpi can be 10x longer than penultimate palpomere. This character is shared also by *Eugeusis*, *Hyperstoma* Wittmer, 1979, and *Lamellipalpus* Maulik, 1921. Similarly as in several other ototretine genera only males of the genus *Lamellipalpodes* are known, and females are assumed to be brachypterous, or apterous, maybe even larviform. Phylogenetic relationships of *Lamellipalpodes* are not known, although the genus is assumed to be closely related to *Lamellipalpus* (Brancucci & Geiser 2009), whose members also have considerably larger mandibles, while those of *Lamellipalpodes* are small.

Due to the external similarities the genus was placed in Drilidae (Wittmer 1944). Drilidae were found to be a polyphyletic assemblage by (Crowson (1972) who considered *Lamellipalpus* a member of predominantly non-luminous fireflies of the subfamily Ototretinae. The placement of closely related *Lamellipalpodes* in the subfamily Ototretinae was supported by Brancucci & Geiser (2009), and recently by Janisova and Bocakova (2013) who clarified the concept of the Ototretinae. Recently, Branham (2010) revived his previous hypothesis and considered Ototretinae genera (including *Lamellipalpodes*) as incertae sedis within elateroids, although the placement of Ototretinae within Lampyridae was robustly supported by analyses of DNA sequences (Bocakova *et al.* 2007, Sagegami-Oba *et al.* 2007, Stanger Hall 2007). Lately, the lampyrid placement of the Ototretinae including related *Lamellipalpus* was corroborated by multigene analyses of huge elateroid dataset (Kundrata *et al.* 2014).

Here, a taxonomic revision of *Lamellipalpodes* is provided and four new species from Nepal, India, Laos, and China are proposed. Descriptions, illustrations, a checklist, and a key to species are provided.

## Acknowledgements

We would like to thank A. Taghavian (MHNP), M. Barclay (BMNH), and E. Spraecher (NHMB) for providing the material in their charge. The type material in the Natural History Museum in London was examined under EU-funded Synthesys project (GB-TAF-1982). This study has been supported by grants PdF\_2014019 (TF) from Palacky University Olomouc (Czech Republic), CZ.1.07/2.3.00/20.0004 (MG) and CZ.1.07/2.3.00/20.0166 (MB, LB) from European Social Fund and the Ministry of Education of the Czech Republic.

## References

- Bocakova, M., Bocak, L., Hunt, T., Teraväinen, M. & Vogler, A.P. (2007) Molecular phylogenetics of Elateriformia (Coleoptera): evolution of bioluminescence and neoteny. *Cladistics*, 23, 477–496.  
<http://dx.doi.org/10.1111/j.1096-0031.2007.00164.x>
- Branham, M. (2010) Lampyridae Latreille, 1817. In: Leschen, R.A., Beutel, R.G. & Lawrence, J.F. (Eds.), *Handbook of Zoology. Part 38. Vol. 2. Morphology and Systematics (Elateroidea, Bostrichiformia, Cucujoidea)*. Walter de Gruyter, Berlin, New York, pp. 141–149.
- Brancucci, M. & Geiser, M. (2009) A revision of the genus Lamellipalpus Maulik, 1921 (Coleoptera, Lampyridae) *Zootaxa*, 2080, 1–20.
- Crowson, R.A. (1972) A review of the classification of Cantharoidea (Coleoptera), with the definition of two new families, Cheoglossidae and Omethidae. *Revista de la Universidad de Madrid*, 21 (82), 35–77.
- Janisova, K. & Bocakova, M. (2013) Revision of the subfamily Ototretinae (Coleoptera: Lampyridae), *Zoologischer Anzeiger*, 252 (1), 1–19.  
<http://dx.doi.org/10.1016/j.jcz.2012.01.001>
- Kawashima, I. (2007) Two new species of the genus Lamellipalpodes (Coleoptera: Lampyridae) from Indochina, Southeast Asia. *Elytra*, 35, 119–128.
- Kundrata, R., Bocakova, M. & Bocak, L. (2014) The comprehensive phylogeny of the superfamily Elateroidea (Coleoptera: Elateriformia), *Molecular Phylogenetics and Evolution*, 76, 162–171.  
<http://dx.doi.org/10.1016/j.ympev.2014.03.012>
- Maulik, S. (1921) New Indian drilid beetles. *Proceedings of Zoological Society London*, 1921, 579–586.
- Wittmer, W. (1944) Catalogue des Drilidae E. Oliv. (Coleoptera-Malacodermata). *Revista de la Sociedad Entomológica Argentina*, 12, 203–221.
- Wittmer, W. (1995) Lampyridae und Omethidae aus dem indo-malaiischen Faunengebiet (Coleoptera). *Mitteilungen der Entomologischen Gesellschaft Basel*, 45 (3), 106–117.
- Sagegami-Oba, R., Takahashi, N. & Oba, Y. (2007) The evolutionary process of bioluminescence and aposematism in cantharoid beetles (Coleoptera: Elateroidea) inferred by the analysis of 18S ribosomal DNA. *Gene*, 400, 104–113.
- Stanger-Hall, K.F., Lloyd, J.E. & Hillis, D.M. (2007) Phylogeny of North American fireflies (Coleoptera: Lampyridae): Implications for the evolution of light signals. *Molecular phylogenetics and evolution*, 45 (1), 33–49.