



<http://dx.doi.org/10.11646/zootaxa.4028.1.7>

<http://zoobank.org/urn:lsid:zoobank.org:pub:D87155DD-C49B-4198-B3C7-5C9A8A8C5037>

New species and additional records of the subgenus *Termidonia* Motschulsky of the genus *Zyras* Stephens in China (Coleoptera, Staphylinidae, Aleocharinae)

ZHU-QI YAN & LI-ZHEN LI¹

Department of Biology, College of Life and Environmental Sciences, Shanghai Normal University, 100 Guilin Road, Xuhui District, Shanghai 200234, P. R. China

¹Corresponding author. E-mail: lomechusini@163.com

Abstract. Two new species are described from Yunnan: *Zyras* (*Termidonia*) *baihuamontis* sp. nov., and *Z. (T.) nabanensis* sp. nov. Additional records of two species of subgenus *Termidonia* Motschulsky, 1860 are reported. Adults of the four species are illustrated, and identification key is provided for Chinese species.

Key words: Aleocharinae, Lomechusini, *Zyras*, *Termidonia*, new species, new record, China.

Introduction

The subgenus *Termidonia* Motschulsky, 1860 currently contains 40 species (Hlaváč *et al.* 2011, Pace 2012), among which seven species, *Zyras* (*Termidonia*) *longwangmontis* Pace, 1986, *Z. (T.) abbreviatus* Fenyés, 1914, *Z. (T.) improltrans* Pace, 2010, *Z. (T.) sexcuspidatus* Assing, 2009, *Z. (T.) viti* Assing, 2009, *Z. (T.) diaoluomontis* Pace, 2012, *Z. (T.) fellowesi* Pace, 2012 are known in China.

Examination of recently collected Chinese material of *Termidonia* revealed two new species from Yunnan, and new locality data for *Z. (T.) fellowesi* from Guangxi and *Z. (T.) longwangmontis* from Shanghai, Jiangsu and Guizhou.

Material and methods

All material treated in the present study is deposited in the Insect Collection of the Shanghai Normal University, Shanghai, China (SNUC).

The following abbreviations are used in the text: BL—body length, from the anterior margin of the head to the posterior margin of the abdominal tergite VIII; FBL—forebody length, from the clypeal anterior margin to the posterior margin of elytra; PL—length of the pronotum along the midline; HW—width of the head across the eyes; PW—maximum width of the pronotum; EL—length of elytra from the apex of the scutellum to the elytral posterior margin; EW—maximum width of the elytra.

Taxonomy

Zyras (*Termidonia*) *baihuamontis* Z.-Q. Yan and L.-Z. Li, new species

(Fig. 1)

Type material. Holotype: China: 1 ♂, labeled 'CHINA: Yunnan Prov., Baoshan City (保山市), Mangkuan Town (芒宽乡), Baihualing (百花岭), 25.16.39 N, 98.47.55 E, alt. 1200 m, 22.iv.2013, Dai, Peng & Song leg.'; HOLOTYPE [red], *Zyras baihuamontis*, Yan & Li det. 2015, (SNUC); **Paratype:** 1 ♀, same label as above-mentioned, PARATYPE [yellow], *Zyras baihuamontis*, Yan & Li det. 2015, (SNUC).

Description. Body length: 9.02–9.22 mm. Coloration: head, elytra, blackish; pronotum darker reddish-brown; abdomen blackish with the posterior margins of the segments reddish-brown; legs and antennae reddish to reddish-brown.