Megataxa 003 (1): 001–383
<https://www.mapress.com/j/mt/>
Copyright © 2020 Magnolia Press

ISSN 2703-3082 (print edition)
MEGATAXA
ISSN 2703-3090 (online edition)

<https://doi.org/10.11646/megataxa.3.1.1>

<http://zoobank.org/urn:lsid:zoobank.org:pub:7A95A1DD-0372-4FAC-BA3B-1896386BC710>

MEGATAXA

3

Grenadiers (Teleostei: Gadiformes: Macrouridae) of Japan and adjacent waters, a taxonomic monograph

NAOHIDE NAKAYAMA

*Department of Marine Biology, School of Marine Science and Technology, Tokai University,
3-20-1 Orido, Shimizu, Shizuoka 424-8610, Japan*

✉ naonakayama@tsc.u-tokai.ac.jp or ✉ gadiformes@gmail.com

🌐 <https://orcid.org/0000-0001-9556-1537>

Magnolia Press
Auckland, New Zealand

NAOHIDE NAKAYAMA

Grenadiers (Teleostei: Gadiformes: Macrouridae) of Japan and adjacent waters, a taxonomic monograph
(*Megataxa* 3)

383 pp.; 30 cm.

5 November 2020

ISBN 978-1-77688-084-3 (Paperback)

ISBN 978-1-77688-085-0 (Online edition)

FIRST PUBLISHED IN 2020 BY
Magnolia Press

P.O. Box 41-383
Auckland 1041
New Zealand

e-mail: magnolia@mapress.com
<https://www.mapress.com/j/mt>

© 2020 Magnolia Press

ISSN 2703-3082 (Print edition)

ISSN 2703-3090 (Online edition)

Table of contents

Abstract.....	4
Introduction.....	5
Historical perspective.....	7
Materials and methods.....	9
Taxonomy.....	15
Family Macrouridae Bonaparte, 1831 (sensu Iwamoto <i>et al.</i> 2015).....	15
Key to genera of grenadiers from Japan and adjacent waters..	15
Genus <i>Asthenomacrus</i> Sazonov & Shcherbachev, 1982	16
<i>Asthenomacrus victoris</i> Sazonov & Shcherbachev, 1982	17
Genus <i>Cetonurus</i> Günther, 1887.....	23
<i>Cetonurus globiceps</i> (Vaillant <i>in</i> Filhol, 1884).....	23
Genus <i>Coelorinchus</i> Giorna, 1809.....	25
Key to species of <i>Coelorinchus</i> from Japan and adjacent waters	26
<i>Coelorinchus anatirostris</i> Jordan & Gilbert <i>in</i> Jordan & Starks, 1904.....	27
<i>Coelorinchus brevirostris</i> Okamura, 1984.....	38
<i>Coelorinchus cingulatus</i> Gilbert & Hubbs, 1920 ..	42
<i>Coelorinchus divergens</i> Okamura & Yatou, 1984	45
<i>Coelorinchus formosanus</i> Okamura, 1963	48
<i>Coelorinchus fuscigulus</i> Iwamoto, Ho & Shao, 2009.....	51
<i>Coelorinchus gilberti</i> Jordan & Hubbs, 1925 ...	55
<i>Coelorinchus hexafasciatus</i> Okamura, 1982....	60
<i>Coelorinchus hige</i> Matsubara, 1943	65
<i>Coelorinchus hubbsi</i> Matsubara, 1936	73
<i>Coelorinchus japonicus</i> (Temminck & Schlegel, 1846)	76
<i>Coelorinchus jordani</i> Smith & Pope, 1906.....	84
<i>Coelorinchus kamoharai</i> Matsubara, 1943.....	87
<i>Coelorinchus kishinouyei</i> Jordan & Snyder, 1900	91
<i>Coelorinchus lanceolatus</i> sp. nov.	94
<i>Coelorinchus leptorhinus</i> Chiou, Shao & Iwamoto, 2004.....	96
<i>Coelorinchus longissimus</i> Matsubara, 1943	98
<i>Coelorinchus macrochir</i> (Günther, 1877).....	100
<i>Coelorinchus matsubarae</i> Okamura, 1982.....	105
<i>Coelorinchus multispinulosus</i> Katayama, 1942...	108
<i>Coelorinchus nox</i> sp. nov.	113
<i>Coelorinchus parallelus</i> (Günther, 1877)	119
<i>Coelorinchus sheni</i> Chiou, Shao & Iwamoto, 2004.....	126
<i>Coelorinchus smithi</i> Gilbert & Hubbs, 1920 ..	128
<i>Coelorinchus tokiensis</i> (Steindachner & Döderlein, 1887)	135

Comments on “ <i>Coelorinchus (Quincuncia) logispinosus</i> ”	139
Genus <i>Coryphaenoides</i> Gunnerus, 1765.....	139
Key to species of <i>Coryphaenoides</i> from Japan and adjacent waters	140
<i>Coryphaenoides acrolepis</i> (Bean, 1884)	141
<i>Coryphaenoides altipinnis</i> Günther, 1877	147
<i>Coryphaenoides armatus</i> (Hector, 1875).....	150
<i>Coryphaenoides asper</i> Günther, 1877	154
<i>Coryphaenoides cinereus</i> (Gilbert, 1896).....	157
<i>Coryphaenoides filifer</i> (Gilbert, 1896).....	164
<i>Coryphaenoides leptolepis</i> Günther, 1877	167
<i>Coryphaenoides longifilis</i> Günther, 1877	179
<i>Coryphaenoides marginatus</i> Steindachner & Döderlein, 1887.....	184
<i>Coryphaenoides microps</i> (Smith & Radcliffe <i>in</i> Radcliffe, 1912).....	190
<i>Coryphaenoides nasutus</i> Günther, 1877	192
<i>Coryphaenoides pectoralis</i> (Gilbert, 1891).....	196
<i>Coryphaenoides rudis</i> Günther, 1878.....	200
<i>Coryphaenoides soyoyae</i> Nakayama & Endo, 2016	203
<i>Coryphaenoides yaquinae</i> Iwamoto & Stein, 1974	205
Genus <i>Hymenocephalus</i> Giglioli, 1884.....	207
Key to species of <i>Hymenocephalus</i> from Japan and adjacent waters	207
<i>Hymenocephalus hachijoensis</i> Okamura, 1970 ...	208
<i>Hymenocephalus lethonemus</i> Jordan & Gilbert <i>in</i> Jordan & Starks, 1904	210
<i>Hymenocephalus longibarbis</i> Günther, 1887... ..	213
<i>Hymenocephalus papyraceus</i> Jordan & Gilbert <i>in</i> Jordan & Starks, 1904.....	215
<i>Hymenocephalus striatissimus</i> Jordan & Gilbert <i>in</i> Jordan & Starks, 1904.....	218
<i>Hymenocephalus yamasakiorum</i> Nakayama, Endo & Schwarzhans, 2015	221
Genus <i>Hymenogadus</i> Gilbert & Hubbs, 1920	222
<i>Hymenogadus gracilis</i> (Gilbert & Hubbs, 1920) ..	223
Genus <i>Kumba</i> Marshall, 1973.....	224
<i>Kumba japonica</i> (Matsubara, 1943).....	225
Genus <i>Kuronezumia</i> Iwamoto, 1974	227
Key to species of <i>Kuronezumia</i> from Japan and adjacent waters	228
<i>Kuronezumia dara</i> (Gilbert & Hubbs, 1916) ..	228
<i>Kuronezumia endoi</i> sp. nov.	231
Genus <i>Lucigadus</i> Gilbert & Hubbs, 1920.....	237
<i>Lucigadus nigromarginatus</i> (Smith & Radcliffe <i>in</i> Radcliffe, 1912).....	237
Genus <i>Malacocephalus</i> Günther, 1862.....	238
<i>Malacocephalus nipponensis</i> Gilbert & Hubbs, 1916.....	239
Genus <i>Nezumia</i> Jordan <i>in</i> Jordan & Starks, 1904	243
Key to species of <i>Nezumia</i> from Japan and adjacent	

waters	243
<i>Nezumia condylura</i> Jordan & Gilbert <i>in</i> Jordan & Starks, 1904.....	243
<i>Nezumia kamoharai</i> Okamura, 1970.....	248
<i>Nezumia proxima</i> (Smith & Radcliffe <i>in</i> Radcliffe, 1912).....	250
<i>Nezumia rara</i> sp. nov.....	256
<i>Nezumia shinoharai</i> Nakayama & Endo, 2012	260
<i>Nezumia spinosa</i> (Gilbert & Hubbs, 1916).....	262
<i>Nezumia tomiyamai</i> (Okamura, 1963).....	264
Comments on Japanese record of <i>Nezumia burragei</i>	266
Comments on Japanese record of <i>Nezumia propinqua</i>	268
Genus <i>Odontomacrurus</i> Norman, 1939.....	268
<i>Odontomacrurus murrayi</i> Norman, 1939.....	268
Genus <i>Pseudocetonurus</i> Sazonov & Shcherbachev, 1982 .	269
<i>Pseudocetonurus</i> sp. cf. <i>septifer</i> Sazonov & Shcherbachev, 1982	270
Genus <i>Pseudonezumia</i> Okamura, 1970.....	270
Key to species of <i>Pseudonezumia</i> from Japan and adjacent waters	272
<i>Pseudonezumia cetonuropsis</i> (Gilbert & Hubbs, 1916)	272
<i>Pseudonezumia japonica</i> Okamura, 1970.....	275
Genus <i>Sphagemacrurus</i> Fowler, 1925	276
<i>Sphagemacrurus pumiliceps</i> (Alcock, 1894)	277
Genus <i>Spicomacrurus</i> Okamura, 1970.....	277
<i>Spicomacrurus kuronumai</i> (Kamohara, 1938).....	278
Genus <i>Trachonurus</i> Günther, 1887	281
<i>Trachonurus villosus</i> (Günther, 1877).....	281
Genus <i>Ventrifossa</i> Gilbert & Hubbs, 1920.....	284
Key to species of <i>Ventrifossa</i> from Japan and adjacent waters	285
<i>Ventrifossa garmani</i> (Jordan & Gilbert <i>in</i> Jordan & Starks, 1904)	285
<i>Ventrifossa johnboborum</i> Iwamoto, 1982.....	289
<i>Ventrifossa macroptera</i> Okamura, 1982	298
<i>Ventrifossa misakia</i> (Jordan & Gilbert <i>in</i> Jordan & Starks, 1904)	301
<i>Ventrifossa nigrodorsalis</i> Gilbert & Hubbs, 1920	306
<i>Ventrifossa rhipidodorsalis</i> Okamura, 1984 ...	307
<i>Ventrifossa saikaiensis</i> Okamura, 1984	310
Funding	312
Acknowledgments.....	312
References.....	313
APPENDIX 1. Additional material examined	330
APPENDIX 2. Comparative material examined	364
APPENDIX 3. Maps of the northwestern Pacific around Japan showing the distribution of each species of grenadier occurring in the area	371

Abstract

The taxonomy of the gadiform fish family Macrouridae (sensu stricto) in the northwestern Pacific off Japan and adjacent waters is critically reviewed on the basis of 7846 specimens. A total of 76 species belonging to 18 genera is recognized, including four new species of the genera *Coelorinchus* (2 species), *Kuronezumia* (1), and *Nezumia* (1). *Coelorinchus lanceolatus* **sp. nov.** superficially resembles *Coe. anatirostris* Jordan & Gilbert *in* Jordan & Starks, 1904, but differs most notably from that species in having a distinctly longer snout (107% of the postrostral length vs. 60–88%). *Coelorinchus nox* **sp. nov.** is closely similar to *Coe. smithi* Gilbert & Hubbs, 1920, but readily differs from the latter in that the occipital scales are covered with short, erect, needle-like spinules in widely divergent, comb-like rows (vs. moderately reclined, keel-like to knife-like spinules in saw-toothed rows). *Kuronezumia endoi* **sp. nov.** is a distinctive species among the genus, and is clearly diagnosed from other congeners in having the highest number of pelvic-fin rays (15 vs. ≤ 14). *Nezumia rara* **sp. nov.** closely resembles *N. tomiyamai* (Okamura, 1963), but they can be distinguished from each other by the combination of a number of morphometric characters, including the orbit-preopercle distance, interorbital width, pelvic-fin length, and length of gill slit. New and reconfirmed synonymies include: *Coe. abbreviatus* Chu & Lo *in* Chu, Chan & Chen, 1963 and *Coe. intermedius* Chu & Lo *in* Chu, Chan & Chen, 1963 with *Coe. multispinulosus* Katayama, 1942; *Coe. productus* Gilbert & Hubbs, 1916 with *Coe. anatirostris*; *Coe. asteroides* Okamura, 1963 with *Coe. hige* Matsubara, 1943; *Coe. sparsilepis* Okamura & Yatou, 1984 with *Coe. parallelus* (Günther, 1877); *Coryphaenoides filamentosus* Okamura, 1970 with *Cor. cinereus* (Gilbert, 1896); *Cor. liocephalus* (Günther, 1887) with *Cor. leptolepis* Günther, 1877; *Cor. spinulosus* (Gilbert & Burke, 1912) with *Cor. acrolepis* (Bean, 1884); and *Ventrifossa fusca* Okamura, 1982 with *V. misakia* (Jordan & Gilbert *in* Jordan & Starks, 1904). Lectotypes are designated for *Coe. parallelus*, *Cor. asper* Günther, 1877, *Cor. liocephalus*, and *Cor. nasutus* Günther, 1877. Previous records of *N. burragei* (Gilbert, 1905) and *N. propinqua* (Gilbert & Cramer, 1897) from Japan represent misidentifications, and these species are eliminated from the list of Japanese grenadiers. *Coryphaenoides armatus* (Hector, 1875) and *Cor. leptolepis* are recorded for the first time from the hadal zone. All species are illustrated, with full descriptions for 14 species. Dichotomous keys to genera and species are also provided. The distribution of each species in the study area is revised based on the specimens examined here, with re-identification of voucher specimens of most previous geographical records.

Key words: Actinopterygii, deep-sea fish, taxonomic review, new species, synonymy, distribution, Japanese Archipelago, northwestern Pacific

FIGURE 1. Map of the northwestern Pacific around Japan showing four tectonic plates and their subduction zones, and major seas, bays, “nada”, ridges, troughs, and trenches in the area.

Introduction

Grenadiers of the family Macrouridae, also known as rattails or whiptails, are among the most dominant deep-sea demersal fishes in terms of species diversity and abundance (Marshall 1979; Gage & Tyler 1992; Iwamoto & Merrett 1997; Merrett & Headrich 1997; Iwamoto & Graham 2001; Shao *et al.* 2008b; Iwamoto *et al.* 2015). This study follows Iwamoto *et al.* (2015) in regarding the three taxa previously considered to be subfamilies of Macrouridae as distinct families, viz., Bathygadidae, Macrouroididae, and Trachyrincidae. Comprising at least 370 valid species in about 27 genera (author’s pers. observ.), the Macrouridae is by far the largest family among the order Gadiformes (596 spp. in 83 gen., fide Eschmeyer & Fong 2011), while many new species still continue to be described (Nelson 2006; Iwamoto *et al.* 2009, 2011; McMillan & Iwamoto 2009, 2014; Nakayama & Endo 2012, 2016a, 2017; Schwarzhans 2014; Nakayama *et al.* 2015a; Nelson *et al.* 2016). Grenadiers are widely distributed in all major oceans, with the highest diversity

in tropical and subtropical areas (Marshall 1965; Marshall & Iwamoto 1973; Iwamoto 1990; Iwamoto *et al.* 2015). Most species are benthopelagic and closely associated with the sea floor, but a few species are midwater dwellers occurring at mesopelagic and bathypelagic depths of the open oceans (Marshall 1965; Marshall & Iwamoto 1973; Iwamoto 1990, 2008b). The vertical distribution of demersal grenadiers is primarily confined to bathyal depths between 200 and 2000 m, with only a few species occurring in much deeper habitats or on the continental shelves (Marshall & Iwamoto 1973; Marshall 1979; Iwamoto 1990; this study). Grenadiers also constitute a large portion of discarded bycatch in slope fisheries (Allain *et al.* 2003; Drazen 2008), while some species are commercially exploited for food (e.g., *Coryphaenoides rupestris* Gunnerus, 1765 and species of *Macrourus* Bloch, 1786).

The Japanese Archipelago is an extensive chain of more than 4000 islands located in the northwestern Pacific off the east coast of Asia. This island arc is developed along margins of four tectonic plates (i.e., Pacific, Eurasia,

FIGURE 2. Map of the northwestern Pacific around Japan showing major currents and straits in the area (modified from Ministry of the Environment 2011: fig. 2).

Okhotsk, and Philippine plates), and is surrounded by four major seas (i.e., Pacific, East China Sea, Okhotsk Sea, and Sea of Japan; Fig. 1). Japan lies in the mid-latitude region, affected by both warm and cold ocean currents (i.e., Kuroshio, Tsushima Current, Oyashio, and Liman Current; Fig. 2). Although Japan is a relatively small country with limited land space ($3.78 \times 10^5 \text{ km}^2$), in terms of marine area, it has the world's sixth largest Exclusive Economic Zone (EEZ; $4.05 \times 10^6 \text{ km}^2$; Fig. 3). Japan's EEZ extends latitudinally for about 3000 km between 17°N and 48°N, in which six marine climatic regimes are recognized (Nishimura 1992; Senou *et al.* 2006; Fujikura *et al.* 2010), viz., subboreal, cool-temperate, middle-temperate, warm-temperate, subtropical, and tropical zones (Fig. 3). These topographical, geological, and physical features form and provide rich habitats for marine fishes in Japanese waters (Matsuura & Kimura 2007; Fujikura *et al.* 2010).

Over the last 120 years, grenadiers of Japan and adjacent waters have been extensively surveyed by many ichthyologists (see the Historical perspective). Consequently, our knowledge of Japanese grenadiers is

relatively rich compared with other regions, with three outstanding reviews or monographs on their taxonomy, viz., Jordan & Gilbert (*in* Jordan & Starks 1904), Gilbert & Hubbs (1916), and Okamura (1970a). However, despite these reviews there remained some ambiguous species requiring further investigation, and several problems regarding the synonymies of nominal taxa made the identification of common species problematic. Some of these issues have been recently resolved (Nakayama & Endo 2012, 2016a; Nakayama *et al.* 2015a, 2015b, 2015c, 2018), but there is still a necessity to provide a comprehensive review of Japanese grenadiers for further development of systematic and associated fields of science. Fortunately, thousands of specimens of Japanese grenadiers have been accumulated in the world's museums as a result of nearly 200 years of collecting efforts. These specimens came from numerous scientific cruises and field trips to local fish markets (see Historical perspective below), and their sampling sites cover most of the continental shelves and slopes along the Japanese Archipelago (Fig. 4). This huge collection provides a

FIGURE 3. Map of the northwestern Pacific around Japan showing marine climatic regimes in the area (after Nishimura 1992: fig. 2) and Japan's Exclusive Economic Zone.

sufficient inventory to elucidate the geographic range of each grenadier species in the northwestern Pacific around Japan. The purposes of this study are to provide a taxonomic review of grenadiers of Japan and adjacent waters, and to clarify the distributional pattern of each species in the study area. A detailed analysis of the zoogeography of Japanese grenadiers will be published separately.

Historical perspective

In the Pisces volume of *Fauna Japonica*, Coenraad J. Temminck (1778–1858) and Hermann Schlegel (1804–1884) described *Macrourus japonicus* (= *Coelorinchus japonicus*) based on specimens collected from the East China Sea off Nagasaki (Temminck & Schlegel 1846). These specimens were collected by Philipp F. von Siebold (1796–1866), a German physician and naturalist, and his assistant, Heinrich Bürger (1804–1854), during 1823–1830 (Boeseman 1947, 1997; Yamaguchi & Machida 2003).

As far as we know, this is the first scientific treatment of grenadiers from Japan as well as the northwestern Pacific. After then, published records of Japanese grenadiers were almost absent from the literature until Albert Günther (1830–1914) described eight new species based on the collection made by the British research vessel HMS *Challenger* (Günther 1877). A decade later, Günther (1887), in the “*Challenger*” Report, provided more thorough descriptions of the specimens. During the late 19th and early 20th centuries, other European and American ichthyologists also described and/or reported grenadiers from Japanese waters (Steindachner & Döderlein 1887; Jordan & Snyder 1900; Doflein 1906; Smith & Pope 1906; Franz 1910; Jordan & Thompson 1914).

The knowledge of Japanese grenadiers was greatly enhanced when Japan's deep-sea fish fauna was explored by the United States fisheries steamer *Albatross*. David S. Jordan (1851–1931) and Charles H. Gilbert (1859–1928) were the first to review the taxonomy of Japanese grenadiers, and their study was based primarily on the *Albatross* specimens collected in 1900 (Jordan & Gilbert

FIGURE 4. Map of the northwestern Pacific around Japan showing localities where grenadier specimens were collected (each symbol may indicate more than one capture). Red circles represent the localities of specimens examined in this study; orange circles represent those based only on literature records where voucher specimens were not re-identified in this study.

in Jordan & Starks 1904). Excluding bathygadids, they recognized 20 species in 10 genera, of which eight species were new to science. An additional *Albatross* collection of Japanese grenadiers came from the 1906 North Pacific Expedition, of which Gilbert participated as the Naturalist-in-Charge (Dunn 1997). This second collection was studied by Gilbert and his student, Carl L. Hubbs (1894–1979). Based on these *Albatross* specimens, Gilbert & Hubbs (1916) published a landmark monograph of grenadier taxonomy, providing a further review of the Japanese species [including descriptions of six new species of macrourids (*sensu stricto*)] and a synopsis of all the then-known genera worldwide.

Subsequent to Gilbert & Hubbs' (1916) contribution, Japanese ichthyologists continued to report grenadiers over the next half century. The major contributions in this period were made by Toshiji Kamohara (1901–1972) and Kiyomatsu Matsubara (1906–1968). Kamohara was one of the first professors who served at Kochi University, and eagerly studied marine fishes of southern Japan,

particularly those from deep waters and coral reefs (Matsuura 1997; Nakayama 2016). During his 45-year career in ichthyology, Kamohara reported 30 species of macrourid grenadiers from Kochi Prefecture (Kamohara 1964), including one new species and three new Japanese records. Kamohara's contemporary (Nakayama 2016), Matsubara, is considered a pioneer in the field of phylogenetic systematics of Japanese fishes (Matsuura 1997). In his early career at the Research Institute for Natural Resources (= "Sigenkagaku Kenkyusyo"), Matsubara described many new species of Japanese fishes, eight of which belong to the family Macrouridae (Matsubara 1936, 1943; Matsubara & Asano 1943).

While the earlier contributions were valuable, it was Osamu Okamura (1933–2008; Fig. 5) who established the foundation for our current knowledge of Japanese grenadiers. He started a taxonomic study of the family at Kochi University under the supervision of Kamohara. After he graduated, he became a Ph.D. candidate of Kyoto University, pursuing his study of grenadiers under

Matsubara's supervision (Okamura 1963a, 1963b). The taxonomic part of his dissertation was published as a monograph entitled *Fauna Japonica, Macrourina (Pisces)* (Okamura 1970a), in which 45 species belonging to 11 genera were recognized (excluding bathygadids). Although his career initially started as an Assistant Professor of Matsubara's laboratory in 1963, he succeeded Kamohara's position at Kochi University after Kamohara's retirement in 1965. During his employment at Kochi (1965–1996), Okamura organized and participated in numerous research cruises to collect deep-sea demersal fishes, establishing a huge collection of grenadiers from Japanese waters (and also from other countries). Many of these specimens were reported in *Fishes of the Kyushu-Palau Ridge and Tosa Bay* (Okamura *et al.* eds. 1982) and *Fishes of the Okinawa Trough and the Adjacent Waters I* (Okamura & Kitajima eds. 1984). These two publications included descriptions of 11 new species of grenadiers, which brought the total number of Japanese macrourids to 61 (Okamura 1984b, 1988). However, information on most of his specimens remained unpublished, and those specimens contributed a large portion of the material examined in this study.

FIGURE 5. Osamu Okamura (1933–2008) at his laboratory in Kochi University, Japan, in 1981. [Photo: H. Endo]

Over the past three decades, many Japanese researchers (including taxonomists, ecologists, and fisheries biologists) conducted a large number of trawl surveys around the archipelago (e.g., Shinohara *et al.* 1996, 2001, 2005, 2009, 2011, 2014; Shinohara & Matsuura 1997; Suetsugu & Ohta 2005; Kitagawa *et al.* 2008; Furuhashi *et al.* 2010). These extensive efforts covered previously poorly explored areas, such as the Pacific Ocean off southern Honshu

including the Shichito-Iojima Ridge, Hyuga-nada, deep slopes, and midwater depths. As a result of the dedication of these researchers, thousands of grenadier specimens were deposited in various Japanese institutions. Some new species and new records have been reported based on the newly acquired specimens (Endo & Okamura 1992; Endo *et al.* 1994; Nakayama & Endo 2012, 2016a; Nakayama *et al.* 2015a, 2015b, 2015c, 2018). However, most of the new material remained unpublished. The author expresses sincere gratitude to all who collected scientific specimens of Japanese grenadiers on which this present publication is largely based.

Materials and methods

Study area. The Pacific off Japan, East China Sea, Sea of Japan, and southern Okhotsk Sea (Fig. 1).

Counts and measurements. Methods for taking counts and making measurements generally follow procedures described by Iwamoto (1970, 1979), Iwamoto & Sazonov (1988), Iwamoto & Williams (1999), Nakayama *et al.* (2015c), and Nakayama & Endo (2017), and are redefined in the following paragraphs (characters marked with an asterisk are preliminary measurements and are not used in detailed morphometric comparisons).

First dorsal-fin rays—The bases of all segmented rays are counted, including the last two rays; the Roman numerals “II” refer to the two spinous rays (= pseudospines; sensu Endo 2002), the first ray is rudimentary (often scarcely discernible without a microscope) and closely attached to the elongate second spinous ray; the Arabic numbers that follow refer to the segmented rays (Iwamoto 1970; Iwamoto & Sazonov 1988; Iwamoto & Williams 1999).

Pectoral- and pelvic-fin rays—All rays are counted on both sides of the body; the lowercase “i” of pectoral-fin rays refers to the rudimentary or short splint-like uppermost ray that is closely attached to the well-developed second ray (Iwamoto 1970; Iwamoto & Sazonov 1988; Iwamoto & Williams 1999).

Branchiostegal rays—The count is often difficult to determine without probing or dissection. The posterior four rays are always attached laterally on the epihyal and ceratohyal, with the base of the anteriormost one being enlarged; the remaining two or three (rarely four) rays are generally small and slender, and are attached anteriorly and medially on the ceratohyal; thus only the anterior rays attached medially on the ceratohyal need to be counted to determine the total number (Iwamoto 1979).

Gill rakers on first and second arches—All gill rakers, including rudimentary ones, of the first (outermost) and second arches are counted; counts of both outer and inner series are given and separated by a slash mark (Iwamoto & Sazonov 1988; Iwamoto & Williams 1999).

Longitudinal scales—Longitudinal scales are counted from the anterior end of the lateral line over a

FIGURE 6. Diagrammatic illustrations of (A) *Ventrifossa garmani*, (B) *Coelorinchus fuscigulus*, (C) *Nezumia proxima*, (D–E) *Coe. japonicus*, (F) *Coryphaenoides leptolepis*, and (G) *Malacocephalus occidentalis* (after Iwamoto 1970: fig. 3) showing methods for making measurements of grenadiers. Abbreviations: 1D–2D—interdorsal length; 1D base—length of first dorsal-fin base; 1D height—height of first dorsal fin; Anus–A—anus–anal distance; Barbel—barbel length; BD–1D—body depth at first dorsal-fin origin; BD–A—body depth at anal-fin origin; BW–P—body width over pectoral-fin bases; CD–40A—caudal depth at base of 40th anal-fin ray; Gill slit—length of gill slit; Head (HL)—head length; HW–preop.—head width over preopercles; Internasal—internasal width; Interorb.—interorbital width; Isth.–V—isthmus–pelvic distance; Isth.–A—isthmus–anal distance; Isth.–anus—isthmus–anus distance; Lat. nasal. r.—length of lateral nasal ridge; Light organ—length of light organ; Nostril—length of posterior nostril; Occipital w.—occipital width; Orb.—orbit diameter; Orb.–preop.—orbit–preopercle distance; Pectoral—pectoral-fin length; Pelvic—pelvic-fin length; Postorb.—postorbital length; Postrostral (PRL)—postrostral length; Pre–1D—predorsal length; Pre–A—preanal length; Pre–anus—preanus length; Preoral—preoral length; Pre–V—prepelvic length; Rictus—length of rictus; Snout—snout length; Snout w.—snout width; Suborb.—suborbital width; Suborb. r.—length of suborbital ridge; Suborb. shelf—suborbital shelf width; Terminal sc.—length of terminal snout scute; Upp. jaw—upper-jaw length; V–A—pelvic–anal distance; V–anus—pelvic–anus distance. [Illustrations: N. Nakayama]

distance equal to the predorsal length. The first scale is located immediately posterior to the hind margin of the posttemporal. This count corresponds to the number of lateral-line scales (over a distance equal to the predorsal length) in previous studies (e.g., Iwamoto 1970; Iwamoto & Sazonov 1988; Iwamoto & Williams 1999). I prefer to use the term “longitudinal scales” rather than “lateral-line scales”, because a grooved lateral line is partly interrupted in some species.

Transverse scale rows—Transverse scale rows are counted below the origin and midbase of the first dorsal fin and the origin of the second dorsal fin, and above the origin of the anal fin; half-scale counts (0.5) are given when the uppermost or lowermost scale(s) is(are) considerably smaller than others of the series (Iwamoto & Sazonov 1988; Iwamoto & Williams 1999).

Cephalic sensory pores—All pores on each cephalic sensory canal are counted.

Pyloric caeca—All distal tips of pyloric caeca are counted (Iwamoto & Sazonov 1988).

Rete-gas gland complexes—All rete-gas gland complexes in the swim bladder are counted (see Iwamoto 1978: fig. 3).

Total length (TL)—Measurement from snout tip to posteriormost extent of tail; a plus sign (+) indicates that a distal portion of the tail tip was missing or that a pseudocaudal was present (Iwamoto & Sazonov 1988; Iwamoto & Williams 1999).

Head length (HL)—Measurement from snout tip to posterodorsal angle of opercle, excluding opercular membrane (Fig. 6A; Iwamoto & Sazonov 1988; Iwamoto & Williams 1999).

FIGURE 7. Diagrammatic illustrations of *Coryphaenoides leptolepis* showing the cephalic sensory system. Abbreviations: Ant. nost.—anterior nostril; Infraorb. c.—infraorbital canal; Infraorb. p.—infraorbital pores; Mand. c.—mandibular canal; Mand. p.—mandibular pores; Post. nost.—posterior nostril; Postorb. c.—postorbital canal; Postorb. p.—postorbital pores; Preop. c.—preopercular canal; Preop. p.—preopercular pores; Occip. c.—occipital canal; Occip. p.—occipital pores; Supraorb. c.—supraorbital canal; Supraorb. p.—supraorbital pore; Supratemp. c.—supratemporal canal; Supratemp. p.—supratemporal pores. [Illustrations: N. Nakayama]

Snout length—Measurement from snout tip to anteriormost rim of orbit (Fig. 6A; Iwamoto & Sazonov 1988; Iwamoto & Williams 1999).

Orbit diameter—Greatest diameter of orbit, often oblique (Fig. 6A; Iwamoto 1970; Iwamoto & Sazonov 1988).

Postorbital length—Measurement from posteriormost rim of orbit to posterodorsal angle of opercle, excluding opercular membrane (Fig. 6A; Iwamoto & Williams 1999).

Postrostral length (PRL)—Measurement from anteriormost rim of orbit to posterodorsal angle of opercle, excluding opercular membrane (Fig. 6A; Iwamoto & Sazonov 1988).

Orbit-preopercle distance—Oblique measurement from posteroventral rim of orbit to posteroventral angle of preopercle (Fig. 6A; Iwamoto & Sazonov 1988; Iwamoto & Williams 1999).

Suborbital width—Least width of suborbital space (Fig. 6A; Iwamoto & Sazonov 1988; Iwamoto & Williams 1999).

**Suborbital shelf width*—Least width of suborbital shelf (= dorsal half of suborbital area above infraorbital ridge; Fig. 6B).

Upper-jaw length—Measurement from anteriormost tip of premaxillary to posteriormost end of maxillary (Fig.

6E; Iwamoto & Sazonov 1988; Iwamoto & Williams 1999).

Length of rictus—Measurement from anteriormost tip of premaxillary to lateral corner of mouth (Fig. 6E; Iwamoto & Williams 1999).

Length of premaxillary tooth band—Greatest measurement of premaxillary tooth band.

Preoral length—Median ventral measurement from snout tip to (but not including) upper lip (Fig. 6E; Iwamoto 1970; Iwamoto & Sazonov 1988; Iwamoto & Williams 1999).

Length of terminal snout scute—Longitudinal measurement of terminal snout scute in dorsal view (Fig. 6D).

Length of lateral nasal ridge—Measurement from snout tip to posteriormost end of lateral nasal ridge, including terminal snout scute (Fig. 6B, D).

Length of suborbital ridge—Measurement from snout tip to posteriormost end of preopercular ridge, including terminal snout scute (Fig. 6B).

Distance between tip and lateral angle of snout—Measurement from snout tip to lateral angle of snout.

Snout width—Transverse measurement between posteriormost ends of lateral nasal ridges (= posterolateral angles of nasals) or lateral angles of snout (Fig. 6D, F).

Internasal width—Least transverse measurement between nasal fossae (= outer margins of supranarial

ridges; Fig. 6F; Iwamoto & Sazonov 1988; Iwamoto & Williams 1999).

Interorbital width—Least transverse measurement between bony orbits (Fig. 6F; Iwamoto 1970; Iwamoto & Sazonov 1988; Iwamoto & Williams 1999).

Occipital width—Least transverse measurement between inner margins of parietal ridges (Fig. 6D; Nakayama *et al.* 2015c).

**Head width over preopercles*—Greatest transverse measurement across preopercles (including bony scutes of suborbital ridges, if any present; Fig. 6E–F).

**Greatest head width*—Greatest transverse measurement of head, usually across preopercles (including bony scutes of suborbital ridges, if any present; Iwamoto & Williams 1999).

Body width over pectoral-fin bases—Transverse measurement across upper pectoral-fin bases (Fig. 6D).

**Greatest body width*—Greatest transverse measurement of body, usually across upper pectoral-fin bases.

Body depths at first dorsal- and anal-fin origins—Body depths below first dorsal-fin origin and above anal-fin origin (Fig. 6B; Iwamoto & Sazonov 1988; Iwamoto & Williams 1999).

Caudal depth at base of 40th anal-fin ray—Vertical measurement of tail at base of 40th anal-fin ray.

Prepelvic, preanal, and predorsal lengths—Measurement from snout tip to extreme base of outermost or anteriormost ray of each fin (Fig. 6B–C; Iwamoto & Sazonov 1988; Iwamoto & Williams 1999).

Preanus length—Measurement from snout tip to center of anal opening (Fig. 6C).

Isthmus-pelvic, isthmus-anus, and isthmus-anal distances—Measurements from anterior end of scaled area on chest to outer pelvic-fin base, center of anal opening, and anal-fin origin respectively (Fig. 6C; Iwamoto 1970; Iwamoto & Sazonov 1988).

Pelvic-anus and pelvic-anal distances—Measurements from outer pelvic-fin base to center of anal opening and anal-fin origin respectively (Fig. 6C; Iwamoto 1970; Iwamoto & Sazonov 1988; Iwamoto & Williams 1999).

Anus-anal distance—Measurement from center of anal opening to anal-fin origin (Fig. 6C).

**Length of light organ*—Measurement from anterior end of ventral light organ to anterior margin of anus (Fig. 6E).

Pelvic- and pectoral-fin lengths—Measurement from extreme base of uppermost, outermost, or anteriormost ray to farthest tip of each fin; fleshy filaments, if any, included (Fig. 6A; Iwamoto 1970).

Height of first dorsal fin—Measurement from fin origin to distal tip of second spinous ray, excluding fleshy filament (Fig. 6A).

Length of first dorsal-fin base—Measurement from fin origin to posterior base of last ray (Fig. 6A).

Interdorsal length—Measurement from posterior base of last ray of first dorsal fin to origin of second dorsal fin (Fig. 6A; Iwamoto 1970; Iwamoto & Sazonov 1988; Iwamoto & Williams 1999).

Length of gill slit—Greatest diameter of outermost gill slit (Fig. 6G; Iwamoto 1970; Iwamoto & Sazonov 1988; Iwamoto & Williams 1999). To make this measurement, the gill membrane may need to be cut anterior to the anteriormost branchiostegal ray to expand the gill opening, particularly in species where the membranes are broadly connected across the isthmus.

Length of posterior nostril—Greatest diameter of posterior nostril (Fig. 6B; Iwamoto & Sazonov 1988; Iwamoto & Williams 1999).

Length of olfactory bulb—Greatest longitudinal measurement of olfactory bulb including filaments, dissection of nasal skin required.

Barbel length—Measurement of free portion of chin barbel, taken from posterior insertion to distal tip (Fig. 6A; Iwamoto 1970; Iwamoto & Sazonov 1988; Iwamoto & Williams 1999).

Length of pyloric caecum—Measurement from base to tip of longest pyloric caecum.

All measurements were taken point-to-point with calipers to the nearest 0.1 mm, except for total length, which was measured with 1 mm precision rulers. Head, total, and postrostral lengths are expressed as HL, TL, and PRL respectively. While HL is regularly used as a standard of comparison in grenadiers, PRL is alternatively used here for the species of *Coelorinchus* and *Hymenocephalus*. This is because of an ontogenetic change in the relative length of the snout in *Coelorinchus* (see Nakayama *et al.* 2015c) or a fragile nature of the head bones in *Hymenocephalus*, viz., the median nasal rostrum of *Hymenocephalus* is so weak that the tip of the snout is easily broken when trawled, or is regularly bent to the side in alcohol preserved specimens due to dehydration. Counting scale rows and examination of the cephalic sensory pores were facilitated by a temporal stain with a Cyanine Blue solution (Akihito *et al.* 1993; Saruwatari *et al.* 1997). Meristic and morphometric data given in the text are based on only the specimens examined in this study unless otherwise stated.

Morphological terminology. The terminology of the cephalic sensory system generally follows Nakayama & Endo (2012: fig. 3), and is demonstrated in Fig. 7.

The terminology of the bony ridges and modified scales on the head (Fig. 8) is that of Iwamoto (1978), but definitions of the lateral nasal, infraorbital, and preopercular ridges follow Okamura (1970b: fig. 36). In the species of *Coelorinchus*, the infraorbital ridge is tightly connected with both the lateral nasal and preopercular ridges, forming a long, continuous, longitudinal ridge (= suborbital ridge; usually stout and bony). The border of the lateral nasal and infraorbital ridges is distinct, as they are often separated from each other by a very

FIGURE 8. Heads of (A–B) *Coelorinchus hige* and (C–D) *Coryphaenoides microps* showing positions of the ridges and modified scutes on the head. Abbreviations: Infraorb. r.—infraorbital ridge; Lat. angle sc.—scute at lateral angle of snout; Lat. nas. r.—lateral nasal ridge; Lead. edge—leading edge of snout; Med. rostr. r.—median rostral ridge; Occip. sc.—occipital scute; Par. r.—parietal ridge; Postorb. r.—postorbital ridge; Posttemp. sc.—posttemporal scute; Preop. r.—preopercular ridge; Suborb. r.—suborbital ridge; Supran. r.—supranarial ridge; Supraorb. r.—supraorbital ridge; Terminal sc.—terminal snout scute. (A, C) Lateral and (B, D) ventral views. [Photos: N. Nakayama]

narrow naked gap where a tiny free neuromast is usually present; however, the border between the infraorbital and preopercular ridges is difficult to determine.

The anterolateral fusion of the lateral and medial processes of the nasal bone is an important character in *Coelorinchus* classification. Methods for determining whether or not the two processes are fused follow Iwamoto & Williams (1999) and Iwamoto *et al.* (2009). Namely, if the lateral nasal ridge yields at any point when lightly pressed with one's fingernail, a gap exists between the lateral and medial processes (see Okamura 1970b: fig. 32B' and D'); this state is described herein as “lateral nasal ridge incompletely supported by nasal bone”. On the other hand, if the lateral nasal ridge feels solid when pressed, the two processes are fully joined anterolaterally (see Okamura 1970b: fig. 32A' and C'), and it is described herein as “lateral nasal ridge completely supported by nasal bone”. Radiographs are also helpful to examine the presence/absence of the fusion. Alternatively, the skin below the ridge can be incised to confirm the shape of the bone (see Iwamoto & Graham 2001: fig. 52).

The term occipital scales refers to head scales between the parietal ridges; morphology of the occipital scales is generally described from those on the dorsal midline of the head.

The term ventral striae refers to areas of fine (microscopic), parallel, black lines extending on ventral surfaces of the chest, shoulder girdle, and belly, and along each side of the isthmus, commonly found in the species of *Hymenocephalus*, *Hymenogadus*, and *Spicomacrurus*; this structure is also found on the gular membrane in the species of the former two genera.

The term periproct refers to the area of black naked skin surrounding the anus and urogenital opening.

Scanning electron microscopy. In addition to a standard binocular microscope, a scanning electron microscope (SEM) was used to examine the fine structure of the body scales. Methods for SEM preparation follow Roberts (1993). SEM images provided by the author were taken at the Center for Advanced Marine Core Research, Kochi University.

Deposition of specimens. For the species distributed in Japan and adjacent waters, a total of 7846 specimens (including those listed in Appendix 1) deposited in the following institutions was examined in the course of this study:

AMS	Australian Museum, Sydney, Australia.
ASIZP	Biodiversity Research Center, Academia Sinica, Taipei, Taiwan.

BMNH	Natural History Museum, London, UK.		
BSKU	Laboratory of Marine Biology, Faculty of Science and Technology, Kochi University, Kochi, Japan.	RMNH	Naturalis Biodiversity Center, Leiden, Netherlands.
CAS	California Academy of Sciences, San Francisco, USA.	SFU	Shanghai Ocean University, Shanghai, China.
CAS-SU	Stanford University, Stanford, USA (currently deposited at CAS; also known as SU).	SHMNH	Saito Ho-on Kai Museum of Natural History, Sendai, Japan (currently deposited at NSMT).
CBM	Natural History Museum and Institute, Chiba, Japan.	SKFC	Sigenkagaku Fish Collection, Research Institute for Natural Resources, Tokyo, Japan (some specimens currently deposited at FAKU).
CM	Carnegie Museum of Natural History, Pittsburgh, USA (currently deposited at FMNH).	SNFR	Seikai National Fisheries Research Institute, Fisheries Research Agency, Nagasaki, Japan.
CMNFI	Canadian Museum of Nature, Ottawa, Canada.	USNM	National Museum of Natural History, Smithsonian Institution, Washington D.C., USA.
CSIRO	Australian National Fish Collection, Commonwealth Scientific and Industrial Research Organization, Hobart, Australia.	ZMMGU	Zoological Museum, Moscow State University, Moscow, Russia.
FAKU	Kyoto University Fish Collection, Kyoto University, Japan (separately deposited at the Kyoto University Museum, Kyoto, and the Maizuru Fisheries Research Station, Maizuru).	ZMUB	Museum of Zoology, University of Bergen, Bergen, Norway.
FFNU	Faculty of Fisheries, Nagasaki University, Nagasaki, Japan.	ZMUC	Zoological Museum, University of Copenhagen, Copenhagen, Denmark.
FMNH	Field Museum of Natural History, Chicago, USA.	ZMUT	University Museum, University of Tokyo, Tokyo, Japan.
FRLM	Fisheries Research Laboratory, Mie University, Tsu, Japan.		
FSFRL	National Research Institute of Far Seas Fisheries, Fisheries Research Agency, Shizuoka, Japan (currently deposited at NSMT).		
HUMZ	Hokkaido University Museum, Hokkaido University, Hakodate, Japan.		
KAUM	Kagoshima University Museum, Kagoshima University, Kagoshima, Japan.		
KPM	Kanagawa Prefectural Museum of Natural History, Odawara, Japan.		
KSHS	Kochi Senior High School, Kochi, Japan (currently deposited at NSMT).		
MCZ	Museum of Comparative Zoology, Harvard University, Cambridge, USA.		
MSM	Marine Science Museum, Tokai University, Shizuoka, Japan.		
MUFS	Faculty of Agriculture, University of Miyazaki, Miyazaki, Japan.		
MZS	Musée Zoologique de l'Université Louis Pasteur, Strasbourg, France.		
NSMT	National Museum of Nature and Science, Tsukuba, Japan.		
OCF	Okinawa Churashima Research Center, Okinawa Churashima Foundation, Motobu, Japan.		
OMNH	Osaka Museum of Natural History, Osaka, Japan.		

Additional comparative material (Appendix 2) is deposited at AMS, ANSP (Academy of Natural Sciences, Philadelphia, Pennsylvania, USA), ASIZP, BPBM (Bernice P. Bishop Museum, Honolulu, USA), BSKU, CAS, CSIRO, FAKU, FMNH, HUMZ, IORAS (P.P. Shirshov Institute of Oceanology, Russian Academy of Sciences, Moscow, Russia), MNHN, MZB, [Museum Zoologicum Bogoriense, Division of Zoology, Research Center for Biology, Indonesian Institute of Sciences (LIPI), Cibinong, Indonesia], NSMT, NTM (Museum and Art Gallery of the Northern Territory, Darwin, Australia), SNFR, USNM, WAM (Western Australian Museum, Perth, Australia), ZMH (Zoologisches Institut und Museum, Universität Hamburg, Hamburg, Germany), ZMMGU, ZMUB, and ZMUC. For other institutional abbreviations used in the text see Eschmeyer (1998) and Fricke & Eschmeyer (2020).

Other notes. The taxonomic section includes six species not reported within Japan's EEZ but known from the southern East China Sea off Taiwan and the Sea of Japan off Korea, viz., *Coelorinchus leptorhinus* Chiou, Shao & Iwamoto, 2004, *Coryphaenoides microps* (Smith & Radcliffe in Radcliffe, 1912), *Lucigadus nigromarginatus* (Smith & Radcliffe in Radcliffe, 1912), *Pseudocetonurus* sp. cf. *septifer* Sazonov & Shcherbachev, 1982, *Sphagemacrurus pumiliceps* (Alcock, 1894), and *Ventrifossa nigrodorsalis* Gilbert & Hubbs, 1920. The keys provided here are designed for use in identifying genera and species known from Japan and adjacent

waters only. Taxa marked with an asterisk are keyed out twice in the same key. Synonym lists for each species are generally confined to the original descriptions, important taxonomic works, and references regarding previous Japanese records of grenadiers. For further literature information see Eschmeyer (1998) and Fricke *et al.* (2020). In the Material examined sections, specimens are generally listed south to north, and west to east; an asterisk denotes lots from which counts and measurements were not taken. The sex of specimens, if indicated, was determined by direct examination of the gonads through an abdominal incision. Data given in the Size sections indicate the maximum size known for each species. Maps for Figs. 1–4 and Appendix 3 were generated using the Generic Mapping Tools (GMT ver. 4.5.9; see Wessel *et al.* 2013), with bathymetric data obtained from ETOPO1 (Amante & Eakins 2009). For the spellings *Coelorinchus* (a correction of its original spelling, viz., *Cælorinchus* Giorna, 1809) and *Macrourus* (rather than “*Macrurus*”) see the Remarks of the former genus.

Taxonomy

Family Macrouridae Bonaparte, 1831 (sensu Iwamoto *et al.* 2015)

[Japanese name: Sokodara-ka]

Diagnosis. Two dorsal fins well separated; first dorsal fin high and short-based; second dorsal fin long-based, poorly developed throughout its length; anal fin similar to, but usually much deeper than second dorsal fin; pelvic fin well developed with 5 or more soft rays; tail greatly elongate, without either caudal fin or caudal peduncle (pseudocaudal occasionally present and variously developed); mouth usually inferior to subterminal (rarely terminal); outermost gill slits restricted by skin folds attached to upper and lower ends of gill arch; gill rakers tubercular, not slender or lath-like; posttemporal fossa absent; body scales almost always ctenoid.

Remarks. This study confirms the occurrence of 76 grenadier species belonging to 18 genera from Japan and adjacent waters, of which 70 species in 15 genera are recognized within Japan’s EEZ.

Key to genera of grenadiers from Japan and adjacent waters

- 1a Infraorbital and preopercular ridges tightly adjoined, forming continuous bony ridge; preopercular ridge elongate posteroventrally and sharply pointed at posteroventral angle; snout long, protruding well beyond upper jaw, acutely pointed in lateral view in most species *Coelorinchus* (25 spp.)
- 1b Infraorbital ridge not connected with preopercular ridge, separated by distinct gap; preopercular ridge not elongate

- posteroventrally nor sharply pointed at posteroventral angle; snout relatively short, moderately to scarcely protruding beyond upper jaw, conical to broadly rounded in lateral view 2
- 2a Anus (or periproct, if developed) abutting or only slightly separated from anal-fin origin 3
- 2b Anus (or periproct, if developed) distinctly separated by several scale rows from anal-fin origin 11
- 3a Gular region covered with net-like pattern or striation of black epithelium; broad areas of fine (microscopic), parallel, black lines (ventral striae) extending on ventral surfaces of chest, shoulder girdle, and belly, and along each side of isthmus; small lens-like structure of light organ on chest 4
- 3b Gular region uniformly pale, dusky, dark, or blackish; ventral striae absent; no lens-like structure of light organ on chest anterior to pelvic fins 6
- 4a Gular region covered with net-like pattern of black epithelium; nasal bones stout, forming 3 horizontally flattened projections; outer pelvic-fin ray flattened, with narrow ribbon-like flange on free distal portion *Spicomacrus* (*S. kuronumai*)
- 4b Gular region covered with striated pattern of black epithelium; nasal bones weak, usually paper-thin, not forming horizontal projections; outer pelvic-fin ray not flattened distally 5
- 5a Second spinous ray of first dorsal fin smooth along its leading edge; trunk moderately deep, laterally compressed in cross section *Hymenocephalus* (6 spp.)
- 5b Second spinous ray of first dorsal fin serrated along its leading edge; trunk shallow, almost cylindrical in cross section *Hymenogadus* (*H. gracilis*)
- 6a Ventral aspects of body well forward so that anal-fin origin distinctly anterior to hind margin of first dorsal-fin base; scales on infraorbital ridge modified into coarsely spinulated, adherent scutes **Sphagemacrus* (*S. pumiliceps*)
- 6b Ventral aspects of body not well forward, and anal-fin origin below or posterior to hind margin of first dorsal-fin base; scales on infraorbital ridge normal or only slightly thickened 7
- 7a Second spinous ray of first dorsal fin smooth along its leading edge; periproct large, occupying more than half of distance between inner pelvic-fin bases to anal-fin origin; scales along anterior part of anal-fin base enlarged *Trachonurus* (*T. villosus*)
- 7b Second spinous ray of first dorsal fin serrated along its leading edge; periproct small, its anterior margin far posterior to midway between inner pelvic-fin bases and anal-fin origin; scales along anterior part of anal-fin base not especially enlarged 8
- 8a Head massive, soft, inflated, and broad; scales along anterior part of second dorsal-fin base enlarged *Cetonus* (*C. globiceps*)
- 8b Head not inflated, moderately compressed laterally; scales along anterior part of second dorsal-fin base not especially enlarged 9
- 9a Periproct absent or rudimentary; reticulate structure absent on unexposed portion of body scales; branchiostegal rays 6 *Coryphaenoides* (15 spp.)

- 9b Periproct narrow but distinct; reticulate structure broadly developed on unexposed portion of body scales; branchiostegal rays 7–8 10
- 10a Dorsal surface of snout broadly naked anterior to line connecting lateral angles of snout; underside of head almost completely naked; posterior margin of upper jaw extending to vertical through hind rim of orbit or beyond; spinules on body scales arranged in widely spaced subparallel to slightly divergent rows..... *Asthenomacrus* (*A. victoris*)
- 10b Dorsal surface of snout fully scaled; naked areas on underside of head confined to area above upper lip and anterior portions of mandibular rami; posterior margin of upper jaw not reaching vertical through hind rim of orbit; spinules on body scales arranged in quincunx order
.....*Pseudonezumia* (2 spp.)
- 11a Premaxillary teeth fang-like, arranged in 1 distinct row; chin barbel absent; trunk strongly compressed.....
.....*Odontomacrus* (*O. murrayi*)
- 11b Premaxillary teeth not fang-like, relatively small, arranged in 2 or more rows; chin barbel present, variously developed; trunk moderately compressed 12
- 12a Second spinous ray of first dorsal fin smooth along its leading edge; mandibular teeth arranged in 1 or 2 rows 13
- 12b Second spinous ray of first dorsal fin variously serrated along its leading edge; mandibular teeth arranged in narrow to wide band..... 15
- 13a Mandibular teeth arranged in 2 irregular rows, with inner row slightly enlarged
.....**Ventrifossa* (in part; subgenus *Atherodus*: *V. macroptera*)
- 13b Mandibular teeth arranged in 1 distinct row..... 14
- 14a Anterior dermal window of light organ large, bean-shaped, laterally expanded; mandibular teeth widely spaced canines; lower portion of gill membranes narrowly scaled along branchiostegal rays*Malacocephalus* (*M. nipponensis*)
- 14b Anterior dermal window of light organ small and circular; mandibular teeth small, not enlarged, tightly arranged; branchiostegal membranes entirely naked.....
.....*Pseudocetonurus* (*P. sp. cf. septifer*)
- 15a Dorsal surface of snout broadly naked anterior to line connecting lateral angles of snout.....*Kumba* (*K. japonica*)
- 15b Dorsal surface of snout completely or almost completely scaled 16
- 16a Scales on infraorbital ridge modified into coarsely spinulated, adherent scutes..... 17
- 16b Scales on infraorbital ridge not especially modified 18
- 17a Anus closer to pelvic-fin bases than to anal-fin origin; ventral aspects of body not positioned forward, and anal-fin origin below or posterior to hind margin of first dorsal-fin base....
.....*Nezumia* (7 spp.)
- 17b Anus closer to anal-fin origin than to pelvic-fin bases; ventral aspects of body positioned forward so that anal-fin origin distinctly anterior to hind margin of first dorsal-fin base.....
.....**Sphagemacrus* (*S. pumiliceps*)
- 18a Posterior margin of upper jaw extending beyond vertical through midorbit; posterior end of premaxillary tooth band extending beyond hind margin of maxillary process.....
.....**Ventrifossa* (in part; subgenera *Sokodara* and *Ventrifossa*: 7 spp.)
- 18b Posterior margin of upper jaw not or barely reaching vertical through midorbit; posterior end of premaxillary tooth band falling short of hind margin of maxillary process..... 19
- 19a Reticulate structure present on unexposed portion of body scales; body uniformly dark to blackish; posterior dermal window of light organ scarcely discernible externally
.....*Kuronezumia* (2 spp.)
- 19b Reticulate structure absent on body scales; body pale to dusky; posterior dermal window of light organ prominent, much larger than anterior window
.....*Lucigadus* (*L. nigromarginatus*)

Genus *Asthenomacrus* Sazonov & Shcherbachev, 1982

[New Japanese name: Narakuno-sokodara-zoku]

Asthenomacrus Sazonov & Shcherbachev, 1982:2 (type species: *Asthenomacrus victoris* Sazonov & Shcherbachev, 1982, by original designation).

Diagnosis. Anus immediately anterior to anal-fin origin, surrounded by narrow but distinct periproct. Ventral light organ absent. Infraorbital ridge not connected with preopercular ridge, separated by distinct gap. Pelvic fin with 7–9 soft rays, its origin distinctly anterior to vertical through pectoral-fin base. Leading edge of second spinous ray of first dorsal fin weakly serrated. Head bones weak, flexible. Mouth large, posterior margin of upper jaw extending beyond vertical through hind rim of orbit. Nasal fossa not greatly enlarged. Chin barbel present. Teeth in narrow tapered bands in both jaws, none especially enlarged. Underside of head almost completely naked, including gular and branchiostegal membranes; dorsal surface of snout broadly naked anterior to line connecting lateral angles of snout; no modified scales along head ridges; tip and lateral angles of snout lacking prominent scutes. Body scales covered with short, erect, needle-like spinules in widely separated, subparallel to slightly divergent rows; buttresses of scale spinules narrowly developed; reticulate structure developed over entire surface of unexposed portion. Scales along second dorsal and anal fins not enlarged. Grooved lateral line complete. Cephalic sensory pores present on underside of head, scarcely discernible. Pyloric caeca very short, thick, 7–10. Branchiostegal rays 7, rarely 8. Body lacking prominent silvery reflections when fresh. [Modified from Iwamoto & Orlov (2008).]

Remarks. *Asthenomacrus* appears to be most similar to *Macrosamia* Merrett, Sazonov & Shcherbachev, 1983, and they share the following combination of features: branchiostegal rays 7 or rarely 8; anus immediately anterior to anal-fin origin; no ventral light organs; pelvic-fin base distinctly anterior to vertical through pectoral-fin base;

first dorsal fin serrated along its leading edge; underside of head almost completely naked; dorsal surface of snout broadly naked anteriorly. *Asthenomacrurus* differs notably from *Macrosmia* in having a small nasal fossa (vs. greatly enlarged) and fewer pelvic-fin rays (7–9 vs. 11–12) and pyloric caeca (7–10 vs. 15–18). [Data for *Macrosmia* are from Merrett *et al.* (1983), Merrett & Iwamoto (2000), and Iwamoto *et al.* (2015).]

The genus currently comprises two bathyal species: *A. victoris* Sazonov & Shcherbachev, 1982, the type species widely distributed in the Indo-West Pacific including Japan, plus a single record from the Mid-Atlantic Ridge; *A. fragilis* (Garman, 1899) known only from the eastern central Pacific. Iwamoto & Orlov (2008) recently provided a redescription of *A. victoris* based on all the then-known specimens, giving a detailed comparison with *A. fragilis*.

***Asthenomacrurus victoris* Sazonov & Shcherbachev, 1982**

[New Japanese name: Narakuno-sokodara]

(Figs. 9–15; Appendix 3-1A)

Asthenomacrurus victoris Sazonov & Shcherbachev, 1982:3, fig. 1 (original description; holotype: ZMMGU P.15349, from Mid-Indian Ridge, in 3530 m; 2 paratypes from Western Australia and Japan); Iwamoto & Orlov 2008:126 (redescription; comparison with *A. fragilis*).

Asthenomacrurus [sic] *victoris*: Nakabo & Kai 2013:1874 (comments on Japanese record).

Diagnosis. Differs from its only congener, *A. fragilis*, in having the following combination of features: pelvic-fin rays 7–8; preoral length 26–40% HL; barbel length 11–17% HL; cephalic sensory pores not prominent; pyloric caeca 8–10; gill rakers on first arch (outer/inner) 8–12/12–16, on second arch 11–15/11–15. [Modified from Iwamoto & Orlov (2008).]

Material examined. 19 specimens. **Holotype of *Asthenomacrurus victoris*:** ZMMGU P.15349 (female, 32.9 mm HL, 245+ mm TL), Mid-Indian Ridge, western Indian Ocean, 2.7667°S, 65.6833°E, 3530 m, R/V *Vityaz*, cr. 31, sta. 4634, coll. V.M. Makushok, 12 Feb. 1960. **Paratype of *A. victoris*:** ZMMGU P.16012 (1 male, 41.9 mm HL, 223+ mm TL), off Cape Leeuwin, Western Australia, Naturalist Plateau, southeastern Indian Ocean, 33.7000°S, 110.8833°E, 2320–2350 m, R/V *Dmitriy Mendeleev*, cr. 16, coll. D.P. Andrianov. **Non-types:** **Japan:** BSKU 47934 (1 female, 41.1 mm HL, 192+ mm TL), BSKU 98223 (1 female, 28.9 mm HL, 135+ mm TL), off Shikoku, 32.2372°N, 134.0220°E, 2565–2647 m, R/V *Tansei-maru*, cr. KT-90-13, sta. T6, beam trawl, 3 Sept. 1990; NSMT-P 98028 (3 males, 28.5–31.8 mm HL, 142+–176+ mm TL), southeast of Boso Peninsula, 34.7267°N, 140.7400°E, 2089–2096 m, R/V *Tansei-maru*, cr. KT-99-

06, sta. 15, 3-m ORE beam trawl, 22 Mar. 1999; BSKU 85666 (1 female, 45.9 mm HL, 197+ mm TL), BSKU 85667 (1 male, 44.1 mm HL, 265 mm TL), off Boso Peninsula, 34.3567°N, 140.2190°E, 2480–2516 m, R/V *Tansei-maru*, cr. KT-94-14, sta. 14-2, beam trawl, coll. H. Endo, 2 Sept. 1998; BSKU 45545 (1 female, 43.3 mm HL, 270 mm TL), off Kinkazan, 38.2355°N, 142.4238°E, 2167–2229 m, R/V *Tansei-maru*, cr. KT-89-03, sta. SR93, beam trawl, coll. O. Okamura and H. Endo, 11 Mar. 1989; KPM-NI 18570 (1 female, 35.1 mm HL, 181+ mm TL), KPM-NI 18571 (1 male, 32.8 mm HL, 202+ mm TL), off Tanega-shima Island, 29.8507°N, 130.9281°E, 1678–1769 m, R/V *Tansei-maru*, cr. KT-07-01, sta. YT-3, 3-m beam trawl, coll. T. Sato, 24 Feb. 2007. **Australia:** CSIRO H5304-13 (1 female, 44.2 mm HL, 204+ mm TL), off Victoria, 38.5667°S, 149.4667°E, 1954–1979 m; CSIRO H5304-38 (2 males, 41.2–42.6 mm HL, 190+–227+ mm TL), off Victoria, Tasman Sea, 38.5667°S, 149.4667°E, 1954–1979 m; CSIRO H5313-05 (1 male, 46.5 mm HL, 229+ mm TL), off Tasmania, Tasman Sea, 39.8000°S, 149.1000°E, 1946–1973 m. **Porcupine Seabight:** BMNH 1996.8.12.2–3 (2 sex undetermined, 26.5–29.6 mm HL, 144+–145+ mm TL), 50.4500°N, 12.9833°W, 2470–2500 m, *Discovery*, sta. 5141101, coll. N.R. Merrett, 29 Mar. 1982. **Mid-Atlantic Ridge:** ZMUB 19710 (1 female, 38.1 mm HL, 230 mm TL), 42.7667°N, 29.2667°W, 3036 m, R/V *G.O. Sars*, cr. MAR-ECO, supersta. 46, local sta. 372, 29-m otter trawl, 11 Jul. 2004.

Counts and measurements. Based on 19 specimens (26.5–46.5 mm HL, 135+–270 mm TL). Counts: first dorsal-fin rays II,8–10; pectoral-fin rays i17–i21; pelvic-fin rays 7–8; gill rakers on first arch (outer/inner) 8–12/12–16, on second arch 12–15/11–15; longitudinal scales 28–39; transverse scale rows below first dorsal-fin origin 6–9.5, below first dorsal-fin midbase 4.5–7.5, below second dorsal-fin origin 6–9; pyloric caeca 9–10.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 27–38 (36–57); orbit diameter 20–27 (29–39); postorbital length 44–51 (62–73); postrostral length 65–76; orbit–preopercle distance 38–47 (53–67); suborbital width 15–20 (21–29); upper-jaw length 35–42 (51–60); length of rictus 31–37 (41–53); length of premaxillary tooth band 20–24 (29–34); preoral length 26–40 (37–58); distance between tip and lateral angle of snout 17–26 (26–39); snout width 24–37 (32–54); internasal width 22–33 (29–49); interorbital width 23–41 (31–58); body width over pectoral-fin bases 31–47 (41–70); body depth at first dorsal-fin origin 67–83 (100–120); body depth at anal-fin origin 53–80 (74–101); prepelvic length 91–108 (120–156); preanus length 124–140 (175–203); preanal length 126–145 (182–211); isthmus–pelvic distance 27–32 (38–48); isthmus–anus distance 59–71 (82–100); isthmus–anal distance 63–78 (90–109); pelvic–anal distance 33–55 (48–77); pelvic-fin length 54–78 (77–114); pectoral-fin length 53–77 (81–110); predorsal

FIGURE 9. Fresh specimens of *Asthenomacrus victoris*. (A) BSKU 45545, 43.3 mm HL, 270 mm TL, off Kinkazan, Miyagi Pref., Pacific, 2167–2229 m depth (reversed); (B) BSKU 85666, 45.9 mm HL, 197+ mm TL, off Boso Peninsula, Chiba Pref., Pacific, 2480–2516 m depth; (C) BSKU 47934, 41.1 mm HL, 192+ mm TL, off Shikoku, Pacific, 2565–2647 m depth. Lateral views. [Photos: BSKU]

length 100–113 (137–164); height of first dorsal fin 71–83 (98–121); length of first dorsal-fin base 17–26 (24–36); interdorsal length 21–40 (28–53); length of gill slit 15–22 (21–32); length of posterior nostril 6–13 (9–18); barbel length 11–16 (17–24).

Supplementary description based on Japanese specimens. A full description of *A. victoris* is unnecessary here, because most of its characters have been detailed by previous authors (Sazonov & Shcherbachev 1982; Iwamoto & Orlov 2008). The following description therefore focuses on poorly known features based on the Japanese specimens examined, emphasizing their

variation. General features of the species are shown in Figs. 9–10.

Body scales small, thin, highly deciduous, covered with short, erect, needle-like spinules in widely separated, subparallel to slightly divergent, comb-like rows (Fig. 11A–B); scales on dorsum below interdorsal space vertically elongate, covered with 5–6 spinule rows (about 3 rows in smaller KPM specimens); middle row slightly enlarged, covered with 5–7 spinules (about 4 spinules in KPM specimens); height of spinules gradually increasing posteriorly, with last spinule in each row extending well beyond posterior scale margin; buttresses of spinules

FIGURE 10. Preserved specimens of *Asthenomacrurus victoris*. (A–C) BSKU 45545, 43.3 mm HL, 270 mm TL, off Kinkazan, Miyagi Pref., Pacific, 2167–2229 m depth; (D) ZMMGU P.15349, holotype, 32.9 mm HL, 245+ mm TL, Mid-Indian Ridge, western Indian Ocean, 3530 m depth; (E) ZMMGU P.16012, paratype, 41.9 mm HL, 223+ mm TL, Western Australia, southeastern Indian Ocean, 2320–2350 m depth. (A, D–E) Lateral views; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

poorly developed; reticulate structure scattered over entire unexposed portion. Scales on isthmus longitudinally elongate, smaller than those on dorsum, covered with 1–3 rows of spinules; middle row distinctly longer than other rows (Fig. 11C–D). Predorsal scales covered with single row of spinules. Scales beneath gill covers and posterior to pectoral and pelvic fins cycloid. In KPM-NI 018571 (32.8 mm HL), scales on posterior half of belly also lacking spinules. Body fully scaled except for fins and periproct.

Head scales similar to those on body, but spinules much shorter and more erect. Scales on gill cover, postorbital canal, and posterior part of nape somewhat larger, covered with about 3 divergent rows of spinules.

Other scales, especially those on dorsal surface of head, small and longitudinally elongate, covered with single row of spinules (Fig. 11E–F), giving striated appearance to head surfaces (Fig. 12). Neither tubercles nor modified scales on tip and lateral angles of snout and head ridges.

Dorsal and lateral surfaces of head mostly covered with scales; naked region confined to top of snout anterior to lateral angles (Figs. 12–13). Underside of head completely naked from snout tip to above posterior margin of upper jaw; ventral surface of preopercle naked; exposed portion of interopercle naked, but narrowly scaled posteriorly; mandibular rami, and gular and branchiostegal membranes lacking scales.

FIGURE 11. Scanning electron micrographs showing scales of *Asthenomacrus victoris*. BSKU 45545, 43.3 mm HL. Scales (A–B) from the dorsum below the interdorsal space; (C–D) from the chest posterior to the isthmus; (E–F) from the interorbital area. (A, C, E) Views from above; (B, D, F) oblique views. [Photos: N. Nakayama]

Cephalic sensory canals somewhat inflated when fresh (Fig. 9) but shrunken in most preserved specimens, giving pointed appearance to snout (see also Iwamoto & Orlov 2008: fig. 2A). Due to this, general appearance of head often significantly changed after dehydration in alcohol.

FIGURE 12. Dorsal surface of the snout of *Asthenomacrus victoris* showing squamation and spinulation on the head scales. BSKU 45545, 43.3 mm HL. Preserved condition, stained with Cyanine Blue solution. [Photo: N. Nakayama]

Small inconspicuous sensory pores present along mandibular and infraorbital canals (Fig. 13). No pores on preopercular, supraorbital, occipital, postorbital, and supratemporal canals. Lateral line complete and not interrupted throughout.

Color when fresh (Fig. 9). Ground color of head and body uniformly dark brown in almost intact specimens (Fig. 9A); this dark color still prominent even when body scales completely missing (Fig. 9B), but ground color becoming dusky pink when epidermis badly damaged

(Fig. 9C). Eye, gill cover, naked areas on head, gill membranes, and abdomen dull black; pectoral fin very dark; other fins uniformly dusky, but second spinous ray of first dorsal fin blackish.

Size. To about 27 cm TL (BSKU 45545, 270 mm TL, off Kinkazan, Japan).

Sexual dimorphism. Compared with females, males have a wider posterior nostril [8.5–12.6% HL ($n = 9$) vs. 6.3–9.1 ($n = 8$); Fig. 14A], a larger olfactory bulb (7.9–11.7% HL vs. 4.8–8.0%; Fig. 14B), and more olfactory lamellae [16–20 (mean 17.6, $n = 5$) vs. 14–15 (mean 15.4, $n = 5$)]. In addition, the distal portion of the lamella in males is notably fimbriated, giving a concave appearance to the olfactory bulb (Fig. 15A). By contrast, that in females is simple or only weakly frilled, and the bulb is somewhat flat in appearance (Fig. 15B). These differences are emphasized in adult specimens, but obscure in juveniles of less than 32.5 mm HL.

Distribution. Sporadically recorded from the Indian and West Pacific Oceans, and Mid-Atlantic Ridge, at depths of 1642–3530 m (Sazonov & Shcherbachev 1982; Parin *et al.* 1993; Iwamoto & Graham 2001; Iwamoto *et al.* 2004; Iwamoto & Orlov 2008). In Japan, known from off the Pacific coasts at depths of 1678–3460 m (Appendix 3-1A). Rare.

Remarks. *Asthenomacrus victoris* was originally described by Sazonov & Shcherbachev (1982) based on three specimens collected from the Mid-Indian Ridge (type locality), Western Australia, and Japan (off Ibaraki Pref.). The holotype and the Australian paratype are deposited in the Zoological Museum of Moscow State University (ZMMGU), and the author had an opportunity to examine the two specimens. According to Sazonov & Shcherbachev (1982), the holotype (Fig. 10D) and “the larger of the paratypes” (= ZMMGU P.16012; Fig. 10E) were immature females. The holotype was confirmed

FIGURE 13. Diagrammatic illustrations showing the cephalic sensory system and naked areas (gray) of *Asthenomacrus victoris*. BSKU 45545, 43.3 mm HL. (A) Lateral, (B) dorsal, and (C) ventral views of the head and trunk. [Illustrations: N. Nakayama]

as a female, with well-developed ovaries in which there were large oocytes of about 0.6–1.0 mm in diameter. As typical of females, it also had a small olfactory bulb (7.0% of HL) consisting of 14 simple lamellae. The sex of the Australian paratype could not be determined with certainty, but its gonads seemed to be immature testes. In addition, its olfactory bulb was greatly enlarged (9.5% of HL), consisting of 17 fimbriated lamellae, which further suggests that it represents a male.

Prior to this study, *A. victoris* was represented by 12 specimens in museum collections, most of which are in poor condition (Iwamoto & Orlov 2008). Additional specimens from Japan are well preserved, which allowed documentation of some squamation features of the species for the first time (see the Supplementary description). Although the scales of *A. victoris* are highly deciduous and easily missing when trawled, most scales are well preserved in the 43.3 mm HL specimen examined (BSKU 45545; Figs. 9A, 10A–C). In *A. victoris*, the dorsum scales below the interdorsal space are covered with short, erect, needle-like spinules in widely spaced parallel rows, and the morphology is similar to those of *Kumba* Marshall, 1973. In contrast, the scales on the dorsal surface of the head are covered with a single row of erect spinules (Fig. 11E–F) that gives a striated appearance to the area (Fig. 12). This characteristic striation is somewhat similar to that found in *Mesovagus* Nakayama & Endo, 2016 (replacement name for *Mesobius* Hubbs & Iwamoto, 1977). In addition, specimens of *A. victoris* are fragile and easily damaged when trawled, which caused an apparent difference in coloration (see the Supplementary description and Fig. 9).

In the original description of *A. victoris*, Sazonov & Shcherbachev (1982) described that “there are small pores in the ... preopercular canal (2 in the lower part)”. However, the number was not confirmed due to their poor condition. Additional specimens from Japan revealed that the preopercular canal has not only two (or rarely three) pores on its lower part, but also a single pore on its upper end (Fig. 13A).

Sexual dimorphism of olfaction is well known in bathypelagic fishes such as ceratoid anglerfishes and several gonostmatids (*Cyclothone* spp.), viz., males are macrosmatic with large modified olfactory organs, whereas those of females are poorly developed (e.g., Marshall 1967, 1979; Herring 2002). Although a few grenadiers are exclusively confined to the mid-water depth, most species are bathypelagic and closely associated with the sea floor. Marshall (1967), who studied the olfactory organs of various bathypelagic fishes, concluded that both sexes of bathypelagic grenadiers are microsmatic. He (1979:409) also noted that “in the bathypelagic fish fauna olfactory dimorphism seems to be confined to the halosaurs”.

FIGURE 14. Scatter diagrams showing relationships of the (A) length of the posterior nostril and (B) length of the olfactory bulb (as % HL) to HL (mm) in *Asthenomacrus victoris*. Circles and squares indicate males and females respectively.

Sexual dimorphism of the olfactory organ in grenadiers was first suggested by Merrett *et al.* (1983), who wrote that “males of several macrourid species are found to be macrosmatic (e.g., *Trachyrinchus trachyrinchus* (Risso, 1810); *Bathygadus melanobranchus* Vaillant, 1888; *Coryphaenoides (Coryphaenoides) güntheri* (Vaillant, 1888) and *Nezumia aequalis* (Günther, 1887)—personal observation, N.R.M.)”. Unfortunately, they did not specify which characters show the sexually dimorphic modification. The results of this study confirmed such a dimorphism expressed in *A. victoris* (see the Sexual dimorphism entry above).

Iwamoto & Orlov (2006) redescribed *Pseudonezumia fragellicauda* (Koefoed, 1927) (as *Paracetonurus fragellicauda*) based on its four syntypes and 44 specimens collected from the northeastern Atlantic and the Madagascar Plateau in the southwestern Indian Ocean. Two of these specimens collected from the Porcupine Seabight (BMNH 1996.8.12.2–3, 26.5–29.6 mm HL, 144+–145+ mm TL) are re-identified here as *A. victoris* by having a broad naked area on the dorsal surface of the snout (vs. completely scaled in *P. fragellicauda*) and short, thick pyloric caeca (vs. slender and long).

FIGURE 15. Olfactory bulbs of *Asthenomacrus victoris*. (A) BSKU 85667, male, 44.1 mm HL; (B) BSKU 85666, female, 45.9 mm HL. [Photos: N. Nakayama]

Genus *Cetonurus* Günther, 1887

[Japanese name: Dango-hige-zoku]

Cetonurus Günther, 1887:143 (as subgenus of *Macrurus*; type species: *Coryphaenoides crassiceps* Günther, 1878, by original designation).

Diagnosis. Anus immediately anterior to anal-fin origin, surrounded by narrow but distinct periproct. Ventral light organ absent. Infraorbital ridge not connected with preopercular ridge, separated by distinct gap. Pelvic fin with 9–13 soft rays, its origin distinctly anterior to vertical through pectoral-fin base. Leading edge of second spinous ray of first dorsal fin weakly serrated. Head massive, soft, bones weak. Snout high, its tip above horizontal through midorbit. Chin barbel rudimentary. Teeth in narrow tapered bands in both jaws, none especially enlarged. Body scales covered with long, erect, needle-like spinules in quincunx order; buttresses of scale spinules well developed; reticulate structure developed over entire surface of unexposed portion. Scales along anterior part of second dorsal fin enlarged. Grooved lateral line interrupted, occurring as short segments. No modified scales along head ridges; tip and lateral angles of snout lacking prominent scutes; head almost completely scaled; scaly patches usually present on gular and branchiostegal membranes. Cephalic sensory canals broad, without open pores. Pyloric caeca 9–15. Branchiostegal rays 7. Body dark to gray overall, lacking

prominent silvery reflections when fresh. [Modified from Sazonov & Shcherbachev (1985).]

Remarks. The genus *Cetonurus* comprises two valid species: *C. crassiceps* (Günther, 1878), the type species known from the South Atlantic, Kermadec Islands (type locality), and Hawaii; and *C. globiceps* (Vaillant in Filhol, 1884) widespread in tropical and temperate waters of the Atlantic and Indo-West Pacific including Japan. A nominal species, *C. robustus* Gilbert & Hubbs, 1916 originally described from Japan, was synonymized with *C. globiceps* by Marshall (1973). Sazonov & Shcherbachev (1985) provided a synopsis of the genus.

Cetonurus globiceps (Vaillant in Filhol, 1884)

[Japanese name: Dango-hige]

(Figs. 16–17; Appendix 3-1B)

Macrurus globiceps Vaillant in Filhol, 1884:183 (original description; lectotype: MNHN 1886-92, from Spanish Sahara, 23°57'N, 17°12'W, in 1400–1435 m; 8 paralectotypes); Eschmeyer 1998:653 (lectotype designation).

Cetonurus globiceps: Goode & Bean 1896:411 (new combination); Marshall 1973:613, fig. 35 (*C. robustus* a junior synonym); Shao *et al.* 2008b: table 2 (3 spec. listed from southeastern Taiwan and South China Sea; first record from Taiwan); Iwamoto *et al.* 2015:45, fig. 4 (brief description; 5 spec. from Taiwan and Philippines).

Cetonurus robustus Gilbert & Hubbs, 1916:207, pl. 11, fig. 2 [original description; holotype: USNM 76870, from “off the east coast of central Hondo” (off Cape Shionomisaki), *Albatross* sta. 4971, in 649 ftn (1187 m); 5 paratypes from *Albatross* sta. 4791 and 4973]; Okada & Matsubara 1938:453 (in key; Japan); Matsubara 1955:1317 (in key; Japan; new Japanese name: “Dango-hige”); Matsubara 1965:510 (compiled; Japan); Okamura 1970a:110, pl. XXIV, text-fig. 46 (description based on 1 spec. from Molucca Sea); Okamura 1970b: table 1 (listed; Japan); Tominaga & Uyeno 1981:488 (listed; Japan); Okamura 1984b:94, pl. 345, fig. E (compiled); Okamura 1988:94, pl. 345, fig. E (compiled); Nakabo 1993:357 (in key; Japan); Nakabo 2000:421 (in key; Japan); Nakabo 2002:420 (in key; Japan); Nakabo & Kai 2013:497 (in key; Japan); Motomura 2020:38 (listed; Japan).

Diagnosis. Differs from its only congener, *C. crassiceps*, in having the following combination of features: orbit diameter 21–34% HL; premaxillary teeth in 3–4 rows, with outer series slightly enlarged; mandibular teeth in 2–4 rows throughout; tail fully scaled along anal-fin base; transverse scale rows below first dorsal-fin origin 13–19 (usually 16–18), below second dorsal-fin origin 10–19 (usually 14–15). [From Sazonov & Shcherbachev (1985).]

Material examined. 7 specimens. **Holotype of *Cetonurus robustus*:** USNM 76870 (57.5 mm HL, 280+ mm TL), off Cape Shionomisaki, Wakayama Pref., Kii Peninsula, Japan, 33.3917°N, 135.5667°E, 649 ftn (1187

FIGURE 16. Preserved specimen of *Cetonurus globiceps*. USNM 76870, holotype of *C. robustus*, 57.5 mm HL, 280+ mm TL, off Cape Shionomisaki, Wakayama Pref., Pacific, 1187 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

FIGURE 17. Scanning electron micrographs showing a body scale (from the dorsum below the interdorsal space) of *Cetonurus globiceps*. USNM 76871, paratype of *C. robustus*, 56.6 mm HL. (A) View from above; (B) oblique view. [Photos: N. Nakayama]

m), *Albatross* sta. 4971, 9-ft. Agassiz beam trawl, 30 Aug. 1906. **Paratypes of *C. robustus*:** USNM 76874 (1, ca. 29 mm HL, 113+ mm TL), off Cape Shionomisaki, 33.4042°N, 135.5083°E, 600 ftm (1098 m), *Albatross* sta. 4973, 9-ft Agassiz beam trawl, 30 Aug. 1906; CAS-SU 22923 (2, 48.0–60.3 mm HL, 241+–250+ mm TL), USNM 76871 (2, 46.9–59.6 mm HL, 261+–280+ mm TL), off Cape Shionomisaki, 33.3917°N, 135.5667°E, 649 ftm (1187 m), *Albatross* sta. 4971, 9-ft Agassiz beam trawl, 30 Aug. 1906. **Non-type: Japan:** HUMZ 146791 (1, 42.2 mm HL, 142+ mm TL), west of Amami-oshima Island, Okinawa Trough, 28.5548°N, 127.1660°E, 996–999 m, 16 Jul. 1994.

Counts and measurements. Based on 7 specimens (ca. 29–60.3 mm HL, 113+–280+ mm TL). Counts:

first dorsal-fin rays II,9–11; pectoral-fin rays i14–i19; pelvic-fin rays 9–10; gill rakers on first arch (outer/inner) 5–12/13–14, on second arch 12–14/11–14; longitudinal scales 59–70; transverse scale rows below first dorsal-fin origin 14–17, below first dorsal-fin midbase 16–18, below second dorsal-fin origin 12–13.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 33–38 (47–55); orbit diameter 23–27 (32–39); postorbital length 45–51 (65–70); postrostral length 69–73; orbit–preopercle distance 47–52 (68–72); suborbital width 22–30 (32–41); upper-jaw length 29–33 (41–45); length of rictus 23–25 (33–36); length of premaxillary tooth band 11–14 (16–19); preoral length 24–34 (33–47); distance between tip and lateral angle of snout 22–28 (32–41); snout width 38–44

(54–65); internasal width 33–43 (47–63); interorbital width 40–41 (56–59); body width over pectoral-fin bases 34–43 (49–59); body depth at first dorsal-fin origin 69–80 (94–114); body depth at anal-fin origin 64–68 (92–97); prepelvic length 98–111 (142–159); preanus length 114–126 (166–181); preanal length 120–130 (175–186); isthmus–pelvic distance 28–33 (41–45); isthmus–anus distance 49 (71); isthmus–anal distance 55 (78–81); pelvic–anal distance 28–31 (40–45); anus–anal distance 7 (10–11); pectoral-fin length 36 (52); predorsal length 100–104 (140–151); height of first dorsal fin 54 (77); length of first dorsal-fin base 17–20 (24–27); interdorsal length 28–38 (41–53); length of gill slit 9–12 (13–17); length of posterior nostril 4–6 (6–8); barbel length 3–5 (4–7).

Size. Attains at least 51 cm TL (Sazonov & Shcherbachev 1985).

Distribution. Widespread in tropical and temperate waters of the world's oceans at depths of 860–4621 m (Sazonov & Shcherbachev 1985; Iwamoto & Graham 2001). In Japanese waters, known only from off the Kii Peninsula and northern Okinawa Trough at depths of 996–1187 m (Appendix 3-1B). Very rare in Japan, but seemingly less rare in other parts of its range.

Remarks. *Cetonurus globiceps* is quite rare in Japanese waters, and its occurrence was previously based on only the type specimens of *C. robustus* Gilbert & Hubbs, 1916, a junior synonym of the former species. The latter species was originally described from six specimens collected from off Cape Shionomisaki, Wakayama Pref. An additional specimen from the Okinawa Trough (HUMZ 146791, heavily damaged) represents the first record of *C. globiceps* from the East China Sea and the second confirmed record of the species from Japanese waters. For further morphological information see Gilbert & Hubbs (1916; original description of *C. robustus*).

Genus *Coelorinchus* Giorna, 1809

[Japanese name: Tojin-zoku]

Coelorinchus Giorna, 1809:179 [type species: *Lepidoleprus caelorhincus* Risso, 1810 (= *Coelorinchus caelorhincus*), by subsequent designation by Fowler (1936)].

[?] *Krohnii* Cocco, 1844:22 (type species: *Krohnii filamentosus* Cocco, 1844, by monotypy).

[?] *Coccolus* Bonaparte, 1846:47, 97 (type species: *Coccolus annectens* Bonaparte, 1846, by monotypy).

Paramacrurus Bleeker, 1874:370 (type species: *Lepidoleprus australis* Richardson, 1839, by original designation).

Oxymacrurus Bleeker, 1874:370 (type species: *Macrurus japonicus* Temminck & Schlegel, 1846, by original designation).

Abyssicola Goode & Bean, 1896:417 (type species: *Macrurus macrochir* Günther, 1877, by original designation).

Quincuncia Gilbert & Hubbs, 1920:432 (as subgenus of *Coelorhynchus*; type species: *Coelorhynchus argentatus* Smith & Radcliffe in Radcliffe, 1912, by original designation).

Oxygadus Gilbert & Hubbs, 1920:515 (as subgenus of *Coelorhynchus*; type species: *Macrurus parallelus* Günther, 1877, by original designation).

Mahia McCann & McKnight, 1980:53 (type species: *Mahia matamua* McCann & McKnight, 1980, by original designation).

Diagnosis. Infraorbital and preopercular ridges tightly adjoined, forming continuous bony ridge; preopercular ridge elongate posteroventrally, sharply pointed at angle. Snout long, acute in lateral view in most species, tipped with stout scute. Scales on head ridges usually coarsely modified. Pelvic-fin rays almost always 7 (rarely 6); pelvic fin inserted below or posterior to vertical through pectoral-fin base. Leading edge of second spinous ray of first dorsal fin smooth (a few denticles occasionally found in distal portion in small individuals). Anus usually abutting or only slightly separated from anal-fin origin; periproct absent or rudimentary. Light organ present in most species. Chin barbel present and variously developed. No gill rakers on outer side of first gill arch (a few rudimentary rakers rarely found in large adults). Body scales large, spinules variable in shape; reticulate structure absent on unexposed portion (except for *C. immaculatus* Sazonov & Iwamoto, 1992). Grooved lateral line complete. Cephalic sensory canals without open pores. Branchiostegal rays 7. Color variable, some species with prominent body markings.

Remarks. Following Art. 32.5.2 of the ICZN (1999) and comments by Iwamoto (2008a), the original spelling of the generic name “*Caelorinchus*” is corrected to “*Coelorinchus*”; other subsequent spellings, including “*Caelorinchus*”, “*Coelorhynchus*”, and “*Caelorhynchus*”, are all considered here as incorrect subsequent spellings [see also Art. 33.3 of the ICZN (1999)]. *Coelorinchus* is by far the largest genus of the family, including about 120 valid species, with many species awaiting description (Iwamoto & Williams 1999; Iwamoto *et al.* 2006, 2015; Nakayama & Endo 2017). Of these, 25 species are herein recorded from Japan and adjacent waters. According to a recent molecular phylogeny (Roa-Varón & Ortí 2009), *Macrourus* Bloch, 1786 [“*Macrurus*” appeared in early literature but should be regarded as an incorrect subsequent spelling of this name; see also Art. 33.3 of the ICZN (1999)] is deeply nested within *Coelorinchus*, suggesting that the two genera are synonymous. In terms of morphology, *Coelorinchus* differs from *Macrourus* in that the second spinous ray of the first dorsal fin is smooth along its leading edge (vs. serrated in *Macrourus*), but otherwise they are very similar. A more in-depth study is therefore required to revise their taxonomic status.

Key to species of *Coelorinchus* from Japan and adjacent waters

- 1a** Anus distinctly separated from anal-fin origin, situated about midway between inner pelvic-fin bases and anal-fin origin ..
..... **2**
- 1b** Anus immediately anterior to or only slightly separated from anal-fin origin..... **3**
- 2a** Mandibular teeth in 2 irregular rows; lateral nasal ridge soft, incompletely supported by nasal bone; suborbital ridge distinctly curved anteriorly in lateral view; body scales covered with moderately erect, knife-like to keel-like spinules in widely divergent rows..... *C. macrochir*
- 2b** Mandibular teeth in narrow band; lateral nasal ridge stout, completely supported by nasal bone; suborbital ridge straight from tip of snout to preopercle in lateral view; body scales covered with greatly reclined, needle-like spinules in tightly packed parallel rows..... **5**
- 3a** Light organ externally discernible, its anterior margin extending to line connecting outer pelvic-fin bases or beyond (duct connecting anterior and posterior fossae often difficult to distinguish externally in species of the *C. hubbsi* group; see the Relationships of *C. brevirostris*) **4**
- 3b** Light organ either discernible or not evident externally; if discernible, its anterior margin not reaching (usually far posterior to) line connecting outer pelvic-fin bases..... **14**
- 4a** Lateral nasal ridge completely supported by nasal bone; anterior margin of light organ extending to line connecting outer pelvic-fin bases or only slightly beyond; snout relatively short, length 50–66% PRL **5**
- 4b** Lateral nasal ridge incompletely supported by nasal bone; anterior margin of light organ extending well beyond line connecting outer pelvic-fin bases and almost reaching isthmus; snout relatively long, 60–90% PRL..... **6**
- 5a** Small rudimentary dark spot between lateral line and pectoral-fin base (often difficult to distinguish or absent); anterior margin of light organ extending anteriorly beyond line connecting outer pelvic-fin bases; anus only slightly separated from anal-fin origin; barbel length 12–19% PRL ..
..... *C. jordani*
- 5b** Prominent large dark blotch between lateral line and pectoral-fin base; anterior margin of light organ situated on line connecting outer pelvic-fin bases; anus situated about midway between inner pelvic-fin bases and anal-fin origin; barbel length 7–13% PRL *C. kishinouyei*
- 6a** Anterior rays of second dorsal fin distinctly shorter than those of anal fin; interdorsal space usually greater than first dorsal-fin base length; premaxillary teeth in long, tapered band, with posterior margin of tooth band almost reaching lateral corner of mouth **7**
- 6b** Anterior rays of second dorsal fin about equal in height to those of anal fin; interdorsal space usually less than first dorsal-fin base length; premaxillary teeth in short, uniformly wide band, with posterior margin of tooth band falling far short of lateral corner of mouth..... **10**
- 7a** Underside of head posterior to vertical through midorbit heavily scaled, but mandibular rami mostly naked anteriorly; spinules on body scales trihedral *C. formosanus*
- 7b** Underside of head including mandibular rami usually completely naked (a few scales or rudimentary scaly patches occasionally found on ventral surfaces of preopercles); spinules on body scales needle-like or narrowly triangular...
..... **8**
- 8a** Short black hair-like papillae (not free neuromasts) densely scattered over underside of head; spinules on body scales narrowly triangular; length of gill slit 26–29% PRL
..... *C. kamoharai*
- 8b** Short black hair-like papillae (not free neuromasts) absent on underside of head; spinules on body scales needle-like; length of gill slit 20–26% PRL..... **9**
- 9a** Spinules on body scales arranged in subparallel to narrowly divergent rows; dark body markings not prominent, often difficult to distinguish; height of first dorsal fin 58–70% PRL; barbel length 11–18% PRL *C. longissimus*
- 9b** Spinules on body scales arranged in quincunx order; dark vermiculations or small irregular spots prominently marked dorsally on body; height of first dorsal fin 81–100% PRL; barbel length 16–24% PRL *C. multispinulosus*
- 10a** Body pale overall, lacking prominent markings; second spinous ray of first dorsal fin greatly elongate, height of first dorsal fin 139–220% PRL; when viewed dorsally lateral contour of snout broadly convex anteriorly, giving spade-shaped appearance to snout; chin barbel short, length 4–10% PRL; spinules on scales anterior to light organ rudimentary .
..... *C. brevirostris*
- 10b** Body with prominent dark saddles, diagonal bars or blotch present on body, or if markings absent, body uniformly dark overall; first dorsal fin high, but second spinous ray not especially elongate, height of first dorsal fin 80–130% PRL (except for *C. cingulatus*, 107–156% PRL); when viewed dorsally, distal portion of snout narrowly attenuated, giving sharply pointed appearance to snout (except for juveniles); chin barbel moderately developed, length 9–19% PRL; spinules on scales anterior to light organ not rudimentary
..... **11**
- 11a** Longitudinally elongate dark blotch between lateral line and pectoral fin, extending from above pectoral-fin base to below about second dorsal-fin origin; narrow dark stripe extending mid-laterally from pectoral-fin base to middle of tail or beyond..... *C. hubbsi*
- 11b** Body color different from above..... **12**
- 12a** Prominent dark blotch between lateral line and pectoral-fin base; fine dark vermiculations over dorsal surface of head (uniformly dark brown in preserved specimens)
..... *C. matsubarae*
- 12b** No prominent dark blotch between lateral line and pectoral-fin base; fine dark vermiculations absent on dorsal surface of head **13**
- 13a** Second spinous ray of first dorsal fin greatly elongate (except for juveniles), height of first dorsal fin 107–156% PRL; snout length 79–86% PRL; dorsal profile of snout distinctly concave in lateral view; barbel length 10–13% PRL *C. cingulatus*

- 13b Second spinous ray of first dorsal fin not elongate, height of first dorsal fin 92–106% PRL; snout length 60–72% PRL; dorsal profile of snout almost straight to gently concave in lateral view; barbel length 13–17% PRL *C. fuscigulus*
- 14a Body scales covered with keel-like spinules in divergent rows; every spinule row complete, extending to posterior scale margin, giving harsh appearance to body surface; middle row not especially high and enlarged compared with adjacent rows; light organ externally discernible; anus slightly separated from anal-fin origin by a few scale rows 15
- 14b Body scales covered with shield-like spinules in parallel to subparallel rows; lateral rows often incomplete, not extending to posterior scale margin; middle row notably higher, giving striated appearance to body surface; no external evidence of light organ; anus abutting anal-fin origin 24
- 15a Underside of head mostly or completely scaled 16
- 15b Underside of head almost entirely naked (oval scaly patches present on lower surfaces of preopercles in *C. hexafasciatus* and *C. tokiensis*) 20
- 16a About 6 prominent dark saddles on trunk and tail; lateral nasal ridge incompletely supported by nasal bone *C. sheni*
- 16b Dark saddles usually absent on body, if present, not prominent and restricted to tail; lateral nasal ridge completely supported by nasal bone 17
- 17a Underside of snout broadly naked (small to moderate scaly patches often present); internasal width 19–24% PRL, interorbital width 27–31% PRL *C. leptorhinus*
- 17b Underside of snout almost completely scaled, naked area confined along ventromedial line and/or narrow band above upper lip; internasal width 26–37% PRL, interorbital width 31–42% PRL 18
- 18a Occipital scales usually covered with single row of spinules (rarely 2–3 rows) *C. japonicus*
- 18b Occipital scales covered with 4 or more rows of spinules 19
- 19a Spinules on occipital scales needle-like, erect, completely free from one another in each row, forming comb-like ridges; head and body uniformly dark brown *C. nox* sp. nov.
- 19b Spinules on occipital scales keel-like (rarely knife-like), moderately reclined, basally adjoined from one another in various degrees in each row, forming saw-toothed ridges; in Japanese specimens, head and body darker dorsally, but distinctly paler ventrally *C. smithi*
- 20a Several dark saddles on trunk and tail; lower surfaces of preopercles scaled; lateral nasal ridge incompletely supported by nasal bone; outer series of premaxillary tooth band enlarged 21
- 20b No prominent dark saddles on trunk and tail; lower surfaces of preopercles completely or almost completely naked; lateral nasal ridge completely supported by nasal bone; outer series of premaxillary tooth band not enlarged 22
- 21a Snout relatively short, length 57–70% PRL *C. hexafasciatus*
- 21b Snout relatively long, length 70–85% PRL *C. tokiensis*
- 22a Spinules on occipital scales needle-like, forming comb-like ridges; nasal fossa usually naked (a few scales rarely found anteroventrally); interdorsal length 51–71% PRL *C. hige*
- 22b Spinules on occipital scales keel-like, forming saw-toothed ridges; nasal fossa usually heavily scaled (only sparsely scaled in some specimens); interdorsal length 33–54% PRL 23
- 23a Snout relatively short, length 60–88% PRL; length of terminal snout scute 7–18% PRL; height of first dorsal fin 85–107% PRL *C. anatirostris*
- 23b Snout very long, length 107% PRL; length of terminal snout scute 27% PRL; height of first dorsal fin 84% PRL *C. lanceolatus* sp. nov.
- 24a Underside of head almost entirely naked; nasal fossa usually lacking scales; free neuromasts on underside of head modified into prominent, knob-like or flap-like projections *C. gilberti*
- 24b Underside of head almost completely scaled; nasal fossa usually heavily scaled; free neuromasts on underside of head not especially enlarged 25
- 25a Occipital scales covered with divergent rows of spinules *C. divergens*
- 25b Occipital scales covered with single row of spinules *C. parallelus*

Coelorinchus anatirostris Jordan & Gilbert in Jordan & Starks, 1904

[Japanese name: Nezumi-hige]

(Figs. 18–20, 21A, 22–25; Table 1; Appendix 3-1C)

Coelorhynchus anatirostris Jordan & Gilbert in Jordan & Starks, 1904:619, unnumbered fig. (original description; holotype: USNM 51471, from Misaki); Jordan *et al.* 1913:418, fig. 388 (listed; Japan; new Japanese name: “Nezumi-hige”); Schmidt 1931:155 (brief description; 3 spec. from Nagasaki); Kamohara 1934a:54 (listed; Kochi); Okada & Matsubara 1938:450 (in key; Japan); Kamohara 1938:73 (1 spec. from Kochi Pref.); Kamohara 1950:277 (listed; Kochi and Wakayama Pref.); Kuroda 1951:391 (listed; Suruga Bay); Kamohara 1952:99 (spec. from Kochi Pref.); Kuroda 1952:177 (2 spec. from Suruga Bay); Matsubara 1955:1313 (in key; Japan); Kamohara 1958:73 (listed; Kochi Pref.); Okada *et al.* 1959:83 (listed; Kumano-nada); Kamohara 1964:95 (listed; Kochi Pref.); Matsubara 1965:506 (compiled; Japan); Okamura 1970a:186, pl. VIII, text-fig. 80 (description; 72 spec. from Pacific off southern Japan from Heta to Tosa Bay; *C. productus* a junior synonym of *C. anatirostris*); Kataoka & Tomida 1981:78 (listed; Mie Pref.); Tominaga & Uyeno 1981:488 (listed; Japan); Okamura 1982:171, 352, fig. 102 (brief description; 2 spec. from Tosa Bay; photo based on BSKU 29586); Ozawa 1983:13 (listed; off Makurazaki, Kagoshima Pref., East China Sea).

Coelorhynchus antirostris [sic]: Franz 1910:27 (misspelled; 1 spec. from Aburatsubo); Matsubara *et al.* 1951:42 (misspelled; listed; Mie Pref.).

Coelorhynchus productus Gilbert & Hubbs, 1916:175, pl. 9, fig. 1 [original description; holotype: USNM 76865, from “Suruga

Gulf" (= Suruga Bay), *Albatross* sta. 5059; 6 paratypes from Suruga Bay]; Okada & Matsubara 1938:450 (in key; Japan; new Japanese name: "Tengu-hige"); Kamohara 1950:277 (listed; Kochi and Wakayama Pref.); Kuroda 1951:391 (listed; Suruga Bay); Kamohara 1952:99 (spec. from Kochi Pref.); Matsubara 1955:1313 (in key; Japan); Kobayashi 1956:72 (listed; Enshu-nada); Kamohara 1958:73 (listed; Kochi Pref.); Kamohara 1964:95 (listed; Kochi Pref.); Matsubara 1965:506 (compiled; Japan).

Coelorhynchus (Oxymacrus) anatirostris: Okamura 1970b: table 1 (listed; Japan).

Coelorrhinchus anatirostris: Yatou 1984:233, 368, fig. 165 [brief description; 9 spec. from Okinawa Trough; photo based on BSKU 37101 (from Mimase fish market)]; Okamura 1984b:97, pl. 83, fig. E (compiled); Okamura 1988:97, pl. 83, fig. E (compiled); Iwamoto 1990:141, fig. 346 (synopsis); Okamura 1997:127, fig. 11 (compiled); Shao *et al.* 2008b: table 2 (5 spec. listed from northeastern and southwestern Taiwan); Nakabo & Kai 2013:511 (in key; Japan); Fukui *et al.* 2015:182, fig. 9 (13 spec. from off Kuro-shima Island, Kagoshima Pref., East China Sea); Iwamoto *et al.* 2015:50 (brief description; 20 spec. from northeastern and southwestern Taiwan, and South China Sea); Motomura 2020:38 (listed; Japan).

Coelorrhinchus productus: Yatou 1984:233, 369, fig. 166 (brief description; 6 spec. from Okinawa Trough; photo based on BSKU 29638); Okamura 1988:435, pl. 373, fig. I (compiled); Iwamoto 1990:179, fig. 406 (synopsis); Okamura 1997:127, fig. 12 (compiled); Shao *et al.* 2008b: table 2 (5 spec. listed from northeast of Taiwan); Nakabo & Kai 2013:511 (in key; Japan); Iwamoto *et al.* 2015:66 (brief description; 5 spec. from northeastern Taiwan); Motomura 2020:38 (listed; Japan).

Coelorrhinchus productus: Nakabo 1993:370 (in key; Japan); Shinohara & Matsuura 1997:291 (listed; Suruga Bay); Nakabo 2000:434 (in key; Japan); Shinohara *et al.* 2001:304 (listed; Tosa Bay); Nakabo 2002:434 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Chiou *et al.* 2004a:44, fig. 11 (brief description; 2 spec. from Da-xi; first record from Taiwan); Shinohara *et al.* 2005:417 (12 spec. listed from Ryukyu Islands); Suetsugu & Ohta 2005: table 2 (listed; southeast of Boso Peninsula).

Coelorrhinchus anatirostris: Nakabo 1993:370 (in key; Japan); Shao 1993:168, fig. 37-6 (compiled; Taiwan; photo spec. is *C. multispinulosus*); Shinohara & Matsuura 1997:290 (listed; Suruga Bay); Nakabo 2000:434 (in key; Japan); Shinohara *et al.* 2001:304 (26 spec. listed from Tosa Bay); Nakabo 2002:434 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Shinohara *et al.* 2005:416 (11 spec. listed from Ryukyu Islands); Senou *et al.* 2006:420 (listed; Sagami Sea); Furuhashi *et al.* 2010: table 2 (268 spec. listed from northern Okinawa Trough).

Coelorrhinchus (Oxymacrus) anatirostris: Chiou *et al.* 2004a: table 1 (listed; Taiwan).

Coelorrhinchus (Oxymacrus) productus: Chiou *et al.* 2004a: table 1 (listed; Taiwan).

Coelorrhinchus gilberti (not Jordan & Hubbs 1925): Shinohara *et al.* 2005:416 (3 spec. listed from Ryukyu Islands).

Diagnosis. Light organ externally represented by short, narrow, naked streak immediately anterior to anus, its anterior margin falling far short of line connecting inner pelvic-fin bases. Underside of head almost completely naked; dorsal surface of snout fully scaled, except narrow naked clefts along each side of median rostral ridge. Nasal fossa usually heavily scaled. Snout long, sharply pointed, length 60–88% PRL, becoming proportionally shorter with growth; dorsal profile of snout slightly concave in lateral view. Terminal snout scute short, broad, arrowhead-shaped, dorsoventrally flattened, length 7–18% PRL. Scales on median rostral ridge covered with divergent rows of spinules. Lateral nasal ridge completely supported by nasal bone, not strongly convex when viewed dorsally. Anus slightly separated from anal-fin origin. Premaxillary teeth small, conical, arranged in short, uniformly wide band, none especially enlarged; posterior margin of tooth band falling far short of lateral corner of mouth. Body scales covered with short, reclined, keel-like spinules in narrowly divergent saw-toothed ridges; every spinule row complete, extending to posterior scale margin; spinules in each row greatly overlapping, closely adjoined to one another, but generally free at tips; buttresses narrowly developed. Occipital scales covered with short, reclined, keel-like spinules in narrowly divergent, saw-toothed rows. Orbit diameter 45–52% PRL; postorbital length 50–56% PRL; orbit–preopercle distance 52–60% PRL; height of first dorsal fin 85–107% PRL; interdorsal length 33–54% PRL; pyloric caeca 21–26. Free neuromasts on underside of head tiny, immaculate, difficult to distinguish; dark hair-like papillae absent on underside of head. Body darker dorsally, paler ventrally, without either blotches or saddles; underside of head dusky; lips white; oral cavity blackish; gular and branchiostegal membranes pale; pelvic fins dusky.

Material examined. 56 specimens. **Holotype of *Coelorhynchus anatirostris*:** USNM 51471 (93.7 mm HL, 395 mm TL), Misaki, Kanagawa Pref., Sagami Bay, Japan, date unknown. **Holotype of *Coelorhynchus productus*:** USNM 76865 (79.9 mm HL, 304 mm TL), off Fuji River, Shizuoka Pref., Suruga Bay, Japan, 35.0917°N, 138.6639°E, 197–297 ftm (361–544 m), *Albatross* sta. 5059, 9-ft *Albatross-Blake* beam trawl, 13 Oct. 1906. **Paratypes of *C. productus*:** USNM 76872 (1, 55.6 mm HL, 233+ mm TL), off Fuji River, Suruga Bay, 35.1014°N, 138.6722°E, 221–293 ftm (405–536 m), *Albatross* sta. 5066, 9-ft *Albatross-Blake* beam trawl, 15 Oct. 1906; CAS-SU 22977 (2, 68.2–68.2 mm HL, 260+–286 mm TL), USNM 76873 (3 of 4, 53.0–58.5 mm HL, 186+–243+ mm TL, counts and measurements taken on 2 of 3), collected with holotype. **Non-types: Japan:** BSKU 29744 (1, 80.2 mm HL, 269+ mm TL), north-northeast of Miyako-jima Island, Okinawa Trough, 25.9833°N, 125.8500°E, 430 m, F/V *Yuryo-maru*, No. 8, sta. 2-T3, bottom trawl, coll. H. Maeda, 10 Oct. 1979; BSKU 34351 (1, 58.1 mm HL, 199+ mm TL), northwest of Kume-

FIGURE 18. Fresh specimens of *Coelorinchus anatirostris*. (A) BSKU 109032, 90.6 mm HL, 354+ mm TL, Okinawa Trough, East China Sea, 351–389 m depth; (B) BSKU 109033, 79.7 mm HL, 318+ mm TL, collected with BSKU 109032; (C–E) BSKU 102003, 73.1 mm HL, 283+ mm TL, Mimase fish market, Kochi, Pacific. (A–C) Lateral views; (D) dorsal and (E) ventral views of the head and trunk. [Photos: N. Nakayama]

jima Island, Okinawa Trough, 27.0083°N, 125.8417°E, 410–420 m, F/V *Yuryo-maru*, No. 8, sta. 5-T3, bottom trawl, coll. K. Uchida, 10 Dec. 1979; BSKU 29586 (1, 94.3 mm HL, 359+ mm TL), northwest of Miyako-jima Island, Okinawa Trough, 25.7917°N, 124.5117°E, 500–550 m, F/V *Yuryo-maru*, No. 8, sta. 1-T9, bottom trawl, coll. T. Kitajima, 15 Sept. 1979; BSKU 32516 (1, 91.6 mm HL, 318+ mm TL), northwest of Miyako-jima Island, Okinawa Trough, 25.7917°N, 124.5117°E, 500–550 m, F/

V *Yuryo-maru*, No. 8, sta. 1-T9, bottom trawl, coll. M. Hirose and T. Kume, 10 Oct. 1979; BSKU 29638 (1, 68.7 mm HL, 226+ mm TL), northwest of Miyako-jima Island, Okinawa Trough, 25.7900°N, 124.3333°E, 600 m, F/V *Yuryo-maru*, No. 8, sta. 1-T13, bottom trawl, coll. T. Kitajima, 20 Sept. 1979; BSKU 106770 (1, 72.7 mm HL, 278+ mm TL), BSKU 106821 (1, 62.5 mm HL, 219+ mm TL), BSKU 106823 (1, 63.8 mm HL, 250+ mm TL), BSKU 106824 (1, 55.7 mm HL, 210+ mm TL), southeast

FIGURE 19. Holotype of *Coelorhynchus anatirostris* (= *Coelorinchus anatirostris*). USNM 51471, 93.7 mm HL, 395 mm TL, Sagami Bay off Misaki, Kanagawa Pref., Pacific. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

FIGURE 20. Preserved specimen of *Coelorhynchus anatirostris*. USNM 76865, holotype of *Coelorhynchus productus*, 79.9 mm HL, 304 mm TL, Suruga Bay, Shizuoka Pref., Pacific, 544–361 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

of Shimokoshiki-jima Island, East China Sea, 31.5658°N, 129.8915°E, 380 m, F/V *Maruko-maru*, tr. 1, bottom trawl, coll. N. Nakayama *et al.*, 24 Apr. 2012; BSKU 106807 (1, 91.3 mm HL, 370+ mm TL), BSKU 106808 (1, 68.2 mm HL, 260+ mm TL), BSKU 106809 (1, 75.3 mm HL, 281+ mm TL), BSKU 106811 (1, 62.1 mm HL, 240 mm TL), southeast of Shimokoshiki-jima Island, East China Sea, 31.5652°N, 129.8878°E, 407 m, F/V *Maruko-maru*, tr. 3, bottom trawl, coll. N. Nakayama *et al.*, 24 Apr. 2012; BSKU 106777 (1, 58.9 mm HL, 230+ mm TL), BSKU

106779 (1, 57.1 mm HL, 191+ mm TL), BSKU 106794 (1, 55.4 mm HL, 208+ mm TL), BSKU 106796 (1, 51.0 mm HL, 195+ mm TL), BSKU 106797 (1, 68.4 mm HL, 259+ mm TL), southeast of Shimokoshiki-jima Island, East China Sea, 31.5560°N, 129.8847°E, 415 m, F/V *Maruko-maru*, tr. 4, bottom trawl, coll. N. Nakayama *et al.*, 24 Apr. 2012; KAUM-I. 32206 (1, 85.5 mm HL, 338+ mm TL), off Nomaie, Kagoshima, 31.5000°N, 129.8833°E, 370–400 m, trawl, coll. M. Yamashita and Y. Ohashi, 10 Sept. 2010; BSKU 109032 (1, 90.6 mm HL, 354+ mm

TABLE 1. Morphometric measurements and meristic counts* for *Coelorinchus anatirostris* and *C. lanceolatus* **sp. nov.**

Species	<i>C. anatirostris</i>				<i>C. lanceolatus</i>
Nominal species	<i>C. anatirostris</i>		<i>C. productus</i>		–
Type status	HT	NT	HT	PT	HT
Number of specimens	<i>n</i> = 1	<i>n</i> = 46	<i>n</i> = 1	<i>n</i> = 5	<i>n</i> = 1
Measurements (mm)					
TL	395	191+–427	304	186+–286	255+
HL	93.7	51.0–106	79.9	53.0–68.2	79.2
% of HL (% of PRL)					
Snout length	43 (73)	38–46 (60–84)	47 (88)	44–47 (79–86)	52 (107)
Orbit diameter	28 (48)	26–31 (45–52)	25 (48)	26–28 (48–50)	23 (47)
Postorbital length	32 (55)	28–34 (50–56)	29 (54)	28–29 (52–53)	26 (53)
Postrostral length	59 (–)	55–64 (–)	53 (–)	54–56 (–)	49 (–)
Orbit–preopercle distance	34 (58)	29–37 (52–60)	30 (56)	29–31 (52–57)	27 (55)
Suborbital width	15 (25)	12–16 (22–27)	14 (25)	11–12 (21–23)	11 (23)
Upper-jaw length	24 (41)	21–27 (37–45)	21 (39)	20–22 (37–41)	18 (38)
Length of rictus	17 (29)	15–20 (26–33)	14 (27)	14–16 (26–29)	– (–)
Length of PMX tooth band	8 (13)	6–10 (11–17)	7 (14)	7–8 (13–15)	8 (16)
Preoral length	38 (64)	32–43 (50–79)	43 (81)	41–42 (74–77)	49 (101)
Length of terminal snout scute	– (–)	5–10 (7–18)	– (–)	8–9 (14–16)	13 (27)
Length of lateral nasal ridge	30 (50)	23–33 (37–61)	34 (63)	31–34 (56–62)	39 (81)
Length of suborbital ridge	94 (159)	89–96 (142–175)	93 (174)	92–95 (166–173)	94 (194)
Snout width	30 (51)	26–32 (44–58)	28 (53)	26–31 (49–56)	25 (51)
Internasal width	19 (32)	17–20 (28–35)	18 (34)	16–18 (28–32)	16 (33)
Interorbital width	22 (37)	19–25 (35–43)	21 (39)	19–21 (34–38)	19 (38)
Occipital width	– (–)	8–10 (14–18)	– (–)	9 (16)	– (–)
Body width over P. bases	43 (73)	37–53 (67–87)	42 (78)	37–41 (68–74)	37 (77)
Body depth at 1D. origin	54 (92)	40–58 (73–99)	48 (90)	42–48 (76–87)	42 (86)
Body depth at A. origin	42 (71)	35–54 (61–91)	45 (85)	37–39 (66–72)	37 (77)
Pre-V. length	111 (188)	106–120 (184–213)	114 (214)	108–113 (199–203)	110 (227)
Pre-anus length	– (–)	132–157 (229–261)	– (–)	135 (247)	135 (278)
Pre-A. length	156 (263)	138–168 (239–280)	148 (278)	142–146 (257–266)	142 (292)
Isthmus–V. distance	32 (54)	23–36 (41–57)	28 (53)	23–29 (42–52)	22 (46)
Isthmus–anus distance	– (–)	47–75 (83–119)	– (–)	– (–)	39 (80)
Isthmus–A. distance	– (–)	54–86 (94–136)	– (–)	61–63 (112–117)	47 (96)
V.–A. distance	49 (82)	33–58 (57–92)	40 (74)	34–40 (62–75)	33 (69)
Anus–A. distance	– (–)	6–11 (10–19)	– (–)	– (–)	– (–)
V. length	40 (67)	31–44 (53–76)	34 (63)	33–37 (60–67)	34 (69)
P. length	42 (70)	34–48 (60–79)	37 (69)	37–40 (69–71)	– (–)
Pre-1D. length	115 (194)	106–115 (177–205)	109 (204)	107–112 (196–204)	104 (214)
Height of 1D.	59 (100)	48–65 (85–107)	52 (96)	52–55 (95–101)	41 (84)
Length of 1D. base	21 (35)	16–24 (29–41)	17 (32)	17–21 (31–37)	14 (30)
Interdorsal length	30 (50)	19–32 (33–54)	25 (46)	21–27 (37–50)	16 (34)
Length of gill slit	9 (16)	9–13 (15–22)	– (–)	9–10 (17)	9 (19)
Length of posterior nostril	9 (16)	8–12 (15–19)	9 (16)	7–8 (14–15)	7 (15)
Barbel length	7 (12)	5–12 (9–18)	– (–)	4–6 (8–11)	6 (13)
Counts					
1D. rays	II,9	II,8–10	II,8	II,7–9	II,8
P. rays	i17	i15–i19	i16	i16–i18	i18
V. rays	7	7	7	6–7	7
Outer GR on first arch	0	0–3	–	0	0
Inner GR on first arch	8	6–8	6	6–7	8
Outer GR on second arch	6	5–7	6	5–6	6
Inner GR on second arch	7	6–8	6	6–7	9
Longitudinal scales	28	26–37	32	32–35	41

...Continued on the next page

TABLE 1. (Continued)

Species	<i>C. anatirostris</i>				<i>C. lanceolatus</i>
Nominal species	<i>C. anatirostris</i>		<i>C. productus</i>		—
Type status	HT	NT	HT	PT	HT
Number of specimens	<i>n</i> = 1	<i>n</i> = 46	<i>n</i> = 1	<i>n</i> = 5	<i>n</i> = 1
Scales below 1D. origin	5.5	4.5–6.5	5	4.5–5.5	4.5
Scales below 1D. midbase	3.5	3.5–5.5	4.5	4.5	3.5
Scales below 2D. origin	4.5	4–6.5	4.5	4–5	5
Scales above A. origin	—	15–19	—	—	15

*Abbreviations: 1D.—first dorsal fin; 2D.—second dorsal fin; A.—anal fin; GR—gill rakers; HT—holotype; NT—non-type; P.—pectoral fin; PMX—premaxillary; PT—paratype; V.—pelvic fin

FIGURE 21. Preserved juvenile specimens of two species of *Coelorinchus*. (A) *C. anatirostris*, BSKU 50891, 21.7 mm HL, 88+ mm TL, Numazu fish market, Shizuoka Pref., Pacific; (B) *C. hige*, BSKU 50881, 30.0 mm HL, 108+ mm HL, off Nomaieke, Kagoshima Pref., East China Sea. Lateral views. [Photos: N. Nakayama]

TL), BSKU 109033 (1, 79.7 mm HL, 318+ mm TL), BSKU 109034 (1, 56.1 mm HL, 207+ mm TL), BSKU 109038 (1, 89.9 mm HL, 382+ mm TL), west of Sanposone, 32.2872°N, 129.0732°E, 351–389 m, T/V *Nagasaki-maru*, cr. N365, sta. B4, 3-m ORE beam trawl, coll. N. Nakayama, 19 Nov. 2012; BSKU 109019 (1, 62.2 mm HL, 256+ mm TL), west of Okikasayama Bank, 32.1503°N, 129.0072°E, 499–504 m, T/V *Nagasaki-maru*, cr. N365, sta. C5, 3-m ORE beam trawl, coll. N. Nakayama, 19 Nov. 2012; NSMT-P 65673 (1, 74.4 mm HL, 310+ mm TL), west of Yaku-shima Island, East China Sea, 30.4467°N, 128.2367°E, 499–500 m, FRV *Yoko-maru*, sta. D9, bottom trawl, coll. T. Kubodera *et al.*, 10 Nov. 2002; BSKU 29821 (1, 86.4 mm HL, 358+ mm TL), BSKU 34038 (1, 96.7 mm HL, 388+ mm TL), west of Yaku-shima Island, Okinawa Trough, 30.4433°N, 128.0177°E, 410–430 m, F/V *Yuryo-maru*, No. 8, sta. 3-T7, bottom trawl, coll. H. Maeda, 28 Oct. 1979; BSKU 27989 (1, 106 mm HL, 383+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 28.8333°N, 127.2333°E, 700–740 m, F/V *Ryoan-maru*, No. 28, tr. 59, bottom trawl, coll. Y. Kinoshita and S. Hagino, 16 Mar. 1978; BSKU 27927 (1, 94.9 mm HL, 377+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 28.7500°N, 127.1167°E, 520–542 m,

F/V *Ryoan-maru*, No. 28, tr. 58, bottom trawl, coll. Y. Kinoshita and S. Hagino, 16 Mar. 1978; BSKU 112539 (1, 100 mm HL, 399+ mm TL), off Aki, Tosa Bay, 380 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 12 Mar. 2014; BSKU 70897 (1, 85.6 mm HL, 359 mm TL), Tosa Bay, 33.2250°N, 133.6933°E, 440–460 m, FRV *Kotaka-maru*, otter trawl, 24 Aug. 2000; BSKU 71581 (1, 79.2 mm HL, 317+ mm TL), BSKU 97956 (1, 83.5 mm HL, 318+ mm TL), BSKU 97957 (1, 85.1 mm HL, 326+ mm TL), BSKU 97958 (1, 73.3 mm HL, 309+ mm TL), Tosa Bay, 400 m, FRV *Kotaka-maru*, otter trawl, 9 May 2000; BSKU 77981 (1, 79.5 mm HL, 292+ mm TL), BSKU 77982 (1, 81.3 mm HL, 314+ mm TL), Tosa Bay, 33.2000°N, 133.6317°E, 370–380 m, FRV *Kotaka-maru*, otter trawl, 8 Sept. 1998; BSKU 43815 (1, 78.9 mm HL, 310+ mm TL), BSKU 43832 (1, 51.1 mm HL, 206+ mm TL), Tosa Bay, 400 m, FRV *Kotaka-maru*, otter trawl, 11 Jun. 1987; BSKU 4385 (1, 98.4 mm HL, 427 mm TL), Mimase fish market, bottom trawl, coll. T. Kamohara, 19 Dec. 1954; BSKU 37101 (1, 77.0 mm HL, 301+ mm TL), Mimase fish market, bottom trawl, coll. Y. Machida, 20 Apr. 1982; BSKU 102701 (1, 58.5 mm HL, 204+ mm TL), Mimase fish market, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama, 21 Feb. 2010; BSKU 102691 (1, 78.4

FIGURE 22. Scanning electron micrographs showing scales of *Coelorinchus anatirostris*. (A–B) BSKU 4385, 98.4 mm HL; (C–F), BSKU 97958, 73.3 mm HL. (A–D) Scales from the dorsum below the interdorsal space; (E–F) from the top of the head between the parietal ridges. (A, C, E) Views from above; (B, D, F) oblique views. [Photos: N. Nakayama]

mm HL, 312 mm TL), Mimase fish market, F/V *Kosei-maru*, tr. 3, bottom trawl, coll. N. Nakayama, 21 Feb. 2010; BSKU 102003 (1, 73.1 mm HL, 283+ mm TL), Mimase fish market, F/V *Seiryō-maru*, bottom trawl, 7 Dec. 2009; BSKU 110034 (1, 74.2 mm HL, 294+ mm

TL), Suruga Bay, 34.7127°N, 138.4553°E, 262–434 m, F/V *Hinode-maru*, sta. 1, bottom trawl, coll. N. Nakayama and R. Misawa, 23 Apr. 2013; *BSKU 50891 (1, 21.7 mm HL, 88+ mm TL), Numazu fish market, coll. S. Ohashi, 15 Feb. 2009. **Taiwan:** ASIZP 65573 (1, 98.2 mm HL,

352+ mm TL), Gwu-san Island, Yilan, 606–653 m, bottom trawl, coll. H.-M. Yeh, 17 Apr. 2003. **Chesterfield and Bellona Plateaus:** CAS 86496 (1, 81.9 mm HL, 288+ mm TL), 19.6808°S, 158.7683°E, 600–615 m, *MUSORSTOM* 5, sta. CC383, 21 Oct. 1986.

FIGURE 23. Scatter diagrams showing relationships of the (A) snout length and (B) barbel length (as % PRL) to PRL (mm) in *Coelorinchus anatirostris* (circles) and *C. productus* (squares). Arrows indicate holotypes.

Redescription based on Japanese specimens.

General features are shown in Figs. 18–20, 21A. Counts and measurements are given in Table 1. Body deepest at first dorsal-fin origin, gradually tapering to long, laterally compressed tail. Trunk short, moderately compressed, width over pectoral-fin bases 1.0–1.3 in depth below first dorsal-fin origin. Head large, HL about 3.3–4.3 in TL. Snout long, protruding greatly beyond upper jaw, length usually 1.4–1.8 times as long as orbit diameter (1.2 times in BSKU 4385); dorsal contour of snout slightly concave in lateral view; lateral nasal ridge completely supported by nasal bone; anterolateral margins of snout moderately convex when viewed dorsally. Orbit large, greatest diameter 1.0–1.2 in postorbital length. Interorbital space almost flat, width 1.1–1.5 in orbit diameter. Mouth moderate in size, inferior, protrusible, upper-jaw length 1.0–1.4 in orbit diameter; posterior margin of maxilla extending to vertical through midorbit or beyond, but not

reaching hind margin of orbit; lateral corner of mouth moderately restricted by skin folds; lips thin, slightly papillose near tooth bands. Suborbital region divided by longitudinal bony ridge passing from tip of snout to posteroventral angle of opercle; its upper half almost vertical, lower half sharply inclined mesially. Preopercle large, posterior margin inclined, forming moderately angular lobe at posteroventral corner. Subopercle prolonged ventrally as slender flap; posterior margin deeply concave. Gill membranes broadly connected across, and attached mesially to isthmus, with slight posterior free fold. Outer gill slit moderately restricted by skin folds, length 2.3–3.2 in orbit diameter. Gill rakers small, low, tubercular, armed with short, fine spines; no rakers on outer side of first arch (a few rudimentary rakers occasionally found in large specimens) and inner side of fourth arch; gill filaments moderately long. Chin barbel slender, tapering into filamentous tip; length usually 2.6–5.6 in orbit diameter (6.5 in USNM 76872, paratype of *C. productus*).

Anus slightly separated from anal-fin origin by a few scale rows. Light organ externally represented by short, narrow, naked streak anterior to anus; its anterior margin reaching about midway between anus and outer pelvic-fin bases, and falling far short of line connecting inner pelvic-fin bases. Pyloric caeca 24–26, rete-gas gland complexes 4 (based on BSKU 106823 and 109038).

Teeth small, sharp, conical, gently incurved, in narrow bands in both jaws. Premaxillary teeth in short, uniformly wide band, with about 6–7 tooth rows near symphysis; posterior margin of tooth band falling far short of lateral corner of mouth; teeth becoming progressively smaller inwardly, and none especially enlarged. Mandibular teeth in broad tapered band, with about 5 tooth rows near symphysis, none especially enlarged; band narrowing abruptly, and its posterior margin extending beyond lateral corner of mouth. All teeth deeply embedded in thick layer of gum papillae.

Body scales large, thin, not deciduous, covered with narrowly divergent rows of short, reclined, keel-like spinules (Fig. 22A–D); in 90.1 mm HL specimen (BSKU 109032), those on dorsum below interdorsal space with 7–9 spinule rows; middle row longest with about 6–8 spinules, but not especially high or enlarged compared with adjacent rows; every spinule row complete throughout; spinules forming angle of about 40° to scale surface; spinules greatly overlapping, closely adjoined to one another, but distal tips usually free, forming saw-toothed ridges that give spiny appearance to body surfaces; spinules increasing in height posteriorly, with last in each row extending slightly beyond posterior scale margin (last spinule rarely falling short of scale margin in large specimens; Fig. 22A–B); buttresses of spinules narrowly developed; no reticulate structures on unexposed portion. Body fully scaled except for fins and ventral light organ.

FIGURE 24. Variation of the snout shape in *Coelorinchus anatirostris*. [Photos: N. Nakayama]

Scales on head ridges coarsely modified; those on median rostral ridge covered with divergent rows of spinules. Terminal snout scute short, broad, arrowhead-shaped, dorsoventrally flattened, its length 2.8–6.5 in orbit diameter. Supraoccipital scute stout, enlarged, armed with single row of spinules; posttemporal scutes thickened, but not especially enlarged. Occipital scales covered with short, reclined, keel-like spinules in narrowly divergent rows (Fig. 22E–F); spinules slightly adjoined basally to one another, and increasing in height posteriorly; scales on dorsal surfaces of snout posterior to leading edges armed with divergent rows or small cluster of short, erect, knife-like spinules. Other scales on head variable in size, generally similar to those on body, but spinules more erect; those on opercle, preopercle, and postorbital and supratemporal canals largest. Nasal fossa usually heavily scaled anteroventrally (scales rarely absent especially in

small specimens). Dorsal surfaces of snout fully scaled; naked clefts along each side of median rostral ridge only slightly developed. Underside of head almost completely naked; small scaly patches often present on ventral surfaces of preopercles and above posterior margins of lower jaws.

No open pores along cephalic sensory canals. Free neuromasts on ventral surface of head tiny, poorly marked, not discernible. Anterior nostril small and oval; posterior large, bean-shaped. Lateral line complete, extending from upper margin of opercle to tip of tail.

Origins of first dorsal and pelvic fins on about same vertical; first dorsal fin moderately high, its height 2.3–3.4 times as long as its base length; second spinous ray not elongated, smooth along its leading edge (1–2 rudimentary denticles rarely present distally); its tip extending to second dorsal-fin origin or beyond when

laid back. Interdorsal space 0.9–1.5 times as long as first dorsal-fin base length. Second dorsal fin originating above or posterior to anal-fin origin. Pectoral-fin base slightly anterior to vertical through pelvic-fin base. Outer pelvic-fin ray slightly prolonged, filamentous at tip.

FIGURE 25. Scatter diagrams showing relationships of the (A) orbit diameter, (B) postorbital length, and (C) orbit–preopercle distance (as % PRL) to PRL (mm) in *Coelorinchus anatirostris* (circles) and *C. weberi* (squares). Arrows indicate holotypes.

Color when fresh (Fig. 18). Dorsum body purplish gray, ventral half whitish without prominent silvery reflection; abdomen generally pale even in ventral view; opercle iridescent soon after capture; first dorsal darker apically, but otherwise paler; pectoral and pelvic

fins dusky; distal margin of anal fin narrowly blackish posteriorly.

Color in alcohol (Figs. 19–20, 21A, 24). Head and body darker dorsally, paler ventrally; abdomen dark violet, but chest paler; lips white; oral and gill cavities blackish; gill rakers and filaments light tan, arches pallid; gular and branchiostegal membranes pale; first dorsal fin paler basally, darker distally, second spinous ray blackish; second dorsal fin immaculate; anal fin paler anteriorly, but prominently darker posteriorly; pectoral and pelvic fins generally dusky.

Size. To about 43 cm TL (BSKU 4385, 427 mm TL, off Kochi, Japan).

Distribution. Distributed in Japan and Taiwan as well as in the tropical southwestern Pacific including Vanuatu and the Chesterfield Islands (Iwamoto & Merrett 1997; Merrett & Iwamoto 2000; Shao *et al.* 2008b; Nakayama 2019). Depth range 110–1160 m. In Japan and adjacent waters, known from off the Pacific coasts of southern Japan northward to the Miura Peninsula (139.62°E), Okinawa Trough, south of Jeju Island in Korea, and the northern South China Sea (Appendix 3-1C). One of the most abundant species of *Coelorinchus* found at upper slope depths off southern Japan.

Nomenclatural discussion. *Coelorinchus anatirostris* was originally described from a single 395 mm TL specimen collected from Misaki, Sagami Bay, Japan (Fig. 19). Subsequently, Gilbert & Hubbs (1916) described *C. productus* based on seven specimens collected from Suruga Bay, Japan (186+–304 mm TL; Fig. 20), suggesting its close relationship with *C. anatirostris*. According to Gilbert & Hubbs (1916), *C. productus* is distinguished from *C. anatirostris* by having the following combination of features: snout longer, sharply pointed in dorsal view (vs. duckbill-shaped in *C. anatirostris*); orbit smaller, 1.8 in snout length, slightly less than distance from pelvic-fin base to anus (vs. 1.5); barbel shorter, about 6 in orbit diameter (vs. almost equal to half eye); maxillary 4.6 in HL (vs. 4); gill membranes without free fold; body scales with 3–5 spinule rows (vs. 6–9); scales atop head not similar to those on body, with 1–3 spinule rows.

The taxonomic status of *C. productus* has been controversial over the past few decades. Okamura (1970a), who examined paratypes of this species and many Japanese specimens of *C. anatirostris*, concluded that the two species are synonymous, considering that the differences reported by Gilbert & Hubbs (1916) are simply attributed to size-related or individual variations. However, Yatou (*in* Okamura & Kitajima 1984) resurrected *C. productus* and distinguished the two species by the combination of the barbel length (1/4–1/3 times as long as orbit diameter in *C. productus* vs. 1/3 times or more in *C. anatirostris*) and preoral length (HL 2.4–2.6 times as long as preoral length vs. 2.6–3.0 times). Subsequently, many authors followed Yatou's (1984) conclusion (see the synonym list),

but Fukui *et al.* (2010) re-synonymized the species with *C. anatirostris* based on selected measurements and an analysis of 16s rRNA sequences. Most recently, Iwamoto *et al.* (2015) continued to recognize *C. productus* as valid, but noted that the problem needs further investigation.

An examination here of the types of both *C. anatirostris* and *C. productus* and an additional 46 specimens from Japan reconfirmed their conspecificity. All measurements of the holotype of *C. productus* fall within, or only slightly outside of the range of variation of *C. anatirostris* (Table 1), except for the snout length (88% PRL vs. 60–84%) and preoral length (81% PRL vs. 50–79%). However, these subtle differences are of little significance when referring to ontogenetic variation of the snout in *C. anatirostris*. In *C. anatirostris*, the snout becomes shorter and broader with growth (Fig. 23A), giving significant change to the head appearance (Fig. 24). Furthermore, the shape of the snout is variable even in a comparison with similar-sized specimens (see Fig. 24: BSKU 77981 vs. BSKU 77982; BSKU 4385 vs. BSKU 27989). These variations influenced other measurements, because the head length was traditionally used as a standard of comparison in grenadiers.

As noted by Gilbert & Hubbs (1916), type specimens of *C. productus* have a somewhat shorter barbel than the holotype of *C. anatirostris* (8–11% PRL vs. 12%). However, the barbel length relative to HL tends to become longer with growth (Fig. 23B), and the difference is unclear when the variation of *C. anatirostris* is considered (9–18% PRL). Unfortunately, the distal portion of the chin barbel is slightly damaged in the holotype of *C. productus*.

Gilbert & Hubbs (1916) distinguished *C. productus* from *C. anatirostris* in having “scales with 3 to 5, instead of 6 to 9, divergent spinous ridges, those on the top of the head not similar to those on body, bearing but one to three ridges”. However, the number of spinule rows tends to increase with growth in grenadiers (Iwamoto & Merrett 1997:509) and the character should be used only for comparisons of similar-sized specimens. Specimens in the type series of *C. productus* are significantly smaller than the holotype of *C. anatirostris* (53.0–79.9 mm HL vs. 93.7 mm HL), and the differences in squamation appear to reflect size-related variation.

Meristic data also show no significant differences between the two species. Therefore, this study supports Fukui *et al.*'s (2010) conclusion that *C. productus* is a junior synonym of *C. anatirostris*.

Comments on type specimens. In the original description of *C. productus*, Gilbert & Hubbs (1920) designated six specimens collected from the *Albatross* sta. 5059 and 5066 as paratypes of the species. However, seven specimens being referable to *C. anatirostris* collected from these stations were found among the USNM and CAS fish collections. Of these, the smallest one from sta. 5059 (USNM 76873, 1 of 4, 93+ mm TL) is herein considered a non-type, because it is obviously

smaller than the size range of the *C. productus* paratypes given in the original description (190–286 mm TL).

Remarks. Okamura (1970b: fig. 14N) provided a figure of the premaxillary tooth band of *C. anatirostris*, where the outermost series was illustrated as being distinctly enlarged. His figure seems to be diagrammatic, considering that the specimens examined in this study lack greatly enlarged teeth on the upper jaw. In *C. anatirostris*, the premaxillary teeth are arranged in a short, uniformly wide band, with tooth size gradually decreasing inward. In addition, inner rows are deeply embedded in a thick layer of gum papillae, which gives a larger appearance to the outermost series.

Chiou *et al.* (2004b; as *C. productus*) reported two Taiwanese specimens (ASIZP 61326 and ASIZP 61327, 278–296 mm TL) as having 11–12 pelvic-fin rays. However, the number is almost always seven (rarely six) in *Coelorinchus*, and a brief examination of the voucher specimens confirmed that Chiou *et al.*'s (2004b) counts were erroneous (those specimens have seven pelvic-fin rays). Chiou *et al.* (2004b) also noted their specimens as having 3.5–4 scale rows below the first dorsal-fin origin and the length of the light organ 2.8–3 times as long as the orbit diameter. Although their scale counts are slightly lower than those of *C. anatirostris* in this study (4.5–6.5), the difference appears to reflect different methods of counting. Further, the length of the light organ given by Chiou *et al.* (2004b) should be read as the length is 2.8–3 in the orbit diameter.

Coelorinchus anatirostris has a disjunct distribution between the northwestern Pacific from Japan to Taiwan and the southwestern Pacific. A single specimen collected from the Chesterfield and Bellona Plateaus (CAS 86496, 81.9 mm HL, 288+ mm TL) was examined to compare with the northwestern Pacific population. As pointed out by Iwamoto & Merrett (1997:489), the New Caledonian specimen has a distinctly paler color, but otherwise they are closely similar to each other. A further in-depth comparison should be done with more specimens from the southeastern Pacific.

Relationships. *Coelorinchus anatirostris* belongs to a group of *Coelorinchus* species characterized by the following combination of features: light organ short, its anterior margin falling far short of pelvic-fin bases; lateral nasal ridge completely supported by nasal bone, not strongly convex when viewed dorsally; premaxillary teeth in short, uniformly wide band; underside of head completely to broadly naked; body scales covered with widely divergent rows of spinules, and every row complete throughout; buttresses of body-scale spinules scarcely to only narrowly developed; no prominent markings on body (Nakayama & Endo 2017; this study). It corresponds with one of two subgroups of the *C. japonicus* group as redefined by Nakayama & Endo (2017), comprising the following eight species in which the underside of the head is completely or almost completely naked: *C. anatirostris*;

C. goobala Iwamoto & Williams, 1999 from the eastern Indian Ocean off Australia; *C. hige* Matsubara, 1943 from Japan to Taiwan (a senior synonym of *C. asteroides* Okamura, 1963); *C. lanceolatus* **sp. nov.** from Japan; *C. okamurai* Nakayama & Endo, 2017 from the Timor Sea; *C. radcliffei* Gilbert & Hubbs, 1920 from the Philippines and Indonesia; *C. shcherbachevi* Iwamoto & Merrett, 1997 from the tropical southwestern Pacific; and *C. weberi* Gilbert & Hubbs, 1920 from the Philippines. For convenience of comparison, this group is called here the *C. anatirostris* group.

Coelorinchus chilensis Gilbert & Thompson in Thompson, 1916, known from the southeastern Pacific, is superficially similar to the aforementioned species, but differs notably in having the lateral nasal ridge incompletely supported by the nasal bone, and in lacking any prominent external evidence of the light organ anterior to the anus.

Comparisons. Among the *C. anatirostris* group, *C. anatirostris* is closely related to *C. weberi*, with which it shares most of the diagnostic features used to distinguish the species from its congeners. Gilbert & Hubbs (1920:378, table 3) assumed that *C. weberi* is a possible sibling species of either *C. anatirostris* or *C. productus* (the last of which was then considered valid, but herein regarded as a junior synonym of *C. anatirostris*; see the Nomenclatural discussion). *Coelorinchus weberi* is currently known only from the holotype collected at the Babuyan Channel off northern Luzon, the Philippines. In their key to species of the genus (Gilbert & Hubbs 1920:425), *C. anatirostris* runs to the couplet j, where it is substantially separated from *C. weberi* by having a larger orbit, with its diameter being about equal to the postorbital length (vs. 1.35 in postorbital) and 1.9 in the snout length (vs. 1.5–1.8). An examination of the holotype of *C. weberi* (USNM 78225, 86.7 mm HL) confirmed their differences in the orbit diameter (45–52% PRL in *C. anatirostris* vs. 44% in *C. weberi*; Fig. 25A) and postorbital length (50–56% PRL vs. 58%; Fig. 25B) as well as the orbit–preopercle distance (52–60% PRL vs. 62%; Fig. 25C). However, the two species differ most notably from each other by coloration of the cephalic neuromasts. In *C. anatirostris*, the free neuromasts on the underside of the head are difficult to distinguish as they are unpigmented, whereas each neuromast of *C. weberi* is marked with two small black dots, and easily discernible without a microscope. The latter species will be fully redescribed in a forthcoming paper by the present author.

Coelorinchus shcherbachevi is another species that is closely similar to *C. anatirostris*, and is known only from four large specimens (118–132 mm HL, 435+–550+ mm TL) collected from the Loyalty Ridge, New Caledonia (Iwamoto & Merrett 1997; Merrett & Iwamoto 2000) and an additional record from the western Coral Sea (Last *et al.* 2014). The holotype (MNHN 1996-966, 126+ mm HL) and one of the paratypes (CAS 90995, 123

mm HL) were examined in the course of this study. Its distinctness from *C. anatirostris* was well documented in the original description given by Iwamoto & Merrett (1997). *Coelorinchus anatirostris* is clearly distinguished from *C. shcherbachevi* by having keel-like spinules on the body scales (vs. broadly triangular, somewhat pyramidal in *C. shcherbachevi*) and poorly developed sensory organs on the underside of the head (vs. dark hair-like papillae densely scattered, with prominent flap-like black neuromasts). As discussed by Iwamoto & Merrett (1997), the number of spinule rows on the body scales also appears to be a good diagnostic character between the two species, especially when similar-sized specimens are compared. In the Japanese specimens of *C. anatirostris* examined, the number increases with size, with one of the largest specimens (BSKU 4385, 98.4 mm HL) having eight rows of spinules (Fig. 22A). However, the *C. shcherbachevi* specimens examined had only four or five rows despite their large size, and the number is considerably fewer than in the *C. anatirostris* specimens examined. This study also confirmed that the two species are further distinguished from each other by the orbit diameter (45–52% PRL in *C. anatirostris* vs. 43–44% in *C. shcherbachevi*), orbit–preopercle distance (52–60% PRL vs. 61–63%), upper-jaw length (37–45% PRL vs. 45–47%), interorbital width (34–43% PRL vs. 42–43%), isthmus–pelvic distance (41–57% PRL vs. 59–63%), height of first dorsal fin (85–107% PRL vs. 80%), length of gill slit (15–22% PRL vs. 23–28%), and coloration (paler ventrally vs. uniformly dark). However, these differences should be verified when more specimens of *C. shcherbachevi* become available, because ontogenetic variations of that species have been poorly understood.

Among Japanese congeners, *C. anatirostris* is most similar to *C. lanceolatus* **sp. nov.**, but is readily distinguished from the latter by its shorter snout (60–88% PRL vs. 107%) and higher first dorsal fin (85–107% PRL vs. 84%). *Coelorinchus anatirostris* is also likely to be confused with *C. hige* (a senior synonym of *C. asteroides* Okamura, 1963; see the Nomenclatural discussion on the latter species), and they occur together in the northwestern Pacific (Appendix 3-1C and 3-2E). It differs primarily from *C. hige* in that the occipital scales are covered with keel-like spinules in narrowly divergent, saw-toothed rows (vs. needle-like spinules in widely divergent, comb-like rows) and the nasal fossa is usually scaled ventrally (vs. completely or almost completely naked). A further comparison between the two species is discussed under the Relationships and comparisons of *C. hige*.

Coelorinchus brevirostris Okamura, 1984

[Japanese name: Tsumari-hige]

(Figs. 26–27, 28A–B; Appendix 3-1D)

Coelorinchus brevirostris Okamura, 1984a:225, fig. 160 (original

description; holotype: BSKU 29562, from Okinawa Trough, 25°47.4'N, 124°23.4'E, in 600 m; new Japanese name: "Tsumari-hige"); Okamura 1988:453, pl. 373, fig. G (compiled); Okamura 1997:126, fig. 4 (compiled); Shao *et al.* 2008b: table 2 (6 spec. listed from northeastern and eastern Taiwan and South China Sea); Nakabo & Kai 2013:507 (in key; Japan); Iwamoto *et al.* 2015:51 (brief description; 21 spec. from northeastern and eastern Taiwan and South China Sea); Motomura 2020:38 (listed; Japan).

Caelorinchus brevirostris: Nakabo 1993:367 (in key; Japan); Nakabo 2000:431 (in key; Japan); Nakabo 2002:431 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Chiou *et al.* 2004b:301, figs. 4–5 (redescription; 11 spec. from Da-xi; first record from Taiwan); Shinohara *et al.* 2005:416 (listed; Ryukyu Islands); Furuhashi *et al.* 2010: table 2 (1 spec. listed from northern Okinawa Trough).

Diagnosis. Ventral light organ externally represented by long dark line, terminated with 1 fossa at each end, one immediately anterior to anus, the other on chest just posterior to isthmus (duct connecting each fossa often difficult to distinguish). Underside of head completely naked except along lateral nasal ridges; dorsal surface of snout mostly scaled, but prominent naked clefts present along each side of median rostral ridge. Snout long, sharply pointed, length 67–83% PRL; dorsal profile of snout straight in lateral view; terminal scute short, moderately broad, arrowhead-shaped, dorsoventrally flattened, length 14–17% PRL; distal portion of snout broadly spade-shaped when viewed dorsally; lateral nasal ridge incompletely supported by nasal bone. Interdorsal space shorter than first dorsal-fin base length. Anal fin originating posterior to vertical through second dorsal-fin origin. Second spinous ray of first dorsal fin greatly elongate, height of fin 139–220% PRL. Second dorsal fin well developed, its rays as long as opposites of anal fin. Anus immediately anterior to anal-fin origin. Premaxillary teeth small, slender, conical in short, uniformly wide band, none especially enlarged; posterior end of tooth band falling far short of lateral corner of mouth. Body scales covered with short, reclined, needle-like spinules in subparallel to slightly divergent rows; spinules in each row overlapping, but free from one another, with last spinule extending well beyond posterior scale margin; buttresses narrowly developed. Spinules on scales anterior to light organ rudimentary. Orbit diameter 43–47% PRL; postorbital length 52–56% PRL; upper-jaw length 36–38% PRL; preoral length 58–72% PRL; barbel length 4–10% PRL. Transverse scale rows below first dorsal-fin midbase 3.5–5.5. Body pale overall, lacking prominent blotches or saddles; scale pockets gently outlined by dark brown; underside of head dusky, heavily peppered; lips pale; oral cavity blackish; gular membrane almost immaculate, but branchiostegal membranes moderately dark; first dorsal fin blackish; pelvic fin dark, but outermost ray paler distally.

Material examined. 7 specimens. **Holotype** of

Coelorinchus brevirostris: BSKU 29562 (54.2 mm HL, 204+ mm TL), northwest of Miyako-jima Island, Okinawa Pref., Okinawa Trough, Japan, East China Sea, 25.7900°N, 124.3900°E, 600 m, F/V *Yuryo-maru*, No. 8, sta. 1-T13, bottom trawl, coll. T. Kitajima, 20 Sept. 1979. **Non-types: Japan:** FFNU P-00640 (2, 52.0–52.6 mm HL, 212+–224 mm TL), east of Okinohira Bank, East China Sea, 32.1757°N, 129.4868°E, 476 m, T/V *Nagasaki-maru*, cr. N218, sta. B5-4, 3-m ORE beam trawl, coll. J. Hashimoto, 1 Aug. 2006; FAKU 84522 (1, 52.6 mm HL, 221+ mm TL), west of Yaku-shima Island, Okinawa Trough, 30.4133°N, 128.2300°E, 499–500 m, FRV *Yokomaru*, sta. D9, tr. T-2, 10 Nov. 2002. **Taiwan:** ASIZP 61352 (1, 36.2 mm HL, 147+ mm TL), Da-xi fish market, Yilan, bottom trawl, 1 Jun. 2001; ASIZP 61350 (1, 53.1 mm HL, 181+ mm TL), Da-xi fish market, Yilan, bottom trawl, 30 Jun. 2000. **South China Sea:** ASIZP 66806 (1, 47.1 mm HL, 181+ mm TL), northeast of Dong-sha Atoll, 21.6701°N, 117.7195°E, 509 m, beam trawl, coll. H.-M. Yeh *et al.*, 17 Aug. 2005.

Counts and measurements. Based on 7 specimens (36.2–54.2 mm HL, 147+–224 mm TL). Counts: first dorsal-fin rays II, 7–9; pectoral-fin rays i15–i19; pelvic-fin rays 7; gill rakers on first arch (outer/inner) 0/6–8, on second arch 5/7–8; longitudinal scales 29–32; transverse scale rows below first dorsal-fin origin 4.5–6.5, below first dorsal-fin midbase 3.5–5.5, below second dorsal-fin origin 4.5–6.5, above anal-fin origin 15.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 41–46 (67–83); orbit diameter 24–27 (43–47); postorbital length 30–33 (52–56); postrostral length 56–61; orbit–preopercle distance 28–33 (50–55); suborbital width 12–14 (21–26); upper-jaw length 20–22 (36–38); length of rictus 15–16 (26–29); length of premaxillary tooth band 6–9 (11–15); preoral length 35–40 (58–72); length of terminal snout scute 8–10 (14–17); length of lateral nasal ridge 28–30 (48–54); length of suborbital ridge 90–92 (148–165); snout width 29–32 (50–55); internasal width 19–20 (31–35); interorbital width 19–21 (32–36); occipital width 9–10 (14–17); body width over pectoral-fin bases 39–46 (68–81); body depth at first dorsal-fin origin 43–49 (75–83); body depth at anal-fin origin 38–43 (66–77); prepelvic length 113–125 (193–224); preanus length 158 (270); preanal length 162–164 (279–291); isthmus–pelvic distance 25–29 (43–51); isthmus–anus distance 74 (126); isthmus–anal distance 66–79 (114–135); pelvic–anal distance 42–53 (73–90); anus–anal distance 6 (10); pelvic-fin length 37–44 (64–75); pectoral-fin length 40–44 (67–74); predorsal length 104–112 (180–190); height of first dorsal fin 82–125 (139–220); length of first dorsal-fin base 16–20 (27–34); interdorsal length 10–16 (18–28); length of gill slit 12–13 (20–22); length of posterior nostril 6–10 (10–17); barbel length 3–6 (4–10).

Supplementary description based on Japanese

FIGURE 26. Fresh specimen of *Coelorinchus brevirostris*. FAKU 84522, 52.7 mm HL, 221+ mm TL, west of Yaku-shima Island, Okinawa Trough, East China Sea, 499–500 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: Kyoto University Museum]

FIGURE 27. Holotype of *Coelorinchus brevirostris*. BSKU 29562, 54.2 mm HL, 204+ mm TL, Okinawa Trough, East China Sea, 600 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

specimens. *Coelorinchus brevirostris* has been well described by Okamura (1984a) and Chiou *et al.* (2004b) that can be referred to for details. Three additional specimens collected from the northern Okinawa Trough (FAKU 84522 1, 52.6 mm HL, 221+ mm TL; FFNU P-640, 2, 52.0–52.6 mm HL, 212+–224 mm TL) allowed to document previously unknown features regarding squamation. General features of the species are shown in Figs. 26–27.

Body scales moderate in size, thin, highly deciduous, covered with short, reclined, needle-like spinules in subparallel to slightly divergent rows (Fig. 28A); those on dorsum below interdorsal space with about 6 rows

of spinules; every spinule row complete, extending to posterior scale margin; middle row not especially high or enlarged; spinules in each row greatly overlapping, but free from one another; height of spinules gradually increasing posteriorly, with tips of last spinules extending well beyond posterior scale margin; lateral buttresses narrowly developed; no reticulate structures on unexposed portion. Spinules on chest scales anterior to light organ rudimentary (Fig. 28B).

Size. To about 22 cm TL (FFNU P-640, 1 of 2, 224 mm TL, Okinawa Trough, Japan).

Distribution. Known only from the Okinawa Trough, Taiwan Strait, and northern South China Sea, at

FIGURE 28. Scanning electron micrographs showing body scales of three species of the *Coelorinchus hubbsi* group. (A–B) *C. brevirostris*, FFNU P-640, 52.6 mm HL; (C–D) *C. cingulatus*, BSKU 98963, 60.4 mm HL; (E–F) *C. fuscigulus*, BSKU 106760, 41.9 mm HL. (A, C, E) Scales from the dorsum below the interdorsal space; (B, D, F) from the chest between the isthmus and the anterior tip of the light organ. Views from above. [Photos: N. Nakayama]

depths of 455–1185 m (Chiou *et al.* 2004b; Shao *et al.* 2008a, 2008b; this study; Appendix 3-1D). Very rare in Japanese waters.

Remarks. *Coelorinchus brevirostris* was originally described from a single specimen collected from the Okinawa Trough, southern Japan, in 600 m. The holotype (Fig. 27) is in a somewhat poor condition, with all body scales missing and the tip of the first dorsal fin broken. Chiou *et al.* (2004b) redescribed the species based on 11 specimens obtained at the Da-xi fish market in northeastern Taiwan, and newly documented that the second spinous ray of the first dorsal fin is greatly elongated. Subsequently, Furuhashi *et al.* (2010) reported two Japanese specimens (FFNU P-00640) collected from the Okinawa Trough off Nagasaki Pref., which represents the northernmost record for the species. An additional specimen from the west of Yaku-shima Island in the Okinawa Trough was also found among the fish collection of the Kyoto University Museum (Fig. 26).

Relationships. *Coelorinchus brevirostris* falls in the group of *Coelorinchus* species as defined by Iwamoto *et al.* (2009:43–44) to include the following 10 species (this group is called here the *C. hubbsi* group; see also Nakayama *et al.* 2020): *C. brevirostris*, *C. cingulatus* Gilbert & Hubbs, 1920 from Japan to the South China Sea (see also the Remarks of that species); *C. cylindricus* Iwamoto & Merrett, 1997 from the Norfolk Ridge and New Zealand; *C. gladius* Gilbert & Cramer, 1897 from Hawaii; *C. fuscigulus* Iwamoto, Ho & Shao, 2009 from Japan to Taiwan; *C. hubbsi* Matsubara, 1936 from Japan to Taiwan; *C. matsubarai* Okamura, 1982 from the Kyushu-Palau Ridge and the Emperor Seamounts; *C. melanosagmatus* Iwamoto & Anderson, 1999 from the Indian Ocean; *C. multifasciatus* Sazonov & Iwamoto, 1992 from the Sala y Gomez Ridge; and *C. spilonotus* Sazonov & Iwamoto, 1992 from the Nazca and Sala y Gomez Ridges and Hawaii. These species are distinctive within the genus in having a somewhat cylindrical body, a long light organ with two external fossae (of which one is immediately anterior to the anus and the other is on the chest immediately posterior to the isthmus), a well-developed second dorsal fin (as high as the anal fin), a narrow interdorsal space (less than the first dorsal-fin base length), an incomplete bony support for the lateral nasal ridge, the absence of scales on the underside of the head except for overlapping scales along the lateral nasal ridges (small scaly patches occasionally found above the posterior margins of the upper jaw), and body scales covered with needle-like spinules in subparallel to slightly divergent rows.

Coelorinchus acutirostris Smith & Radcliffe in Radcliffe, 1912 distributed in the Philippines, Australia, and New Caledonia agrees generally with the above-mentioned features, but its lateral nasal ridge is completely supported by the nasal bone. It further differs from all species of the *C. hubbsi* group in having body scales covered in widely

divergent rows of erect spinules (vs. subparallel to slightly divergent rows of reclined spinules).

Coelorinchus melanobranchus Iwamoto & Merrett, 1997, known from New Caledonia, is also similar in many aspects to the members of the *C. hubbsi* group. However, it has very different physiognomy associated with a bluntly pointed snout (47–57% PRL vs. $\geq 60\%$), a short terminal scute (4–8% PRL vs. $\geq 12\%$), a narrow interorbital space (27–32% PRL vs. $\geq 32\%$), and a long chin barbel (33–42% PRL vs. $\leq 19\%$).

This group almost corresponds with Okamura's (1984a:227) “*hubbsi* complex”, although the latter also includes *C. acutirostris*.

Comparisons. *Coelorinchus brevirostris* differs most notably from all other species of the *C. hubbsi* group in lacking prominent body markings on the body (vs. dark saddles, blotches, or stripes present) and in having rudimentary spinules on the chest scales anterior to the anterior fossa of the light organ (vs. generally similar to those on the dorsum). It also differs from all except *C. cingulatus* by having a greatly elongate second spinous ray of the first dorsal fin (vs. not elongate). *Coelorinchus brevirostris* further differs from *C. cingulatus* in having shorter terminal scute (14–17% PRL vs. 20–26%) and chin barbel (4–10% PRL vs. 10–13%), and prominent naked clefts along each side of the median rostral ridge (vs. only narrowly developed). The shape of the snout is also significantly different between the two species. In *C. brevirostris*, its dorsal contour is almost straight in lateral view, whereas it is obviously concave in *C. cingulatus*. When viewed dorsally, the anterolateral margins are broadly convex in *C. brevirostris*, giving a spade-shaped appearance to the snout. In contrast, *C. cingulatus* has a more attenuated snout in dorsal view, with the leading edges distinctly concave anteriorly.

Coelorinchus cingulatus Gilbert & Hubbs, 1920

[Japanese name: Kurakake-sokodara]

(Figs. 28C–D, 29; Appendix 3-1E)

Coelorhynchus cingulatus Gilbert & Hubbs, 1920:480, fig. 15 [original description; holotype: USNM 78221, from “China Sea near Taiwan”, *Albatross* sta. 5317, in 230 fth (421 m); 1 paratype from “China Sea off northern Luzon”]; Chen 1951:210 (listed; Taiwan).

Coelorinchus cingulatus: Okamura 1984a:229, 366, fig. 161 (first record from Japan; brief description; 1 spec. from Okinawa Trough; BSKU 28658; new Japanese name: “Kurakake-sokodara”); Okamura 1988:453, pl. 373, fig. H (compiled); Shao *et al.* 2008b: table 2 (17 spec. listed from northeastern and southwestern Taiwan, and South China Sea); Ho & Shao 2011:36 (type catalog); Nakabo & Kai 2013:508 (in key; Japan); Iwamoto *et al.* 2015:52 (brief description; 29 spec. from northeastern and southwestern Taiwan, and Philippines); Motomura 2020:38 (listed; Japan).

Caelorinchus cingulatus: Nakabo 1993:367 (in key; Japan); Shao 1993:168 (compiled; Taiwan); Nakabo 2000:431 (in key; Japan); Nakabo 2002:431 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Shinohara *et al.* 2005:416 (listed; Ryukyu Islands).

Caelorinchus dorsalis (not Gilbert & Hubbs 1920): Shao 1993:168 (compiled; Taiwan) [fide Chiou *et al.* (2004a: table1)].

Caelorinchus (Paramacrurus) cinguletus [sic]: Chiou *et al.* 2004a: table 1 (listed; Taiwan).

Diagnosis. Ventral light organ externally represented by long dark line, terminated with 1 fossa at each end, one immediately anterior to anus, the other on chest just posterior to isthmus (duct connecting each fossa often difficult to distinguish). Underside of head completely naked except along lateral nasal ridges; dorsal surface of snout almost fully scaled, naked clefts only narrowly developed along each side of median rostral ridge. Snout long, sharply pointed, length 79–86% PRL; dorsal profile of snout distinctly concave in lateral view; terminal scute long, slender, spearhead-shaped, length 20–26% PRL; distal portion of snout narrowly attenuated when viewed dorsally; lateral nasal ridge incompletely supported by nasal bone. Interdorsal space less than first dorsal-fin base length. Anal fin originating posterior to vertical through second dorsal-fin origin. Second spinous ray of first dorsal fin greatly elongate, height of fin 123–156% PRL in adults. Second dorsal fin well developed, its rays as long as opposites of anal fin. Anus immediately anterior to anal-fin origin. Premaxillary teeth small, slender, conical in short, uniformly wide band, none especially enlarged; posterior end of tooth band falling far short of lateral corner of mouth. Body scales covered with short, reclined, needle-like spinules in subparallel to slightly divergent rows; spinules in each row overlapping, but free from one another, with last spinule extending well beyond posterior scale margin; buttresses scarcely developed. Spinules on scales anterior to light organ not rudimentary. Orbit diameter 41–47% PRL; postorbital length 54–61% PRL; upper-jaw length 35–43% PRL; preoral length 66–80% PRL; barbel length 10–13% PRL. Transverse scale rows below first dorsal-fin midbase 4–5.5. Body with several dark markings including prominent diagonal stripes on trunk and faint elongate blotches on anterior portion of tail; longitudinal dark stripe running along lower margin of postorbital canal; scale pockets on dorsal half of body outlined by dark brown; underside of head dusky, heavily peppered; lips pale; oral cavity blackish; gular and branchiostegal membranes dusky to dark; first dorsal fin uniformly dark, but second spinous ray paler; pelvic fin dark, but outermost ray paler distally.

Material examined. 13 specimens. **Holotype of *Coelorhynchus cingulatus*:** USNM 78221 (40.0 mm HL, 127+ mm TL), northern South China Sea, 21.6000°N, 117.4500°E, 230 ftm (421 m), *Albatross* sta. 5317, 12-ft Tanner beam trawl, 5 Nov. 1908. **Paratype of *C.***

***cingulatus*:** USNM 78233 (1, 41.2 mm HL, 146+ mm TL), South China Sea off northern Luzon, Hermanos Island, Philippines, 18.5708°N, 121.8542°E, 224 ftm (410 m), *Albatross* sta. 5325, 12-ft Tanner beam trawl, 12 Nov. 1908. **Non-types: Japan:** BSKU 28658 (1, 60.6 mm HL, 232 mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.9000°N, 127.7167°E, 250 m, F/V *Ryoan-maru*, No. 28, tr. 78, bottom trawl, coll. Y. Kinoshita and S. Hagino, 21 Mar. 1978; FAKU 204486 (1, 28.2 mm HL, 105+ mm TL), Mimase fish market, F/V *Tsukasa-maru*, bottom trawl, coll. N. Nakayama 4 Mar. 2017, bottom trawl. **Taiwan:** KAUM-I. 44654 (1, 67.8 mm HL, 224+ mm TL), off Kaohsiung, Pingtung, 400 m, trawl, coll. M. Matsunuma and H.-C. Ho, 1 Jul. 2011; BSKU 98963 (1, 60.4 mm HL, 191+ mm TL), BSKU 98964 (1, 34.8 mm HL, 122+ mm TL), Dong-gang fish market, Pingtung, bottom trawl, coll. H.-C. Ho, 16 Nov. 2007; *FAKU 204447 (1, 45.3 mm HL, 179+ mm TL), Dong-gang fish market, Pingtung, bottom trawl, coll. N. Nakayama, 27 Feb. 2017. **South China Sea:** BSKU 106847 (1, 62.8 mm HL, 224+ mm TL), BSKU 106848 (1, 60.6 mm HL, 232+ mm TL), off Swatow, 21.6867°N, 117.5183°E, 415–437 m, R/V *Hakuho-maru*, cr. KH-73-02, sta. 44-1, 3-m beam trawl, 19 Mar. 1973. **Philippines:** ASIZP 68017 (1, 59.0 mm HL, 227+ mm TL), east of Luzon Island, 14.7665°N, 123.6553°E, 356–367 m, French type beam trawl, coll. Y.-C. Liao and K.-T. Shao, 26 May 2007; ASIZP 67859 (1, 61.0 mm HL, 227+ mm TL), east of Luzon Island, 15.9908°N, 121.8447°E, 422–431 m, French type beam trawl, coll. Y.-C. Liao and K.-T. Shao, 20 May 2007; ASIZP 68394 (1 of 2, 61.3 mm HL, 211+ mm TL), east of Luzon Island, 15.9908°N, 121.8447°E, 422–431 m, French type beam trawl, coll. Y.-C. Liao and K.-T. Shao, 20 May 2007.

Counts and measurements. Based on 12 specimens (28.2–67.8 mm HL, 105+–232+ mm TL). Counts: first dorsal-fin rays II,8–9; pectoral-fin rays i16–i20; pelvic-fin rays 7; gill rakers on first arch (outer/inner) 0/7, on second arch 5–6/7–9; longitudinal scales 31–37; transverse scale rows below first dorsal-fin origin 5.5–6.5, below first dorsal-fin midbase 4–5.5, below second dorsal-fin origin 4–6, above anal-fin origin 13.5–16.5.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 44–46 (79–86); orbit diameter 22–27 (41–47); postorbital length 30–34 (54–61); postrostral length 54–57; orbit–preopercle distance 29–34 (53–60); suborbital width 11–15 (21–27); upper-jaw length 19–24 (35–43); length of rictus 14–19 (26–34); length of premaxillary tooth band 7–9 (13–17); preoral length 37–44 (66–80); length of terminal snout scute 11–14 (20–26); length of lateral nasal ridge 31–34 (55–63); length of suborbital ridge 90–96 (160–174); snout width 25–31 (44–57); internasal width 17–20 (30–37); interorbital width 20–21 (35–38); occipital width 8–10 (15–19); body width over pectoral-fin bases 31–45 (56–82); body depth at first

FIGURE 29. Fresh specimen of *Coelorinchus cingulatus*. (A–C) FAKU 204486, 28.2 mm HL, 105+ mm TL, Mimase fish market, Kochi Pref., Pacific; (D) FAKU 204447, 45.3 mm HL, 179+ mm TL, Dong-gang fish market, southwestern Taiwan, South China Sea. (A, D) Lateral views (A, reversed); (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

dorsal-fin origin 39–52 (71–93); body depth at anal-fin origin 32–41 (58–74); prepelvic length 101–117 (183–212); preanus length 135–154 (245–276); preanal length 139–161 (252–292); isthmus–pelvic distance 24–32 (44–58); isthmus–anus distance 62–75 (111–134); isthmus–anal distance 64–80 (116–142); pelvic–anal distance 42–53 (75–95); anus–anal distance 3–8 (5–15); pelvic-fin length 33–39 (61–70); pectoral-fin length 32–45 (57–83); predorsal length 104–109 (186–199); height of first dorsal fin 59–85 (107–156); length of first dorsal-fin base 16–18 (27–32); interdorsal length 4–12 (8–22); length of gill slit 10–13 (18–24); length of posterior nostril 8–11 (15–20); barbel length 5–7 (10–13).

Size. To about 23 cm TL (BSKU 28658, 232 mm TL, Okinawa Trough, Japan).

Distribution. Known from Tosa Bay and the East and South China Seas at depths of 250–1211 m (Gilbert & Hubbs 1920; Shao *et al.* 2008a, 2008b; this study; Appendix 3-1E). Very rare in Japanese waters.

Remarks. *Coelorinchus cingulatus* was originally described from the northern South China Sea by Gilbert & Hubbs (1920), to which the reader can refer for further morphological details. This species was first reported

from Japanese waters by Okamura (1984a) based on a single specimen collected from the Okinawa Trough (BSKU 28658).

Iwamoto & Merrett (1997) and Merrett & Iwamoto (2000) recorded this species from the New Caledonian region and off the east coast of Australia. However, Iwamoto & Graham (2001:442) subsequently re-identified these specimens as *Coelorinchus* sp. cf. *cingulatus*, noting that “... their specimens, though closely similar, did not entirely agree with the description of *C. cingulatus* Gilbert & Hubbs, 1920, especially in regards to certain body markings ...” A further study is therefore necessary to elucidate the taxonomic status of the southwestern Pacific population.

Comparisons. *Coelorinchus cingulatus* is a distinctive species of the *C. hubbsi* group (see the Relationships of *C. brevirostris*). Its body markings, consisting of somewhat broken, diagonal bands or stripes, readily diagnose the species from other members of this group. Similar markings are also found in *C. cylindricus* Iwamoto & Merrett, 1997 known from the Norfolk Ridge and New Zealand, *C. fuscigulus* Iwamoto, Ho & Shao, 2009 from the East China Sea, *C. melanosagmatus*

Iwamoto & Anderson, 1999 from the Indian Ocean, and *C. multifasciatus* Sazonov & Iwamoto, 1992 from the Sala y Gomez Ridge; however, their markings consist of regularly arranged distinct saddles. It further differs from all except *C. brevirostris* Okamura, 1984 in that the second spinous ray of the first dorsal fin is distinctly elongate in adult specimens [height of first dorsal fin 123–156% PRL (107% in a 28.2 mm HL spec.; FAKU 204486) vs. not elongate, usually $\leq 122\%$]. *Coelorinchus hubbsi* Matsubara, 1936 also has a high first dorsal fin (85–130% PRL, usually $\leq 122\%$), but its second spinous ray is not especially elongated.

***Coelorinchus divergens* Okamura & Yatou, 1984**

[Japanese name: Hanare-hige]

(Figs. 30, 31A–B; Appendix 3-1F)

Coelorinchus divergens Okamura & Yatou, 1984:237, fig. 169 (original description; holotype: BSKU 26865, from Okinawa Trough; 1 paratype from Okinawa Trough; new Japanese name: “Hanare-hige”); Okamura 1988:453, pl. 373, fig. J (compiled); Shao *et al.* 2008b: table 2 (15 spec. listed from northeastern and southwestern Taiwan, and South China Sea); Nakabo & Kai 2013:509 (in key; Japan); Iwamoto *et al.* 2015:53, fig. 5 (brief description; 17 spec. from northeastern and eastern Taiwan, South China Sea, and Philippines); Motomura 2020:38 (listed; Japan).

Coelorinchus divergens: Nakabo 1993:368 (in key; Japan); Nakabo 2000:432 (in key; Japan); Nakabo 2002:432 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Shinohara *et al.* 2005:416 (listed; Ryukyu Islands).

Diagnosis. No external evidence of light organ anterior to anus. Underside of head, including ventral surfaces of preopercles, and dorsal surface of snout almost completely and uniformly scaled; nasal fossa scaled anteroventrally. Snout long, sharply pointed, length 64–69% PRL; dorsal profile of snout almost straight in lateral view; terminal scute moderately short, diamond-shaped, dorsoventrally flattened; lateral nasal ridge incompletely supported by nasal bone. Anus abutting anal-fin origin. Premaxillary teeth small, conical, arranged in short, uniformly wide band, with outer series slightly enlarged; posterior margin of tooth band falling far short of lateral corner of mouth. Body scales covered with stout, narrowly trihedral, shield-like spinules in subparallel ridge-like rows; middle row enlarged and higher, giving striated appearance to body surface; lateral rows often incomplete, not extending to posterior scale margin; buttresses broadly developed. Transverse scale rows below first dorsal-fin origin 5.5–6. Occipital scales covered with widely divergent, crest-like rows of keel-like spinules; no prominent naked area atop head between parietal ridges. Modified scales on head ridges prominently spiny and thickened. Free neuromasts on underside of head not especially enlarged. Upper-

jaw length 45–47% PRL; height of first dorsal fin 58–74% PRL. Head and body uniformly dark brown; orbit narrowly edged in black; all fins blackish.

Material examined. 3 specimens. **Holotype of *Coelorinchus divergens*:** BSKU 26865 (98.1 mm HL, 358+ mm TL), north-northwest of Amami-oshima Island, Kagoshima Pref., Okinawa Trough, Japan, East China Sea, 29.7850°N, 128.4467°E, 1000 m, F/V *Yuryo-maru*, No. 8, tr. 17, bottom trawl, coll. Y. Kinoshita and S. Hagino, 3 Feb. 1978. **Paratype of *C. divergens*:** BSKU 33464 (1, 81.6 mm HL, 315+ mm TL), west of Yaku-shima Island, Okinawa Trough, 30.2067°N, 128.2833°E, 780–810 m, F/V *Yuryo-maru*, No. 8, sta. 3, tr. T1, bottom trawl, coll. H. Maeda, 26 Oct. 1979. **Non-type: South China Sea:** ASIZP 66084 (1, 73.7 mm HL, 214+ mm TL), 20.7167°N, 117.5500°E, 954 m, otter trawl, coll. P.-F. Lee, 19 Aug. 2005.

Counts and measurements. Based on 3 specimens (73.7–98.1 mm HL, 214+–358+ mm TL). Counts: first dorsal-fin rays II, 7–8; pectoral-fin rays i17–i18; pelvic-fin rays 7; gill rakers on first arch (outer/inner) 0/7–9, on second arch 6/8–9; longitudinal scales 28–42; transverse scale rows below first dorsal-fin origin 5.5–6, below first dorsal-fin midbase 4–5.5, below second dorsal-fin origin 5–5.5.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 39–41 (64–69); orbit diameter 25–27 (41–44); postorbital length 34–36 (56–59); postrostral length 60–62; orbit–preopercle distance 35–37 (56–61); suborbital width 14–15 (22–25); upper-jaw length 28–28 (45–47); length of rictus 21–22 (34–37); length of premaxillary tooth band 12–15 (20–24); preoral length 32–36 (52–60); length of terminal snout scute 7 (12); length of lateral nasal ridge 26–29 (42–48); length of suborbital ridge 91–97 (148–162); snout width 26–31 (43–52); internasal width 20–22 (32–35); interorbital width 24–26 (40–42); occipital width 8 (14); body width over pectoral-fin bases 37–46 (62–75); body depth at first dorsal-fin origin 48 (80); body depth at anal-fin origin 37 (63); prepelvic length 103–109 (168–183); preanus length 140 (234); preanal length 143 (239); isthmus–pelvic distance 26–32 (44–52); isthmus–anal distance 63 (105); pelvic–anal distance 38 (63); pelvic-fin length 34–43 (56–70); pectoral-fin length 35–42 (58–67); predorsal length 107–112 (177–181); height of first dorsal fin 35–46 (58–74); length of first dorsal-fin base 13–15 (22–24); interdorsal length 13–14 (22–23); length of gill slit 12–14 (20–22); length of posterior nostril 7–9 (12–15); barbel length 7–9 (11–15).

Size. To about 36 cm TL (BSKU 26865, holotype, 358+ mm TL).

Distribution. Restricted to Japan and Taiwan. So far known only from the Okinawa Trough and northern South China Sea at depths of 515–1110 m (Shao *et al.* 2008a, 2008b; this study; Appendix 3-1F). Very rare.

Remarks. *Coelorinchus divergens* was described

FIGURE 30. Holotype of *Coelorinchus divergens*. BSKU 26865, 98.1 mm HL, 358+ mm TL, Okinawa Trough, East China Sea, 1000 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

from two specimens collected from the Okinawa Trough (Fig. 30), and subsequently recorded from the East and South China Seas off Taiwan (Shao *et al.* 2008a, 2008b; Iwamoto *et al.* 2015). For further morphological details see the original description given by Okamura & Yatou (1984). Despite intensive collecting efforts, no additional specimens have been collected from Japanese waters.

Relationships. *Coelorinchus divergens* falls into a group of *Coelorinchus* species characterized by the following combination of features: no external evidence of light organ anterior to anus; lateral nasal ridge incompletely supported by nasal bone; premaxillary teeth in short, uniformly wide band; underside of head entirely or almost entirely scaled; body scales covered with shield-like spinules in parallel to subparallel rows, with middle row enlarged and some rows incomplete; buttresses of body-scale spinules broadly developed; modified scales on head ridges prominently spiny and thickened; body uniformly dark brown to swarthy. This group was originally recognized by Iwamoto *et al.* (2006:437), and is called here the *C. parallelus* group. Apart from *C. divergens*, nine valid species belong to this group: *C. acanthiger* Barnard, 1925 from southern Africa across the Indian Ocean to Australia and New Zealand; *C. amirantensis* Iwamoto, Golani, Baranes & Goren, 2006 from the Seychelles and the Mascarene Ridge in the Indian Ocean; *C. braueri* Barnard, 1925 from South Africa; *C. kermadecus* Jordan & Gilbert in Jordan & Starks, 1904 from the southwestern Pacific; *C. mycterismus* McMillan & Paulin, 1993 from New Zealand and off the Pacific and Indian Ocean coasts of Australia; *C. parallelus*

(Günther, 1877) from Japan, Taiwan, and New Caledonia (a senior synonym of *C. sparsilepis* Okamura & Yatou, 1984; see the Nomenclatural discussion of *C. parallelus*); *C. spinifer* Gilbert & Hubbs, 1920 from the Gulf of Tomini, Indonesia, and the South China Sea off Taiwan; *C. trachycarus* Iwamoto, McMillan & Shcherbachev, 1999 from the southwestern Pacific and off the southern and western coasts of Australia; and *C. yurii* Iwamoto, Golani, Baranes & Goren, 2006 from the Seychelles and the Mascarene Ridge in the Indian Ocean.

Comparisons. *Coelorinchus divergens* readily differs from *C. acanthiger* in having fewer scales below the first dorsal-fin origin (5.5–6 vs. 7–10) and in lacking a prominent naked area atop the head between the parietal ridges; from *C. amirantensis* and *C. yurii* in that the underside of the head is densely scaled (vs. somewhat sparsely scaled); from *C. braueri* and *C. yurii* in having a lower first dorsal fin (height of first dorsal fin 58–74% PRL vs. >98%); from *C. mycterismus* in that the dorsal contour of the snout is almost straight (vs. distinctly concave); and from *C. trachycarus* in that the ventral surfaces of preopercles are heavily scaled (vs. broadly naked). *Coelorinchus divergens* also differs from all except *C. kermadecus*, *C. mycterismus*, *C. parallelus*, and *C. spinifer* in that the nasal fossa is heavily scaled (vs. completely or generally naked); all except *C. acanthiger*, *C. braueri*, *C. parallelus*, and *C. yurii* in having a distinctly shorter snout (64–69% PRL vs. >74%); and from all except *C. kermadecus* and *C. parallelus* in having a somewhat larger mouth (upper-jaw length 45–47% PRL vs. <45%).

FIGURE 31. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of two species of the *Coelorinchus occa* group. (A–B) *C. divergens*, BSKU 33464, paratype, 81.6 mm HL; (C–D) *C. parallelus*, BMNH 1887.12.7.65, lectotype, 93.2 mm HL; (E–F) *C. parallelus*, BSKU 28441, holotype of *C. sparsilepis*, 73.3 mm HL. (A, C, E) Views from above; (B, D, F) oblique views. [Photos: (A–B, E–F) NSMT; (C–D) N. Nakayama]

Coelorinchus divergens may be confused with *C. parallelus*, with which it occurs sympatrically in the East and South China Seas. However, these two species are readily distinguished by two squamation characters. The first character concerns spinulation of the occipital scales, and *C. divergens* readily differs from *C. parallelus* in having occipital scales covered with divergent rows of spinules (vs. uniserial). The second character involves the shape and arrangement of the body-scale spinules; although both species have prominent buttresses at each base of the spinules, those of *C. divergens* are less developed than those of *C. parallelus* (Fig. 31A–B vs. C–F); consequently, the spinules of *C. divergens* are narrowly triangular in appearance, whereas those of *C. parallelus* are broadly triangular. *Coelorinchus divergens* further differs from *C. parallelus* by having a relatively larger mouth (upper-jaw length 45–47% PRL vs. 36–46%) and a shorter snout (64–69% PRL vs. 67–89%), although these ranges are slightly overlapping.

***Coelorinchus formosanus* Okamura, 1963**

[Japanese name: Taiwan-sokodara]

(Figs. 32–33, 34A; Appendix 3-2A)

Coelorhynchus formosanus Okamura, 1963b:37, figs. 1–2 [original description; holotype: FAKU 35836, from “Tashi” (Da-xi); 4 paratypes from Taiwan]; Okamura 1970a:161, pl. XXXIV, text-figs. 65–66 (description; re-examination of type specimens; new Japanese name: “Taiwan-sokodara”); Tominaga & Uyeno 1981:488 (listed; Japan).

Coelorhynchus (Quincuncia) formosanus: Okamura 1970b: table 1 (listed; Japan).

Coelorinchus longissimus (not Matsubara 1943): Yatou 1984:223, 366, fig. 159 (brief description; Okinawa Trough; in part, BSKU 26581).

Coelorinchus formosanus: Okamura 1984b:96, pl. 82, fig. J (compiled); Okamura 1988:96, pl. 82, fig. J (compiled); Iwamoto 1990:158, fig. 372 (synopsis); Shao *et al.* 2008b: table 2 (9 spec. listed from northeastern and southwestern Taiwan, and South China Sea); Kim *et al.* 2009b:112, fig. 6 (brief description; 2 spec. from Korea; new Korean record); Ho & Shao 2011:36 (type catalog); Nakabo & Kai 2013:506 (in key; Japan); Iwamoto *et al.* 2015:54 (brief description; 38 spec. from northeastern Taiwan); Motomura 2020:38 (listed; Japan).

Coelorinchus formosanus: Nakabo 1993:366 (in key; Japan); Shao 1993:168 (compiled; Taiwan); Nakabo 2000:430 (in key; Japan); Nakabo 2002:430 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Shinohara *et al.* 2005:416 (listed; Ryukyu Islands).

Coelorinchus (Quincuncia) formosanus: Chiou *et al.* 2004a: table 1 (listed; Taiwan).

Diagnosis. Light organ tubular, completely covered with scales, externally represented by long black streak

extending from immediately anterior to anus to chest just posterior to isthmus. Underside of snout completely naked except for overlapping scales along lateral nasal ridges; underside of head posterior to level of midorbit heavily scaled, but anterior portions of mandibular rami mostly naked; top of snout broadly naked along each side of median rostral ridge. Snout long, sharply pointed, length 66–84% PRL, its dorsal profile slightly concave in lateral view; terminal scute short, arrowhead-shaped, dorsoventrally flattened, length 7–18% PRL. Lateral nasal ridge incompletely supported by nasal bone. Anus abutting anal-fin origin. Premaxillary teeth small, slender, conical in moderately wide, long, tapered band, with outer series slightly enlarged; posterior margin of tooth band almost reaching lateral corner of mouth. Body scales covered with short, spike-like, trihedral spinules in irregularly quincunx order to widely divergent rows; buttresses well developed. Second dorsal fin distinctly lower than anal fin. Upper-jaw length 45–51% PRL; pectoral-fin length 69–75% PRL; height of first dorsal fin 74–85% PRL; interdorsal length 35–50% PRL; length of gill slit 21–26% PRL; barbel length 20–26% PRL. Dorsal half of body with faint, dark, irregular blotches; lower half of head and body silvery white when viewed laterally; median nasal bone blackish; underside of head generally pale; black hair-like papillae scarcely developed on naked areas on head; lips dusky; oral cavity pale; gular and branchiostegal membranes heavily peppered with large melanophores; chest dark; first dorsal fin dusky, but second spinous ray prominently blackish; pelvic fin darker distally.

Material examined. 23 specimens. **Holotype of *Coelorhynchus formosanus*:** FAKU 35856 (72.8 mm HL, 221+ mm TL), Da-xi fish market, Yilan, northwestern Taiwan, bottom trawl, 6 Dec. 1961. **Paratypes of *C. formosanus*:** FAKU 35859 (1, 62.3 mm HL, 238+ mm TL), FAKU 35860 (1, 57.8 mm HL, 212+ mm TL), Da-xi fish market, Yilan, bottom trawl, 1958–1961; *FAKU 35857 (1, 46.7 mm HL, 167+ mm TL), *FAKU 35858 (1, 55.4 mm HL, 195+ mm TL), Kaoshiung, South China Sea, bottom trawl, 1958–1961. **Non-types: Japan:** BSKU 159 (1, 69.0 mm HL, 272+ mm TL), Mimase fish market, bottom trawl, 21 Mar. 1951. **Taiwan:** BSKU 108963 (1, 79.9 mm HL, 318+ mm TL), Da-xi fish market, Yilan, 25 Jul. 2012; BSKU 98972 (1, 57.9 mm HL, 216+ mm TL), Da-xi fish market, Yilan, coll. H.-C. Ho, 8 Nov. 2007; BSKU 50363 (1, 51.3 mm HL, 215 mm TL), BSKU 50466 (1, 37.6 mm HL, 128+ mm TL), BSKU 50468 (1, 53.2 mm HL, 189+ mm TL), BSKU 50469 (1, 38.9 mm HL, 152+ mm TL), BSKU 50471 (1, 51.8 mm HL, 210+ mm TL), BSKU 50472 (1, 54.3 mm HL, 200+ mm TL), BSKU 50473 (1, 59.2 mm HL, 214+ mm TL), BSKU 50474 (1, 57.4 mm HL, 217+ mm TL), BSKU 50488 (1, 38.6 mm HL, 158 mm TL), Da-xi fish market, Yilan, coll. T. Yamakawa and R. Asaoka, 23 Apr. 2012; *BSKU 99863 (1, ca. 72 mm HL, 302 mm TL), BSKU 99866 (1, 66.1 mm HL, 216+ mm TL), Da-xi fish market, Yilan, coll.

M.-Y. Lee, 30 Jun. 2009; KAUM-I. 44655 (1, 69.7 mm HL, 204+ mm TL), off Kaohsiung, Yilan, 400 m, trawl, coll. M. Matsunuma and H.-C. Ho, 1 Jul. 2011; BSKU 29993 (1, 52.5 mm HL, 184+ mm TL), BSKU 29994 (1, 40.7 mm HL, 155+ mm TL), Dong-gang, Pingtung, otter trawl, coll. T. Kanayama, 6–7 Dec. 1978; *FAKU 204426 (1, 367+ mm TL), Dong-gang fish market, Pingtung, bottom trawl, coll. N. Nakayama, 27 Feb. 2017.

Counts and measurements. Based on 19 specimens (37.6–79.9 mm HL, 128+–318+ mm TL). Counts: first dorsal-fin rays II,8–10; pectoral-fin rays i15–i17; pelvic-fin rays 7; gill rakers on first arch (outer/inner) 0/7–9, on second arch 6–7/8–9; longitudinal scales 36–45; transverse scale rows below first dorsal-fin origin 4.5–6.5, below first dorsal-fin midbase 3.5–5, below second dorsal-fin origin 3.5–5, above anal-fin origin 11.5–14.5.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 40–46 (66–84); orbit diameter 23–27 (41–47); postorbital length 30–36 (54–59); postrostral length 54–61; orbit–preopercle distance 33–37 (58–66); suborbital width 10–12 (17–20); upper-jaw length 25–30 (45–51); length of rictus 20–24 (36–43); length of premaxillary tooth band 17–22 (30–36); preoral length 33–39 (55–72); length of terminal snout scute 4–10 (7–18); length of lateral nasal ridge 26–34 (43–61); length of suborbital ridge 93–99 (157–180); snout width 22–25 (38–45); internasal width 18–21 (31–38); interorbital width 21–24 (37–43); occipital width 6–9 (10–15); body width over pectoral-fin bases 32–44 (58–81); body depth at first dorsal-fin origin 47–57 (82–104); body depth at anal-fin origin 41–47 (70–86); prepelvic length 100–109 (171–195); preanus length 141–152 (249–270); preanal length 144–156 (256–273); isthmus–pelvic distance 30–39 (53–70); isthmus–anus distance 71–83 (126–146); isthmus–anal distance 73–87 (130–153); pelvic–anal distance 40–54 (72–94); anus–anal distance 2–4 (4–7); pelvic-fin length 26–35 (47–61); pectoral-fin length 38–44 (69–75); predorsal length 102–110 (178–200); height of first dorsal fin 41–47 (74–85); length of first dorsal-fin base 17–20 (29–36); interdorsal length 20–28 (35–50); length of gill slit 12–14 (21–26); length of posterior nostril 6–8 (9–14); barbel length 11–15 (20–26).

Size. To about 32 cm TL (BSKU 108963, 318+ mm TL, Da-xi, northern Taiwan).

Distribution. Restricted to the northwestern Pacific off Taiwan, Japan, and Korea. Known from the northern South China Sea, Okinawa Trough, Tosa Bay, and Sea of Japan off the Korean Peninsula (Chu & Lo *in Chu et al.* 1963; Shao *et al.* 2008a, 2008b; Kim *et al.* 2009b; this study; Appendix 3-2A). Depth range 158–400 m, plus a single capture in 985–1110 m (ASIZP 65595; Iwamoto *et al.* 2015). One of the most common grenadiers found in fish markets in Taiwan, but rare in Japanese waters.

Remarks. *Coelorinchus formosanus* was described by Okamura (1963b) based on five specimens collected

from northeastern Taiwan (Fig. 33). The original description is excellent and can be referred to for details. In the same year, Chu & Lo (*in Chu et al.* 1963) described *C. abbreviatus* and *C. intermedius* from the northern part of the East China Sea. Although Okamura (1970a) subsequently regarded these species synonymous, *C. abbreviatus* and *C. intermedius* are considered as junior synonyms of *C. multispinulosus* Nakayama, 1942 in this study (see the Nomenclatural discussion of *C. multispinulosus*).

Relationships. *Coelorinchus formosanus* belongs to the *C. argentatus* group as defined by Nakayama *et al.* (2020), which includes the following 13 species: *C. argentatus* Smith & Radcliffe *in* Radcliffe, 1912 from the Philippines, New Caledonia and northern Australia; *C. denticulatus* Regan, 1921 from the western Indian Ocean; *C. formosanus*; *C. gaesorhynchus* Iwamoto & Williams, 1999 from Western Australia; *C. kamoharai* Matsubara, 1943 and *C. longissimus* Matsubara, 1943 from Japan to Taiwan; *C. mayiae* Iwamoto & Williams, 1999 from the Arafura Sea; *C. multispinulosus* from Japan to Vietnam; *C. pardus* Iwamoto & Williams, 1999 from the Arafura Sea; *C. posteromaculatus* Nakayama, Prokofiev & Kawai, 2020 from the eastern Indian Ocean; *C. quincunciatus* Gilbert & Hubbs, 1920 from the Philippines; *C. semaphoreus* Iwamoto & Merrett, 1997 from New Caledonia and Queensland, Australia; and *C. thompsoni* Gilbert & Hubbs, 1920 from the Philippines. These species are distinguished from other congeners by having the following combination of features (Nakayama *et al.* 2020): light organ long, extending from anus to chest just posterior to isthmus; no interspace between anus and anal-fin origin; lateral nasal ridge incompletely supported by nasal bone; premaxillary teeth in long tapered band, with posterior margin of tooth band almost reaching lateral corner of mouth; body scales covered with short, erect spinules in quincunx order or subparallel to slightly divergent rows; modified scales on head ridges not prominently spinulated; dorsal surfaces of snout broadly naked along each side of median rostral ridge; second dorsal fin poorly developed, its height distinctly lower than that of anal fin; dark prominent markings, such as spots, blotches, stripes, and vermiculations, usually present dorsally on trunk and tail; lower half of body overlain with silvery reflections (often indistinct in preserved specimens). The above definition generally agrees with the original concept of the subgenus *Quincuncia* Gilbert & Hubbs, 1920 that initially comprised *C. argentatus* (type species), *C. quincunciatus* and *C. thompsoni* (see Nakayama *et al.* 2020).

McCann & McKnight (1980) described *C. campbellicus* from the southeast of New Zealand, classifying the species under *Quincuncia*. This species is excluded from the *C. argentatus* group, because of its short light organ [43% HL (fide McCann & McKnight 1980:57) vs. about 70% in species of the *C. argentatus*

FIGURE 32. Fresh specimen of *Coelorinchus formosanus*. FAKU 204426, 367+ mm TL, Dong-gang fish market, southwestern Taiwan, South China Sea. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

FIGURE 33. Holotype of *Coelorhynchus formosanus* (= *Coelorinchus formosanus*). FAKU 35856, 72.8 mm HL, 221+ mm TL, Da-xi fish market, northeastern Taiwan, East China Sea. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

group]. McCann & McKnight (1980:57) also described *C. campbellicus* as having “anterior photophore covered by scales, posterior ovoid, naked”, but the light organ of the *C. argentatus* group is completely covered with scales. The species seems to be closely related to *C. kaiyomaru* Arai & Iwamoto, 1979, widely distributed in temperate waters in the Southern Hemisphere.

Comparisons. *Coelorinchus formosanus* is distinctive among the *C. argentatus* group in that the underside of the head is heavily scaled posteriorly, but almost completely naked anteriorly (vs. either almost entirely scaled or naked). It further differs from all except large specimens of *C. argentatus* in having body scales

covered with short, spike-like, trihedral spinules with well-developed buttresses (Fig. 34A vs. spinules needle-like or triangular, and buttresses poorly developed, Fig. 34B–D).

In Japanese waters, *C. formosanus* might be confused with *C. longissimus* due to their similar appearance and coloration. In addition to the above-mentioned features, *C. formosanus* is readily distinguished from *C. longissimus* by having a longer barbel (20–26% PRL vs. 11–18%) and pectoral fin (69–75% PRL vs. 55–64%), higher first dorsal fin (74–85% PRL vs. 58–70%), and body-scale spinules arranged in more or less quincunx order (vs. subparallel narrowly divergent rows). Large individuals of *C. longissimus* occasionally have rudimentary scaly

FIGURE 34. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of four species of the *Coelorinchus argentatus* group. (A) *C. formosanus*, BSKU 99863, ca. 72 mm HL; (B) *C. kamoharai*, BSKU 93575, 59.6 mm HL; (C) *C. longissimus*, BSKU 40448, 95.9 mm HL; (D) *C. multispinulosus*, BSKU 99273, 69.9 mm HL. Views from above. [Photos: N. Nakayama]

patch(es) on the ventral surfaces of the preopercles, but the mandibular rami are always naked, except the posterior margins where a few scales are rarely present (vs. usually heavily scaled in *C. formosanus*).

***Coelorinchus fuscigulus* Iwamoto, Ho & Shao, 2009**

[Japanese name: Shima-sokodara]

(Figs. 28E–F, 35–36; Appendix 3-2B)

Coelorinchus cingulatus (not Gilbert & Hubbs 1920): Shao *et al.* 2008b: table 2 (listed; in part, 1 spec. from northeastern Taiwan, 400–600 m).

Coelorinchus cylindricus (not Iwamoto & Merrett 1997): Shao *et al.* 2008b: table 2 [1 spec. listed from northeastern Taiwan (as first record from Taiwan)].

Coelorinchus fuscigulus Iwamoto, Ho & Shao, 2009:40, figs. 1–2 [original description; holotype: ASIZP 70169, from Ta-shi (= Da-xi), northeastern Taiwan; 9 paratypes from north

eastern Taiwan]; Nakabo & Kai 2013:508 (in key; Japan; new Japanese name “Shima-sokodara”); Iwamoto *et al.* 2015:55 (brief description based on type specimens); Motomura 2020:38 (listed; Japan).

Diagnosis. Ventral light organ externally represented by long dark line, terminated with 1 fossa at each end, one immediately anterior to anus, the other on chest just posterior to isthmus (duct connecting each fossa often difficult to distinguish). Underside of head completely naked except along lateral nasal ridges; dorsal surface of snout mostly scaled, but prominent naked clefts present along each side of median rostral ridge; occipital area between parietal ridges without prominent naked area. Snout long, sharply pointed, length 60–72% PRL; dorsal profile of snout almost straight to gently concave in lateral view; terminal scute short, arrowhead-shaped, dorsoventrally flattened, length 14–21% PRL; distal portion of snout narrowly attenuated when viewed

dorsally; lateral nasal ridge incompletely supported by nasal bone. Interdorsal space less than first dorsal-fin base length. Anal fin originating posterior to vertical through second dorsal-fin origin. First dorsal fin high, but second spinous ray not especially elongate, height of fin 80–106% PRL. Second dorsal fin well developed, its rays as long as opposites of anal fin. Anus immediately anterior to anal-fin origin. Premaxillary teeth small, slender, conical in short, uniformly wide band, with outer series slightly enlarged; posterior end of tooth band falling far short of lateral corner of mouth. Body scales covered with short, reclined, needle-like spinules in subparallel to slightly divergent rows; spinules in each row greatly overlapping, but free from one another, with last spinule extending well beyond posterior scale margin; buttresses scarcely developed. Orbit diameter 40–43% PRL; postorbital length 58–61% PRL; orbit–preopercle distance 55–61% PRL; upper-jaw length 41–46% PRL; preoral length 50–64% PRL; body depth at first dorsal-fin origin 84–90% PRL; isthmus–anal distance 132–158%; length of gill slit 20–23% PRL; barbel length 13–17% PRL. Transverse scale rows below first dorsal-fin midbase 5.5–7.5. Spinules on scales anterior to light organ not rudimentary. About 8–11 dark saddles on body (often faint in large specimens); scale pockets outlined in black; underside of head dark to dusky, heavily peppered; lips pale; oral cavity, and gular and branchiostegal membranes prominently blackish; first dorsal fin dark, with white apical tip; pelvic fin dark, but outermost ray paler distally.

Material examined. 13 specimens. **Holotype of *Coelorinchus fuscigulus*:** ASIZP 70169 (76.8 mm HL, 322+ mm TL), Da-xi fish market, Yilan, northeastern Taiwan, bottom trawl, coll. H.-C. Ho, 23 Mar. 2007. **Paratypes of *C. fuscigulus*:** ASIZP 63249 (1, 56.1 mm HL, 238 mm TL), Da-xi fish market, Yilan, bottom trawl, coll. H.-C. Ho, 21 Mar. 2004; ASIZP 70168 (1, 77.4 mm HL, 300+ mm TL), CAS 228337 (2, ex. ASIZP 70168, 69.3–74.1 mm HL, 283+–303+ mm TL), Da-xi fish market, Yilan, bottom trawl, coll. H.-C. Ho, 29 Jan. 2007; ASIZP 66922 (1, 66.5 mm HL, 286 mm TL), off northeastern Taiwan, Okinawa Trough, 24.8500°N, 122.0333°E, 536 m, R/V *Ocean Researcher I*, sta. CP248, French-type beam trawl, coll. H.-M. Yeh *et al.*, 28 Aug. 2004; ASIZP 66973 (1, 72.8 mm HL, 291 mm TL), Nan-fang-ao fish market, Yilan, bottom trawl, coll. H.-C. Ho, 26 Jan. 2007; CAS 228338 [1, ex. CAS 224583 (in part), 65.8 mm HL, 293+ mm TL], Nan-fang-ao fish market, Yilan, bottom trawl, coll. H.-C. Ho, 30 Aug. 2005; ASIZP 63193 (1, 53.7 mm HL, 226 mm TL), CAS 224492 (1, ex. ASIZP 63193, 44.4 mm HL, 189 mm TL), off Senkaku Island, Okinawa Trough, 25.75°N, 123.48°E, bottom trawl, coll. H.-C. Ho, 24 Apr. 2004. **Non-types: Japan:** KAUM-I. 55534 (1, 65.0 mm HL, 265+ mm TL), northwest of Kuro-shima Island, 30.9877°N, 129.5725°E, 300–400 m, coll. Y. Fukui and M. Matsunuma, 16 Jul. 2013; BSKU 106760 (1, 41.9 mm HL, 174+ mm TL), southeast of Shimokoshiki-jima

Island, East China Sea, 31.5658°N, 129.8915°E, 380 m, F/V *Maruko-maru*, tr. 1, bottom trawl, coll. N. Nakayama *et al.*, 24 Apr. 2012. **Taiwan:** *BSKU 108968 (1, 67.2 mm HL, 295+ mm TL), Da-xi fish market, Yilan, bottom trawl, coll. N. Nakayama and M.-Y. Lee, 25 Jul. 2012.

Counts and measurements. Based on 12 specimens (41.9–77.4 mm HL, 174+–322+ mm TL). Counts: first dorsal-fin rays II, 8–9; pectoral-fin rays i16–i18; pelvic-fin rays 7; gill rakers on first arch (outer/inner) 0/7–9, on second arch 5–6/7–9; longitudinal scales 30–36; transverse scale rows below first dorsal-fin origin 6.5–8.5, below first dorsal-fin midbase 5.5–7.5, below second dorsal-fin origin 6–8, above anal-fin origin 21.5–22.5; pyloric caeca 19.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 37–42 (60–72); orbit diameter 24–26 (40–43); postorbital length 34–38 (58–61); postrostral length 58–63; orbit–preopercle distance 33–38 (55–61); suborbital width 12–14 (20–23); upper-jaw length 25–27 (41–46); length of rictus 18–21 (31–36); length of premaxillary tooth band 10–13 (16–21); preoral length 31–38 (50–64); length of terminal snout scute 9–12 (14–21); length of lateral nasal ridge 25–30 (40–52); length of suborbital ridge 89–92 (143–158); snout width 18–28 (31–48); internasal width 17–21 (26–36); interorbital width 20–23 (33–38); occipital width 9–11 (15–18); body width over pectoral-fin bases 36–48 (61–78); body depth at first dorsal-fin origin 49–56 (84–90); body depth at anal-fin origin 40–48 (66–79); prepelvic length 103–112 (169–185); preanus length 150–165 (245–270); preanal length 144–169 (243–277); isthmus–pelvic distance 29–37 (49–59); isthmus–anus distance 80–96 (134–152); isthmus–anal distance 79–100 (132–158); pelvic–anal distance 45–68 (75–111); anus–anal distance 4 (6–7); pelvic-fin length 36–44 (58–74); pectoral-fin length 39–45 (64–76); predorsal length 106–114 (178–187); height of first dorsal fin 50–66 (80–106); length of first dorsal-fin base 17–23 (28–38); interdorsal length 10–14 (15–24); length of gill slit 12–14 (20–23); length of posterior nostril 7–9 (11–15); barbel length 8–11 (13–17).

Description of Japanese specimens. General features are shown in Figs. 35–36. Body slender, deepest below first dorsal-fin origin, progressively tapering into long, laterally compressed tail. Trunk subcylindrical in cross section, width across pectoral-fin bases about 1.2 in depth below first dorsal-fin origin. Head large, shallow, HL about 4.1–4.2 in TL. Snout long, sharply pointed, protruding well beyond upper jaw, length 1.7 times as long as orbit diameter; dorsal profile almost straight in lateral view, but gently upturned near tip; ventral surface slightly inflated when fresh, but concave in preserved specimens; lateral nasal ridge incompletely supported by nasal bone; anterolateral margins of snout slightly convex when viewed dorsally. Orbit large, oval in outline, greatest diameter about 1.4–1.5 in postorbital length. Interorbital space broad, slightly concave in preserved

FIGURE 35. Fresh specimens of *Coelorinchus fuscigulus*. (A–C) BSKU 106760, 41.9 mm HL, 174+ mm TL, southeast of Shimokoshiki-jima Island, Kagoshima Pref., East China Sea, 380 m depth; (D) BSKU 108968, 67.2 mm HL, 295+ mm TL, Da-xi fish market, northeastern Taiwan, East China Sea. (A, D) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: (A–C) N. Nakayama; (D) M.-Y. Lee]

specimen, width 1.2 in orbit diameter. Mouth moderate in size, inferior, protrusible; upper-jaw length about equal to orbit diameter; anterior margin of upper lip below anterior nostril; posterior margin of maxilla extending to below hind 1/3 of orbit; lateral corner of mouth slightly restricted by lip folds; lips slightly papillose near teeth. Suborbital region narrow, sharply divided in dorsal and ventral parts by longitudinal ridge passing from tip of snout to posterior angle of preopercle. Preopercle large, moderately lobed posteroventrally; hind margin oblique, forming angle of about 60° to horizontal axis of head and body. Interopercle completely hidden behind preopercle. Subopercle produced as slender tip posteroventrally. Gill membranes thick, broadly connected across isthmus, with narrow posterior free fold. Gill opening narrow, not reaching forward to vertical through hind margin of orbit. Outer gill slit moderately restricted by skin folds, length 1.8–2.1 in orbit diameter. Gill rakers developed into small, spinous tubercles, but those on epibranchial somewhat pad-like; no rakers on outer side of first arch and inner side of fourth arch. Chin barbel short, terminating as fine tip, length 2.6–2.7 in orbit diameter.

Anus immediately anterior to anal-fin origin. Two dermal fossae of light organ on midventral line of abdomen; anterior fossa oval, situated midway between isthmus and line connecting inner bases of pelvic fin; posterior window small, scarcely discernible, located just anterior to anus.

Teeth small, conical, gently incurved, in bands in both jaws; outer series slightly enlarged, followed by progressively smaller teeth medially. Premaxillary band short, almost uniformly broad, about 5-teeth wide anteriorly, extending to posterior 2/3 of rictus. Mandibular teeth in narrow tapered band, about 4-teeth wide near symphysis; posterior margin of tooth band nearly reaching lateral corner of mouth. All teeth deeply embedded in gum papillae.

Body scales moderately large, thin, somewhat deciduous; those on dorsum below interdorsal space covered with short, slender, needle-like spinules in 6–7 subparallel to slightly divergent, ridge-like rows (Fig. 28E); middle row longest but spinules not especially high or enlarged; spinules reclined about 25° to scale surface and greatly overlapping; spinules in each row

FIGURE 36. Holotype of *Coelorinchus fuscigulus*. ASIZP 70169, 76.8 mm HL, 322+ mm TL, Da-xi fish market, northeastern Taiwan, East China Sea. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

not adjoined to one another, forming reclined comb-like ridges; height of spinules increasing posteriorly, with last in each row extending well beyond posterior scale margin; lateral buttresses of spinules poorly developed; no reticulate structures on unexposed portion. Scales on isthmus much smaller than those on dorsum, covered with much thicker spinules arranged in 4–6 subparallel rows (Fig. 28F). Scales becoming cycloid on light organ, beneath gill cover, and posterior to pectoral and pelvic fins. Body uniformly scaled, but all fins naked.

Scales on head ridges thickened, weakly spinulated; those on median rostral ridge covered with divergent rows of spinules. Terminal snout scute short, arrowhead-shaped, dorsoventrally flattened, its length 2.0 in orbit diameter. Supraoccipital scute rudimentary; posttemporal scutes thickened, elongated, armed with 1–2 rows of spinules. Scales on head generally similar to those on body, but variable in size and spinulation. Occipital scales covered with long, erect, needle-like spinules in narrowly divergent rows; spinules in each row free from one another, forming comb-like ridges, and height of spinules gradually increasing posteriorly; scales on dorsal surfaces of snout posterior to leading edges armed with small cluster of similar spinules. Other scales on head similar to those on body, but spinules arranged in parallel to subparallel rows; those on gill cover and between parietal and postorbital ridges largest. Nasal fossa narrowly scaled ventrally. Dorsal surfaces of snout mostly scaled, but narrow Y-shaped clefts along each side of median rostral ridge naked. Nape between occipital ridges without prominent naked area. Underside of head completely naked, except for characteristic overlapping scales along lateral nasal ridges.

No open pores along cephalic sensory canals. Small, elongated free neuromasts serially arranged along ventral surfaces of infraorbital canals (prominently marked when

fresh). Anterior nostril small and circular; posterior large, elongated slit; septum between nostrils forming anteriorly opened hood. Grooved lateral line well developed, not interrupted throughout.

Origins of first dorsal and pelvic fins slightly posterior to vertical through pectoral-fin base; first dorsal fin moderately high, its height 2.8 times as long as its base length; second spinous ray not elongated, almost entirely smooth along its leading edge (small fine spine present near tip in BSKU 106760); its tip extending to base of 7th–9th rays of second dorsal fin when laid back. Interdorsal space equal to half length of first dorsal-fin base. Origins of second dorsal and anal fins on about same vertical. Outer pelvic-fin ray prolonged as fine distal tip, reaching posterior fossa of light organ or beyond when laid back.

Color when fresh (Fig. 35). Ground color of head and body purplish gray dorsally, paler ventrally; abdomen tinged with blue in ventral view; anterior and posterior fossae of light organ marked with dark blue; about 8 broad dark saddles on trunk and tail; anteriormost saddle located at level of first dorsal-fin origin; second saddle faint, diagonal, extending from second dorsal-fin origin to pectoral-fin base; bands farther posterior generally broader and prominent; upper part of opercle much darker than surroundings; lips pale; underside of head generally dusky, heavily peppered, but lower preopercles paler; branchiostegal membranes dull black behind gill apparatus; first dorsal fin dark with white apical tip; pectoral fin pale with blackish uppermost ray; pelvic fin dark, but outermost ray white; second dorsal and anal fin pale to dusky; small intense black spot at origin of first dorsal fin and each base of pectoral and pelvic fins.

Color in alcohol (Fig. 36). Almost same as in fresh condition; oral and gill cavities blackish; gill rakers and arches dusky, filaments yellowish brown.

Size. To about 30 cm TL (CAS 228337, paratype, 303+ mm TL, Da-xi, northern Taiwan).

Distribution. Restricted to Japan and Taiwan (Appendix 3-2B). So far known only from the Okinawa Trough and northern South China Sea at depths of 380–536 m (Iwamoto *et al.* 2009; this study). Very rare in Japanese waters.

Remarks. The original description given by Iwamoto *et al.* (2009) is excellent and can also be referred to for details. *Coelorinchus fuscigulus* was originally described from 10 specimens collected from the East China Sea (Fig. 36), some of which were initially reported as *C. cylindricus* Iwamoto & Merrett, 1997 by Shao *et al.* (2008b: table 2). Interestingly, *C. fuscigulus* was not recorded by previous Japanese ichthyologists, despite their extensive collecting efforts in the East China Sea (e.g., Okamura & Kitajima 1984; Okamura 1985; Yamada 1986; Shinohara *et al.* 2005; Yamada *et al.* 2007; Furuhashi *et al.* 2010). Iwamoto *et al.* (2009) indicated the collection site of CAS 224492 (paratype, 44.4 mm HL) as “25°75'N, 123°48'E, Diaoyutai Archipelago, Yilan, Taiwan” [sic], but the coordinates fall inside Japan’s EEZ (it should also be read as 25.75°N, 123.48°E). Accordingly, Nakabo & Kai (2013) subsequently proposed a Japanese name “Shima-sokodara” for the species. Two additional Japanese specimens are herein recorded from the East China Sea off Kagoshima Pref., one of which from off Shimokoshiki-jima Island (BSKU 106760, 41.9 mm HL; Fig. 35A–C) represents the northernmost record and the smallest specimen of *C. fuscigulus*.

The presence of dark saddles dorsally on the body is one of the important diagnostic features of *C. fuscigulus*. While these saddles are prominent in small specimens (Fig. 35A), they are often indistinct in large adults (Fig. 35D).

Relationships and comparisons. *Coelorinchus fuscigulus* is one of 10 species of the *C. hubbsi* group (see the Relationships of *C. brevirostris*), and closely resembles the following three species: *C. cylindricus* Iwamoto & Merrett, 1997 known from New Caledonia and New Zealand; *C. melanosagmatus* Iwamoto & Anderson, 1999 widely distributed in the Indian Ocean; and *C. multifasciatus* Sazonov & Iwamoto, 1992 represented by only the holotype collected from the Sala y Gomez Ridge. They share the presence of similar body markings consisting of more than six dark saddles on the body, and the absence of a large dark blotch between the lateral line and the pectoral-fin base. *Coelorinchus fuscigulus* is distinguished from these three species by having more scales below the midbase of the first dorsal fin (5.5–7.7 vs. 4.5, 4–5, and 5 in *C. cylindricus*, *C. melanosagmatus*, and *C. multifasciatus* respectively) and a greater body depth at the first dorsal-fin origin (84–90% PRL vs. 77%, 62–83%, and 81% respectively). It further differs from *C. cylindricus* in having a smaller orbit (40–43% PRL vs. 46%), longer postorbital (58–61% PRL vs. 55%) and upper-jaw (41–46% PRL vs. 36%) lengths, a greater

orbit–preopercle distance (55–61% PRL vs. 50%), and a wider outer gill slit (20–23% PRL vs. 18%). It also differs from *C. melanosagmatus* in having a shorter snout (60–72% PRL vs. 74–95%), greater orbit–preopercle (55–61% PRL vs. 48–54%) and isthmus–anal (132–158% PRL vs. 90–118%) distances, and a longer chin barbel (13–17% PRL vs. 7–13%). The holotype of *C. multifasciatus* (ZMMGU P.18117, 32.3 mm HL; examined here) is much smaller than the *C. fuscigulus* specimens examined (41.9–77.4 mm HL), which makes comparisons of their morphometric characters difficult. However, *C. fuscigulus* differs notably from that species in lacking prominent naked areas along each side the median rostral ridge and atop the head between the parietal ridges (vs. present in *C. multifasciatus*).

Coelorinchus gilberti Jordan & Hubbs, 1925

[Japanese name: Oni-hige]

(Figs. 37–39; Appendix 3-2C)

Macrurus parallelus (not Günther 1877): Günther 1877:439 (original description; in part, 2 paralectotypes, BMNH 1887.12.7.67–68, off Eno-shima Island, Sagami Bay, *Challenger* sta. 232); Günther 1887:125 (description; in part, 2 paralectotypes, BMNH 1887.12.7.67–68, off Eno-shima Island, Sagami Bay, *Challenger* sta. 232).

Coelorhynchus japonicus (not Temminck & Schlegel 1846): Jordan & Thompson 1914:306 [2 spec. from Misaki; FMNH 57268 (ex. CM 6207a)].

Coelorhynchus gilberti Jordan & Hubbs, 1925:327 [original description; holotype: FMNH 59401 (ex. CM 7960), from “fish-market at Shizuoka”].

Coelorhynchus gilberti: Okada & Matsubara 1938:451 (in key; Japan); Kamohara 1950:277 (listed; Kochi and Wakayama Pref.; Japanese name: “Oni-hige”); Kuroda 1951:392 (listed; Suruga Bay); Kamohara 1952:99 (spec. from Kochi Pref.); Matsubara 1955:1313 (in key; Japan); Kamohara 1958:73 (listed; Kochi Pref.); Kobayashi & Ikeda 1962:20, unnumbered fig. (spec. from Shirikishinai, Pacific off southern Hokkaido); Kamohara 1964:96 (listed; Kochi Pref.); Ueno 1965:19 (listed; Pacific off Hokkaido); Okamura 1970a:201, pl. XLIV, text-fig. 87 (description; 23 spec. from Pacific off Japan from Hokkaido to Misaki); Maruyama 1971:33 (listed; Iwate Pref.); Ueno 1971:101 (listed; Hokkaido); Tominaga & Uyeno 1981:488 (listed; Japan); Okamura 1982:181, 354, fig. 107 (brief description; 8 spec. from Kyushu-Palau Ridge and Tosa Bay; photos based on BSKU 29453 and BSKU 29509); Shiogaki 1982:34 (listed; Aomori Pref.); Sawada 1983:113, 196, fig. 65 (brief description; 4 spec. from Pacific off Tohoku; photo based on HUMZ 87121).

Coelorhynchus kanetomoi Matsubara & Asano, 1943:95 (original description; holotype: SKFC 2, from Misaki, in ca. 700 m; new Japanese name: “Syôno-fue”); Matsubara 1955:1314 (in key; Japan).

[?] *Coelorhynchus japonicus* (not Temminck & Schlegel 1846):

- Ueno 1965:19 (listed; Pacific off Hokkaido); Maruyama 1971:33 (listed; Iwate Pref.); Ueno 1971:101 (listed; Hokkaido).
- [?] *Coelorhynchus parallelus* (not Günther 1877): Ueno 1965:19 (listed; Pacific off Hokkaido); Ueno 1971:101 (listed; Hokkaido).
- Coelorhynchus (Oxymacrus) gilberti*: Okamura 1970b: table 1 (listed; Japan).
- Coelorhynchus longicephalus* (not Okamura 1982): Okamura 1982:177, fig. 105 [original description; in part, BSKU 26102; Kyushu-Palau Ridge; see Nakayama *et al.* (2015c)].
- Coelorinchus gilberti*: Okamura 1984b:97, pl. 345, fig. J (compiled); Okamura 1988:97, pl. 345, fig. J (compiled); Maeda & Maruyama 1991:366 (listed; Hokkaido); Shiohaki *et al.* 1992:29 (listed; Aomori Pref.); Amaoka *et al.* 1995:104, fig. 155 (compiled; northern Japan); Suzuki & Kataoka 1997:84, pl. 34, fig. 184 (brief description; 1 spec. from off Kii Peninsula); Okamura 1997:127, fig. 19 (compiled); Goto 2000:6 (19 spec. from Iwate Pref.); Shinohara *et al.* 2009:708 (6 spec. listed from Pacific off Tohoku); Nakabo & Kai 2013:509 (in key; Japan); Ikeda & Nakabo 2015:321, pl. 67, fig. 11 (brief description; spec. from Wakayama Pref.); Amaoka *et al.* 2020:164, fig. 222 (listed; Hokkaido); Motomura 2020:38 (listed; Japan).
- [?] *Coelorinchus japonicus* (not Temminck & Schlegel 1846): Maeda & Maruyama 1991:366 (listed with a question mark; Hokkaido); Amaoka *et al.* 2020:164, fig. 223 (listed; Hokkaido).
- [?] *Coelorinchus parallelus* (not Günther 1877): Maeda & Maruyama 1991:366 (listed with a question mark; Hokkaido).
- Caelorinchus gilberti*: Nakabo 1993:368 (in key; Japan); Shinohara *et al.* 1996:168 (4 spec. listed from Pacific off Tohoku; 4 spec.); Shinohara & Matsuura 1997:290 (listed; Suruga Bay); Nakabo 2000:432 (in key; Japan); Shinohara *et al.* 2001:304 (2 spec. listed from Tosa Bay); Nakabo 2002:432 (in key; Japan); Maeda & Tsutsui 2003:488 (listed; Hokkaido); Shiohaki *et al.* 2004:55 (listed; Pacific off Aomori Pref.); Suetsugu & Ohta 2005: table 3 (listed; Enshu-nada); Senou *et al.* 2006:420 (listed; Sagami Sea); Kitagawa *et al.* 2008:41, unnumbered fig. (brief description; spec. from Pacific off Tohoku); Matsuura *et al.* 2009:24 (1 spec. from Miyagi Pref.); Amaoka *et al.* 2011:131, fig. 166 (compiled; Hokkaido).
- [?] *Caelorinchus japonicus* (not Temminck & Schlegel 1846): Maeda & Tsutsui 2003:488 (listed with a question mark; Hokkaido).
- [?] *Caelorinchus parallelus* (not Günther 1877): Maeda & Tsutsui 2003:488 (listed with a question mark; Hokkaido).
- [?] *Caelorinchus (Oxygadus) gilberti*: Chiou *et al.* 2004a: table 1 (questionable; listed without voucher spec.; Taiwan).
- Caelorinchus anatirostris* (not Jordan & Gilbert in Jordan & Starks 1904): Matsuura *et al.* 2009:24 (1 spec. from Miyagi Pref.; NSMT-P 74846).
- Caelorinchus* sp.: Matsuura *et al.* 2009:24 (in part, NSMT-P 74239; 1 spec. from Iwate Pref.).

Diagnosis. No external evidence of light organ anterior to anus. Underside of head almost completely naked; narrow scaly patches usually present on mandibular rami and areas below infraorbital and preopercular ridges; dorsal surface of snout fully scaled, except for narrow naked clefts along each side of median rostral ridge; nasal fossa usually lacking scales. Snout long, sharply pointed, length 67–81% PRL; dorsal profile of snout slightly concave in lateral view; terminal scute moderately short, diamond-shaped, dorsoventrally flattened; lateral nasal ridge incompletely supported by nasal bone. Anus abutting anal-fin origin. Premaxillary teeth small, conical, arranged in short, uniformly wide band, none especially enlarged; posterior end of tooth band falling far short of lateral corner of mouth. Body scales covered with stout, trihedral, shield-like spinules in parallel ridge-like rows; middle row enlarged, giving striated appearance to body surface; lateral rows often incomplete, not extending to posterior scale margin; buttresses broadly developed. Scales on head ridges coarsely modified, highly spiny, and thickened. Occipital scales covered with short, erect, trihedral spinules in narrowly divergent ridge-like rows. Free neuromasts on underside of head modified into prominent, knob-like or flap-like projections; short, dark, hairy papillae densely scattered over ventral surface of head except for mandibular rami. Head and body uniformly dark; orbit narrowly edged in black; lips, oral cavity, and all fins blackish.

Material examined. 30 specimens. **Holotype of *Caelorhynchus gilberti*:** FMNH 59401 (ex. CM 7960, 137 mm HL, 478+ mm TL), fish market in Shizuoka, Shizuoka Pref., Japan, coll. D.S. Jordan, Oct.–Nov. 1922. **Paralectotypes of *Macrurus parallelus*** (designated herein): BMNH 1887.12.7.67–68 (2, 63.8–107 mm HL, 221+–355+ mm TL), off Enoshima Island, Sagami Bay, 35.1833°N, 139.4667°E, 345 fms (631 m), *Challenger* sta. 232, trawl and dredge, 12 Mar. 1875. **Non-types: Japan:** *BSKU 44480 (1, 39.7 mm HL, 150+ mm TL), Tosa Bay, 700 m, FRV *Kotaka-maru*, 26 Jan. 1988; *BSKU 44839 (1, 50.6 mm HL, 186+ mm TL), Tosa Bay, 32.9886°N, 133.5978°E, 700–720 m, FRV *Kotaka-maru*, otter trawl, 21 Jun. 1988; BSKU 99459 (1, 60.4 mm HL, 198+ mm TL), Tosa Bay, 33.1900°N, 133.6883°E, 542–585 m, FRV *Kotaka-maru*, otter trawl, coll. H. Endo, 22 Jan. 1998; BSKU 44792 (1, 52.6 mm HL, 185+ mm TL), BSKU 44793 (1, 49.8 mm HL, 160+ mm TL), Tosa Bay, 33.0517°N, 133.6152°E, 700 m, FRV *Kotaka-maru*, 24 May 1988; BSKU 46145 (1, 117 mm HL, 416+ mm TL), Tosa Bay, 33.0400°N, 133.6098°E, 675–708 m, FRV *Kotaka-maru*, otter trawl, 22 May 1989; BSKU 29509 (1, 85.6 mm HL, 259+ mm TL), Tosa Bay, 32.9167°N, 133.5500°E, 700 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, cr. 4-Ky, sta. T20, long line, coll. O. Okamura *et al.*, 21 Dec. 1979; BSKU 30494 (1, 40.0 mm HL, 139+ mm TL), Tosa Bay, 32.9667°N, 133.5333°E, 605 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No.

2, bottom trawl, coll. O. Okamura *et al.*, 21 Dec. 1979; BSKU 29453 (1, 177 mm HL, 626+ mm TL), off Kochi, 32.4582°N, 133.4675°E, 745–810 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. 4-L4, long line, coll. O. Okamura *et al.*, 21 Dec. 1979; BSKU 45516 (1, 193 mm HL, 653+ mm TL), off Tosa-Shimizu, long line, 14 Jul. 1988; KPM-NI 18243 (1, 120 mm HL, 425+ mm TL), off Shima Peninsula, Kumano-nada, 34.0144°N, 136.9054°E, 767–771 m, R/V *Tansei-maru*, cr. KT-04-06, sta. KN-1(2), 3-m ORE beam trawl, coll. T. Sato, 1 May 2004; FRLM 3579 (1, 129 mm HL, 455+ mm TL), FRLM 3580 (1, 130 mm HL, 478 mm TL), off Mie, Kumano-nada, long line, coll. Fisheries Research Institute of Mie Pref., 18 May 1982; FRLM 8964 (1, 138 mm HL, 522+ mm TL), Kumano-nada, coll. K. Suzuki, date unknown; FRLM 3577 (1, 108 mm HL, 392+ mm TL), off Mikizaki, Kumano-nada, long line, coll. Fisheries Research Institute of Mie Pref., 18 May 1982; NSMT-P 97154 (1, 74.0 mm HL, 236+ mm TL), southeast of Miyake-jima Island, Shichito-Iojima Ridge, 34.1340°N, 139.6717°E, 565–644 m, R/V *Tansei-maru*, cr. KT-89-06, sta. ME11, 3-m ORE beam trawl, 19 May 1989; NSMT-P 101440 (1, 79.2 mm HL, 217+ mm TL), off Oshima Island, Izu Islands, 621–710 m, R/V *Hakuho-maru*, cr. KH-89-test-III, 4-m ORE beam trawl, 5 Sept. 1989; BSKU 101982 (1, 28.5 mm HL, 107+ mm TL), off Iwaki, 36.9162°N, 141.5995°E, 615–633 m, FRV *Wakataka-maru*, sta. 49, coll. H. Endo, 4 Nov. 1995; BSKU 97953 (1, 83.5 mm HL, 313+ mm TL), off Onahama, 450 m, FRV *Wakataka-maru*, sta. G-450, coll. D. Kitagawa, 16 Oct. 1997; NSMT-P 77845 (1, 162 mm HL, 620 mm TL), off Hitachi, 36.5300°N, 141.1450°E, 557–561 m, FRV *Wakataka-maru*, sta. H-550, bottom trawl, coll. H. Saito *et al.*, 14 Nov. 2006; NSMT-P 74691 (1, 149 mm HL, 535 mm TL), off Kesennuma, date unknown; NSMT-P 48953 (1, 130 mm HL, 480 mm TL), NSMT-P 48954 (1, 116 mm HL, 433+ mm TL), off Kuji, 40.0015°N, 142.3657°E, 611–617 m, FRV *Wakataka-maru*, sta. 15, bottom trawl, coll. G. Shinohara, 26 Oct. 1995; BSKU 30589 (1, 125 mm HL, 400+ mm TL), Kyushu-Palau Ridge, 27.9000°N, 134.6500°E, 910 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, cr. Ky, sta. T-12, bottom trawl, coll. O. Okamura *et al.*, 18 Dec. 1979; BSKU 26016 (1, 175 mm HL, 631+ mm TL), Kyushu-Palau Ridge, 27.9000°N, 134.6417°E, 720 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. 2-1, bottom trawl, coll. Y. Kobayashi *et al.*, 14 Feb. 1978; *BSKU 114872 (1, 107 mm HL, 416 mm TL), east-southeast of Cape Erimomisaki, 41.8170°N, 143.7875°E, 576 m, FRV *Wakataka-maru*, sta. H-550, otter trawl, coll. R. Misawa, 6 Jun. 2014; NSMT-P 76191 (1, 130 mm HL, 487+ mm TL), off Hokkaido, T/V *Oshoro-maru*, sta. OST-2, coll. T. Yanagimoto, 16 Sept. 2005.

Counts and measurements. Based on 27 specimens (28.5–193 mm HL, 107+–653+ mm TL). Counts: first dorsal-fin rays II,7–8; pectoral-fin rays i16–i19; pelvic-fin rays 6–7; gill rakers on first arch (outer/inner) 0–2/7–

10, on second arch 6–7/8–9; longitudinal scales 35–45; transverse scale rows below first dorsal-fin origin 5.5–8, below first dorsal-fin midbase 4.5–7, below second dorsal-fin origin 5–7, above anal-fin origin 17–23.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 40–45 (67–81); orbit diameter 23–28 (39–47); postorbital length 30–37 (52–62); postrostral length 56–61; orbit–preopercle distance 31–39 (55–65); suborbital width 11–15 (18–25); upper-jaw length 22–27 (39–44); length of rictus 17–23 (30–39); length of premaxillary tooth band 8–12 (14–20); preoral length 36–45 (59–80); length of terminal snout scute 7–11 (12–20); length of lateral nasal ridge 28–34 (47–60); length of suborbital ridge 92–97 (153–174); snout width 24–31 (39–55); internasal width 17–22 (28–36); interorbital width 21–23 (37–41); occipital width 8–10 (13–18); body width over pectoral-fin bases 31–47 (54–83); body depth at first dorsal-fin origin 41–58 (73–96); body depth at anal-fin origin 30–48 (53–81); prepelvic length 99–116 (167–193); preanus length 127–155 (225–255); preanal length 130–162 (230–266); isthmus–pelvic distance 22–34 (38–57); isthmus–anus distance 47–75 (83–123); isthmus–anal distance 50–82 (89–135); pelvic–anal distance 30–52 (53–87); anus–anal distance 3–7 (4–12); pelvic-fin length 27–44 (46–73); pectoral-fin length 31–40 (55–66); predorsal length 105–111 (177–196); height of first dorsal fin 30–41 (52–68); length of first dorsal-fin base 12–17 (21–28); interdorsal length 10–24 (18–41); length of gill slit 10–14 (18–24); length of posterior nostril 5–10 (9–16); barbel length 3–12 (6–20).

Size. Attains at least 65 cm TL (BSKU 45516, 653+ mm TL, off Tosa-Shimizu, Kochi, Japan).

Distribution. Japan and the Emperor Seamounts (Iwamoto & Okamoto 2015; this study; Appendix 3-2C). In Japan, known from the upper continental slope along the Pacific coasts from Hokkaido to Shikoku and the Kyushu-Palau Ridge. Depth range 400–930 m. Moderately common in the Pacific off northern Honshu (Tohoku), but rare in the west of the Boso Peninsula (139.89°E).

Remarks. For further morphological information see the original description (Jordan & Hubbs 1925) and Okamura (1970a). *Coelorinchus gilberti* is common in the Pacific off northern Japan, dominating museum specimens of the genus collected from the area.

While Shao (1993) recorded *C. gilberti* from Taiwan, it seems to represent a different species of *Coelorinchus*. Shao's (1993) specimen is the only record of *C. gilberti* from Taiwan, but unfortunately, it is presumed lost (Chiou *et al.* 2004a: table 1). The occurrence of *C. gilberti* in Taiwanese waters was therefore not verified in this study.

Nomenclatural discussion. *Coelorinchus gilberti* was described from a single specimen obtained at a fish market in Shizuoka, Shizuoka Pref., Japan (Fig. 38). Subsequently, Matsubara & Asano (1943) described *C. kanetomoi* based on a single specimen collected from off

FIGURE 37. Fresh specimens of *Coelorinchus gilberti*. (A) BSKU 29453, 177 mm HL, 626+ mm TL, off Kochi Pref., Pacific., 745–810 m depth; (B–C) BSKU 114872, 107 mm HL, 416 mm TL, east-southeast of Cape Erimomisaki, Hokkaido, Pacific, 576 m depth; (E) BSKU 46145, 117 mm HL, 416+ mm TL, Tosa Bay, Kochi Pref., Pacific, 675–708 m depth; (F) BSKU 44839, 50.6 mm HL, 186+ mm TL, Tosa Bay, 700–720 m depth; (G) BSKU 44480, 39.7 mm HL, 150+ mm TL, Tosa Bay, 500 m depth. (A–B, E–G) Lateral views; (C) dorsal and (D) ventral views of the head and trunk. [Photos: (A, E–G) BSKU; (B–D) N. Nakayama]

FIGURE 38. Holotype of *Caelorhynchus gilberti* (= *Coelorinchus gilberti*). FMNH 59401, 137 mm HL, 478+ mm TL, fish market in Shizuoka (probably from Suruga Bay), Pacific. (A) Lateral view (snout broken and bent upwards); (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

FIGURE 39. Scanning electron micrographs showing a body scale (from the dorsum below the interdorsal space) of *Coelorinchus gilberti*. NSMT-P 76191, 130 mm HL. (A) View from above; (B) oblique view. [Photos: N. Nakayama]

Misaki, Kanagawa Pref., Japan (probably from Sagami Bay). The holotype of *C. kanetomoi* was registered in the Sigenkagaku Fish Collection of the Research Institute for Natural Resources, Tokyo (SKFC 2), but apparently lost during WWII (see also the Comments on holotype of *C. hige*). Okamura (1970a) later included *C. kanetomoi* in the synonymy of *C. gilberti* without explanation.

According to Matsubara & Asano (1943:98) and Matsubara (1955:1313–1314), *C. kanetomoi* can be distinguished from *C. gilberti* by having the following combination of features: spinules of body scales arranged

in 8–11 convergent rows (vs. 5 parallel rows in *C. gilberti*); snout length twice as long as orbit diameter (vs. less than two times as long as orbit); interorbital width only slightly greater than interdorsal space (1.33 times vs. about 1.5 times as long as first dorsal-fin base length); pelvic-fin length about equal to pectoral-fin length (vs. shorter than pectoral-fin length); and two narrow scaly patches below infraorbital ridge (vs. underside of head entirely naked). However, the holotype of *C. kanetomoi* (586 mm TL, fide Matsubara & Asano 1943:95) is much larger than that of *C. gilberti* (478+ mm TL), and the differences in squamation

appear to reflect ontogenetic or individual variation. To a certain extent, the number of spinule rows increases with size in grenadiers (Iwamoto & Merrett 1997:509) and also differs on different parts of the body. In addition, the scaly patches on the underside of the head are usually absent in smaller specimens of *C. gilberti*. The differences in the measurements are also less pronounced or even negligible when considering the range of variation of the *C. gilberti* examined: snout length 1.53–1.91 times as long as orbit diameter; interorbital width 1.33–1.84 times as long as first dorsal-fin base length; and pectoral-fin length 0.87–1.30 in pelvic-fin length. Consequently, no significant differences were found between the two nominal species, and *C. kanetomoi* is confirmed as a junior synonym of *C. gilberti*.

Relationships. *Coelorinchus gilberti* belongs to a group of *Coelorinchus* species characterized by the following combination of features: no external evidence of light organ anterior to anus; lateral nasal ridge incompletely supported by nasal bone; premaxillary teeth in short, uniformly wide band; underside of head entirely or almost entirely naked; body scales covered with shield-like spinules in parallel to subparallel rows, with middle row largest and some rows incomplete; buttresses of spinules broadly developed; scales on head ridges coarsely modified, highly spiny, and thickened; body uniformly dark brown to swarthy; lips and oral cavity blackish. This group is called here the *C. occa* group, in which the following eight valid species are included: *C. anisacanthus* Sazonov, 1994 from the Emperor Seamounts; *C. aratrum* Gilbert, 1905 and *C. doryssus* Gilbert, 1905 from Hawaii; *C. gilberti*; *C. labiatus* (Köhler, 1896) from the northeastern Atlantic; *C. mediterraneus* Iwamoto & Ungaro, 2002 from the Mediterranean; *C. obscuratus* McMillan & Iwamoto, 2009 from New Zealand to New Caledonia; and *C. occa* (Goode & Bean, 1885) from the western Atlantic. This group is closely related to the *C. parallelus* group (see the Relationships of *C. divergens*), with which it is considered to comprise the subgenus *Oxygadus* Gilbert & Hubbs, 1920 (type species *C. parallelus*). The *C. parallelus* group readily differs from the *C. occa* group in having the underside of the head completely or almost completely scaled (vs. completely or mostly naked).

Comparisons. *Coelorinchus gilberti* is readily distinguished from other species of the *C. occa* group except *C. occa* by the presence of short, dark, hairy papillae densely scattered over the ventral surface of the head except the mandibular rami (vs. absent). It further differs from all but *C. occa* in having prominent flap-like (knob-like in well-preserved specimens) black neuromasts serially arranged on the underside of the head. In the other species, the neuromasts are relatively small, unpigmented, and scarcely discernible without a microscope. *Coelorinchus gilberti* is readily distinguished from *C. mediterraneus* in that naked areas along each

side of the median rostral ridge are only narrowly developed (vs. broadly developed). It also differs from *C. anisacanthus*, *C. doryssus*, *C. labiatus*, and *C. mediterraneus* by head squamation: in *C. gilberti*, scales on dorsal surfaces of the head are strongly spinulated, which gives a spiny appearance to these areas, whereas in the latter four species, the spinulation is less pronounced, giving more or less smooth texture to the surfaces. *Coelorinchus gilberti* differs from *C. occa* in that scaly patches are only narrowly developed (or even absent) on the mandibular rami and areas below the infraorbital and preopercular ridges (vs. broadly developed in *C. occa*).

***Coelorinchus hexafasciatus* Okamura, 1982**

[Japanese name: Musuji-sokodara]

(Figs. 40–41, 42A–B; Appendix 3-2D)

Coelorhynchus hexafasciatus Okamura, 1982:173, fig. 104 (original description; holotype: BSKU 31443, from Kyushu-Palau Ridge, 26°45.0'N, 135°19.0'E, in 336 m; 31 paratypes from Kyushu-Palau Ridge, in 685–910 m; new Japanese name: “Musuji-sokodara”).

Coelorinchus hexafasciatus: Okamura 1984b:97, pl. 83, fig. G (compiled); Okamura 1988:97, pl. 83, fig. G (compiled); Okamura 1997:127, fig. 16 (compiled); Nakabo & Kai 2013:512 (in key; Japan); Iwamoto *et al.* 2015:56, fig. 6 (brief description; 1 spec. from Da-xi; first record from Taiwan); Motomura 2020:38 (listed; Japan).

Coelorinchus hexfasciatus [sic]: Iwamoto 1990:159, fig. 374 (synopsis).

Coelorinchus hexafasciatus: Nakabo 1993:371 (in key; Japan); Nakabo 2000:435 (in key; Japan); Nakabo 2002:435 (in key; Japan).

Diagnosis. Light organ externally represented by short, narrow, naked streak immediately anterior to anus, its anterior margin falling far short of line connecting inner pelvic-fin bases. Underside of head completely naked except for oval scaly patches on ventral surfaces of preopercles; naked clefts along each side of mid rostral ridge scarcely developed. Snout long, deep, length 57–70% PRL, its dorsal profile almost straight in lateral view; tip of snout not sharply pointed, with 3 bluntly pointed scutes; lateral nasal ridge incompletely supported by nasal bone. Anus slightly separated from anal-fin origin. Premaxillary teeth in long, tapered band, with outer series prominently enlarged; posterior margin of tooth band almost reaching lateral corner of mouth. Body scales covered with short, reclined, keel-like spinules in widely divergent, fine saw-toothed ridges; every spinule row complete, extending to posterior scale margin; spinules in each row overlapping, but generally free from one another, gradually increasing in height posteriorly; buttresses poorly developed. Occipital scales covered with narrowly divergent rows of spinules. Preoral length

41–58% PRL; barbel length 10–18% PRL. About 6–7 dark saddles on body; underside of head dusky; lips pale; oral cavity blackish; gular region dusky, branchiostegal membranes blackish posteriorly.

Material examined. 46 specimens. **Holotype of *Coelorhynchus hexafasciatus*:** BSKU 30443 (150 mm HL, 546+ mm TL), Kyushu-Palau Ridge, Japan, 26.7500°N, 135.3167°E, 336 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. T8, bottom trawl, coll. Y. Kobayashi *et al.*, 17 Dec. 1979. **Paratypes of *C. hexafasciatus*:** BSKU 30440 (1, 62.6 mm HL, 213+ mm TL), BSKU 30441 (1, 73.9 mm HL, 274 mm TL), BSKU 30442 (1, 73.9 mm HL, 275 mm TL), collected with holotype; BSKU 29533 (1, 131 mm HL, 462+ mm TL), Kyushu-Palau Ridge, 26.7500°N, 135.3167°E, 336 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, tr. 9, bottom trawl, coll. O. Okamura *et al.*, 17 Dec. 1979; BSKU 29427 (1, 97.6 mm HL, 334+ mm TL), BSKU 29452 (1, 167 mm HL, 541+ mm TL), Kyushu-Palau Ridge, 27.9178°N, 134.7460°E, 685–710 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. 4-T8, bottom trawl, coll. O. Okamura *et al.*, 19 Dec. 1978; BSKU 32149 (1, 131 mm HL, 460+ mm TL), BSKU 32150 (1, 91.9 mm HL, 298+ mm TL), BSKU 32151 (1, 73.6 mm HL, 253+ mm TL), Kyushu-Palau Ridge, 27.9178°N, 134.7460°E, 685–710 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, tr. T8, bottom trawl, coll. O. Okamura *et al.*, 19 Dec. 1979; BSKU 29486 (1, 76.9 mm HL, 247+ mm TL), BSKU 29506 (1, 115 mm HL, 418 mm TL), Kyushu-Palau Ridge, 27.9000°N, 134.6667°E, 697–700 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, tr. 12, bottom trawl, coll. O. Okamura *et al.*, 18 Dec. 1979; BSKU 30472 (1, 70.1 mm HL, 218+ mm TL), BSKU 30473 (1, 75.3 mm HL, 233+ mm TL), Kyushu-Palau Ridge, 27.9000°N, 134.6667°E, 697–700 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. T11, bottom trawl, coll. Y. Kobayashi *et al.*, 18 Dec. 1979; BSKU 26106 (1, 90.3 mm HL, 291+ mm TL), BSKU 26107 (1, 89.1 mm HL, 337 mm TL), BSKU 26108 (1, 86.0 mm HL, 275+ mm TL), BSKU 26109 (1, 85.7 mm HL, 313 mm TL), BSKU 26110 (1, 88.3 mm HL, 298+ mm TL), BSKU 26111 (1, 74.7 mm HL, 252+ mm TL), Kyushu-Palau Ridge, 27.9033°N, 134.6583°E, 700 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. 2-4, bottom trawl, coll. Y. Kobayashi *et al.*, 20 Feb. 1978; BSKU 26025 (1, 76.5 mm HL, 245+ mm TL), BSKU 26026 (1, 74.8 mm HL, 255+ mm TL), Kyushu-Palau Ridge, 27.9000°N, 134.6550°E, 650 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. 2-2, tr. 9, bottom trawl, coll. Y. Kobayashi *et al.*, 15 Feb. 1978; BSKU 26070 (1, 118 mm HL, 416+ mm TL), BSKU 26071 (1, 69.3 mm HL, 224+ mm TL), BSKU 26072 (1, 96.4 mm HL, 335+ mm TL), BSKU 26073 (1, 75.1 mm HL, 257+ mm TL), BSKU 26074 (1, 70.6 mm HL, 256 mm TL), BSKU 26075 (1, 74.0 mm HL, 258+ mm TL), BSKU 26076 (1, 82.1 mm HL, 300 mm TL), Kyushu-Palau Ridge, 27.9167°N, 134.6500°E, 700 m, F/

Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. 2-3, bottom trawl, coll. Y. Kobayashi *et al.*, 20 Feb. 1978; BSKU 30590 (1, 50.8 mm HL, 175+ mm TL), BSKU 30591 (1, 79.6 mm HL, 263+ mm TL), BSKU 30592 (1, 119 mm HL, 440 mm TL), Kyushu-Palau Ridge, 27.9000°N, 134.6500°E, 910 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, cr. Ky, sta. T-12, bottom trawl, coll. O. Okamura *et al.*, 18 Dec. 1979. **Non-types: Japan:** HUMZ 95116 (1, 156 mm HL), west of Amami-oshima, Okinawa Trough, 28.4033°N, 127.7567°E, 735–770 m, bottom trawl, coll. K. Amaoka, 9 Nov. 1981; BSKU 42885 (1, 122 mm HL, 416+ mm TL), Kyushu-Palau Ridge, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, date unknown; BSKU 40980 (1, 91.1 mm HL, 320+ mm TL), BSKU 40981 (1, 53.5 mm HL, 181+ mm TL), Kyushu-Palau Ridge, 27.9100°N, 134.6667°E, 697–1320 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, bottom trawl, coll. O. Okamura *et al.*, 18 Dec. 1979; HUMZ 71890 (1, 174 mm HL, 632+ mm TL), HUMZ 71891 (1, 203 mm HL, 706+ mm TL), HUMZ 71892 (1, 185 mm HL, 659+ mm TL), HUMZ 71893 (1, 127 mm HL), HUMZ 71897 (1, 108 mm HL), Shichito-Iojima Ridge, 26.1667°N, 140.9167°E, 500–1200 m, coll. Japan Marine Fishery Resources Research Center (JAMARC), 30 Oct. 1977; NSMT-P 102809 (1, 137 mm HL, 533+ mm TL), NSMT-P 102810 (1, 86.9 mm HL, 325+ mm TL), off Hachijo-jima Island, Shichito-Iojima Ridge, F/V *Ryosei-maru*, 23 Apr. 2008; NSMT-P 64134 (1, 182 mm HL, 635+ mm TL), off Hachijo-jima Island, Shichito-Iojima Ridge, F/V *Taiyo-maru*, 6 Jun. 2001; FAKU 133888 (1, 93.5 mm HL, 354 mm TL), off Nii-jima Island, 20 Dec. 2010. **Taiwan:** ASIZP 71202 (1, 133 mm HL, 494+ mm TL), Da-xi fish market, Yilan, 24.9400°N, 121.9000°E, coll. H.-C. Ho, 27 Mar. 2004.

Counts and measurements. Based on 46 specimens (50.8–203 mm HL, 175+–706+ mm TL). Counts: first dorsal-fin rays II,8–10; pectoral-fin rays i16–i20; pelvic-fin rays 7; gill rakers on first arch (outer/inner) 0–3/7–11, on second arch 6–8/8–11; longitudinal scales 35–49; transverse scale rows below first dorsal-fin origin 6.5–9.5, below first dorsal-fin midbase 5–8, below second dorsal-fin origin 6–10, above anal-fin origin 26–30; pyloric caeca 54.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 37–42 (57–70); orbit diameter 22–28 (35–45); postorbital length 34–41 (57–66); postrostral length 59–64; orbit–preopercle distance 37–47 (62–75); suborbital width 12–15 (20–24); upper-jaw length 27–38 (45–61); length of rictus 21–31 (35–49); length of premaxillary tooth band 17–23 (27–37); preoral length 26–35 (41–58); length of lateral nasal ridge 20–33 (32–56); length of suborbital ridge 94–99 (147–163); snout width 23–28 (37–46); internasal width 17–21 (27–35); interorbital width 22–26 (35–43); occipital width 10 (15–16); body width over pectoral-fin bases 32–56 (53–88); body depth at first dorsal-fin origin 52–71 (85–112); body depth at anal-fin origin 40–60 (66–

FIGURE 40. Fresh specimens of *Coelorinchus hexafasciatus*. (A) FAKU 133888, 93.5 mm HL, 354 mm TL, Tokyo Bay, Pacific; (B) HUMZ 71893, 127 mm HL, 416+ mm TL, Shichito-Iojima Ridge, Pacific, 500–1200 m depth; (C) BSKU 29533, paratype, 131 mm HL, 462+ mm TL, Kyushu-Palau Ridge, Pacific, 336 m depth; (D) BSKU unknown, Kyushu-Palau Ridge. Lateral views. [Photos: (A) Y. Kai; (B) HUMZ; (C–D) BSKU]

FIGURE 41. Holotype of *Coelorhynchus hexafasciatus* (= *Coelorinchus hexafasciatus*). BSKU 30443, 150 mm HL, 546+ mm TL, Kyushu-Palau Ridge, Pacific, 336 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

95); prepelvic length 100–117 (162–188); preanus length 114–148 (186–233); preanal length 118–156 (193–248); isthmus–pelvic distance 27–38 (46–60); isthmus–anal distance 62–74 (98–116); pelvic–anal distance 33–69 (54–110); anus–anal distance 5–10 (8–17); pelvic-fin length 30–50 (48–82); pectoral-fin length 40–50 (66–80); predorsal length 103–112 (165–182); height of first dorsal fin 42–53 (69–88); length of first dorsal-fin base 15–25 (25–40); interdorsal length 13–24 (21–39); length of gill slit 15–20 (24–33); length of posterior nostril 7–17 (12–28); barbel length 6–11 (10–18).

Size. Attains at least 71 cm TL (HUMZ 71891, 706+ mm TL, Shichito-Iojima Ridge, Japan).

Distribution. Known only from the Kyushu-Palau Ridge, Shichito-Iojima Ridge, and Okinawa Trough, at depths of 366–1320 m (Appendix 3-2D). Common at the Kyushu-Palau Ridge, but rare in other parts of its range.

Remarks. *Coelorinchus hexafasciatus* was described from 32 specimens collected from the Kyushu-Palau Ridge (Figs. 40C–D, 41). The reader can refer to the original description (Okamura 1982) for further morphological details. Recently, Iwamoto *et al.* (2015) reported an additional specimen from the Da-xi fish market in northern Taiwan. This species was also obtained from the middle Okinawa Trough and the Shichito-Iojima Ridge (see Nakayama *et al.* 2015c:258).

Relationships. *Coelorinchus hexafasciatus* belongs to the *C. tokiensis* group as redefined by Nakayama *et al.* (2015c). In addition to *C. hexafasciatus*, this group includes the following five species: *C. celaenostoma* McMillan & Paulin, 1993 and *C. osipullus* McMillan & Iwamoto, 2009 from New Caledonia to New Zealand;

C. quadricristatus (Alcock in Wood-Mason & Alcock, 1891) from the Andaman Sea; *C. sheni* Chiou, Shao & Iwamoto, 2004 from Japan and Taiwan; and *C. tokiensis* (Steindachner & Döderlein, 1887) from Japan and the South China Sea. The above six species are separable from other congeners by having the following combination of features (modified from Nakayama *et al.* 2015c): light organ externally represented by short naked streak anterior to anus, its anterior margin falling far short of line connecting inner pelvic-fin bases; dorsal profile of snout almost straight in lateral view; lateral nasal ridge incompletely supported by nasal bone; premaxillary teeth in long tapered band, with outer series enlarged; posterior end of premaxillary tooth band almost reaching lateral corner of mouth; body scales covered with keel-like spinules in widely divergent, saw-toothed ridges; series of dark saddles on body; and oral cavity blackish.

Comparisons. *Coelorinchus hexafasciatus* differs most notably from other species of the *C. tokiensis* group except *C. tokiensis* in having the underside of the head completely naked except oval scaly patches on the ventral surfaces of the preopercles. The species is further distinguished from *C. celaenostoma* by having a shorter snout (57–70% PRL vs. 86–94%; preoral length 41–58% PRL vs. 71–80%), about 6–7 dark saddles on the body (vs. ca. 5), pale lips (vs. black), and lacking a bluish band completely encircling the trunk (vs. present); from *C. osipullus* by having a shorter barbel (10–18% PRL vs. 19–26%) and fewer saddles on the body (ca. 6–7 vs. ca. 9–11); and from *C. quadricristatus* and *C. sheni* by having a bluntly pointed tip of the snout (vs. sharply pointed). *Coelorinchus hexafasciatus* is most similar to the sympatric *C. tokiensis*, but differs in having a shorter

FIGURE 42. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of three species of the *Coelorinchus tokiensis* group. (A–B) *C. hexafasciatus*, BSKU 29452, paratype, 167 mm HL; (C–D) *C. sheni*, CAS 215541, 112 mm HL; (E–F) *C. tokiensis*, BSKU 26104, paratype of *C. longicephalus*, 202 mm HL. (A, C, E) Views from above; (B, D, F) oblique views. [Photos: (A–B, E–F) N. Nakayama; (C–D) NSMT]

snout (57–70% PRL vs. 70–85%) with negative allometry (vs. positive; Nakayama *et al.* 2015c).

***Coelorinchus hige* Matsubara, 1943**

[Japanese name: Kata-hige]

(Figs. 21B, 43–45; Table 2; Appendix 3-2E)

Coelorhynchus hige Matsubara, 1943:144, fig. 6 (original description; holotype: SKFC 6, from Suruga Bay off Heta; new Japanese name: “Kata-hige”); Kuroda 1951:391 (listed; Suruga Bay); Matsubara 1955:1313 (in key; Japan); Okamura 1970a:193 [description after Matsubara (1943)]; Tominaga & Uyeno 1981:488 (listed; Japan); Ozawa 1983:13 (listed; off Makurazaki, Kagoshima Pref., East China Sea).

Coelorhynchus asteroides Okamura, 1963a:21, fig. 1 [original description; holotype: FAKU 23801, from off Owase (misspelled as Owashi); 7 paratypes from Heta, Owase and Mimase; new Japanese name: “Hoshihige”]; Kamohara 1964:95 (listed; Kochi Pref.); Okamura 1970a:189, pl. XLI, text-figs. 81–82 [description; 8 spec. from Suruga Bay, Owase (misspelled as Owashi), and Mimase]; Kuroda 1971:56 [listed from Suruga Bay after Okamura (1970a)]; Tominaga & Uyeno 1981:488 (listed; Japan).

Coelorhynchus (Oxymacrus) asteroides: Okamura 1970b: table 1 (listed; Japan).

Coelorhynchus (Oxymacrus) hige: Okamura 1970b: table 1 (listed; Japan).

Coelorinchus asteroides: Yatou 1984:235, 369, fig. 167 (brief description; 10 spec. from Okinawa Trough; photo based on BSKU 29833); Okamura 1984b:97, pl. 83, fig. F (compiled); Okamura 1988:97, pl. 83, fig. F (compiled); Okamura 1997:127, fig. 13 (compiled); Shao *et al.* 2008b: table 2 (3 spec. listed from northeastern and southwestern Taiwan); Nakabo & Kai 2013:511 (in key; Japan); Iwamoto *et al.* 2015:50 (brief description; 8 spec. from northeastern and southwestern Taiwan, and Philippines); Motomura 2020:38 (listed; Japan).

Coelorinchus asteroides: Nakabo 1993:370 (in key; Japan); Shinohara & Matsuura 1997:290 (listed; Suruga Bay); Nakabo 2000:434 (in key; Japan); Shinohara *et al.* 2001:304 (listed; Tosa Bay); Nakabo 2002:434 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Chiou *et al.* 2004a:43, fig. 9 (brief description; 2 spec. from Da-xi and Dong-gang; first record from Taiwan); Shinohara *et al.* 2005:416 (listed; Ryukyu Islands); Furuhashi *et al.* 2010: table 2 (39 spec. listed from northern Okinawa Trough).

Coelorinchus (Oxymacrus) asteroides: Chiou *et al.* 2004a: table 1 (listed; Taiwan).

Diagnosis. Light organ externally represented by short, narrow, naked streak immediately anterior to anus, its anterior margin falling far short of line connecting inner pelvic-fin bases. Underside of head almost completely naked; dorsal surface of snout fully scaled, except narrow naked clefts along each side of median rostral ridge. Nasal

fossa usually naked. Snout long, sharply pointed, length 57–83% PRL, becoming proportionally shorter with growth; dorsal profile of snout almost straight in lateral view. Terminal snout scute short, broad, arrowhead-shaped, dorsoventrally flattened, length 11–18% PRL. Scales on median rostral ridge covered with radiating rows of spinules. Lateral nasal ridge completely supported by nasal bone, not strongly convex when viewed dorsally. Anus slightly separated from anal-fin origin. Premaxillary teeth small, conical, arranged in short, uniformly wide band, none especially enlarged; posterior margin of tooth band falling far short of lateral corner of mouth. Body scales covered with short, reclined, keel-like spinules in narrowly divergent, saw-toothed ridges; every spinule row complete, extending to posterior scale margin; spinules in each row greatly overlapping, closely adjoined to one another, but generally free at tips; buttresses scarcely developed. Occipital scales covered with short, erect, recurved, needle-like spinules in widely divergent comb-like rows. Height of first dorsal fin 71–106% PRL; interdorsal length 51–70% PRL; pyloric caeca 47–49. Body darker dorsally, paler ventrally; no obvious body markings in adults, but juveniles with small dark spots on posterior part of tail; underside of head dusky; lips white; oral cavity blackish; gular and branchiostegal membranes pale; pelvic fins dusky.

Material examined. 30 specimens. **Holotype of *Coelorhynchus asteroides*:** FAKU 23801 (85.5 mm HL, 376+ mm TL), Owase fish market, Mie Pref., Kumano-nada, Japan, bottom trawl, 24 Oct. 1954. **Paratypes of *C. asteroides*:** FAKU 29115 (1, 76.2 mm HL, 266+ mm TL), Mimase fish market, bottom trawl, date unknown; FAKU 21513 (1, 60.2 mm HL, 269+ mm TL), FAKU 24909 (1, 67.5 mm HL), Owase fish market, Kumano-nada, bottom trawl, date unknown. **Non-types: Japan:** BSKU 106771 (1, 66.1 mm HL, 268 mm TL), BSKU 106822 (1, 61.2 mm HL, 255+ mm TL), southeast of Shimokoshiki-jima Island, East China Sea, 31.5658°N, 129.8915°E, 380 m, F/V *Maruko-maru*, tr. 1, bottom trawl, coll. N. Nakayama *et al.*, 24 Apr. 2012; BSKU 50881 (1, 30.0 mm HL, 108+ mm TL), off Nomaie, Kagoshima, East China Sea, F/V *Maruko-maru*, bottom trawl, coll. M. Yamashita and Y. Ohashi, 10 Sept. 2010; BSKU 109039 (1, 84.2 mm HL, 351+ mm TL), BSKU 109040 (1, 88.2 mm HL, 401+ mm TL), east of Kasayama Bank, 32.3884°N, 129.0505°E, 304–312 m, T/V *Nagasaki-maru*, cr. N365, sta. A3, 3-m ORE beam trawl, coll. N. Nakayama, 19 Nov. 2012; SNFR 14555 (1, 110 mm HL, 461+ mm TL), East China Sea, 31.4372°N, 128.3418°E, 233–303 m, F/V *Choun-maru*, No. 6, sta. C6-j, tr. 119, bottom trawl, coll. T. Jintoku, 29 Sept. 2009; BSKU 29833 (1, 74.0 mm HL, 299+ mm TL), BSKU 33819 (1, 76.1 mm HL, 330 mm TL), BSKU 33820 (1, 70.7 mm HL, 297 mm TL), BSKU 33821 (1, 72.4 mm HL, 300+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 30.4815°N, 127.8432°E, 320–360 m, F/V *Yuryo-maru*, No. 8, sta. 3-T8, bottom trawl,

coll. H. Maeda, 28 Oct. 1979; BSKU 28286 (1, 75.5 mm HL, 289+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.7500°N, 127.5000°E, 325–360 m, F/V *Ryoan-maru*, No. 28, tr. 70, bottom trawl, coll. Y. Kinoshita and S. Hagino, 19 Mar. 1978; *BSKU 27032 (1, 70.2 mm HL, 223+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.9267°N, 127.7483°E, 360 m, F/V *Yuryo-maru*, No. 8, tr. 20, bottom trawl, coll. Y. Kinoshita and S. Hagino, 4 Feb. 1978; BSKU 29773 (1, 69.7 mm HL, 297+ mm TL), northwest of Okinawa-jima Island, Okinawa Trough, 28.7000°N, 125.9333°E, 320 m, F/V *Yuryo-maru*, No. 8, sta. 2-T7, bottom trawl, coll. M. Hirose and T. Kume, 12 Oct. 1979; BSKU 29032 (1, 71.2 mm HL, 222+ mm TL), Mimase fish market, 4 Nov. 1978; BSKU 879 (1, 47.1 mm HL, 198+ mm TL), Mimase fish market, bottom trawl, coll. T. Kamohara, 16 Jan. 1951; BSKU 9684 (1, 67.7 mm HL, 297 mm TL), BSKU 9685 (1, 67.1 mm HL, 274+ mm TL), Mimase fish market, bottom trawl, coll. T. Kamohara, 17 Mar. 1949; BSKU 275 (1, 63.1 mm HL, 264 mm TL), BSKU 276 (1, 58.1 mm HL, 256 mm TL), BSKU 277 (1, 64.4 mm HL, 263+ mm TL), BSKU 278 (1, 65.4 mm HL, 291 mm TL), BSKU 279 (1, 58.2 mm HL, 254+ mm TL), BSKU 280 (1, 57.2 mm HL, 241+ mm TL), Mimase fish market, bottom trawl, coll. T. Kamohara, 20 Feb. 1950; BSKU 9683 (1, 63.4 mm HL, 276+ mm TL), Mimase fish market, bottom trawl, coll. T. Kamohara, 30 Jan. 1951; BSKU 112434 (1, 83.6 mm HL, 336+ mm TL), BSKU 112451 (1, 72.0 mm HL, 310+ mm TL), off Susaki, Tosa Bay, 320–380 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 9 Mar. 2014.

Redescription based on Japanese specimens.

General features are shown in Figs. 21B, 43–44. Counts and measurements are given in Table 2. Body deepest at first dorsal-fin origin, progressively tapering to long, laterally compressed tail. Trunk short, moderately compressed, width over pectoral-fin bases 1.1–1.3 in depth below first dorsal-fin origin. Head large, HL about 3.6–4.5 in TL. Snout long, protruding greatly beyond upper jaw, length 1.3–1.7 times as long as orbit diameter; dorsal contour of snout almost straight in lateral view; lateral nasal ridge completely supported by nasal bone; anterolateral margins of snout moderately convex when viewed dorsally. Orbit large, its greatest diameter 0.9–1.3 in postorbital length. Interorbital space flat, width 1.1–1.5 in orbit diameter. Mouth moderate in size, inferior, protrusible, upper-jaw length 1.0–1.3 in orbit diameter; posterior margin of maxilla extending to below midorbit or beyond, but not reaching hind margin of orbit; lateral corner of mouth moderately restricted by skin folds; lips thin, slightly papillose near tooth bands. Suborbital region divided by longitudinal bony ridge passing from tip of snout to posteroventral angle of opercle; its upper half almost vertical, lower half sharply inclined mesially. Preopercle large, posterior margin inclined, forming moderately angular lobe at posteroventral corner.

Subopercle prolonged ventrally as slender flap; posterior margin deeply concave. Gill membranes broadly connected across, and attached mesially to isthmus, with slight posterior free fold. Outer gill slit moderately restricted by skin folds, length 2.0–3.4 in orbit diameter (1.8 in largest specimen; SNFR 14555, 101 mm HL). Gill rakers small, low, tubercular, armed with short, fine spines; no rakers on outer side of first arch (rudimentary raker occasionally found in larger specimens) and inner side of fourth arch; gill filaments moderately long. Chin barbel slender, tapering into fine tip; length 2.1–4.7 in orbit diameter.

Anus slightly separated from anal-fin origin by 1–2 scale rows. Light organ externally represented by short, narrow, naked streak, extending from anus forward to about midway between anus and outer pelvic-fin bases, its anterior margin falling far short of line connecting inner pelvic-fin bases. Pyloric caeca 47–47 (based on 3 paratypes of *C. asteroides* and BSKU 106771), rete-gas gland complexes 2 (based on BSKU 106771).

Teeth small, sharp, conical, gently incurved, in narrow bands in both jaws. Premaxillary teeth in short, uniformly wide band, with about 6 tooth rows near symphysis; posterior margin of tooth band falling far short of lateral corner of mouth; teeth becoming progressively smaller inwardly. Mandibular teeth uniformly small, in broad tapered band, with about 5–6 tooth rows across widest point near symphysis; band abruptly narrowing posteriorly, its posterior margin extending beyond lateral corner of mouth. All teeth deeply embedded in thick layer of gum papillae.

Scales on body large, thin, not deciduous, covered with short, reclined, keel-like spinules in narrowly divergent rows (Fig. 45A–D); in 88.2 mm HL specimen (BSKU 109040), those on dorsum below interdorsal space with 8–9 rows of spinules; middle row longest with about 8–9 spinules, but not especially high or enlarged compared with adjacent rows; every spinule row complete throughout; spinules forming angle of about 40° to scale surface; spinules greatly overlapping, closely adjoined to one another, but their distal tips usually free, forming saw-toothed ridges; spinules increasing in height posteriorly, but last spinule in each row barely extending beyond posterior scale margin; buttresses of spinules scarcely developed; no reticulate structures on unexposed portion. Body fully scaled except for fins and light organ.

Scales on head ridges stout, thickened; those on median rostral ridge covered with radiating rows of spinules. Terminal snout scute short, broad, arrowhead-shaped, dorsoventrally flattened, its length 2.8–4.4 in orbit diameter. Supraoccipital scute stout, enlarged, armed with single row of spinules; posttemporal scutes similar to that on supraoccipital, but spinules arranged in multiple rows. Occipital scales covered with short, erect, needle-like spinules in widely divergent rows (Fig. 45E–F); spinules recurved posteriorly, but scarcely imbricated, independent

FIGURE 43. Fresh specimens of *Coelorinchus hige*. (A) BSKU 109039, 84.2 mm HL, 351+ mm TL, Okinawa Trough, East China Sea, 304–312 m depth; (B) BSKU 109040, 88.2 mm HL, 401+ mm TL, collected with BSKU 109039; (C–E) BSKU 112451, 72.0 mm HL, 310+ mm TL, Mimase fish market, Kochi Pref., Pacific. (A–C) Lateral views (D) dorsal and (E) ventral views of the head and trunk. [Photos: N. Nakayama]

from one another, forming comb-like ridges that increase in height posteriorly; scales on dorsal surfaces of snout posterior to leading edges armed with cluster of short, erect, needle-like spinules. Other scales on head variable in size, generally similar to those on body, but spinules more erect and spinule rows more divergent; those on opercle, preopercle, and postorbital and supratemporal canals largest. Nasal fossa usually naked (a few tiny scales

occasionally present anteroventrally). Dorsal surfaces of snout almost completely scaled; pair of small naked areas posterior to lateral nasal ridges and along each side of median rostral ridge. Underside of head almost completely naked; a few small scales often present on lower portion of preopercle and above posterior margins of lower jaws.

No open pores along cephalic sensory canals. Free neuromasts on ventral surface of head poorly marked,

TABLE 2. Morphometric measurements and meristic counts* for *Coelorinchus hige*.

Nominal species	<i>C. hige</i>	<i>C. asteroides</i>		
Type status	HT**	HT	PT	NT
Number of specimens	<i>n</i> = 1	<i>n</i> = 1	<i>n</i> = 3	<i>n</i> = 25
Measurements (mm)				
TL	256	376	266–269	108–461
HL	62.9	85.5	60.2–76.2	30.0–110
% of HL (% of PRL)				
Snout length	39	39 (63)	39–42 (61–70)	37–46 (57–83)
Orbit diameter	28	28 (45)	28–30 (46–49)	26–31 (43–52)
Postorbital length	33	35 (56)	32–33 (53–55)	29–37 (48–58)
Postrostral length	–	62 (–)	60–63 (–)	55–65 (–)
Orbit–preopercle distance	–	40 (64)	35–38 (59–64)	32–41 (55–64)
Suborbital width	14	15 (23)	12–14 (21–22)	11–16 (20–25)
Upper-jaw length	25	26 (42)	23–26 (38–44)	22–29 (37–45)
Length of rictus	–	21 (34)	16–19 (27–33)	16–21 (27–35)
Length of PMX tooth band	–	12 (19)	8–11 (13–19)	8–12 (14–20)
Preoral length	–	27 (44)	32–34 (51–57)	25–39 (39–72)
Length of terminal snout scute	–	– (–)	– (–)	7–10 (11–18)
Length of lateral nasal ridge	–	24 (39)	25–29 (40–48)	22–32 (34–58)
Length of suborbital ridge	–	95 (153)	93–98 (148–165)	91–97 (141–175)
Snout width	–	29 (47)	28–30 (45–50)	27–31 (42–53)
Internasal width	–	21 (33)	18–20 (31–34)	18–20 (30–35)
Interorbital width	25	24 (39)	22–25 (37–41)	21–25 (35–41)
Occipital width	–	– (–)	– (–)	8–11 (13–18)
Body width over P. bases	–	– (–)	46 (77)	35–53 (59–84)
Body depth at 1D. origin	–	63 (101)	56–59 (93–94)	43–61 (72–98)
Body depth at A. origin	–	53 (84)	47–50 (79–80)	35–50 (58–87)
Pre-V. length	–	118 (189)	111–114 (176–192)	103–125 (178–200)
Pre-anus length	–	– (–)	– (–)	132–150 (221–258)
Pre-A. length	–	160 (257)	154–155 (247–258)	138–164 (232–274)
Isthmus–V. distance	30	39 (63)	35–39 (58–62)	28–43 (47–67)
Isthmus–anus distance	–	– (–)	– (–)	53–71 (89–117)
Isthmus–A. distance	–	– (–)	– (–)	59–80 (100–134)
V.–A. distance	–	48 (77)	43–46 (72–74)	33–49 (56–86)
Anus–A. distance	5	– (–)	– (–)	5–10 (9–18)
V. length	41	30 (48)	31–38 (51–60)	33–48 (58–79)
P. length	47	36 (58)	40–43 (66–69)	37–57 (64–94)
Pre-1D. length	–	117 (188)	112–116 (183–194)	108–117 (173–209)
Height of 1D.	51	52 (83)	54–59 (90–94)	46–64 (71–106)
Length of 1D. base	19	22 (36)	19 (31–32)	15–23 (26–39)
Interdorsal length	38	43 (70)	33–42 (56–67)	29–41 (51–69)
Length of gill slit	–	12 (20)	11–12 (19)	11–16 (18–24)
Length of posterior nostril	–	7 (12)	8–9 (13–14)	7–11 (12–17)
Barbel length	11	9 (15)	– (–)	6–13 (11–22)
Counts				
1D. rays	II,8	II,9	II,8–10	II,8
P. rays	i17	i17	i15–i19	i16
V. rays	7	7	7	7
Outer GR on first arch	–	0	0–3	
Inner GR on first arch	7	8	6–8	6
Outer GR on second arch	–	6	5–7	6
Inner GR on second arch	–	7	6–8	6
Longitudinal scales	–	28	26–37	32
Scales below 1D. origin	–	5.5	4.5–6.5	5

...Continued on the next page

TABLE 2. (Continued)

Nominal species	<i>C. hige</i>	<i>C. asteroides</i>		
Type status	HT**	HT	PT	NT
Number of specimens	<i>n</i> = 1	<i>n</i> = 1	<i>n</i> = 3	<i>n</i> = 25
Scales below 1D. midbase	–	3.5	3.5–5.5	4.5
Scales below 2D. origin	5	4.5	4–6.5	4.5
Scales above A. origin	–	15	17	13.5–19

*Abbreviations: 1D.—first dorsal fin; 2D.—second dorsal fin; A.—anal fin; GR—gill rakers; HT—holotype; NT—non-type; P.—pectoral fin; PMX—premaxillary; PT—paratype; V.—pelvic fin

**Based on Matsubara (1943)

FIGURE 44. Preserved specimen of *Coelorinchus hige*. FAKU 23801, holotype of *Coelorhynchus asteroides*, 85.5 mm HL, 376+ mm HL, Owase fish market, Mie Pref., Pacific. (A) Lateral, (B) dorsal, and (C) ventral views. [Photos: N. Nakayama]

not discernible. Anterior nostril small and oval; posterior elongated slit. Grooved lateral line complete throughout.

Origins of first dorsal and pelvic fins on about same vertical; pectoral-fin base slightly anterior to this level. First dorsal fin moderately high, its height 2.0–3.6 times as long as its base length; second spinous ray not elongated, smooth along its leading edge (1–2 rudimentary denticles rarely present distally); its tip usually extending to second dorsal-fin origin or beyond when laid back. Interdorsal space 1.4–2.4 (usually ≥ 1.7) times as long as first dorsal-fin base length. Second dorsal fin originating posterior to

vertical through anal-fin origin (above base of 5th–10th anal-fin rays). Outer pelvic-fin ray slightly prolonged, with filamentous tip.

Color when fresh (Fig. 43). Head and body purplish gray dorsally, whitish ventrally; silvery reflection slightly developed over lateral surface of body; abdomen pale in lateral and ventral views; opercle and dorsum body iridescent soon after capture; first dorsal fin darker distally, but paler basally; pectoral fin dark dusky; pelvic fin dusky; distal margin of anal fin narrowly black posteriorly.

Color in alcohol (Figs. 21B, 44). Head and body

generally light brown dorsally, paler ventrally; no obvious body markings in adults, but juveniles with small dark spots on posterior part of tail (Fig. 21B); underside of head dusky; abdomen dark violet in ventral view, but chest paler; lips white; oral and gill cavities blackish; gill rakers and filaments pale, arches dusky; gular and branchiostegal membranes pale; first dorsal fin uniformly dark, but second spinous ray blackish; second dorsal fin immaculate; anal fin paler anteriorly, but prominently darker posteriorly; pectoral and pelvic fins generally dusky; outer pelvic-fin rays paler distally.

Size. To about 46 cm TL (SNFR 14555, 416+ mm TL, Okinawa Trough, Japan).

Distribution. Restricted to Japan and Taiwan (Appendix 3-2E). Known from off the Pacific coasts of southern Japan northward to the Izu Peninsula (138.96°E), Okinawa Trough, and northern South China Sea, at depths of 233–420 m. Common in the East China Sea, but seemingly rare in the Pacific off southern Japan.

Comments on holotype. *Coelorinchus hige* was described from a single specimen collected from Suruga Bay off Heta, Shizuoka Pref., Japan. It is one of the six species of grenadiers described by Matsubara (1943), whose holotypes were originally registered in the Sigenkagaku Fish Collection of the Research Institute for Natural Resources, Tokyo. These species are: *C. kamoharai* (holotype SKFC 4; ex. Matsubara's personal number 2498); *C. hige* (SKFC 6); *C. longissimus* (SKFC 5; ex. 1592); *C. vermicularis* (SKFC 3; ex. 2495) (a junior synonym of *C. multispinulosus* Katayama, 1942); *Lionurus abei* (SKFC 7; ex. 4909) [a junior synonym of *Nezumia proxima* (Smith & Radcliffe in Radcliffe, 1912)]; and *L. japonicus* (SKFC 8; ex. 1951) (= *Kumba japonica*). Unfortunately, the building of the institute was destroyed by a bombing raid in 1945, and these holotypes are now believed to be lost (Eschmeyer 1998, Fricke *et al.* 2020). These types, if still extant, are most likely to be present in the ichthyological collection of the Maizuru Fisheries Research Station, Kyoto University, Maizuru, where Matsubara worked after the war. However, none were found in the FAKU collection, despite thorough surveys by the present author in 2009–2010 and 2016–2018.

In the original description of *C. hige*, Matsubara (1943) provided an excellent description of the holotype, with an illustration of a body scale taken from the dorsum below the interdorsal space (Matsubara 1943: fig. 6). Although the holotype is apparently lost, its diagnostic features were adequately detailed in Matsubara's description, viz., snout length 39% HL; orbit diameter 28% HL; interorbital width 38% HL; postorbital length 33% HL; pelvic–anus distance 33% HL; interdorsal length 38% HL; barbel length 11% HL; teeth in bands in both jaws, none especially enlarged; premaxillary tooth band moderately broad throughout its length; short light organ immediately anterior to anus, its length as long as pupil diameter; body scales covered with 7–9 divergent

rows of spinules; nasal fossa lacking scales; underside of snout almost completely naked; second dorsal-fin origin far posterior to vertical through anal-fin origin and above base of eighth anal-fin ray. Subsequently, Matsubara (1955:1313), in his key to species of Japanese grenadiers, noted that *C. hige* has five scales below the midbase of the first dorsal fin.

Nomenclatural discussion. Although the name *C. hige* is available under the ICZN (1999), many recent authors did not mention the presence of this species (e.g., Okamura 1984b; Nakabo 2002; Nakabo & Kai 2013). Iwamoto (1990) is one of the few authors who referred to *C. hige*, implying its close similarity with *C. asteroides* Okamura, 1963. In his key to species of *Coelorinchus*, *C. hige* keys out together with *C. asteroides* at 27a, but no further couplet is provided to distinguish the two species (Iwamoto 1990:130).

Coelorinchus asteroides was described by Okamura (1963a) based on eight specimens collected from off the Pacific coasts of southern Japan, and all subsequent authors have considered the species valid (e.g., Okamura 1970a, 1984b; Yatou 1984; Iwamoto 1990; Nakabo 1993, 2000, 2002; Nakabo & Kai 2013; Iwamoto *et al.* 2015). The holotype (Fig. 44) and two of the paratypes (FAKU 21513 and 29115) were examined in the course of this study. Unfortunately, five paratypes (FAKU 7057, 24909, 25194, 33910, and 33981) were not found in the FAKU collection during the author's visit in 2009. Subsequently, a specimen labeled as a paratype of *C. asteroides* was rediscovered during the 2010 visit. Although it neither bore a catalog number nor had locality data, it is characterized by having the following counts and measurements: first dorsal-fin rays II,8; pectoral-fin rays I,17; pelvic-fin rays 7; inner gill rakers on first arch 6; HL 67.5 mm; snout length 26.0 mm (39% HL); orbit diameter 20.0 mm (30%); postorbital length 22.5 mm (33%); suborbital width 9.3 mm (14%); upper-jaw length 16.1 mm (24%); interorbital width 16.2 mm (24%); interdorsal length 28.2 mm (42%). These data completely match with those given by Okamura (1963a: table 1) for FAKU 24909, a paratype of *C. asteroides* collected from Owase; thus, the specimen is considered here to bear the catalog number of this specimen.

According to Okamura (1963a:25), *C. asteroides* is similar in general appearance to *C. hige*, but differs in having 3.5 scales below the midbase of the first dorsal fin (vs. 5 in *C. hige*), the postorbital length less than the pelvic–anus distance (vs. almost equal), and the scales on the median rostral ridge covered with radiating rows of spinules (vs. "... without divergent carinae of spinules"). Subsequently, Okamura (1970a:154) further distinguished *C. asteroides* from *C. hige* by the absence of scales on the underside of the head (vs. "underside of head with a triangular scaly area at the distal part"). However, these proposed differences appear to be too subtle or obscure to reflect their specific separation.

FIGURE 45. Scanning electron micrographs showing body scales of *Coelorinchus hige*. (A–B) SNFR 14555, 110 mm HL; (C–F), BSKU 27032, 70.2 mm HL. (A–D) Scales from the dorsum below the interdorsal space; (E–F) from the top of the head between the parietal ridges. (A, C, E) Views from above; (B, D, F) oblique views. [Photos: N. Nakayama]

An examination of the available types and additional specimens of *C. asteroides*, along with the original description of *C. hige*, revealed that the two species are not separable morphologically. In the beginning of this study, the additional specimens were treated as *C.*

asteroides for convenience of comparison, because some were already reported under this name by previous authors (Okamura 1984b, 1988, 1997; Yatou 1984). All counts and measurements of the holotype of *C. hige* (based on Matsubara 1943, 1955) fall within the range of variation

of the *C. asteroides* specimens examined, including the numbers of scale rows and the pelvic–anus distance (Table 2). Unfortunately, it is unknown how the spinules on the midrostral scales were arranged in the holotype of *C. hige*, because Matsubara (1943:145) described only the holotype as having the scales “without carinae diverging outward and backward from front of the scales” [sic, probably meaning the spinules are not arranged in distinct rows]. In *C. asteroides*, the midrostral scales are generally covered with radiating rows of spinules that originate slightly anterior to the center of the scale. However, the rows are often irregular in small specimens, and the condition agrees well with Matsubara’s (1943) description. Furthermore, the holotype of *C. hige* was somewhat small, measuring 256 mm TL (fide Matsubara 1943:144). Matsubara (1943) described *C. hige* as having “the underside of the head naked except the small triangular area at the distal part where the several small spinous scales are crowded” [sic]. However, a scaly area such as this was never observed in the *C. asteroides* specimens examined nor in closely related congeners. His description probably refers to the ventral surface of the terminal scute. Consequently, these two species are difficult to differentiate, and *C. hige* is herein regarded as a senior synonym of *C. asteroides*.

Relationships and comparisons. *Coelorinchus hige* belongs to the *C. anatirostris* group (see the Relationships of *C. anatirostris*), and is most closely related to *C. radcliffei* Gilbert & Hubbs, 1920 known from the Philippines and Indonesia. The holotype of the latter species was examined here for comparison. *Coelorinchus hige* agrees with the holotype and the original description of *C. radcliffei*, with no diagnostic differences in counts and measurements, except for the internasal width (30–35% PRL in *C. hige* vs. 27% in *C. radcliffei*). However, according to the original description, *C. radcliffei* has distinctly fewer pyloric caeca (39; Gilbert & Hubbs 1920:501) as compared with *C. hige* (47–49; this study). Although adults of the two species are strikingly similar to each other, *C. hige* is easily diagnosed when young by the presence of several dark spots on the posterior part of the tail [Fig. 21B; vs. absent in *C. radcliffei*, fide Gilbert & Hubbs 1920:502]. The relatively short snout of young *C. hige* is also distinctive: the snout of three small specimens of *C. hige* (30.0–41.0 mm HL) ranged from 42 to 44% HL, whereas *C. radcliffei* of comparable size had a distinctively longer snout of 48% HL (Gilbert & Hubbs 1920: unnumbered table in p. 503, fig. 22, 38.0 mm HL). Furthermore, the terminal snout scute of *C. hige* is shorter than that of *C. radcliffei* (see Gilbert & Hubbs 1920: fig. 23), although the difference is only especially prominent in small specimens. Except for the holotype, no additional specimens of *C. radcliffei* were examined in this study, and further investigations based on more specimens are required to verify the difference between adults of the two species.

Coelorinchus hige superficially resembles *C. goobala* Iwamoto & Williams, 1999 known from the eastern Indian Ocean off northwestern Australia. However, it readily differs from *C. goobala* in that the dorsal surface of the snout is completely scaled except for the narrow naked clefts along each side of the median rostral ridge; by contrast, the snout of *C. goobala* is broadly naked posterior to the lateral nasal ridges. Furthermore, the body scales of *C. hige* are covered with keel-like spinules that greatly overlap and closely adjoin one another to form saw-toothed ridges, whereas in *C. goobala*, the spinules are slender, conical, and individually separated, forming comb-like ridges. Other pronounced differences include the number of pyloric caeca (47–49 in *C. hige* vs. 33–44 in *C. goobala*) and the coloration of the oral cavity (blackish vs. pale). *Coelorinchus hige* is further distinguished from *C. goobala* when young by the presence of dark spots on the posterior part of the tail (Fig. 21B) (vs. absent), and a relatively shorter scute at the tip of the snout [length of terminal snout scute 13–17% PRL (30.0–63.4 mm HL; see also Fig. 21B) vs. 18–33% (54.7–63.5 mm HL; see also Iwamoto & Williams 1999: fig. 11b–b’)].

Because of their similarity in general appearance and sympatric occurrence, in Japanese waters *C. hige* is most likely to be misidentified as *C. anatirostris* Jordan & Gilbert in Jordan & Starks, 1904. However, the two species are easily distinguished by squamation characters. In *C. hige*, the occipital scales are covered with short, erect, recurved, needle-like spinules in widely divergent comb-like rows (Fig. 45E–F), whereas in *C. anatirostris*, spinules are keel-like and arranged in narrowly divergent rows (Fig. 22E–F). The absence of scales on the nasal fossa is also useful to diagnose *C. hige* (vs. usually heavily scaled in *C. anatirostris*), although large adults of *C. hige* often have a few tiny scales, and also the fossa in small specimens of *C. anatirostris* is occasionally naked. In *C. hige*, scales on the median rostral ridge are covered with radiating rows of spinules, whereas in *C. anatirostris*, the spinules are arranged in divergent rows (the spinule rows are often irregular in small specimens of the former). The dorsal contour of the snout is almost straight in *C. hige*, but it is generally concave in *C. anatirostris*. Additionally, *C. hige* has significantly higher counts of pyloric caeca (47–49 vs. 21–26 in *C. anatirostris*) and a relatively wider interdorsal space [51–70% PRL vs. 33–54%; second dorsal fin originating above the bases of 1st–6th anal-fin ray (usually anterior to 5th) vs. 5th–10th (usually posterior to 6th)]. Although the two species are difficult to identify in the field, *C. hige* is readily recognized when young by having several dark spots on the posterior part of the tail (vs. absent in *C. anatirostris*; Fig. 21). The combination of the snout shape, the position of the second dorsal-fin origin, and the presence/absence of scales of the nasal fossa distinguishes larger specimens of the two species in the field.

***Coelorinchus hubbsi* Matsubara, 1936**

[Japanese name: Moyo-hige]

(Figs. 46, 47A; Appendix 3-2F)

Coelorhynchus hubbsi Matsubara, 1936:358, figs. 2–3 [original description; single specimen (whereabouts unknown); type locality: Owase, in 100–200 ftm (183–366 m); new Japanese name: “Moyô-hige”]; Kamohara 1936:934 (listed; Kochi); Okada & Matsubara 1938:450 (in key; Japan); Kamohara 1938:73 (brief description; spec. from Kochi Pref.); Kamohara 1950:276 (listed; Kochi and Wakayama Pref.); Matsubara *et al.* 1951:42 (listed; Mie Pref.); Kamohara 1952:99 (brief description; spec. from Kochi Pref.); Matsubara 1955:1311, pl. 133, fig. 456 (in key; Japan); Kamohara 1958:73 (listed; Kochi Pref.); Okada *et al.* 1959:83 (listed; Kumano-nada); Kamohara 1964:95 (listed; Kochi Pref.); Tataru *et al.* 1965:121 (listed; off Shikoku); Matsubara 1965:505 (compiled; Japan); Okamura 1970a:168, pl. XXXVI, text-fig. 68 (description; biological notes; 118 spec. from Pacific off southern Japan from Choshi to Totoro); Kuroda 1971:56 [listed from Suruga Bay after Okamura (1970a)]; Kataoka & Tomida 1981:78 (listed; Mie Pref.); Tominaga & Uyeno 1981:488 (listed; Japan); Ozawa 1983:13 (listed; off Makurazaki, Kagoshima Pref., East China Sea).

Coelorhynchus (Quincuncia) hubbsi: Okamura 1970b: table 1 (listed; Japan).

Coelorinchus hubbsi: Okamura 1984b:96, pl. 82, fig. L (compiled); Okamura 1988:96, pl. 82, fig. L (compiled); Okamura 1997:126, fig. 5 (compiled); Shao *et al.* 2008b: table 2 (2 spec. listed from northeastern Taiwan); Nakabo & Kai 2013:507 (in key; Japan); Fukui *et al.* 2015:182, fig. 10 (1 spec. from off Kuro-shima Island, Kagoshima Pref., East China Sea); Ikeda & Nakabo 2015:321, pl. 68, figs. 1–6 (brief description; spec. from Wakayama Pref.); Iwamoto *et al.* 2015:57 (brief description; 2 spec. from northeastern Taiwan); Motomura 2020:38 (listed; Japan).

Coelorinchus hubbsi: Nakabo 1993:367 (in key; Japan); Shao 1993:169 (compiled; Taiwan); Suzuki & Kataoka 1997:83, pl. 33, fig. 180 (brief description; 1 spec. from off Owase); Shinohara & Matsuura 1997:290, pl. 2-B (listed; Suruga Bay); Nakabo 2000:431 (in key; Japan); Shinohara *et al.* 2001:304 (76 spec. listed from Tosa Bay); Nakabo 2002:431 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Nakajima 2003:54 (brief description; 1 spec. from Enshu-nada); Shinohara *et al.* 2005:416 (listed; Ryukyu Islands); Iwatsuki *et al.* 2017:32 (listed; Hyuga-nada).

Diagnosis. Ventral light organ externally represented by long dark line, terminated with 1 fossa at each end, one immediately anterior to anus, the other on chest just posterior to isthmus (duct connecting each fossa often difficult to distinguish). Underside of head completely naked except along lateral nasal ridges; dorsal surface of snout almost fully scaled, naked clefts only narrowly developed along each side of median rostral ridge. Snout long, sharply pointed, length 72–85% PRL; dorsal profile

of snout slightly concave in lateral view; terminal scute long, slender, spearhead-shaped, length 20–27% PRL; distal portion of snout narrowly attenuated when viewed dorsally; lateral nasal ridge incompletely supported by nasal bone. Interdorsal space less than first dorsal-fin base length. Anal fin originating posterior to vertical through second dorsal-fin origin. First dorsal fin high, but second spinous ray not especially elongate, height of fin 85–130% PRL. Second dorsal fin well developed, its rays as long as opposites of anal fin. Anus immediately anterior to anal-fin origin. Premaxillary teeth small, slender, conical in short, uniformly wide band, none especially enlarged; posterior end of tooth band falling far short of lateral corner of mouth. Body scales covered with short, reclined, needle-like spinules in subparallel to slightly divergent rows; spinules in each row overlapping, but free from one another, with last spinule extending well beyond posterior scale margin; buttresses scarcely developed. Orbit diameter 45–52% PRL; postorbital length 47–56% PRL; upper-jaw length 38–43% PRL; preoral length 64–79% PRL; barbel length 13–19% PRL. Transverse scale rows below first dorsal-fin midbase 4–6.5. Spinules on scales anterior to light organ not rudimentary. Longitudinally elongate dark blotch between lateral line and pectoral fin, extending from above pectoral-fin base to below about second dorsal-fin origin; narrow dark stripe extending mid-laterally from pectoral-fin base to middle of tail or beyond; scale pockets on dorsal half of body outlined by dark brown; underside of head dark to dusky, heavily peppered; lips pale; oral cavity blackish; gular and branchiostegal membranes dusky to uniformly black; first dorsal fin pale, with large, black, apical blotch; pelvic fin dark, but outermost ray paler distally.

Material examined. 32 specimens. **Japan:** BSKU 77924 (1, 42.8 mm HL, 178+ mm TL), BSKU 97993 (1, 59.5 mm HL, 234+ mm TL), south of Muroto, 33.0790°N, 134.1645°E, 414–513 m, R/V *Tansei-maru*, cr. KT-05-29, sta. K-500, beam trawl, coll. H. Endo *et al.*, 17 Nov. 2005; BSKU 97985 (1, 54.9 mm HL, 232+ mm TL), BSKU 97988 (1, 49.4 mm HL, 188+ mm TL), Tosa Bay, 300 m, FRV *Kotaka-maru*, otter trawl, 16 Jul. 1997; BSKU 42505 (1, 53.9 mm HL, 216+ mm TL), BSKU 42509 (1, 28.8 mm HL, 120+ mm TL), Tosa Bay, FRV *Kotaka-maru*, otter trawl, 14 Jan. 1986; *BSKU 74229 (1, ca. 18 mm HL, 78+ mm TL), Mimase fish market, bottom trawl, 14 Feb. 2005; BSKU 99355 (1, 47.9 mm HL, 193+ mm TL), Mimase fish market, bottom trawl, 11 Jan. 2007; BSKU 75645 (1, 34.0 mm HL, 131+ mm TL), Mimase fish market, bottom trawl, 15 Dec. 2004; BSKU 57945 (1, 44.9 mm HL, 187+ mm TL), Mimase fish market, bottom trawl, 17 Mar. 2002; BSKU 72793 (1, 57.5 mm HL, 238 mm TL), Mimase fish market, bottom trawl, 20 Feb. 2004; BSKU 99468 (1, 53.5 mm HL, 216+ mm TL), Mimase fish market, bottom trawl, 26 Mar. 2005; BSKU 99302 (1, 34.9 mm HL, 138+ mm TL), Mimase

fish market, bottom trawl, 8 Apr. 2007; BSKU 99300 (1, 27.1 mm HL, 110+ mm TL), Mimase fish market, bottom trawl, coll. N. Nakayama, 8 Apr. 2007; BSKU 94810 (1, 40.2 mm HL, 161+ mm TL), Mimase fish market, F/V *Kosei-maru*, bottom trawl, 28 Nov. 2007; BSKU 92269 (1, 40.8 mm HL, 169+ mm TL), Mimase fish market, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama, 9 Nov. 2007; BSKU 99283 (1, 43.4 mm HL, 178+ mm TL), Mimase fish market, F/V *Seiryō-maru*, bottom trawl, coll. N. Nakayama, 14 Nov. 2007; *BSKU 95378 (1, 54.0 mm HL, 195+ mm TL), Mimase fish market, F/V *Seiryō-maru*, bottom trawl, coll. N. Nakayama, 12 Mar. 2008; BSKU 97967 (1, 51.9 mm HL, 210+ mm TL), BSKU 97968 (1, 52.4 mm HL, 216+ mm TL), BSKU 97969 (1, 56.1 mm HL, 232+ mm TL), BSKU 97970 (1, 60.3 mm HL, 231+ mm TL), BSKU 97974 (1, 61.6 mm HL, 244+ mm TL), BSKU 97975 (1, 57.3 mm HL, 243+ mm TL), BSKU 97978 (1, 54.4 mm HL, 226 mm TL), Tosa Bay, 33.1412°N, 133.5388°E, 295–336 m, FRV *Kotaka-maru*, otter trawl, 18 Jul. 1988; BSKU 113676 (1, 46.0 mm HL, 182+ mm TL), BSKU 113677 (1, 51.0 mm HL, 215+ mm TL), BSKU 113679 (1, 52.4 mm HL, 200+ mm TL), BSKU 113680 (1, 48.4 mm HL, 199 mm TL), BSKU 113681 (1, 43.7 mm HL, 174+ mm TL), BSKU 113685 (1, 44.4 mm HL, 184+ mm TL), off Okitsu, Tosa Bay, 280–320 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 24 Jan. 2014; BSKU 110077 (1, 55.3 mm HL, 221+ mm TL), Suruga Bay, 34.7481°N, 138.4652°E, 243–424 m, F/V *Hinode-maru*, sta. 3, bottom trawl, coll. N. Nakayama and R. Misawa, 23 Apr. 2013.

Counts and measurements. Based on 30 specimens (27.1–61.6 mm HL, 110+–244+ mm TL). Counts: first dorsal-fin rays II,8–9; pectoral-fin rays i15–i19; pelvic-fin rays 7; gill rakers on first arch (outer/inner) 0/8–9, on second arch 5–7/8–9; longitudinal scales 30–38; transverse scale rows below first dorsal-fin origin 5.5–8, below first dorsal-fin midbase 4–6.5, below second dorsal-fin origin 5–7, above anal-fin origin 13.5–19.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 42–46 (72–85); orbit diameter 25–30 (45–52); postorbital length 27–31 (47–56); postrostral length 54–59; orbit–preopercle distance 28–31 (49–54); suborbital width 11–14 (21–24); upper-jaw length 21–25 (38–43); length of rictus 16–19 (28–34); length of premaxillary tooth band 7–11 (13–20); preoral length 37–43 (64–79); length of terminal snout scute 11–15 (20–27); length of lateral nasal ridge 29–34 (50–62); length of suborbital ridge 91–96 (157–174); snout width 25–30 (43–52); internasal width 17–23 (31–39); interorbital width 20–24 (36–42); occipital width 9–12 (15–20); body width over pectoral-fin bases 33–48 (58–85); body depth at first dorsal-fin origin 40–54 (69–95); body depth at anal-fin origin 31–44 (53–77); prepelvic length 100–113 (174–201); preanus length 137–158 (238–278); preanal length 139–160 (243–285); isthmus–pelvic distance 25–33 (43–

56); isthmus–anus distance 62–80 (108–143); isthmus–anal distance 66–85 (114–152); pelvic–anal distance 41–58 (72–103); anus–anal distance 2–6 (3–10); pelvic-fin length 33–46 (59–79); pectoral-fin length 39–44 (67–76); predorsal length 107–113 (185–202); height of first dorsal fin 48–71 (85–130); length of first dorsal-fin base 16–22 (29–39); interdorsal length 8–14 (14–25); length of gill slit 10–14 (18–23); length of posterior nostril 6–9 (11–16); barbel length 7–11 (13–19).

Size. To about 26 cm TL (FRLM 8636, 255+ mm TL, Owase, Japan).

Distribution. Restricted to Japan and Taiwan (Appendix 3-2F). Known from the Pacific off southern Japan from Hitachi (36.60°N; Ibaraki Pref.) southward to Cape Ashizurimisaki (133.02°E), off the Izu Islands, and the East and South China Seas off Taiwan, at depths of 240–555 m (Chiou *et al.* 2004b; Shao *et al.* 2008b; this study). Common on the upper continental slope off the Pacific coasts of southern Japan, but rare in the east of the Boso Peninsula (139.89°E) as well as in Taiwan.

Remarks. *Coelorinchus hubbsi* was originally described from a single male specimen (224 mm TL) collected from off Owase, Mie Pref., Japan. In the original description of this species, Matsubara (1936) indicated neither the catalog number of the holotype nor the whereabouts of the specimen. Unfortunately, the holotype is presumed to be destroyed during WWII (see also the Comments on holotype of *C. hige*).

For a full description see Okamura (1970a). *Coelorinchus hubbsi* is moderately common in the Pacific off southern Japan, with a great abundance in Tosa Bay (author's pers. observ.). It is also known from the Senkaku Islands, off northwestern Taiwan (Iwamoto *et al.* 2015). Surprisingly, despite intensive collecting efforts, no specimens of this species have been obtained in the East China Sea by Japanese researchers (e.g., Okamura & Kitajima 1984; Okamura 1985; Yamada 1986; Yamada *et al.* 2007; Shinohara *et al.* 2005; Furuhashi *et al.* 2010).

Relationships and comparisons. *Coelorinchus hubbsi* is a distinctive species of the *C. hubbsi* group (see the Relationships of *C. brevirostris*), and unlikely to be confused with any other congeners except when young. Its characteristic dark stripes on the body (Fig. 46A) readily distinguish the species from the other species of the *C. hubbsi* group. In young specimens of *C. hubbsi*, the horizontal stripes are less pronounced, whereas the saddles are more or less distinct (Fig. 46D). These features may lead to an erroneous identification of the species as *C. matsubarai* Okamura, 1982. However, the pectoral blotch of young *C. matsubarai* is circular and not elongated horizontally, as seen in adults of the species (Fig. 70).

FIGURE 46. Fresh specimens of *Coelorinchus hubbsi* collected from the Mimase fish market, Kochi Pref., Pacific. (A–C) BSKU 95378, 54.0 mm HL, 195+ mm TL; (D) BSKU 74229, ca. 18 mm HL, 78+ mm TL. (A, D) Lateral views; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

FIGURE 47. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of two species of the *Coelorinchus hubbsi* group. (A) *C. hubbsi*, BSKU 97970, 60.3 mm HL; (B) *C. matsubarai*, BSKU 32226, paratype, 75.5 mm HL. Views from above. [Photos: N. Nakayama]

***Coelorinchus japonicus* (Temminck & Schlegel, 1846)**

[Japanese name: Tojin]

(Figs. 48–51, 52A, 53A–B, 54A; Appendix 3-3A)

Macrourus japonicus Temminck & Schlegel, 1846:256, pl. CXII, figs. II, IIa, IIb [original description; in part; lectotype: RMNH 3746, from “bays in provinces Oomura and Shimabara” (Nagasaki Pref.)]; Boeseman 1947:184 (in part; lectotype designation; re-examination of type specimens).

Macrurus japonicus: Günther 1862:395 [brief description; 2 spec. (only heads) from Japan]; Günther 1887:127 [brief description; 1 spec. from off Inoshima (= Enoshima Island), Sagami Bay, Challenger sta. 232]; Steindachner & Döderlein 1887:283 (brief description; 1 spec. probably from Sagami Bay); Doflein 1906:234, unnumbered fig. (spec. from Sagami Bay).

Coelorhynchus japonicus: Jordan & Snyder 1901:120 (listed; Japan; local name “Hige”); Jordan & Gilbert in Jordan & Starks 1904:617 (description; 8 spec. from Misaki and Nagasaki); Franz 1910:26 (12 spec. from Yokohama and Aburatsubo); Jordan *et al.* 1913:418 (listed; Japan; Japanese names: “Hige” or “Tōjin”); Gilbert & Hubbs 1916:178 (supplementary description; 4 spec. from off Cape Shionomisaki and Suruga Bay); Izuka & Matsuura 1920:103 (spec. from “Tōkyō Market”); Jordan & Hubbs 1925:327 (spec. from Misaki); Tanaka 1927:477, fig. 923 (compiled; Japan; standard Japanese name: “Tojin”); Schmidt 1931:155 (brief description; 1 spec. from Nagasaki); Kamohara 1934a:54 (listed; Kochi); Kamohara 1934b:301 (1 spec. from Susaki, Kochi Pref.); Okada & Matsubara 1938:450 (in key; Japan); Kamohara 1938:73 (4 spec. from Kochi Pref.); Kamohara 1950:276, fig. 211 (listed; Kochi and Wakayama Pref.); Tanaka 1951:137, fig. 350 (compiled; Japan); Kuroda 1951:391 (listed; Suruga Bay); Uchinomi 1951:76 (listed; southern part of Kii Peninsula); Kamohara 1952:99 (spec. from Kochi Pref.); Asano *et al.* 1952:97 (listed; Ibaraki Pref.); Kamohara 1955:63, pl. 63, fig. 2 (compiled; Japan); Matsubara 1955:1313 (in key; Japan); Kobayashi 1956:72 (listed; Enshu-nada); Kamohara 1958:73 (listed; Kochi Pref.); Kamohara 1961a:63, pl. 63, fig. 2 (compiled; Japan); Kamohara 1964:96 (listed; Kochi Pref.); Tataru *et al.* 1965:121 (listed; off Shikoku); Matsubara 1965:506 (compiled; Japan); Dotsu & Tomiyama 1967:35 (spec. from Goto Islands); Kamohara 1967:125, pl. 63, fig. 2 (compiled; Japan); Okamura 1970a:183, pl. XL, text-figs. 78–79 (description; biological notes; 93 spec. from Pacific off southern Japan from Sagami Bay to Mimase); Tominaga & Uyeno 1981:488 (listed; Japan); Yatou 1982:181, 353, fig. 106 (description; 8 spec. from Tosa Bay; photo based on BSKU 29937); Ohta 1983: table A (listed with a question mark; Suruga Bay; in situ observation); Akazaki 1984:265 (listed; Aoshima, Miyazaki Pref.).

[?] *Coelorhynchus parallelus* (not Günther 1877): Kuroda 1941:80, fig. 11 (brief description; 2 spec. from Suruga Bay) [re-identified from figure]; Kuroda 1951:392 (listed; Suruga Bay); Kuroda 1962:8 (description of fresh color; 1 spec. from Suruga Bay).

Coelorhynchus (Oxymacrus) japonicus: Okamura 1970b: table 1 (listed; Japan).

Coelorinchus japonicus: Yatou 1984:231, 367, fig. 163 (brief description; 5 spec. from Okinawa Trough; photo based on BSKU 29886); Okamura 1984b:97, pl. 83, fig. D (compiled); Kamohara & Okamura 1985:32, pl. 31, fig. 154 (compiled; Japan); Okamura 1988:97, pl. 83, fig. D (compiled); Iwamoto 1990:162, fig. 378 (synopsis); Hori 1996:28 (listed; Ibaraki Pref.); Suzuki & Kataoka 1997:84, pl. 33, fig. 182 (brief description; 1 spec. from Kumano-nada); Okamura 1997:127, fig. 14 (compiled); Funabashi 1998:85 (listed; Ibaraki Pref.); Youn 2002:539 (questionable; listed; Korea); Shao *et al.* 2008b: table 2 (3 spec. listed from northeastern Taiwan); Nakabo & Kai 2013:510 (in key; Japan); Ikeda & Nakabo 2015:322, pl. 68, fig. 12 (brief description; spec. from Wakayama Pref.); Iwamoto *et al.* 2015:57 (brief description; 7 spec. from northeastern Taiwan); Miyazaki *et al.* 2019: table 1 (1 spec. listed from Tokyo Submarine Canyon); Motomura 2020:38 (listed; Japan).

Caelorinchus japonicus: Nakabo 1993:369 (in key; Japan); Shao 1993:169, fig. 37-7 (compiled; Taiwan; photo spec. is *C. leptorhynchus*); Shinohara & Matsuura 1997:290 (listed; Suruga Bay); Nakabo 2000:433 (in key; Japan); Shinohara *et al.* 2001:305 (listed; Tosa Bay); Nakabo 2002:43 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Nakajima 2003:55, pl. 15, fig. 90 (brief description; 1 spec. from Enshu-nada); Yamaguchi & Machida 2003:240, pl. 26, fig. 97 (comments on type specimens); Shinohara *et al.* 2005:416 (8 spec. listed from Ryukyu Islands); Kim *et al.* 2005:172, fig. 292 (compiled; Korea; photo spec. represents *C. multispinulosus*); Senou *et al.* 2006:420 (listed; Sagami Sea); Shinohara *et al.* 2009:708 (listed; Pacific off Tohoku); Furuhashi *et al.* 2010: table 2 (44 spec. listed from northern Okinawa Trough); Iwatsuki *et al.* 2017:32 (listed; Hyuga-nada).

Caelorinchus (Oxymacrus) japonicus: Chiou *et al.* 2004a: table 1 (listed; Taiwan).

Coelorinchus parallelus (not Günther 1877): Ikeda & Nakabo 2015:321, pl. 68, figs. 7–11 (brief description; spec. from Wakayama Pref.) [re-identified from figures].

Coelorinchus sp.: Ikeda & Nakabo 2015:322, pl. 69, figs. 1–5 (brief description; 1 spec. from Kii Channel off Wakayama Pref.; new Japanese name: “Minabe-sokodara”).

Diagnosis. Light organ externally represented by short, narrow, naked streak immediately anterior to anus, its anterior margin falling far short of line connecting inner pelvic-fin bases. Underside of head almost completely scaled, except for narrow naked area above upper lip and anterior portions of mandibular rami; top of snout fully scaled. Snout long, sharply pointed, length 67–90% PRL; dorsal profile of snout concave in lateral view (rarely almost straight in large specimens); terminal scute moderately short, diamond-shaped, dorsoventrally flattened; lateral nasal ridge completely supported by nasal bone. Anus slightly separated from anal-fin origin. Premaxillary teeth

small, conical, arranged in short, uniformly wide band, with outer series enlarged; posterior margin of tooth band falling far short of lateral corner of mouth. Body scales covered with short, reclined, keel-like spinules in widely divergent, coarsely saw-toothed ridges; every spinule row complete, extending to posterior scale margin; spinules in each row greatly overlapping, increasing in height posteriorly; buttresses narrowly developed. Occipital scales covered with single (rarely 2–3 narrowly divergent) saw-toothed row of keel-like spinules. Transverse scale rows below second dorsal-fin origin 5–7.5. Orbit diameter 39–52% PRL; suborbital width 18–23% PRL; internasal width 26–33% PRL; interorbital width 31–38% PRL; pectoral-fin length 55–71% PRL. Body generally dark, without prominent markings; lips white; oral cavity blackish; gular membrane pale; branchiostegal membranes darker posteriorly; fins blackish.

Material examined. 48 specimens. **Lectotype of *Macrourus japonicus*:** *RMNH 3476 (33+ cm TL, fide Boeseman 1947:184; examination based on only photographs; Fig. 49A), bays in provinces Omura and Shimabara, Nagasaki Pref., Japan, East China Sea, coll. P.F. von Siebold. **Non-types: Japan:** BSKU 106780 (1, 96.0 mm HL, 320+ mm TL), southeast of Shimokoshikijima Island, East China Sea, 31.5560°N, 129.8847°E, 415 m, F/V *Maruko-maru*, tr. 4, bottom trawl, coll. N. Nakayama *et al.*, 24 Apr. 2012; BSKU 26863 (1, 135 mm HL, 421+ mm TL), north-northwest of Amami-oshima Island, Okinawa Trough, 29.7850°N, 128.4467°E, 1000 m, F/V *Yuryo-maru*, No. 8, tr. 17, bottom trawl, coll. Y. Kinoshita and S. Hagino, 3 Feb. 1978; BSKU 27111 (1, 105 mm HL, 322+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.3833°N, 127.5150°E, 650 m, F/V *Yuryo-maru*, No. 8, tr. 21, bottom trawl, coll. Y. Kinoshita and S. Hagino, 5 Feb. 1978; *BSKU 109017 (1, 128 mm HL, 485+ mm TL), west of Okikasayama Bank, 32.1503°N, 129.0072°E, 499–504 m, T/V *Nagasaki-maru*, cr. N365, sta. C5, 3-m ORE beam trawl, coll. N. Nakayama, 19 Nov. 2012; *BSKU 43971 (1, 188 mm HL, 710+ mm TL), off Kamikawaguchi, Tosa Bay, long line, 26 Aug. 1987; *BSKU 115479 (1, 122 mm HL, 403+ mm TL), off Aki, Tosa Bay, 550 m, F/V *Tsukasa-maru*, rod and reel, coll. T. Yamasaki, 6 Jul. 2014; *BSKU 45514 (1, 222 mm HL, 665+ mm TL), off Tosa-Shimizu, long line, 31 Aug. 1988; *BSKU 77919 (1, 34.2 mm HL, 126 mm TL), south of Muroto, 33.0790°N, 134.1645°E, 414–513 m, R/V *Tansei-maru*, cr. KT-05-29, sta. K-500, beam trawl, coll. H. Endo *et al.*, 17 Nov. 2005; BSKU 32426 (1, 127 mm HL, 468+ mm TL), BSKU 32430 (1, 119 mm HL, 410+ mm TL), off Nahari, Tosa Bay, 33.2792°N, 133.8947°E, 450–470 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. 1-L3, long line, coll. Y. Ishida and T. Araya, 17 Nov. 1979; BSKU 69780 (1, 82.8 mm HL, 279+ mm TL), Tosa Bay, 33.1900°N, 133.6883°E, 542–585 m, FRV *Kotaka-maru*, otter trawl, coll. H. Endo, 22 Jan. 1998; BSKU 44419 (1, 106 mm HL, 337+ mm

TL), BSKU 44420 (1, 128 mm HL, 416+ mm TL), BSKU 44423 (1, 117 mm HL, 377+ mm TL), Tosa Bay, 600 m, FRV *Kotaka-maru*, otter trawl, 11 Jan. 1988; BSKU 71583 (1, 38.8 mm HL, 150+ mm TL), Tosa Bay, 400 m, FRV *Kotaka-maru*, otter trawl, 9 May 2000; BSKU 112040 (1, 49.5 mm HL, 172+ mm TL), BSKU 112044 (1, 51.0 mm HL, 163+ mm TL), BSKU 112046 (1, 31.3 mm HL, 116+ mm TL), BSKU 112049 (1, 91.4 mm HL, 285+ mm TL), BSKU 112051 (1, 63.4 mm HL, 228+ mm TL), BSKU 112052 (1, 62.7 mm HL, 177+ mm TL), BSKU 112054 (1, 109 mm HL, 400+ mm TL), BSKU 112070 (1, 45.0 mm HL, 136+ mm TL), BSKU 112074 (1, 99.1 mm HL, 343+ mm TL), Tosa Bay, 33.1428°N, 133.6248°E, 450–490 m, FRV *Kotaka-maru*, 1 Aug. 1988; BSKU 43816 (1, 71.0 mm HL, 247+ mm TL), Tosa Bay, 400 m, FRV *Kotaka-maru*, 11 Jun. 1987; BSKU 65948 (1, 86.6 mm HL, 275+ mm TL), BSKU 97955 (1, 72.4 mm HL, 261+ mm TL), Tosa Bay, 400 m, FRV *Kotaka-maru*, otter trawl, 9 May 2000; BSKU 43977 (1, 90.5 mm HL, 327+ mm TL), Tosa Bay, 500 m, FRV *Kotaka-maru*, otter trawl, 20 Apr. 1987; BSKU 44422 (1, 118 mm HL, 448 mm TL), Tosa Bay, 600 m, FRV *Kotaka-maru*, otter trawl, 11 Jan. 1988; BSKU 23288 (1, 150 mm HL, 486+ mm TL), Mimase fish market, bottom trawl, date unknown; BSKU 110136 (1, 134 mm HL, 487 mm TL), BSKU 110138 (1, 97.7 mm HL, 367+ mm TL), Suruga Bay, 34.7619°N, 138.4811°E, 196–387 m, F/V *Hinode-maru*, sta. 5, bottom trawl, coll. N. Nakayama and R. Misawa, 23 Apr. 2013; BSKU 110084 (1, 80.2 mm HL, 243+ mm TL), BSKU 110085 (1, 77.7 mm HL, 282 mm TL), BSKU 110119 (1, 120 mm HL, 449+ mm TL), BSKU 110120 (1, 122 mm HL, 428 mm TL), Suruga Bay, 34.7489°N, 138.4664°E, 200–450 m, F/V *Hinode-maru*, sta. 6, bottom trawl, coll. N. Nakayama and R. Misawa, 23 Apr. 2013; BSKU 110032 (1, 88.5 mm HL, 334 mm TL), BSKU 110033 (1, 62.6 mm HL, 193+ mm TL), BSKU 110035 (1, 88.0 mm HL, 300+ mm TL), Suruga Bay, 34.7127°N, 138.4553°E, 262–434 m, F/V *Hinode-maru*, sta. 1, bottom trawl, coll. N. Nakayama and R. Misawa, 23 Apr. 2013; KPM-NI 28790 (1, 152 mm HL, 514+ mm TL), ca. 22 km south-southeast of Mie-shima Island, Kumano-nada, 34.0713°N, 136.6453°E, 606–669 m, R/V *Tansei-maru*, cr. KT-10-16, sta. SK-1, 3-m beam trawl, coll. T. Sato, 14 Aug. 2010; FRLM 2535 (1, 130 mm HL, 419+ mm TL), off Mie, Kumano-nada, long line, coll. Fisheries Research Institute of Mie Pref., 16 Apr. 1980; *BSKU 110419 (1, 29.0 mm HL, 112+ mm TL), Suruga Bay, 34.7489°N, 138.4664°E, 200–450 m, F/V *Hinode-maru*, sta. 6, bottom trawl, coll. N. Nakayama and R. Misawa, 23 Apr. 2013; KPM-NI 13883 (1, 140 mm HL, 518+ mm TL), off Zushi, Sagami Bay, 500 m, hook and line, coll. M. Miyazawa, 14 Mar. 2004; BSKU 103027 (1, 137 mm HL, 454+ mm TL), Sagami Bay, 950 m, rod and reel, coll. T. Okamoto, 18 Apr. 2010; KPM-NI 12824 (1, 149 mm HL, 528+ mm TL), off Odawara, Sagami Bay, 450 m, hook and line, 15 Mar. 2003; KPM-NI 12080 (1, 126 mm HL, 468+ mm

FIGURE 48. Fresh specimens of *Coelorinchus japonicus*. (A) BSKU 77919, 34.2 mm HL, 126 mm TL, south of Muroto, Kochi Pref., Pacific, 414–513 m depth; (B) BSKU 106780, 96.0 mm HL, 320+ mm TL, southeast of Shimokoshiki-jima Island, Kagoshima Pref., East China Sea, 415 m depth; (C) BSKU 103027, 137 mm HL, 454+ mm TL, Sagami Bay, Pacific; (D) BSKU 115479, 122 mm HL, 403+ mm TL, Tosa Bay off Aki, Kochi Pref., Pacific, 550 m depth; (E) BSKU 43971, 188 mm HL, 710+ mm TL, Tosa Bay, Pacific. Lateral views. [Photos: (A–D) N. Nakayama; (E) BSKU]

FIGURE 49. Type specimens of *Macrourus japonicus* (= *Coelorinchus japonicus*) collected from Nagasaki Pref., East China Sea. (A) RMNH 3476, lectotype; (B) RMNH D1405, paralectotype; (C) RMNH D1406, paralectotype. (A) Preserved in ethanol; (B–C) stuffed. Lateral views. [Photos: Y. Machida]

TL), off Fukuura, Sagami Bay, 350 m, hook and line, 8 Apr. 2002; KPM-NI 19322 (1, 172 mm HL, 630+ mm TL), west of Nii-jima Island, Izu Islands, 400 m, hook and line, coll. H. Ishikawa, 14 Jun. 2007.

Counts and measurements. Based on 41 specimens (31.3–172 mm HL, 116+–630+ mm TL). Counts: first dorsal-fin rays II,8–10; pectoral-fin rays I,17–20; pelvic-fin rays 7; gill rakers on first arch (outer/inner) 0–1/6–9, on second arch 6–8/7–9; longitudinal scales 37–48; transverse scale rows below first dorsal-fin origin 5.5–8, below first dorsal-fin midbase 4–6.5, below second dorsal-fin origin 5–7.5, above anal-fin origin 17–28.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 40–47 (67–90); orbit diameter 22–30 (39–52); postorbital length 28–36 (49–61); postrostral length 53–60; orbit–preopercle distance 30–38 (52–64); suborbital width 10–14 (18–23); upper-jaw length 23–29 (41–49); length of rictus 17–23 (31–41); length of premaxillary tooth band 11–15 (19–26); preoral length 35–45 (58–84); length of terminal snout scute 5–9 (8–17); length of lateral nasal ridge 25–34 (42–63); length of suborbital ridge 91–97 (152–182); snout width 23–29 (38–52); internasal width 15–19 (26–33); interorbital width 17–22 (31–38); occipital width 7–11 (13–19); body width over pectoral-fin bases 32–48 (57–85); body depth at first dorsal-fin origin 40–55 (72–99); body depth at anal-fin origin 30–47 (53–82); prepelvic length 103–118 (178–207); preanus length 127–147 (221–256); preanal length 130–154 (225–268); isthmus–pelvic distance 22–33 (38–57); isthmus–anus distance 44–68 (81–117); isthmus–anal distance 48–74 (88–127); pelvic–anal distance 25–47 (43–82); anus–anal distance 4–9 (8–16); pelvic-fin length 28–48 (47–83); pectoral-fin length 31–41

(55–71); predorsal length 105–112 (180–202); height of first dorsal fin 35–42 (62–74); length of first dorsal-fin base 12–19 (22–32); interdorsal length 18–30 (32–49); length of gill slit 10–13 (17–23); length of posterior nostril 7–9 (11–16); barbel length 7–11 (12–18).

Size. Attains about 93 cm TL (FRLM 3456, 927+ mm TL, Kumano-nada, Japan).

Distribution. Restricted to Japan and Taiwan. So far known only from the Pacific off southern Japan northward to Hitachi (36.60°N; Ibaraki Pref.) and the Okinawa Trough, at depths of 150–1000 m (Shao *et al.* 2008b; this study; Appendix 3-3A). Very common in the east of the Boso Peninsula (139.89°E).

Comments on type specimens. *Coelorinchus japonicus* was the first grenadier species described from the North Pacific. It was originally described as *Macrourus japonicus* in the Pisces volume of the *Fauna Japonica* written by Coenraad J. Temminck and Hermann Schlegel. This outstanding monograph was published separately in 16 fascicles from 1842 to 1850, and the description of *C. japonicus* (pp. 256–258) appeared in the 15th fascicle published in 1846 (fide Sherborn & Jentink 1895; Yamaguchi & Machida 2003:93). The specimens examined by Temminck and Schlegel were collected by Philipp F. von Siebold and Heinrich Bürger (see also the Introduction). To record fresh coloration, many of Bürger's specimens were drawn by Keiga Kawahara, a Japanese artist who produced 260 figures of fishes in response to Bürger's order (Yamaguchi & Machida 2003:89). The plates and color descriptions of Temminck & Schlegel are based largely on Kawahara's unpublished drawings, which are currently available online (Nagasaki Museum of History and Culture 2015; Naturalis 2020; Wikimedia Commons 2020).

FIGURE 50. Illustrations showing type specimens of *Macrourus japonicus* (= *Coelorinchus japonicus*). (A) Figures provided in the original description (Temminck & Schlegel 1846: pl. CXII, figs. II, IIa, IIb); (B) drawing by Keiga Kawahara. [Photos: (A) Faculty of Science Common Library, Kyoto University; and (B) Naturalis Biodiversity Center, Leiden; these images were cropped from originally digitized versions.]

According to Temminck & Schlegel (1846:257), *C. japonicus* was described from a single specimen (about one ft TL) collected from Nagasaki. This specimen was also illustrated in their monograph (pl. CXII, figs. II, IIa, IIb; see also Fig. 50A), but its catalog number was not indicated in the original description. About a century after the *Fauna Japonica*, Boeseman (1947) re-examined fish specimens collected by von Siebold and Bürger, reviewing the type materials of the species described by Temminck & Schlegel. According to Boeseman (1947:184), there were three specimens referable to *C. japonicus* in the ichthyological collection of the Rijksmuseum

van Natuurlijke Histoire, Leiden (now the Naturalis Biodiversity Center), of which one is preserved in alcohol (RMNH 3476, Fig. 49A) and the other two are stuffed (RMNH 1405 and RMNH 1406, Fig. 49B–C). Boeseman considered these three specimens to be conspecific and the type series of *C. japonicus*, designating RMNH 3467 as the lectotype. The remaining two specimens automatically became paralectotypes under Art. 74.1.3 of the ICZN (1999), although no indication was given by Boeseman. However, as mentioned above, Temminck & Schlegel (1846) noted that their description was based on only a single specimen, and thus Boeseman's treatment

FIGURE 51. Preserved specimen of *Coelorinchus japonicus*. BSKU 45514, 222 mm HL, 665+ mm TL, off Tosa-Shimizu, Kochi Pref., Pacific. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

FIGURE 52. Photographs showing spinulation of the occipital scales in two species of *Coelorinchus*. (A) *C. japonicus*, BSKU 44422, 118 mm HL; (B) *C. smithi*, BSKU 65947, 97.2 mm HL. Preserved specimens. [Photos: N. Nakayama]

of the type materials contradicts Temminck & Schlegel's statement.

Another problem is also included in the description and figures of *C. japonicus* provided by Temminck & Schlegel (1846). Although the *C. japonicus* specimens examined in this study generally agreed with the original description of this species, significant differences were found in coloration. Temminck & Schlegel (1846:258) described the species as having prominent, circular, grayish-blue spots on the lateral surface of the body. The characteristic spots are also depicted in their color figure (pl. CXII, fig. II). However, none of the specimens

examined in this study had such body markings. The color of the first dorsal, pectoral, and pelvic fins is also significantly different. In Temminck & Schlegel's figure, these fins are illustrated as pale except for a dark line along the second spinous ray of the first dorsal fin, whereas in the specimens examined in this study, these fins were generally uniformly dark.

While the author had no opportunity to directly examine the type series of *C. japonicus*, their photographs were provided by Dr. Yoshihiko Machida, a Professor Emeritus of Kochi University (Fig. 49). Based on the photograph (Fig. 49A), RMNH 3467 corresponds

FIGURE 53. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of three species of the *Coelorinchus japonicus* group. (A–B) *C. japonicus*, BSKU 109017, 128 mm HL; (C–D) *C. leptorhinus*, BSKU 99870, 96.5 mm HL; (E–F) *C. smithi*, BSKU 65947, 97.2 mm HL. (A, C, E) Views from above; (B, D, F) oblique views. [Photos: NSMT]

with what many authors have called *C. japonicus*. Unfortunately, both stuffed specimens (RMNH D1405 and RMNH D1406; Fig. 49B–C) are in poor condition, with limited information available on their identification. However, they appear to be a species of the *C. argentatus*

group (see the Relationships of *C. formosanus*) and most likely to be *C. multispinulosus* Katayama, 1942. This suggestion is further supported by Kawahara's drawing for one of the two stuffed specimens (Fig. 50B). His drawing agrees in all respects with fresh specimens of *C.*

FIGURE 54. Preserved juvenile specimens of two species of *Coelorinchus*. (A) *C. japonicus*, BSKU 110419, 29.0 mm HL, 112+ mm TL, Suruga Bay, (B) *C. smithi*, BSKU 71585, 34.5 mm HL, 123+ mm TL, Tosa Bay. Lateral views. [Photos: N. Nakayama]

multispinulosus (Fig. 72), viz., general appearance, body markings, and fin color. Therefore, there is little doubt that the two stuffed specimens do not represent *C. japonicus* but *C. multispinulosus*.

As mentioned above, Temminck & Schlegel (1846:257) stated that the original description of *C. japonicus* was based on a single specimen of about one ft TL. Considering the size of the specimen, it must correspond to the alcohol preserved specimen RMNH 3476 (ca. 33+ cm TL, fide Boeseman 1947:184). Temminck & Schlegel's figures (figs. II, IIa, IIb in pl. CXII) are also considered to be based on this specimen, because they capture the presence of numerous tiny scales on the underside of the head (vs. absent in *C. multispinulosus*). However, it is also clear that they complemented the color of the figure with Kawahara's drawing that illustrates either RMNH 1405 or RMNH 1406 (= *C. multispinulosus*). Therefore, the original description is based substantially on two specimens, which means that Boeseman's (1947) use of the term "lectotype" is correct. For convenience, both RMNH 1405 and RMNH 1406 are recognized here as paralectotypes, although it is still uncertain which specimen was drawn by Kawahara.

Remarks. For further morphological information see Okamura (1970a). *Coelorinchus japonicus* is one of a few grenadier species recorded from the Sea of Japan. Although many authors listed the species from the area (e.g., Kikuchi 1931; Katayama 1940; Mori 1952, 1956; Tsuda 1964), no voucher specimens exist in museum collections, and these records appear to represent misidentified *C. multispinulosus*. These misidentifications are probably due to the fact that the color figure given in the original description exhibits mixed characters of *C. japonicus* and

C. multispinulosus (as discussed above). In addition, prior to Okamura (1970a) who provided a review of all the then-known Japanese grenadiers, *C. japonicus* was often illustrated as having fine dark vermiculations dorsally on the body (e.g., Tanaka 1927: fig. 923; Kamohara 1950: fig. 211; Matsubara 1965: fig. 1434). *Coelorinchus multispinulosus* is occasionally found on the continental shelf north of southern Honshu, dominating the grenadier specimens from the Sea of Japan (based on the museum collections examined).

Ueno (1965, 1971) recorded *C. japonicus* from off the Pacific coast of Hokkaido, but it appears to represent *C. gilberti* Jordan & Hubbs, 1925, a species often found in the area. Maeda & Maruyama (1991), Maeda & Tsutsui (2003), and Nakabo & Kai (2013:1874) also doubted Ueno's records of *C. japonicus*. Similarly, Amaoka *et al.* (2020) recently listed *C. japonicus* from Hokkaido, but their photographed specimen (fig. 223) likely represents *C. gilberti*.

A large individual of *Coelorinchus* sp. reported by Ikeda & Nakabo (2015:70, 322, pl. 69, figs. 1–5; WW 9706, 649 mm TL) is considered here an unusual variant of *C. japonicus*, because their brief description and several figures of the specimen agree well with the diagnosis of the latter species, except that the snout tip is distinctly rounded in dorsal view. The snout is usually sharply pointed in *C. japonicus* when viewed dorsally, but its tip is occasionally rounded in the large specimens examined (Fig. 51), possibly due to damage during early ontogeny. Unfortunately, Ikeda & Nakabo's (2015) specimen has been processed to be transferred to the Wakayama Prefectural Museum of Natural History, Wakayama (WW), and the present author had no access to the specimen in the

course of this study. A direct examination of the voucher specimen is needed to determine the taxonomic status of this ambiguous individual.

Weber (1913) reported *C. japonicus* from Indonesia, including the Flores Sea, Bali Sea, Makassar Strait, and Arafura Sea. Subsequently, Gilbert & Hubbs (1920:488) cited Weber's *C. japonicus* in the synonymy of their new species, *C. acantholepis* Gilbert & Hubbs, 1920.

Okamura (1970a:186) considered *C. japonicus* to be widely distributed in tropical and subtropical waters of the Indo-West Pacific. However, Iwamoto (1990:163) claimed that there were no voucher specimens that would support such a distribution, and this species is regarded here as restricted to the northwestern Pacific off Japan and Taiwan.

Relationships. *Coelorinchus japonicus* belongs to a group of *Coelorinchus* species characterized by the following combination of features: light organ short, extending from anus to far posterior to pectoral-fin bases; lateral nasal ridge completely supported by nasal bone, not strongly convex when viewed dorsally; premaxillary teeth in short, uniformly wide band; underside of head mostly or almost entirely scaled; body scales covered with widely divergent rows of keel-like spinules, and every row complete throughout; buttresses of body-scale spinules scarcely to narrowly developed; no prominent markings on body (Nakayama & Endo 2017; this study). This group corresponds with one of two subgroups of the *C. japonicus* group as redefined by Nakayama & Endo (2017), comprising the following nine species in which the underside of the head is completely or almost completely scaled: *C. charius* Iwamoto & Williams, 1999 from off Western Australia; *C. commutabilis* Smith & Radcliffe in Radcliffe, 1912 from the Philippines; *C. japonicus*; *C. leptorhinus* Chiou, Shao & Iwamoto, 2004 from Taiwan; *C. macrorhynchus* Smith & Radcliffe in Radcliffe, 1912 from the Philippines and Australia; *C. nox* **sp. nov.** described here; *C. sereti* Iwamoto & Merrett, 1997 from the tropical southwestern Pacific; *C. smithi* Gilbert & Hubbs, 1920 from Japan to Australia; *C. supernasutus* McMillan & Paulin, 1993 from Australia and New Zealand. This group is called here the *C. japonicus* group (*sensu stricto*) for convenience of comparison.

Coelorinchus flabellispinis (Alcock, 1894), *C. lasti* Iwamoto & Williams, 1999, and *C. trunovi* Iwamoto & Anderson, 1994, all known from the Indian Ocean, share most of the above-mentioned features. However, the lateral nasal ridge of these three species is incompletely supported by the nasal bone. *Coelorinchus flabellispinis* and *C. lasti* also lack any external evidence of the light organ anterior to the anus.

Comparisons. *Coelorinchus japonicus* is a distinctive species of the *C. japonicus* group (*sensu stricto*) that readily differs from its similar congeners in that the occipital scales are covered with a single row of spinules [rarely 2–3 rows (Fig. 52A) vs. 4 or more widely

divergent rows (Fig. 52B)]. In addition, it differs notably from *C. leptorhinus* in having ventral surfaces of the head almost completely scaled (vs. underside of snout mostly naked); and from *C. supernasutus* in having a distinctly shorter snout (67–90% PRL vs. $\geq 111\%$). Young specimens of *C. japonicus* are easily distinguished from those of *C. charius* and *C. smithi* by the absence of dark saddles on the tail [Fig. 54A vs. present in *C. charius* (see Iwamoto & Williams 1999:135, fig. 9b) and *C. smithi* (Fig. 54B)].

***Coelorinchus jordani* Smith & Pope, 1906**

[Japanese name: Kyushu-hige]

(Figs. 55–56, 57A–B; Appendix 3-3B)

[?] *Coelorhynchus australis* (not Richardson 1893): Ishikawa & Matsuura 1897:26 [listed; spec. from Awa (= Chiba Pref.)]; Jordan & Snyder 1901:120 (listed; Japan; local name: “Hige”).

Coelorhynchus jordani Smith & Pope, 1906:494, fig. 11 (original description; holotype: USNM 55609, from Kagoshima); Jordan *et al.* 1913:419, fig. 389 (listed; Japan; new Japanese name: “Kiusiu-hige”); Gilbert & Hubbs 1916:173 (supplementary description; 24 spec. from Kagoshima market, East China Sea, and Kagoshima Bay; comparison with *C. kishinouyei*); Schmidt 1931:156 (brief description; 11 spec. from Kagoshima); Okada & Matsubara 1938:450 (in key; Japan); Kamohara 1950:277 (listed; Kochi and Wakayama Pref.); Kamohara 1952:99 (spec. from Kochi Pref.); Matsubara 1955:1310 (in key; Japan); Kamohara 1958:73 (listed; Kochi Pref.); Kamohara 1964:95 (listed; Kochi Pref.); Okamura 1970a:175 (description; 31 spec. from Pacific off southern Japan from Suruga Bay to Mimase); Kuroda 1971:56 [listed from Suruga Bay after Okamura (1970a)]; Tominaga & Uyeno 1981:488 (listed; Japan); Ozawa 1983:13 (listed; off Makurazaki, Kagoshima Pref., East China Sea).

Coelorhynchus (Paramacrurus) jordani: Okamura 1970b: table 1 (listed; Japan).

Coelorinchus jordani: Yatou 1984:229, 367, fig. 162 [brief description; 10 spec. from Okinawa Trough; photo based on BSKU 37574 (from Kagoshima Bay)]; Okamura 1984b:96, pl. 83, fig. B (compiled); Yamada 1986:101 (brief description; East China Sea); Okamura 1988:96, pl. 83, fig. B (compiled); Okamura 1997:126, fig. 10 (compiled); Nakabo & Kai 2013:508 (in key; Japan); Iwatsubo & Motomura 2017:59, unnumbered fig. (listed; Kagoshima Bay); Motomura 2020:38 (listed; Japan).

Coelorinchus jordani: Nakabo 1993:368 (in key; Japan); Shinohara & Matsuura 1997:290 (listed; Suruga Bay); Nakabo 2000:432 (in key; Japan); Shinohara *et al.* 2001:305 (listed; Tosa Bay); Nakabo 2002:432 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Shinohara *et al.* 2005:416 (listed; Ryukyu Islands); Yamada *et al.* 2007:315, pl. 14, fig. 6 (brief description; 1 spec.; East China Sea; biological notes); Furuhashi *et al.* 2010: table 2 (1 spec. listed from northern Okinawa Trough); Iwatsuki *et al.* 2017:32 (listed; Hyuga-nada).

Diagnosis. Light organ moderately long, externally represented by narrow black streak immediately anterior to anus (covered with scales), its anterior margin extending forward to line connecting outer pelvic-fin bases, but falling far short of isthmus. Underside completely naked; dorsal surface of snout posterior to lateral nasal ridges broadly naked. Snout moderately long, length 51–65% PRL, its dorsal profile slightly concave in lateral view; terminal scute small but broad, dorsoventrally flattened; lateral nasal ridge completely supported by nasal bone, broadly convex in dorsal view. Length of chin barbel 12–19% PRL. Anus separated from anal-fin origin by several scale rows. Premaxillary teeth small, conical, arranged in short, uniformly wide band, none especially enlarged; posterior margin of tooth band falling far short of lateral corner of mouth. Body scales covered with short, greatly reclined, needle-like spinules in tightly packed parallel rows; spinules in each row greatly overlapping, but free from one another; buttresses absent. Body silvery white but darker dorsally; small dorsoventrally elongate dark spot between lateral line and pectoral-fin base (occasionally absent or indistinct); lips and oral cavity whitish; gular membrane prominently blackish; chest mostly pale; anal fin pale except for dark anterior margin.

Material examined. 31 specimens. **Holotype of *Coelorhynchus jordani*:** USNM 55609 (39.5 mm HL, 172+ mm TL), Kagoshima, Kagoshima Pref., Japan, coll. H.M. Smith, 16 Jun. 1903. **Non-types: Japan:** BSKU 37574 (1, 44.7 mm HL, 207+ mm TL), BSKU 37579 (1, 39.8 mm HL, 189 mm TL), off Furue, Kagoshima Bay, 13 Oct. 1982; BSKU 47208 (1, 52.7 mm HL, 222+ mm TL), off Ibusuki, Kagoshima Bay, 190 m, 24 May 1989; BSKU 50884 (1, 46.7 mm HL, 205+ mm TL), off Nomaike, Kagoshima, East China Sea, F/V *Maruko-maru*, bottom trawl, coll. M. Yamashita and Y. Ohashi, 10 Sept. 2010; BSKU 109023 (1, 38.7 mm HL, 181+ mm TL), BSKU 109024 (1, 34.9 mm HL, 160+ mm TL), BSKU 109027 (1, 27.8 mm HL, 107+ mm TL), BSKU 109042 (1, 42.0 mm HL, 208+ mm TL), east of Kasayama Bank, 32.3884°N, 129.0505°E, 304–312 m, T/V *Nagasaki-maru*, cr. N365, sta. A3, 3-m ORE beam trawl, coll. N. Nakayama, 19 Nov. 2012; BSKU 27725 (1, 48.3 mm HL, 181+ mm TL), north-northwest of Okinawa Island, Okinawa Trough, 28.5000°N, 127.0000°E, 225–235 m, F/V *Ryoan-maru*, No. 28, tr. 51, bottom trawl, coll. Y. Kinoshita and S. Hagino, 12 Mar. 1978; BSKU 92878 (1, 57.1 mm HL, 241+ mm TL), off Kannoura, Mimase fish market, F/V *Kosei-maru*, bottom trawl, 9 Dec. 2007; BSKU 91920 (1, 44.3 mm HL, 227+ mm TL), Mimase fish market, bottom trawl, coll. N. Nakayama, 29 Oct. 2007; BSKU 99349 (1, 50.8 mm HL, 230+ mm TL), BSKU 99350 (1, 59.2 mm HL, 247+ mm TL), Mimase fish market, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama, 9 Dec. 2007; BSKU 87825 (1, 41.8 mm HL, 204+ mm TL), Mimase fish market, bottom trawl, 14 Jan. 2004; BSKU 51770 (1, 53.9 mm HL, 242+ mm TL), BSKU 51772 (1, 64.3 mm

HL, 279+ mm TL), Mimase fish market, bottom trawl, 18 Nov. 1999; BSKU 68922 (1, 55.5 mm HL, 274+ mm TL), Mimase fish market, bottom trawl, 20 Jan. 2004; BSKU 54045 (1, 51.7 mm HL, 217+ mm TL), Mimase fish market, bottom trawl, 4 Sept. 1999; BSKU 36133 (1, 48.8 mm HL, 252+ mm TL), Mimase fish market, bottom trawl, 5 Dec. 1981; BSKU 58380 (1, 55.3 mm HL, 252+ mm TL), Mimase fish market, bottom trawl, 6 Dec. 2001; BSKU 89341 (1, 50.0 mm HL, 222+ mm TL), BSKU 99357 (1, 47.2 mm HL, 193+ mm TL), Mimase fish market, bottom trawl, coll. N. Nakayama, 11 Jan. 2007; BSKU 76501 (1, 46.2 mm HL, 227+ mm TL), Mimase fish market, bottom trawl, coll. N. Nakayama, 30 Nov. 2005; BSKU 112513 (1, 54.1 mm HL, 238+ mm TL), off Okitsu, Tosa Bay, 380 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 11 Mar. 2014; *BSKU 101167 (1, 56.2 mm HL, 275+ mm TL), Mimase fish market, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama, 5 Nov. 2009; BSKU 110421 (1, 32.5 mm HL, 166 mm TL), Suruga Bay, 34.7651°N, 138.4880°E, 200–450 m, F/V *Hinode-maru*, sta. 4, bottom trawl, coll. N. Nakayama and R. Misawa, 23 Apr. 2013; BSKU 111216 (1, 68.2 mm HL, 281+ mm TL), BSKU 111219 (1, 57.1 mm HL, 223+ mm TL), BSKU 111222 (1, 61.6 mm HL, 298+ mm TL), BSKU 111224 (1, 55.0 mm HL, 269+ mm TL), Isshiki fish market, coll. S. Ohashi, 14 Feb. 2010.

Counts and measurements. Based on 30 specimens (27.8–68.2 mm HL, 107+–298+ mm TL). Counts: first dorsal-fin rays II, 8–10; pectoral-fin rays i13–i17; pelvic-fin rays 7–8; gill rakers on first arch (outer/inner) 0/8–10, on second arch 6–8/7–9; longitudinal scales 27–37; transverse scale rows below first dorsal-fin origin 4.5–6, below first dorsal-fin midbase 3.5–5, below second dorsal-fin origin 3.5–5, above anal-fin origin 11.5–17.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 34–40 (51–65); orbit diameter 29–36 (46–56); postorbital length 30–36 (46–55); postrostral length 61–67; orbit–preopercle distance 36–41 (56–66); suborbital width 12–15 (19–22); upper-jaw length 24–29 (36–43); length of rictus 18–23 (28–34); length of premaxillary tooth band 10–13 (16–20); preoral length 28–38 (41–59); length of terminal snout scute 4–8 (7–12); length of lateral nasal ridge 24–28 (35–46); length of suborbital ridge 90–100 (134–162); snout width 28–32 (43–50); internasal width 17–21 (27–33); interorbital width 19–23 (30–36); occipital width 10–12 (15–19); body width over pectoral-fin bases 37–53 (59–79); body depth at first dorsal-fin origin 54–72 (83–115); body depth at anal-fin origin 42–67 (65–102); prepelvic length 106–117 (163–186); preanus length 128–148 (200–235); preanal length 136–160 (211–254); isthmus–pelvic distance 30–41 (47–65); isthmus–anus distance 50–70 (77–111); isthmus–anal distance 56–84 (87–133); pelvic–anal distance 29–47 (46–75); anus–anal distance 7–16 (11–25); pelvic-fin length 37–48 (55–78); pectoral-fin length 45–59 (68–87); predorsal length 111–

119 (168–186); height of first dorsal fin 59–79 (93–122); length of first dorsal-fin base 20–27 (31–42); interdorsal length 29–45 (43–74); length of gill slit 11–14 (17–23); length of posterior nostril 8–13 (13–21); barbel length 7–13 (12–19).

Size. To about 32 cm TL (ZMUT 856, 323+ mm TL, Sagami Bay, Japan).

Development. The early life stages of *C. jordani* were fully described by Fukui *et al.* (2010), with a complete series of eggs and larvae.

Distribution. So far known only from the Pacific off southern Japan northward to Hitachi (36.60°N; Ibaraki Pref.), Kagoshima Bay, and the Okinawa Trough, at depths of 110–550 m (Appendix 3-3B). Common, especially in Kagoshima Bay (Ohtomi 2006).

Remarks. *Coelorinchus jordani* was described by Smith & Pope (1906) based on a single specimen collected from Kagoshima Pref., Japan (Fig. 56). This species has been well described by previous authors, and thus does not need repeating here. For further morphological details

see the original description (Smith & Pope 1906) and Okamura (1970a). It is one of the most abundant demersal fishes in Kagoshima Bay (Ohtomi 2006), but not so in other parts of its range.

Relationships. *Coelorinchus jordani* is one of 12 species closely related to *C. argus* Weber, 1913. According to Iwamoto *et al.* (2015), this complex also includes: *C. dorsalis* Gilbert & Hubbs, 1920; *C. kishinouyei* Jordan & Snyder, 1900; *C. macrolepis* Gilbert & Hubbs, 1920; *C. maculatus* Gilbert & Hubbs, 1920; *C. notatus* Smith & Radcliffe in Radcliffe, 1912; *C. sexradiatus* Gilbert & Hubbs, 1920; *C. thurla* Iwamoto & Williams, 1999; *C. triocellatus* Gilbert & Hubbs, 1920; *C. velifer* Gilbert & Hubbs, 1920; and *C. sp. cf. notatus* (sensu Iwamoto *et al.* 2015). Gilbert & Hubbs (1920) classified most of these species under the subgenus *Paramacrurus* Bleeker, 1874. However, this subgenus was originally established with the type species *C. australis* (Richardson, 1839) that is now considered a member of the *C. fasciatus* group as defined by Iwamoto & Graham (2008). Therefore,

FIGURE 55. Fresh specimens of *Coelorinchus jordani*. (A) BSKU 109042, 42.0 mm HL, 208+ mm TL, Okinawa Trough, East China Sea, 304–312 m depth; (B–D) BSKU 101167, 56.2 mm HL, 275+ mm TL, Mimase fish market, Kochi Pref., Pacific. (A–B) Lateral views; (C) dorsal and (D) ventral views of the head and trunk. [Photos: N. Nakayama]

FIGURE 56. Holotype of *Coelorhynchus jordani* (= *Coelorinchus jordani*). USNM 55609, 39.5 mm HL, 172+ mm TL, Kagoshima, Pref. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

the complex is called here the *C. argus* group, which is defined by the following combination of features: light organ moderately long, extending to or slightly beyond line connecting inner bases of pelvic fins; lateral nasal ridge completely supported by nasal bone; anus slightly separated from anal-fin origin; body scales covered with parallel to slightly divergent rows of greatly reclined, needle-like spinules; dark blotch between lateral line and pectoral-fin base (in most species). The *C. argus* group differs most notably from the *C. fasciatus* group in that the lateral nasal ridge is incompletely supported by the nasal bone.

Comparisons. Among the *C. argus* group, *C. jordani* differs from all but *C. kishinouyei* in that the spinules on the body scales are arranged in tightly packed parallel rows (vs. moderately separated, parallel to slightly divergent rows). The species also differs from all, except *C. argus*, *C. kishinouyei*, *C. macrolepis*, *C. thurla*, and *C. sp. cf. notatus*, in that the second spinous ray of the first dorsal fin is not especially elongate (vs. greatly prolonged; see Iwamoto 1990: figs. 284–292). *Coelorinchus jordani* is most similar to *C. kishinouyei*, the only other species of the group occurring in the Pacific off southern Japan. However, *C. jordani* differs from *C. kishinouyei* in that the dark spot between the lateral line and the pectoral-fin base is small and rudimentary, and even absent in some specimens (vs. large and prominent). Apart from the difference in coloration, *C. jordani* is distinguished from that species by the position of the light organ: in *C. jordani*, the anterior margin of the light organ extends forward beyond a line connecting the outer pelvic-fin bases, whereas that of *C. kishinouyei* only reaches this line. Compared with *C. kishinouyei*, *C. jordani* has a longer barbel [2.6–4.1 in orbit (12–19% PRL) vs. 4.2–

7.9 (7–13%)], and a somewhat smaller orbit [0.8–1.2 in postorbital length (usually 1.0–1.1) vs. 0.8–1.1 (usually 0.8–0.9)]. In addition, dorsal surfaces of the snout posterior to the leading edges are broadly naked in most specimens of *C. jordani*, whereas the areas are variously scaled in *C. kishinouyei* (rarely lacking scales).

***Coelorinchus kamoharai* Matsubara, 1943**

[Japanese name: Ichimonji-hige]

(Figs. 34B, 58–59; Appendix 3-3C)

Coelorhynchus kamoharai Matsubara, 1943:136, figs. 1B, 2B, 4 [original description; holotype: FAKU 2498 (ex. SKFC 4), from Kumano-nada; 14 paratypes from Kumano-nada and off Heta; new Japanese name: “Itimonji-hige”]; Matsubara *et al.* 1951:42 (listed; Mie Pref.); Kuroda 1951:391 (listed; Suruga Bay); Kamohara 1952:99 (spec. from Kochi Pref.); Matsubara 1955:1311, pl. 133, fig. 455 (in key; Japan); Kamohara 1958:73 (listed; Kochi Pref.); Okada *et al.* 1959:83 (listed; Kumano-nada); Kuroda 1962:8 (description of fresh color; 4 spec. from Suruga Bay); Kamohara 1964:95 (listed; Kochi Pref.); Matsubara 1965:505 (compiled; Japan); Okamura 1970a:159, pl. I, fig. b, text-fig. 64 (description; 111 spec. from Pacific off southern Japan from Heta to Kochi); Kataoka & Tomida 1981:78 (listed; Mie Pref.); Tominaga & Uyeno 1981:488 (listed; Japan); Ohta 1983: tables A–B (listed; Suruga Bay); Ozawa 1983:13 (listed; off Makurazaki, Kagoshima Pref., East China Sea); Akazaki 1984:265 (listed; Aoshima, Miyazaki Pref.).

Coelorhynchus (Quincuncia) kamoharai: Okamura 1970b: table 1 (listed; Japan).

Coelorinchus kamoharai: Yatou 1984:221, 365, fig. 157 [brief description; 10 spec. from Okinawa Trough; photo based

FIGURE 57. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of two species of the *Coelorinchus argus* group. (A–B) *C. jordani*, BSKU 101167, 56.2 mm HL; (C–D) *C. kishinouyei*, BSKU 111224, 55.0 mm HL. (A, C) Views from above; (B, D) oblique views. [Photos: NSMT]

on BSKU 36042 (from Mimase fish market)]; Okamura 1984b:96, pl. 82, fig. I (compiled); Okamura 1988:96, pl. 82, fig. I (compiled); Iwamoto 1990:164, fig. 382 (synopsis); Suzuki & Kataoka 1997:83, pl. 33, fig. 179 (brief description; 1 spec. from Kumano-nada); Okamura 1997:126, fig. 6 (compiled); Shao *et al.* 2008b: table 2 (6 spec. listed from northeastern Taiwan); Shinohara *et al.* 2009:708 (listed; Pacific off Tohoku); Nakabo & Kai 2013:506 (in key; Japan); Fukui *et al.* 2015:182, fig. 11 (1 spec. from off Kuro-shima Island, Kagoshima Pref., East China Sea); Ikeda & Nakabo 2015:321, pl. 67, figs. 1–5 (brief description; spec. from Wakayama Pref.); Iwamoto *et al.* 2015:58 (brief description; 9 spec. from northeastern Taiwan and Philippines); Motomura 2020:38 (listed; Japan).

Caelorinchus kamoharai: Nakabo 1993:366 (in key; Japan); Shao 1993:169, fig. 37–8 (compiled; Taiwan; photo spec. is *C. formosanus*?); Shinohara *et al.* 1996:169 (1 spec. listed from Pacific off Tohoku; 1 spec.); Shinohara & Matsuura 1997:290 (listed; Suruga Bay); Nakabo 2000:430 (in key; Japan); Shinohara *et al.* 2001:304 (98 spec. listed from Tosa

Bay); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Nakabo 2002:430 (in key; Japan); Nakajima 2003:54, pl. 15, fig. 87 (brief description; 1 spec. from Enshu-nada); Shinohara *et al.* 2005:416 (listed; Ryukyu Islands); Iwatsuki *et al.* 2017:32 (listed; Hyuga-nada).

Caelorinchus (Quincuncia) kamoharai: Chiou *et al.* 2004a: table 1 (listed; Taiwan).

Diagnosis. Light organ tubular, completely covered with scales, externally represented by long black streak extending from immediately anterior to anus to chest just posterior to isthmus. Underside of head completely naked except for overlapping scales posterior to lateral nasal ridges; dorsal surface of snout broadly naked along each side of median rostral ridge. Snout long, sharply pointed, length 63–77% PRL, its dorsal profile straight in lateral view, with slight humplike rise above nostrils; terminal scute short, arrowhead-shaped, dorsoventrally flattened, length 6–11% PRL. Lateral nasal ridge incompletely supported by nasal bone. Anus abutting anal-fin origin.

Premaxillary teeth small, slender, conical in moderately wide, long, tapered band, with outer series slightly enlarged; posterior margin of tooth band almost reaching lateral corner of mouth. Body scales covered with short, erect, narrowly triangular spinules in quincunx order; buttresses poorly developed. Second dorsal fin distinctly lower than anal fin. Orbit diameter 43–48% PRL; postorbital length 52–59% PRL; upper-jaw length 47–54% PRL; pectoral-fin length 56–74% PRL; height of first dorsal fin 65–90% PRL; interdorsal length 41–58% PRL; length of gill slit 26–29% PRL; barbel length 14–24% PRL. Dorsal half of body with faint, dark, irregular blotches; lower half of head and body silvery white when viewed laterally; median nasal bone blackish; underside of head except mandibular rami densely covered with short, black, hair-like papillae; similar papillae also present over naked areas on dorsal surface of snout; lips blackish; oral cavity pale; gular and branchiostegal membranes heavily peppered with large melanophores; chest prominently dark; first dorsal fin dusky, but second spinous ray blackish; pelvic fin pale proximally, darker distally.

Material examined. 35 specimens. **Paratypes of *Coelorrhynchus kamoharai*:** FAKU 2496 (1, 59.4 mm HL, 237+ mm TL), Owase fish market, Kumano-nada, bottom trawl, coll. K. Matsubara, 4–9 Jan. 1936; FAKU 1593 (1, 57.2 mm HL, 221+ mm TL), *FAKU 1594 (1, 48.7 mm HL, 204+ mm TL), *FAKU 1595 (1, 46.5 mm HL, 200+ mm TL), *FAKU 1596 (1, 53.1 mm HL, 208+ mm TL), Owase fish market, Kumano-nada, bottom trawl, coll. K. Matsubara, 6–9 Dec. 1935. **Non-types: Japan:** BSKU 106803 (1, 59.5 mm HL, 228+ mm TL), BSKU 106804 (1, 62.8 mm HL, 252 mm TL), southeast of Shimokoshiki-jima Island, East China Sea, 31.5560°N, 129.8847°E, 415 m, F/V *Maruko-maru*, tr. 4, bottom trawl, coll. N. Nakayama *et al.*, 24 Apr. 2012; BSKU 106828 (1, 58.0 mm HL, 226 mm TL), southeast of Shimokoshiki-jima Island, East China Sea, 31.5658°N, 129.8915°E, 380 m, F/V *Maruko-maru*, tr. 1, bottom trawl, coll. N. Nakayama *et al.*, 24 Apr. 2012; BSKU 106840 (1, 58.3 mm HL, 215+ mm TL), BSKU 106841 (1, 44.3 mm HL, 184+ mm TL), southeast of Shimokoshiki-jima Island, East China Sea, 31.5675°N, 129.8930°E, 393 m, F/V *Maruko-maru*, tr. 2, bottom trawl, coll. N. Nakayama *et al.*, 24 Apr. 2012; BSKU 112350 (1, 56.8 mm HL, 238+ mm TL), BSKU 112923 (1, 75.8 mm HL, 300 mm TL), BSKU 112924 (1, 72.3 mm HL, 290+ mm TL), BSKU 112925 (1, 50.4 mm HL, 203+ mm TL), BSKU 112926 (1, 66.0 mm HL, 269+ mm TL), off Okitsu, Tosa Bay, 280–320 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 24 Jan. 2014; BSKU 112428 (1, 56.7 mm HL, 226+ mm TL), BSKU 112466 (1, 61.2 mm HL, 256+ mm TL), off Susaki, Tosa Bay, 320–380 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 9 Mar. 2014; BSKU 99361 (1, 32.3 mm HL, 128+ mm TL), Mimase fish market, bottom trawl, 11 Jan. 2007; BSKU 89690 (1, 65.3 mm HL, 272+ mm TL), Mimase fish market, bottom trawl, 15 Jan. 2007; BSKU 99410 (1, 34.6

mm HL, 140+ mm TL), BSKU 99411 (1, 41.0 mm HL, 163+ mm TL), Mimase fish market, bottom trawl, 19 Apr. 2006; BSKU 99238 (1, 64.8 mm HL, 259+ mm TL), BSKU 99239 (1, 48.4 mm HL, 202+ mm TL), BSKU 99240 (1, 54.7 mm HL, 220+ mm TL), Mimase fish market, bottom trawl, coll. N. Nakayama, 31 Oct. 2007; BSKU 97005 (1, 53.2 mm HL, 209+ mm TL), Mimase fish market, F/V *Kosei-maru*, bottom trawl, 11 Dec. 2007; *BSKU 93575 (1, 59.6 mm HL, 248+ mm TL), Mimase fish market, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama, 21 Dec. 2007; *BSKU 94341 (1, 59.0 mm HL, 254+ mm TL), Mimase fish market, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama, 27 Mar. 2008; BSKU 108872 (1, 56.1 mm HL, 219+ mm TL), Mimase fish market, F/V *Kosei-maru*, bottom trawl, 22 Oct. 2012; BSKU 94344 (1, 61.8 mm HL, 239+ mm TL), Mimase fish market, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama, 27 Mar. 2008; BSKU 99284 (1, 49.6 mm HL, 206+ mm TL), Mimase fish market, F/V *Seiryō-maru*, bottom trawl, coll. N. Nakayama, 14 Nov. 2007; BSKU 89490 (1, 39.7 mm HL, 160+ mm TL), Kochi, date unknown; FRLM 34363 (1, 69.2 mm HL, 282+ mm TL), off Owase, Kumano-nada, coll. Iwata, date unknown; BSKU 110036 (1, 60.2 mm HL, 247+ mm TL), Suruga Bay, 34.7127°N, 138.4553°E, 262–434 m, F/V *Hinode-maru*, sta. 1, bottom trawl, coll. N. Nakayama and R. Misawa, 23 Apr. 2013; BSKU 110012 (1, 68.4 mm HL, 264+ mm TL), BSKU 110020 (1, 44.1 mm HL, 174 mm TL), Suruga Bay, 34.7080°N, 138.4587°E, 200–450 m, F/V *Hinode-maru*, sta. 2, bottom trawl, coll. N. Nakayama and R. Misawa, 23 Apr. 2013.

Counts and measurements. Based on 30 specimens (32.3–75.8 mm HL, 128+–300 mm TL). Counts: first dorsal-fin rays II, 8–10; pectoral-fin rays i14–i17; pelvic-fin rays 7; gill rakers on first arch (outer/inner) 0/9–11, on second arch 7–9/9–12; longitudinal scales 34–45; transverse scale rows below first dorsal-fin origin 4.5–6.5, below first dorsal-fin midbase 3.5–5, below second dorsal-fin origin 4–6, above anal-fin origin 11–20; pyloric caeca 10.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 39–44 (63–77); orbit diameter 25–29 (43–48); postorbital length 32–36 (52–59); postrostral length 57–62; orbit–preopercle distance 35–39 (57–66); suborbital width 11–12 (17–21); upper-jaw length 28–33 (47–54); length of rictus 24–27 (41–45); length of premaxillary tooth band 20–24 (32–39); preoral length 32–37 (51–65); length of terminal snout scute 4–7 (6–11); length of lateral nasal ridge 26–31 (41–54); length of suborbital ridge 94–100 (152–175); snout width 21–27 (35–45); internasal width 19–24 (33–39); interorbital width 24–28 (40–47); occipital width 6–8 (10–13); body width over pectoral-fin bases 33–46 (54–74); body depth at first dorsal-fin origin 45–61 (73–99); body depth at anal-fin origin 34–46 (57–78); prepelvic length 101–110 (166–185); preanus length 142–156 (231–268); preanal length 144–159 (236–271); isthmus–pelvic distance 32–40 (53–66); isthmus–anus distance 71–90 (119–146);

FIGURE 58. Fresh specimen of *Coelorinchus kamoharai*. BSKU 94341, 59.0 mm HL, 254+ mm TL, Mimase fish market, Kochi Pref., Pacific. (A) Lateral; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

FIGURE 59. Rediscovered paratype of *Coelorhynchus kamoharai* (= *Coelorinchus kamoharai*). FAKU 2496, 59.4 mm HL, 237+ mm TL, Owase fish market, Mie Pref., Pacific. (A) Lateral, (B) dorsal, and (C) ventral views. Preserved condition. [Photos: N. Nakayama]

isthmus–anal distance 73–93 (121–150); pelvic–anal distance 43–57 (71–97); anus–anal distance 1–4 (2–6); pelvic-fin length 28–40 (46–65); pectoral-fin length 33–46 (56–74); predorsal length 104–110 (170–190); height of first dorsal fin 39–55 (65–90); length of first dorsal-fin base 18–22 (30–38); interdorsal length 25–34 (41–58); length of gill slit 15–18 (26–29); length of posterior nostril 4–7 (7–12); barbel length 8–14 (14–24).

Size. To 30 cm TL (BSKU 112923, 300 mm TL, Tosa Bay, Japan).

Distribution. Restricted to Japan and Taiwan

(Appendix 3-3C). Known from off the Pacific coasts of Japan northward to the Oshika Peninsula (38.32°N), Okinawa Trough, and east of Tsushima Island, at depths of 110–450 m (Shao *et al.* 2008a, 2008b; this study). One of the most abundant *Coelorinchus* found in southern Japan, but rare in the east of the Boso Peninsula (139.89°E).

Remarks. *Coelorinchus kamoharai* was originally described by Matsubara (1943) based on 13 specimens collected from off the Pacific coast of southern Honshu, Japan (Kumano-nada and Suruga Bay off Heta). The type specimens of this species were believed to be lost

during WWII (see the Comments on holotype of *C. hige*), but five paratypes (FAKU 1593–1596, 2496, 46.5–59.4 mm HL, 200+–237+ mm TL; Fig. 59) were rediscovered among the FAKU collection during the author's visit in 2009 (Iwamoto *et al.* 2015:58).

For further morphological information see the original description (Matsubara 1943) and Okamura (1970a). This species is well represented in trawl catches from the Pacific off southern Japan at depths between 200 and 400 m (author's pers. observ.).

Relationships and comparisons. *Coelorinchus kamoharai* belongs to the *C. argentatus* group (see the Relationships of *C. formosanus*) and is most similar to *C. argentatus* Smith & Radcliffe in Radcliffe, 1912 known from the East Indies and Australasian waters. The two species are unique within the group in having short, black, hair-like papillae densely scattered over the underside of the head and the top of the snout. In *C. kamoharai* and *C. argentatus*, the gular membrane is overlain with fine striations or reticulations consisting of small black melanophores; this characteristic pigmentation is not found in other species of the *C. argentatus* group. *Coelorinchus kamoharai* differs most notably from large specimens of *C. argentatus* in having narrowly triangular spinules on the body scales (vs. broadly triangular and somewhat pyramidal). It is further distinguished from *C. argentatus* by having a larger orbit (43–48% PRL vs. 38–42%) and a shorter postorbital length (52–59% PRL vs. 57–64%).

Among Japanese congeners, *C. kamoharai* is closely similar to *C. multispinulosus* Katayama, 1942 in general appearance and coloration, with both species having similar dark spots or blotches dorsally on the body. However, the body markings of *C. kamoharai* are much larger and more obscure than those of *C. multispinulosus* (Fig. 58 vs. Fig. 72), although the dark spots of *C. multispinulosus* are less pronounced when fresh. Spinulation of the body scales also differs between the two species: in *C. kamoharai*, the body scales are covered with short, erect, narrowly triangular spinules, whereas in *C. multispinulosus*, the spinules are much longer and needle-like (Fig. 34B vs. D). It further differs from *C. multispinulosus* in having a longer interdorsal space (41–58% PRL vs. 25–41%) and a wider gill slit (26–29% PRL vs. 21–27%). While their vertical distribution is broadly overlapping at depths between 110 and 400 m, *C. kamoharai* typically occurs on the continental slope at depths below 200 m, where *C. multispinulosus* is rarely distributed (author's pers. observ.).

***Coelorinchus kishinouyei* Jordan & Snyder, 1900**

[Japanese name: Mugura-hige]

(Figs. 57C–D, 60–61; Appendix 3-3D)

Coelorhynchus kishinouyei Jordan & Snyder, 1900:376, pl. XX (original description; holotype: USNM 49395, from Misaki);

Jordan & Snyder 1901:120 (listed; Japan); Jordan & Gilbert in Jordan & Starks 1904:618, unnumbered fig. [after Jordan & Snyder (1900)]; Jordan *et al.* 1913:420, fig. 390 (listed; Japan; new Japanese name: “Mugura-hige”); Gilbert & Hubbs 1916:170 (supplementary description; 7 spec. from Suruga Bay); Izuka & Matsuura 1920:103 [spec. from “Bōshiu” (= Chiba Pref.)]; Kamohara 1931b:544 (listed; Kochi); Kamohara 1934a:54 (listed; Kochi); Okada & Matsubara 1938:450, pl. 110, fig. 2 (in key; Japan); Kamohara 1938:73 (spec. from Kochi Pref.); Kamohara 1950:277 (listed; Kochi and Wakayama Pref.); Matsubara *et al.* 1951:42 (listed; Mie Pref.); Kuroda 1951:391 (listed; Suruga Bay); Uchinomi 1951:76 (listed; southern part of Kii Peninsula); Kamohara 1952:99 (spec. from Kochi Pref.); Kuroda 1952:176, fig. 9 (brief description; 8 spec. from Suruga Bay); Matsubara 1955:1310 (in key; Japan); Kamohara 1958:73 (listed; Kochi Pref.); Okada *et al.* 1959:83 (listed; Kumano-nada); Kamohara 1964:95 (listed; Kochi Pref.); Matsubara 1965:505 (compiled; Japan); Okamura 1970a:172, pl. XXXVII, text-fig. 71 (description; biological notes; 92 spec. from Pacific off southern Japan from Kominato to Mimase); Tominaga & Uyeno 1981:488 (listed; Japan); Ohta 1983: table A (listed; Suruga Bay); Akazaki 1984:265 (listed; Aoshima, Miyazaki Pref.).

Coelorhynchus (Paramacrurus) kishinouyei: Okamura 1970b: table 1 (listed; Japan).

Coelorinchus kishinouyei: Okamura 1984b:96, pl. 83, fig. A (compiled); Okamura 1988:96, pl. 83, fig. A (compiled); Iwamoto 1990:167, fig. 386 (synopsis); Suzuki & Kataoka 1997:83, pl. 33, fig. 181 (brief description; 1 spec. from off Kii Peninsula); Shao *et al.* 2008b: table 2 (15 spec. listed from northeastern and southwestern Taiwan, and South China Sea); Fukui *et al.* 2010:171, figs. 1–2 (descriptions of early life stages; Suruga Bay); Nakabo & Kai 2013:508 (in key; Japan); Ikeda & Nakabo 2015:321, pl. 66, fig. 12 (brief description; 1 spec. from Shingu, Wakayama Pref.); Iwamoto *et al.* 2015:59 (brief description; 55 spec. from northeastern and southwestern Taiwan); Miyazaki *et al.* 2019: fig. 3C, table 1 (1 spec. listed from Tokyo Submarine Canyon); Motomura 2020:38 (listed; Japan).

Coelorinchus kishinouyei: Nakabo 1993:368 (in key; Japan); Shao 1993:169, fig. 37-9 (compiled; Taiwan); Shinohara & Matsuura 1997:291 (listed; Suruga Bay); Nakabo 2000:432 (in key; Japan); Shinohara *et al.* 2001:305 (listed; Tosa Bay); Nakabo 2002:432 (in key; Japan); Nakajima 2003:54 (brief description; 1 spec. from Enshu-nada); Senou *et al.* 2006:420 (listed; Sagami Sea).

[?] *Coelorinchus matsubarae* (not Okamura 1982): Shao 1993:170 (compiled; Taiwan).

Coelorinchus (Paramacrurus) kishinouyei [sic]: Chiou *et al.* 2004a: table 1 (listed; Taiwan).

Diagnosis. Light organ moderately long, externally represented by narrow black streak immediately anterior to anus (covered with scales), its anterior margin situated on (or very slightly anterior to) line connecting outer pelvic-fin bases. Underside almost completely naked; dorsal

surface of snout posterior to lateral nasal ridges usually heavily scaled. Snout moderately long, length 50–66% PRL, its dorsal profile concave in lateral view; terminal scute small but broad, dorsoventrally flattened; lateral nasal ridge completely supported by nasal bone, broadly convex in dorsal view. Length of chin barbel 7–13% PRL. Anus separated from anal-fin origin by several scale rows. Premaxillary teeth small, conical, arranged in short, uniformly wide band, none especially enlarged; posterior margin of tooth band falling far short of lateral corner of mouth. Body scales covered with short, greatly reclined, needle-like spinules in tightly packed parallel rows; spinules in each row greatly overlapping, but free from one another; buttresses absent. Body silvery white but darker dorsally; prominent large dark blotch between lateral line and pectoral-fin base; lips and oral cavity whitish; gular and branchiostegal membranes uniformly dull black; anal fin dark especially along anteroventral margin.

Material examined. 24 specimens. **Holotype of *Coelorhynchus kishinouyei*:** USNM 49395 (68.5 mm HL, 350+ mm TL), Misaki, Kanagawa Pref., Sagami Bay, Japan, coll. K. Okita, date unknown. **Non-types: Japan:** BSKU 840 (1, 49.9 mm HL, 256 mm TL), BSKU 842 (1, 43.3 mm HL, 226 mm TL), Mimase fish market, coll. T. Kamohara, 11 Dec. 1950; BSKU 74207 (1, 49.4 mm HL, 246+ mm TL), BSKU 74208 (1, 51.6 mm HL, 281 mm TL), BSKU 74209 (1, 48.7 mm HL, 244+ mm TL), BSKU 74210 (1, 46.8 mm HL, 219+ mm TL), off Numazu, Suruga Bay, bottom trawl, coll. E. Izuka, 16 Jan. 2005; FAKU 28365 (1, 61.3 mm HL, 311+ mm TL), Miya fish market, Enshu-nada, bottom trawl, coll. O. Okamura, 5–7 Feb. 1959; FAKU M7512 (1, 59.2 mm HL, 320 mm TL), Owase fish market, Kumano-nada, bottom trawl, coll. K.

Matsubara, 4–9 Jan. 1936; *BSKU 110139 (1, 38.5 mm HL, 205 mm TL), Suruga Bay, 34.7651°N, 138.4880°E, 200–450 m, F/V *Hinode-maru*, sta. 4, bottom trawl, coll. N. Nakayama and R. Misawa, 23 Apr. 2013; KPM-NI 10133 (1, 57.0 mm HL, 300+ mm TL), southwest of Boso Peninsula, Uraga Channel, Sagami Bay, 120–220 m, gill net, 25 Mar. 1998; FAKU 34242 (1, 55.3 mm HL, 265+ mm TL), FAKU 34243 (1, 59.7 mm HL, 309 mm TL), off Kurihama, ?Tokyo Bay, bottom trawl, coll. O. Okamura, Apr. 1969; KPM-NI 10131 (1, 61.8 mm HL, 330+ mm TL), east of Izu Peninsula, Sagami Bay, 60–80 m, gill net, 30 Mar. 1998. **Taiwan:** ASIZP 65991 (1, 68.5 mm HL, 318+ mm TL), Da-xi fish market, Yilan, bottom trawl, coll. P.-F. Lee, 7 Mar. 2005; BSKU 99867 (1, 57.1 mm HL, 298+ mm TL), BSKU 99869 (1, 50.5 mm HL, 230+ mm TL), Da-xi fish market, Yilan, coll. M.-Y. Lee, 30 Jun. 2009; KAUM-I. 44657 (1, 43.8 mm HL, 194+ mm TL), KAUM-I. 44658 (1, 50.2 mm HL, 200+ mm TL), KAUM-I. 44659 (1, 47.8 mm HL, 206+ mm TL), off Kaohsiung, 400 m, trawl, coll. M. Matsunuma and H.-C. Ho, 1 Jul. 2011; BSKU 98956 (1, 56.2 mm HL, 237+ mm TL), BSKU 98958 (1, 47.6 mm HL, 237 mm TL), BSKU 98959 (1, 44.7 mm HL, 217+ mm TL), BSKU 98960 (1, 51.9 mm HL, 252+ mm TL), Dong-gang fish market, Pingtung, bottom trawl, coll. H.-C. Ho, 16 Nov. 2007.

Counts and measurements. Based on 23 specimens (43.3–68.5 mm HL, 194+–350+ mm TL). Counts: first dorsal-fin rays II, 8–10; pectoral-fin rays i14–i19; pelvic-fin rays 7; gill rakers on first arch (outer/inner) 0/8–10, on second arch 6–8/7–9; longitudinal scales 27–35; transverse scale rows below first dorsal-fin origin 5–7, below first dorsal-fin midbase 3–5.5, below second dorsal-fin origin 4–5.5, above anal-fin origin 14–19.

FIGURE 60. Fresh specimen of *Coelorhynchus kishinouyei*. BSKU 110139, 38.5 mm HL, 205 mm TL, Suruga Bay, Shizuoka Pref., Pacific, 200–450 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

FIGURE 61. Holotype of *Coelorhynchus kishinouyei* (= *Coelorinchus kishinouyei*). USNM 49395, 68.5 mm HL, 350+ mm TL, Sagami Bay, Pacific. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 34–42 (50–66); orbit diameter 32–39 (50–57); postorbital length 28–34 (45–54); postrostral length 62–68; orbit–preopercle distance 36–43 (55–64); suborbital width 13–15 (20–23); upper-jaw length 24–28 (36–44); length of rictus 18–23 (26–34); length of premaxillary tooth band 10–14 (15–22); preoral length 30–37 (44–59); length of terminal snout scute 3–6 (5–10); length of lateral nasal ridge 24–30 (35–49); length of suborbital ridge 94–102 (138–165); snout width 29–34 (43–54); internasal width 18–21 (26–33); interorbital width 19–23 (27–35); occipital width 9–11 (13–17); body width over pectoral-fin bases 40–51 (61–78); body depth at first dorsal-fin origin 60–76 (95–118); body depth at anal-fin origin 50–74 (78–117); prepelvic length 106–122 (156–189); preanus length 125–150 (191–235); preanal length 137–167 (207–262); isthmus–pelvic distance 31–43 (46–64); isthmus–anus distance 47–62 (75–95); isthmus–anal distance 55–84 (88–132); pelvic–anal distance 31–52 (49–81); anus–anal distance 8–20 (13–32); pelvic-fin length 41–50 (61–74); pectoral-fin length 55–72 (87–107); predorsal length 110–122 (166–187); height of first dorsal fin 66–77 (101–117); length of first dorsal-fin base 20–28 (32–41); interdorsal length 29–46 (47–72); length of gill slit 9–13 (15–20); length of posterior nostril 8–12 (12–18); barbel length 5–9 (7–13).

Size. To 36 cm TL (Okamura 1970a).

Distribution. Restricted to Japan and Taiwan.

Known from off the Pacific coasts of southern Japan from Kochi to Chiba Pref., and the East and South China Seas off Taiwan (Appendix 3-3D). The species has a disjunct distribution between central Japan and Taiwan, and is absent from most of the East China Sea. Depth range of Japanese specimens 60–458 m. Common in Suruga Bay and Taiwan, but rare in other parts of its range.

Remarks. *Coelorinchus kishinouyei* was described by Jordan & Snyder (1900) based on a single specimen collected from Misaki, Kanagawa Pref. Japan (Fig. 61; probably originally from Sagami Bay). The reader can refer to the original description (Jordan & Snyder 1900) and Okamura (1970a) for further morphological details. As noted by Iwamoto *et al.* (2015), the species has a disjunct distribution between Japan and Taiwan, and is absent from the Okinawa Trough in the East China Sea. It is abundant in Taiwan and the Pacific off southern Japan east of Cape Shionomisaki (135.75°E), but seemingly rare between these two areas. So far as I know, only three specimens of this species have been collected in Tosa Bay despite thorough collecting effort by previous researchers.

Coelorinchus kishinouyei falls in the *C. argus* group and is most similar to *C. jordani* Smith & Pope, 1906. The two species are fully compared in the Comparisons section of the latter species.

***Coelorinchus lanceolatus* sp. nov.**

[New Japanese name: Yajiri-hige]

(Figs. 62–63; Table 1; Appendix 3–3E)

Diagnosis. Light organ externally represented by short, narrow, naked streak immediately anterior to anus, its anterior margin falling far short of line connecting inner pelvic-fin bases. Underside of head almost completely naked; dorsal surface of snout fully scaled, except narrow naked clefts along each side of median rostral ridge. Nasal fossa heavily scaled anteroventrally. Snout very long, sharply pointed, length 107% PRL; dorsal profile of snout moderately concave in lateral view. Terminal snout scute long, slender, spearhead-shaped, dorsoventrally flattened, length 27% PRL. Scales on median rostral ridge covered with divergent rows of spinules. Lateral nasal ridge completely supported by nasal bone, almost straight when viewed dorsally, giving sharply pointed and distinctly attenuated appearance to snout. Anus slightly separated from anal-fin origin. Premaxillary teeth small, conical, arranged in short, uniformly wide band, none especially enlarged; posterior margin of tooth band falling far short of lateral corner of mouth. Body scales covered with short, reclined, keel-like spinules in narrowly divergent, saw-toothed ridges; every spinule row complete, extending to posterior scale margin; spinules in each row greatly overlapping, closely adjoined to one another, but generally free at tips; buttresses only narrowly developed. Occipital scales covered with short, erect, keel-like spinules in widely divergent, saw-toothed rows. Postorbital length 53% PRL; preoral length 101% PRL; prepelvic length 227% PRL; preanal length 292% PRL; predorsal length 214% PRL; height of first dorsal fin 84% PRL; interdorsal length 34% PRL. Free neuromasts on underside of head tiny, immaculate, difficult to distinguish; dark hair-like papillae absent on underside of head. Body uniformly pale in preserved specimen; lips white; oral cavity blackish.

Material examined. 1 specimen. **Holotype:** NSMT-P 95590 (79.2 mm HL, 255+ mm TL), east of Muko-jima Island, Ogasawara Islands, Shichito-Iojima Ridge, Japan, 521–536 m, R/V *Tansei-maru*, cr. KT-97-07, sta. MK05, 3-m ORE beam trawl, 7 Jun. 1997.

Description. General features are shown in Fig. 62. Counts and measurements are given in Table 1. Body deepest at first dorsal-fin origin, tapering gradually to long laterally compressed tail. Trunk shallow, width over pectoral-fin bases 1.1 in depth below first dorsal-fin origin. Head large, HL about 3.2 in TL. Snout very long, sharply pointed, protruding greatly beyond upper jaw, length 2.3 times as long as orbit diameter; dorsal contour of snout moderately concave in lateral view; lateral nasal ridge completely supported by nasal bone, almost straight in dorsal view, giving sharply pointed and distinctly attenuated appearance to snout. Orbit large, greatest diameter 1.1 in postorbital length. Interorbital space flat,

width 1.2 in orbit diameter. Mouth moderate in size, inferior, protrusible, upper-jaw length 1.2 in orbit diameter; posterior margin of maxilla reaching vertical through hind 2/5 of orbit; lateral corner of mouth moderately restricted by skin folds; lips thin, slightly papillose near tooth bands. Suborbital region divided by longitudinal bony ridge passing from tip of snout to posteroventral angle of opercle; its upper half almost vertical, lower half sharply inclined mesially. Preopercle large, posterior margin inclined at about 60° angle, forming moderately angular lobe at posteroventral corner. Subopercle prolonged ventrally as slender flap; posterior margin deeply concave. Gill membranes broadly connected across, and attached mesially to isthmus, with slight posterior free fold. Outer gill slit moderately restricted by skin folds, length 2.5 in orbit diameter. Gill rakers small, low, tubercular, armed with short, fine spines; no rakers on outer side of first arch and inner side of fourth arch; gill filaments moderately long. Chin barbel slender, length 3.7 in orbit diameter, tapering into filamentous tip.

Anus only slightly separated from anal-fin origin by 1–2 scale rows. Light organ externally represented by short, narrow, naked streak, extending from anus forward to midway between anus and outer pelvic-fin base, its anterior margin falling far short of line connecting inner pelvic-fin bases.

Teeth small, sharp, conical, gently incurved, in narrow bands in both jaws. Premaxillary teeth in short, uniformly wide band, with about 7 tooth rows across widest point near symphysis, posterior margin of tooth band falling far short of lateral corner of mouth; teeth becoming progressively smaller inwardly. Mandibular teeth in broad tapered band, with about 5 tooth rows near symphysis; mandibular band narrowing gradually to about 3 rows posteriorly, its posterior margin extending beyond lateral corner of mouth; all teeth uniformly small.

Body scales large, thin, not deciduous; those on dorsum below interdorsal space covered with short, reclined, keel-like spinules in 6–7 narrowly divergent rows (Fig. 63A–B); middle row longest with about 7 spinules, but not especially high or enlarged compared with adjacent rows; every spinule row complete throughout; spinules forming angle of about 40° to scale surface; spinules greatly overlapping, closely adjoined to one another, but their distal tips often free, forming saw-toothed ridges that give spiny appearance to body surfaces; spinules increasing in height posteriorly, and tip of last spinule in each row extending slightly beyond posterior scale margin; buttresses of spinules narrowly developed; no reticulate structures on unexposed portion. Body fully scaled except for fins and ventral light organ.

Scales on head ridges coarsely modified; those on median rostral ridge covered with radiating rows of spinules. Terminal snout scute very long, slender, attenuated, spearhead-shaped, dorsoventrally flattened, its length 1.7 in orbit diameter. Supraoccipital and posttemporal scutes stout, but not especially enlarged.

Occipital scales covered with short, erect, keel-like spinules in narrowly divergent, saw-toothed rows (Fig. 63C–D); those on dorsal surfaces of snout posterior to leading edges armed with small cluster of short, erect, needle-like spinules. Other scales on head variable in size, generally similar to those on body, but spinules more erect and more or less needle-like in shape; those on opercle, preopercle, and postorbital and supratemporal canals largest. Nasal fossa heavily scaled anteroventrally. Dorsal surfaces of snout almost fully scaled; naked clefts along each side of median rostral ridge narrowly developed. Underside of head completely naked.

No open pores along cephalic sensory canals. Free neuromasts tiny, poorly marked, not discernible. Anterior nostril small slit; posterior large, oval in shape. Grooved lateral line complete, not interrupted throughout.

Origins of first dorsal and pelvic fins on about same vertical; first dorsal fin moderately high, its height 2.9 times as long as its base length; second spinous ray not elongated, smooth along its leading edge; its tip extending posteriorly to base of 5th ray of second dorsal fin when laid back. Interdorsal space 1.1 times as long as first dorsal-fin base length. Origin of second dorsal fin above base of 2nd ray of anal fin. Tips of pectoral and pelvic fins extending slightly beyond vertical through anal-fin origin when laid back; outer pelvic-fin ray slightly prolonged, filamentous at tip.

Color when fresh. Unknown.

Color in alcohol (Fig. 62). Head and body uniformly light brown; abdomen darker; lips white; oral cavity faded, but probably blackish originally; gill cavity blackish; gill rakers and arches pale, filaments light tan; fins uniformly pale, but second spinous ray of first dorsal fin slightly darker distally.

Size. To at least 26 cm TL.

Distribution. So far known from off the Ogasawara Islands, Shichito-Iojima Ridge, at a depth of 521–536 m (Appendix 3-3E). Very rare.

Etymology. The specific epithet is derived from the Latin adjective *lanceolatus*, meaning “armed with a lance”, in reference to the species’ long snout tipped with a spearhead-shaped scute. The Japanese name, Yajiri-hige is derived from a combination of the Japanese nouns “yajiri”, meaning a spearhead, and “hige”, meaning grenadiers.

Remarks, relationships, and comparisons. *Coelorinchus lanceolatus* **sp. nov.** is represented by only a single specimen collected from the Shichito-Iojima Ridge, Japan. It belongs to the *C. anatirostris* group in having the following combination of features (see the Relationships of *C. anatirostris*): light organ externally represented by short naked streak anterior to anus, its anterior margin not reaching line connecting inner pelvic-fin bases; lateral nasal ridge completely supported by nasal bone; premaxillary teeth in short, uniformly wide band; underside of head completely naked; body scales covered with keel-like spinules in narrowly divergent rows, and every row complete throughout; buttresses of scale spinules narrowly developed; no prominent markings on body. *Coelorinchus lanceolatus* is similar to *C. anatirostris* Jordan & Gilbert in Jordan & Starks, 1904 in many aspects, and the holotype of the former was initially considered to represent an unusual variant of the latter species. Actually, the two species share the most useful diagnostic features distinguishing *C. anatirostris* from its congeners, and no significant differences are found in meristic characters. However, *C. lanceolatus* differs notably from *C. anatirostris* in having a distinctly

FIGURE 62. Holotype of *Coelorinchus lanceolatus* **sp. nov.** NSMT-P 95590, 79.2 mm HL, 255+ mm TL, off Ogasawara Islands, Shichito-Iojima Ridge, Pacific, 521–536 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

FIGURE 63. Scanning electron micrographs showing scales of *Coelorinchus lanceolatus* sp. nov. NSMT-P 95590, holotype, 79.2 mm HL. (A–B) Scales from the dorsum below the interdorsal space; (C–D) from the top of the head between the parietal ridges. (A, C) Views from above; (B, D) oblique views. [Photos: N. Nakayama]

longer snout [107% PRL (Fig. 62) vs. 60–88% (Figs. 18–20, 24)] with a well-developed terminal scute (27% PRL vs. 7–18%). The difference in the snout length is too great to ignore, even considering the variation of *C. anatirostris* in this study, which causes other pronounced differences in measurements, including the preoral length (101% PRL in *C. lanceolatus* vs. 50–81% in *C. anatirostris*), prepelvic length (227% PRL vs. 184–214%), preanal length (292% PRL vs. 239–380%), and predorsal length (214% PRL vs. 177–205%). In addition, *C. lanceolatus* has a relatively lower first dorsal fin than *C. anatirostris* (84% PRL vs. 85–107%). Despite limited study material, there is little doubt that the specimen represents a distinct species previously unknown to science.

The new species is also similar in many aspects with *C. weberi* Gilbert & Hubbs, 1916 known only from the holotype collected from the Philippines. However, *C. lanceolatus* is readily distinguished from the latter species in that free neuromasts on the underside of the head are unpigmented and difficult to distinguish without

a microscope (vs. each neuromast marked with two small black dots). It further differs notably from *C. weberi* by having a longer snout (107% PRL vs. 80%), shorter postorbital (53% PRL vs. 58%), and lower first dorsal fin (84% PRL vs. 99%). A pronounced difference between the two species was also found in the snout shape. In *C. lanceolatus*, the anterolateral margins of the snout are almost straight when viewed dorsally, giving a sharply pointed and distinctly attenuated appearance to the snout, whereas those of *C. weberi* are broadly convex so as to give a spade-shaped appearance to the snout. The latter species will be fully redescribed in a forthcoming paper by the present author.

***Coelorinchus leptorhinus* Chiou, Shao & Iwamoto, 2004**

[No Japanese name]

(Figs. 53C–D, 64–65; Appendix 3-3F)

Caelorinchus tokiensis (not Steindachner & Döderlein 1887): Shao 1993:170 (compiled; Taiwan) [fide Chiou *et al.* 2004a: table 1].
Caelorinchus (Oxymacrus) leptorhinus: Chiou *et al.* 2004a: table 1 (listed; Taiwan).
Caelorinchus leptorhinus Chiou, Shao & Iwamoto, 2004b:299, figs. 1–3 [original description; holotype: ASIZP 61344, from Ta-shi (= Da-xi, northeastern Taiwan); 23 paratypes from northeast of Taiwan].
Caelorinchus leptorhinus: Shao *et al.* 2008b: table 2 (23 spec. listed from northeastern and southwestern Taiwan); Ho & Shao 2011:36 (type catalog); Iwamoto *et al.* 2015:59 (brief description; 54 spec. from northeastern and southwestern Taiwan).

Diagnosis. Light organ externally represented by short, narrow, naked streak immediately anterior to anus, its anterior margin falling far short of line connecting inner pelvic-fin bases. Snout completely scaled dorsally, mostly naked ventrally; underside of head posterior to hind margins of lateral nasal ridges almost completely scaled. Snout very long, sharply pointed, length 69–99% PRL; dorsal profile of snout concave in lateral view; terminal scute short, diamond-shaped, dorsoventrally flattened; lateral nasal ridge completely supported by nasal bone. Anus only slightly separated from anal-fin origin. Premaxillary teeth small, conical, arranged in moderately wide tapered band, with outer series distinctly enlarged; posterior margin of tooth band extending beyond midway of rictus. Body scales covered with short, reclined, keel-like spinules in divergent saw-toothed ridges; every spinule row complete, extending to posterior scale margin; spinules in each row greatly overlapping, increasing in height posteriorly; buttresses barely developed. Occipital scales covered with keel-like spinules in narrowly divergent rows. Orbit diameter 36–41% PRL; suborbital width 14–18% PRL; internasal width 19–24% PRL; interorbital width 27–31% PRL; pectoral-fin length 72–97% PRL; transverse scale rows below first dorsal-fin midbase 5.5–6.5. Head and body uniformly dark brown, but lips pale to dusky; all fins blackish.

Material examined. 11 specimens. **Holotype of *Caelorinchus leptorhinus*:** ASIZP 61344 (234 mm HL, 811+ mm TL), Da-xi fish market, Yilan, northeastern Taiwan, 24.9105°N, 122.0636°E, 400–600 m, beam trawl, coll. M.-L. Qiu, 15 Mar. 1999. **Paratypes of *C. leptorhinus*:** ASIZP 61345 (1, 119 mm HL, 388+ mm TL), Nan-fang-ao fish market, Yilan, 15 Dec. 1999; ASIZP 61347 (3, 171–176 mm HL, 507+–645+ mm TL), Da-xi fish market, Yilan, 24.9105°N, 122.0636°E, bottom trawl, coll. M.-L. Qiu, 30 May 2001. **Non-types: Taiwan:** *BSKU 99870 (1, 96.5 mm HL, 367+ mm TL), Da-xi fish market, Yilan, coll. M.-Y. Lee, 30 Jun. 2009; FAKU 204420 (1, 131 mm HL, 470 mm TL), FAKU 204421 (1, 107 mm HL, 411 mm TL), FAKU 204422 (1, 112 mm HL, 424+ mm TL), FAKU 204423 (1, 96.5 mm HL, 302+ mm TL), FAKU 204424 (1, 80.6 mm HL, 245+ mm TL), Dong-gang fish market, Pingtung, bottom trawl, coll. N. Nakayama, 27 Feb. 2017.

Counts and measurements. Based on 10 specimens (80.6–234 mm HL, 245+–811+ mm TL). Counts: first dorsal-fin rays II,8–9; pectoral-fin rays i16–i19; pelvic-fin rays 7; gill rakers on first arch (outer/inner) 0/8–9, on second arch 6–7/8–9; longitudinal scales 43–51; transverse scale rows below first dorsal-fin origin 5.5–8, below first dorsal-fin midbase 5.5–6.5, below second dorsal-fin origin 6–8, above anal-fin origin 21–22.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 41–50 (69–99); orbit diameter 20–21 (36–41); postorbital length 30–38 (59–65); postrostral length 50–59; orbit–preopercle distance 31–39 (61–68); suborbital width 7–11 (14–18); upper-jaw length 24–31 (47–53); length of rictus 18–26 (35–43); length of premaxillary tooth band 15–20 (28–34); preoral length 27–40 (45–79); length of terminal snout scute 6–9 (11–19); length of lateral nasal ridge 21–34 (35–67); length of suborbital ridge 88–95 (149–189); snout width 18–21 (30–40); internasal width 11–14 (19–24); interorbital width 14–18 (27–31); occipital width 6–7 (11–13); body width over pectoral-fin bases 26–39 (51–66); body depth at first dorsal-fin origin 33–48 (66–80); body depth at anal-fin origin 29–37 (58–65); prepelvic length 103–115 (175–212); preanus length 124–139 (220–256); preanal length 133–147 (234–269); isthmus–pelvic distance 27–35 (50–62); isthmus–anus distance 46–56 (88–106); isthmus–anal distance 55–65 (105–116); pelvic–anal distance 29–40 (56–67); anus–anal distance 8–11 (14–19); pelvic-fin length 27–36 (51–65); pectoral-fin length 37–57 (72–97); predorsal length 86–110 (146–210); height of first dorsal fin 35–37 (60–71); length of first dorsal-fin base 13–17 (25–29); interdorsal length 24–37 (44–63); length of gill slit 9–14 (18–24); length of posterior nostril 7–8 (12–14); barbel length 7–14 (15–24).

Size. Attains at least 81 cm TL (ASIZP 61244, holotype, 811+ mm TL).

Distribution. So far known only from the East and South China Seas off Taiwan at depths of about 100–650 m (Shao *et al.* 2008a, 2008b; Appendix 3-3F). Common.

Comments on type specimens. *Coelorinchus leptorhinus* was originally described from northeastern Taiwan based on 23 specimens collected at the Da-xi fish market, Yilan (Chiou *et al.* 2004b; Fig. 65). Ten paratypes registered in the Academia Sinica, Taipei (ASIZP 61346) were not found among the collection during the author's visit in 2012 and 2017.

Remarks, relationships, and comparisons. Morphology of the species was detailed in the original description given by Chiou *et al.* (2004b). No specimens of *C. leptorhinus* have been collected within Japan's EEZ, although it is commonly found in the Da-xi fish market in northern Taiwan. The species belongs to the *C. japonicus* groups (*sensu stricto*; see the Relationships of *C. japonicus*), and is unlikely to be confused with Japanese congeners in that the underside of the snout is broadly naked (*vs.*

FIGURE 64. Fresh specimen of *Coelorinchus leptorhinus*. FAKU 204420, 130 mm HL, 570 mm TL, Dong-gang fish market, southwestern Taiwan, South China Sea. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

FIGURE 65. Holotype of *Coelorinchus leptorhinus* (= *Coelorinchus leptorhinus*). ASIZP 61344, 234 mm HL, 811+ mm TL, Dong-gang fish market, southern Taiwan, South China Sea. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

completely and uniformly scaled, except for narrow naked areas above upper lip and along ventromedial line). It further differs from Japanese species of the *C. japonicus* group in having a relatively smaller orbit [36–41% PRL vs. 39–52%, 45–52%, and 43–52% in *C. japonicus* (Temminck & Schlegel, 1846), *C. nox* **sp. nov.**, and *C. smithi* Gilbert & Hubbs, 1920 respectively], longer pectoral fin (72–97% PRL vs. 55–71%, 57–69%, and 60–76%), narrower suborbital (14–18% PRL vs. 18–23%, 18–23%, and 19–23%), internasal (19–24% PRL vs. 26–33%, 27–31%, and 28–37%), and interorbital regions (27–31% PRL vs. 31–38%, 39–41%, and 33–42%). This species is also distinguished from *C. nox* and *C. smithi* in having more scales below the first dorsal-fin midbase (5.5–6.5 vs. 3.5–4.5 and 3.5–5.5 in *C. nox* and *C. smithi* respectively).

***Coelorinchus longissimus* Matsubara, 1943**

[Japanese name: Tongari-hige]

(Figs. 34C, 66; Appendix 3-4A)

Coelorhynchus longissimus Matsubara, 1943:140, figs. 1C, 2C, 5 (original description; holotype: FAKU 1592, from Kumano-nada; new Japanese name: “Tongari-hige”); Matsubara 1955:1312, pl. 134, fig. 457 (in key; Japan); Kamohara 1958:73 (listed; Kochi Pref.); Chyung & Kim 1959:8, pl. II, figs. 16–17 (first record from Korea; brief description; 1 spec. from Pusan, Korea); Okada *et al.* 1959:83 (listed; Kumano-nada); Kamohara 1964:95 (listed; Kochi Pref.); Matsubara 1965:506 (compiled; Japan); Okamura 1970a:165, pl. XXXV, text-fig. 67 (description; biological notes; 48 spec. from Pacific off southern Japan from Suruga Bay to Totoro); Kuroda 1971:56 [listed from Suruga Bay after Okamura (1970a)]; Kataoka & Tomida 1981:78 (listed; Mie Pref.); Tominaga & Uyeno 1981:488 (listed; Japan); Ohta 1983: table A (listed; Suruga

Bay); Ozawa 1983:13 (listed; off Makurazaki, Kagoshima Pref., East China Sea).

Coelorhynchus (Quincuncia) longissimus: Okamura 1970b: table 1 (listed; Japan).

Coelorinchus longissimus: Yatou 1984:223, 366, fig. 159 [brief description; 9 spec. from Okinawa Trough; photo based on BSKU 39080 (from Mimase fish market); in part, BSKU 26581 is *C. formosanus*]; Okamura 1984b:96, pl. 82, fig. 1 (compiled); Okamura 1988:96, pl. 82, fig. 1 (compiled); Okamura 1997:126, fig. 9 (compiled); Youn 2002:538 (listed; Korea); Shao *et al.* 2008b: table 2 (3 spec. listed from northeastern and southwestern Taiwan); Nakabo & Kai 2013:506 (in key; Japan); Ikeda & Nakabo 2015:321, pl. 67, figs. 6–10 (brief description; spec. from Wakayama Pref.); Iwamoto *et al.* 2015:60 (brief description; 9 spec. from northeastern and southwestern Taiwan, and South China Sea); Motomura 2020:38 (listed; Japan).

Coelorinchus longissimus: Nakabo 1993:366 (in key; Japan); Shinohara & Matsuura 1997:291 (listed; Suruga Bay); Nakabo 2000:430 (in key; Japan); Shinohara *et al.* 2001:304 (listed; Tosa Bay); Nakabo 2002:430 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Chiou *et al.* 2004a:43, fig. 10 (brief description; 2 spec. from Daxi and Dong-gang; first record from Taiwan); Shinohara *et al.* 2005:416 (listed; Ryukyu Islands); Kim *et al.* 2005:172, fig. 293 (compiled; Korea; photo spec. represents *C. multispinulosus*); Kim *et al.* 2009a:13 (questionable; listed; Jeju Island, Korea); Furuhashi *et al.* 2010: table 2 (45 spec. listed from northern Okinawa Trough); Iwatsuki *et al.* 2017:32 (listed; Hyuga-nada).

Coelorinchus (Quincuncia) longissimus: Chiou *et al.* 2004a: table 1 (listed; Taiwan).

Diagnosis. Light organ tubular, completely covered with scales, externally represented by long black streak extending from immediately anterior to anus to chest just posterior to isthmus. Underside of head completely naked except for overlapping scales posterior to lateral nasal ridges (small scaly patches occasionally found on lower preopercles and posterior ends of mandibular rami); dorsal surface of snout broadly naked along each side of median rostral ridge. Snout long, sharply pointed, length 77–90% PRL, its dorsal profile concave in lateral view; terminal scute short, arrowhead-shaped, dorsoventrally flattened, length 12–22% PRL. Lateral nasal ridge incompletely supported by nasal bone. Anus abutting anal-fin origin. Premaxillary teeth small, slender, conical in moderately wide, long, tapered band, with outer series slightly enlarged; posterior margin of tooth band almost reaching lateral corner of mouth. Body scales covered with short, erect, needle-like spinules in subparallel to narrowly divergent rows; buttresses barely developed. Second dorsal fin distinctly lower than anal fin. Upper-jaw length 42–49% PRL; preoral length 66–81% PRL; pectoral-fin length 55–64% PRL; height of first dorsal fin 58–70% PRL; interdorsal length 27–47% PRL; length of

gill slit 20–25% PRL; barbel length 11–18% PRL. Body darker dorsally, silvery white ventrally; dark prominent blotches faint, often difficult to distinguish (prominently marked only in juveniles); median nasal bone blackish; underside of head generally pale, but mandibular rami blackish; no black hair-like papillae on naked areas on head; lips pale to dusky; oral cavity immaculate; gular and branchiostegal membranes heavily peppered with large melanophores; chest pale to dark; first dorsal fin dusky, but second spinous ray blackish; tip of pelvic fin prominently blackish.

Material examined. 29 specimens. **Japan:** BSKU 19077 (1, 73.0 mm HL, 256 mm TL), off Satsuma Peninsula, Okinawa Trough, 31.2267°N, 129.9750°E, 310 m, FRV *Soyo-maru*, sta. 72d, beam trawl, 12 Feb. 1959; BSKU 106767 (1, 68.2 mm HL, 240+ mm TL), southeast of Shimokoshiki-jima Island, East China Sea, 31.5658°N, 129.8915°E, 380 m, F/V *Maruko-maru*, tr. 1, bottom trawl, coll. N. Nakayama *et al.*, 24 Apr. 2012; BSKU 109052 (1, 48.7 mm HL, 185 mm TL), east of Kasayama Bank, 32.3884°N, 129.0505°E, 304–312 m, T/V *Nagasaki-maru*, cr. N365, sta. A3, 3-m ORE beam trawl, coll. N. Nakayama, 19 Nov. 2012; BSKU 113743 (1, 86.1 mm HL, 313+ mm TL), off Katsurahama, Tosa Bay, 230 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 9 Apr. 2014; BSKU 99262 (1, 88.4 mm HL, 314+ mm TL), Mimase fish market, bottom trawl, 1 Mar. 2006; BSKU 40440 (1, 91.4 mm HL, 327+ mm TL), BSKU 40441 (1, 92.4 mm HL, 345+ mm TL), BSKU 40442 (1, 92.9 mm HL, 339 mm TL), BSKU 40445 (1, 96.2 mm HL, 342+ mm TL), BSKU 40447 (1, 92.6 mm HL, 341+ mm TL), BSKU 40448 (1, 95.9 mm HL, 277+ mm TL), BSKU 40451 (1, 96.6 mm HL, 338+ mm TL), BSKU 40457 (1, 81.0 mm HL, 304 mm TL), Mimase fish market, bottom trawl, 8 Mar. 1985; BSKU 99346 (1, 93.0 mm HL, 314+ mm TL), Mimase fish market, bottom trawl, coll. N. Nakayama, 23 Apr. 2008; BSKU 99235 (1, 81.5 mm HL, 303+ mm TL), BSKU 99237 (1, 58.9 mm HL, 219+ mm TL), Mimase fish market, bottom trawl, coll. N. Nakayama, 31 Oct. 2007; BSKU 55 (1, 76.4 mm HL, 302 mm TL), Mimase fish market, bottom trawl, coll. T. Kamohara, 5 Mar. 1951; BSKU 100497 (1, 78.7 mm HL, 287+ mm TL), Mimase fish market, F/V *Kosei-maru*, bottom trawl, 15 Oct. 2009; BSKU 108871 (1, 77.5 mm HL, 278+ mm TL), Mimase fish market, F/V *Kosei-maru*, bottom trawl, 22 Oct. 2012; BSKU 94450 (1, 71.6 mm HL, 251+ mm TL), Mimase fish market, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama, 14 Apr. 2008; BSKU 93088 (1, 63.4 mm HL, 237 mm TL), BSKU 93090 (1, 89.9 mm HL, 321+ mm TL), Mimase fish market, F/V *Seiryō-maru*, bottom trawl, coll. N. Nakayama, 24 Dec. 2007; BSKU 94387 (1, 51.5 mm HL, 203+ mm TL), Mimase fish market, F/V *Seiryō-maru*, bottom trawl, coll. N. Nakayama, 4 Apr. 2008; BSKU 69291 (1, 56.1 mm HL, 198+ mm TL), BSKU 69295 (1, 55.7 mm HL, 187+ mm TL), Tosa Bay, 33.1450°N, 133.5150°E, 240–

249 m, FRV *Kotaka-maru*, otter trawl, coll. H. Endo, 20 Aug. 1997; BSKU 112928 (1, 98.7 mm HL, 342+ mm TL), off Okitsu, Tosa Bay, 280–320 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 24 Jan. 2014; BSKU 110426 (1, 29.8 mm HL, 113+ mm TL), Suruga Bay, 34.7651°N, 138.4880°E, 200–450 m, F/V *Hinode-maru*, sta. 4, bottom trawl, coll. N. Nakayama and R. Misawa, 23 Apr. 2013; BSKU 110081 (1, 40.7 mm HL, 154 mm TL), BSKU 110082 (1, 38.3 mm HL, 132+ mm TL), Suruga Bay, 34.7489°N, 138.4664°E, 200–450 m, F/V *Hinode-maru*, sta. 6, bottom trawl, coll. N. Nakayama and R. Misawa, 23 Apr. 2013.

Counts and measurements. Based on 29 specimens (29.8–98.7 mm HL, 113+–345+ mm TL). Counts: first dorsal-fin rays II,9–10; pectoral-fin rays i15–i18; pelvic-fin rays 7; gill rakers on first arch (outer/inner) 0/7–8, on second arch 5–6/7–9; longitudinal scales 36–45; transverse scale rows below first dorsal-fin origin 4.5–7, below first dorsal-fin midbase 3.5–5.5, below second dorsal-fin origin 3.5–5.5, above anal-fin origin 12–18.5.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 44–48 (77–90); orbit diameter 23–29 (43–51); postorbital length 28–31 (51–58); postrostral length 53–57; orbit–preopercle distance 31–37 (57–68); suborbital width 10–12 (19–22); upper-jaw length 23–27 (42–49); length of rictus 18–21 (32–38); length of premaxillary tooth band 15–18 (26–33); preoral length 38–45 (66–81); length of terminal snout scute 7–12 (12–22); length of lateral nasal ridge 30–35 (54–66); length of suborbital ridge 96–101 (168–185); snout width 22–27 (39–48); internasal width 17–22 (32–41); interorbital width 21–24 (38–43); occipital width 6–8 (11–14); body width over pectoral-fin bases 28–44 (50–80); body depth at first dorsal-fin origin 37–54 (70–99); body depth at anal-fin origin 32–45 (56–82); prepelvic length 101–115 (183–207); preanus length 130–152 (234–275); preanal length 133–155 (236–281); isthmus–pelvic distance 27–36 (50–65); isthmus–anus distance 55–71 (99–128); isthmus–anal distance 56–79 (103–143); pelvic–anal distance 31–47 (56–88); anus–anal distance 1–4 (2–8); pelvic-fin length 20–32 (36–59); pectoral-fin length 30–34 (55–64); predorsal length 103–109 (184–200); height of first dorsal fin 32–37 (58–70); length of first dorsal-fin base 15–19 (28–35); interdorsal length 15–26 (27–47); length of gill slit 11–14 (20–25); length of posterior nostril 6–9 (10–16); barbel length 6–10 (11–18).

Size. To about 36 cm TL (Okamura 1970a).

Distribution. Known from Japan and Taiwan. Distributed in the Pacific off southern Japan northward to the Boso Peninsula (139.89°E), Okinawa Trough, and northern South China Sea, at depths of 170–450 m (Chiou *et al.* 2004b; Shao *et al.* 2008a, 2008b; this study; Appendix 3–4A). Common.

Comments on holotype. *Coelorinchus longissimus* was described by Matsubara (1943) based on a single

specimen collected from the Kumano-nada, Japan. Unfortunately, the holotype is considered lost during WWII (see the Comments on holotype of *C. hige*).

Remarks, relationships, and comparisons. *Coelorinchus longissimus* has been well described by previous authors, and thus does not need repeating here. The reader can refer to the original description (Matsubara 1943) and Okamura (1970a) for further morphological details. This species belongs to the *C. argentatus* group (see the Relationships of *C. formosanus*), and is most similar to *C. gaesorhynchus* Iwamoto & Williams, 1999 known from the eastern Indian Ocean off Western Australia. The two species are similar to each other in having the absence of prominent markings on the body, such as dark spots, blotches, stripes, or vermiculations (except for small specimens of *C. longissimus*). However, *C. longissimus* is readily distinguished from *C. gaesorhynchus* by having a shorter snout (77–90% PRL vs. 93–100%), preoral (66–81% PRL vs. 85–92%), and terminal snout scute (12–22% PRL vs. 25–26%).

If some scales remain on the body dorsum, the species is unlikely to be confused with other Japanese species of the *C. argentatus* group, viz., *C. formosanus* Okamura, 1963, *C. kamoharai* Matsubara, 1943, and *C. multispinulosus* Katayama, 1942. In *C. longissimus*, spinules on the body scales are arranged in subparallel to narrowly divergent rows (Fig. 34C), whereas those of the other three species are in quincunx order (Fig. 34A–B, D). In addition, the blotched pattern of *C. longissimus* gradually becomes obscure with growth, but it is still discernable or prominently marked in adults of the other species (especially in preserved specimens). *Coelorinchus longissimus* is also distinguished from *C. formosanus* and *C. multispinulosus* in having a lower first dorsal fin (58–70% PRL vs. ≥74%). It further differs from *C. formosanus* in lacking broad scaly patches on the ventral surfaces of the preopercle and the mandibular rami (vs. well developed), and in having a shorter barbel (11–18% PRL vs. 20–26%) and pectoral fin (55–64% PRL vs. 69–75%). Due to their close resemblance, *C. longissimus* may be confused with *C. kamoharai* in the field. However, *C. longissimus* lacks dark hair-like papillae on the underside of the head (vs. well developed in *C. kamoharai*). It is further separable from *C. kamoharai* in having a longer snout (snout length 77–90% PRL vs. 63–77%; preoral length 66–81% PRL vs. 51–65%), a shorter upper jaw (42–49% PRL vs. 47–54%), and a narrower gill slit (20–25% PRL vs. 26–29%).

Coelorinchus macrochir (Günther, 1877)

[Japanese name: Tenaga-dara]

(Figs. 67–69; Appendix 3–4B)

Macrurus macrochir Günther, 1877:438 [original description; holotype: BMNH 1887.12.7.123, from “off Inoshima” (= Eno-shima Island), *Challenger* sta. 232, at a depth of 345

FIGURE 66. Fresh specimens of *Coelorinchus longissimus* collected from the Mimase fish market, Kochi Pref., Pacific. (A) BSKU 94450, 71.6 mm HL, 251+ mm TL; (B–C) BSKU 100497, 78.7 mm HL, 287+ mm TL. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: (A) N. Nakayama; (B–C) S. Yamamoto]

ftm (631 m)]; Günther 1887:48, pl. XXIX, fig. B (further description of holotype).

Abyssicola macrochir: Jordan & Snyder 1900 (spec. from off Tokyo); Jordan & Snyder 1901:119 (listed; Japan); Jordan & Gilbert in Jordan & Starks 1904:607 (3 spec. from Sagami Bay); Franz 1910:26 (1 spec. from Aburatsubo); Jordan *et al.* 1913:414 (listed; Japan; new Japanese name: “Tenagadara”); Gilbert & Hubbs 1916:183 (description; 36 spec. from East China Sea, Hyuga-nada, and Pacific off Tohoku); Schmidt 1931:154 (brief description; 2 spec. from Misaki and Tokyo); Okada & Matsubara 1938:450 (in key; Japan); Kamohara 1938:71 (brief description; 1 spec. from Kochi Pref.); Kamohara 1950:279 (listed; Kochi Pref.); Kuroda 1951:392 (listed; Suruga Bay); Kamohara 1952:98 (3 spec. from Kochi Pref.); Kuroda 1952:177 (brief description; 3 spec. from Suruga Bay); Matsubara 1955:1314 (in key; Japan); Kamohara 1958:73 (listed; Kochi Pref.); Okada *et al.* 1959:83 (listed; Kumano-nada); Kuroda 1962:8, fig. 2 (description of fresh color; 1 spec. from Suruga Bay); Kamohara 1964:96 (listed; Kochi Pref.); Matsubara 1965:507 (compiled; Japan); Okamura 1970a:145, pl. XXXII, text-fig. 58 (description; biological notes; 136 spec. from Pacific off Japan from Hokkaido to Mimase); Okamura 1970b: table 1 (listed; Japan); Ueno 1971:101 (listed; Hokkaido); Kataoka & Tomida 1981:78 (listed; Mie Pref.); Tominaga & Uyeno 1981:488 (listed; Japan); Shiogaki 1982:34 (listed; Aomori Pref.); Okamura 1984a:221, 364, fig. 156 [brief description; 1 spec. from Okinawa Trough; photo probably based on BSKU 22773 (from Tosa Bay)]; Okamura 1984b:95, pl. 82, fig. G (compiled); Okamura 1988:95, pl. 82, fig. G (compiled); Maeda & Maruyama 1991:366 (listed with a question mark; Hokkaido); Nakabo 1993:365 (in key; Japan); Shinohara & Matsuura 1997:290 (listed; Suruga Bay); Goto 2000:6 (3 spec. from Iwate Pref.); Nakabo 2000:429 (in key; Japan); Zama 2001:40 (listed; Miyagi Pref.); Nakabo 2002:429 (in

key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Maeda & Tsutsui 2003:488 (listed; Hokkaido); Shiogaki *et al.* 2004:55 (listed; Pacific off Aomori Pref.); Shinohara *et al.* 2005:416 (listed; Ryukyu Islands); Suetsugu & Ohta 2005: table 2 (listed; southeast of Boso Peninsula); Senou *et al.* 2006:420 (listed; Sagami Sea); Honma *et al.* 2008:68, fig. 4 (brief description; 1 spec. from Niigata); Kitagawa *et al.* 2008:41, unnumbered fig. (brief description; spec. from Pacific off Tohoku); Shinohara *et al.* 2009:707 (8 spec. listed from Pacific off Tohoku); Amaoka *et al.* 2011:131, fig. 167 (compiled; Hokkaido); Nakabo & Kai 2013:505 (in key; Japan); Kawano *et al.* 2014: appendix table (listed; Sea of Japan); Shinohara *et al.* 2014:241 (listed; Sea of Japan); Amaoka *et al.* 2020:165, fig. 224 (listed; Hokkaido).

[?] *Coelorhynchus kishinouyei* (not Jordan & Snyder 1900): Shiogaki 1982:33 (listed; Aomori Pref.; “a large species attaining 1 m TL”).

[?] *Coelorinchus kishinouyei* (not Jordan & Snyder 1900): Shiogaki 1988: table 1 [listed; Pacific off Aomori Pref.; obviously after Shiogaki (1982)]; Shiogaki *et al.* 1992:29 [listed after Shiogaki (1982); Aomori Pref.].

Coelorinchus macrochir: Iwamoto 1990:171, fig. 392 (synopsis); Balanov 2003:483 (first record from Okhotsk Sea; description; 1 spec.); Shao *et al.* 2008b: table 2 (2 spec. listed from northeastern Taiwan; first record from Taiwan); Kim *et al.* 2009b:110, fig. 5 (brief description; 7 spec. from Korea and Japan; new Korean record); Iwamoto *et al.* 2015:61, fig. 7 (brief description; 2 spec. from northeastern Taiwan); Motomura 2020:38 (listed; Japan).

Coelorinchus macrochir: Shinohara *et al.* 1996:169 (2 spec. listed from Pacific off Tohoku); Shinohara *et al.* 2001:305 (listed; Tosa Bay).

[?] *Coelorinchus kishinouyei* (not Jordan & Snyder 1900): Shiogaki *et al.* 2004:55 [listed after Shiogaki (1982); Pacific off Aomori Pref.]; Shinohara *et al.* 2009:708 [listed after Shiogaki (2004);

Pacific off Tohoku].

Caelorinchus sp.: Matsuura *et al.* 2009:24 (1 spec. from Fukushima Pref.; NSMT-P 73686).

Coryphaenoides sp.: Matsuura *et al.* 2009:24 (1 spec. from Miyagi Pref.; NSMT-P 74215).

Diagnosis. Anus situated about midway between inner pelvic-fin bases and anal-fin origin. Light organ small, located immediately anterior to anus. Head fully scaled. Snout moderately long, deep, bluntly pointed, length 40–52% PRL; dorsal profile of snout convex in lateral view; tip of snout situated above horizontal through midorbit; terminal scute small, tubercular; suborbital ridge distinctly curved anteriorly; lateral nasal ridge incompletely supported by nasal bone. Premaxillary teeth small, conical, arranged in narrow band (about 3 rows), with outer series enlarged; posterior margin of tooth band extending beyond midway of rictus; mandibular teeth in 2 irregular rows, with inner series distinctly enlarged. Body scales covered with short, moderately erect, knife-like to keel-like spinules in widely divergent rows; spinules in each row overlapping, but free from one another; buttresses narrowly developed. Occipital scales covered with short, erect, needle-like spinules in widely divergent, comb-like rows. Tip of pectoral fin extending well beyond vertical through anal-fin origin; pectoral-fin length 88–104% PRL. Body darker dorsally, paler ventrally (lateral surfaces of head and body broadly silvery-white in small specimens); no prominent markings in adult (several dark bars and spots present in juveniles); fins blackish.

Material examined. 22 specimens. **Holotype of *Macrurus macrochir*:** BMNH 1887.12.7.123 (142 mm HL, 657+ mm TL), off Eno-shima Island, Kanagawa Pref., Sagami Bay, Japan, 35.1833°N, 139.4667°E, 345 fm (631 m), *Challenger* sta. 232, trawl and dredge, 12 Mar. 1875. **Non-types: Japan:** BSKU 22773 (1, 97.7 mm HL, 512 mm TL), Tosa Bay, 32.9467°N, 133.5567°E, 714–735 m, R/V *Hakuho-maru*, cr. KH-74-3, sta. B5-1, beam trawl, 24 Jul. 1974; *BSKU 110140 (1, 30.5 mm HL, 100+ mm TL), Suruga Bay, 34.7651°N, 138.4880°E, 200–450 m, F/V *Hinode-maru*, sta. 4, bottom trawl, coll. N. Nakayama and R. Misawa, 23 Apr. 2013; KPM-NI 34639 (1, 139 mm HL, 689+ mm TL), Suruga Bay off Izu Peninsula, 710 m, coll. H. Masuda, 24 Aug. 1995; KPM-NI 34659 (1, 141 mm HL, 707+ mm TL), east of Futo, Sagami Bay, 700 m, F/V *Yusaku-maru*, coll. H. Masuda, 11 Feb. 1997; *BSKU 19160 (1, 59.0 mm HL, 300+ mm TL), off Mito, 36.2717°N, 140.9467°E, 355–370 m, FRV *Soyo-maru*, sta. 18, beam trawl, 25 Jul. 1960; NSMT-P 78249 (5, ex. SHMNH-P 19325, 62.6–75.6 mm HL, 248+–345+ mm TL, counts and measurements taken on 2 of 5), off Arahama, coll. M. Atsumi, Apr. 1933; NSMT-P 73686 (1, 60.7 mm HL, 295+ mm TL), off Onahama, coll. H. Kakuda (or H. Sumida), Mar. 1932; NSMT-P 74215 (1, 65.7 mm HL, 324+ mm TL), off Natori, coll. T. Kikuta, 15 May 1932; NSMT-P 73515 (5, ex. SHMNH-P 25782,

51.8–57.8 mm HL, 248+–294+ mm TL, counts and measurements taken on 3 of 5), southeast off Onahama, coll. H. Sumida, 3 Jan. 1932; NSMT-P 77837 (1, 120 mm HL, 588+ mm TL), off Hitachi, 36.5300°N, 141.1450°E, 557–561 m, FRV *Wakataka-maru*, sta. H-550, bottom trawl, coll. H. Saito *et al.*, 14 Nov. 2006; NSMT-P 77846 (1, 110 mm HL, 614+ mm TL), east of Oshika Peninsula, 38.4033°N, 142.0500°E, 423 m, FRV *Wakataka-maru*, sta. E-425, bottom trawl, coll. G. Shinohara *et al.*, 4 Nov. 2006; HUMZ 206465 (1, 128 mm HL, 643 mm TL), south of Tomakomai, Hokkaido, 42.1670°N, 141.6503°E, 725 m, T/V *Oshoro-maru*, bottom trawl, coll. T. Tada and M. Ono, 29 Sept. 2009; HUMZ 206304 (1, 134 mm HL, 679+ mm TL), south of Tomakomai, Hokkaido, 42.1267°N, 141.6133°E, 650–750 m, T/V *Oshoro-maru*, bottom trawl, coll. F. Tashiro and M. Ono, 29 Sept. 2009.

Counts and measurements. Based on 15 specimens (52.3–142 mm HL, 249+–707+ mm TL). Counts: first dorsal-fin rays II,9–11; pectoral-fin rays i15–i19; pelvic-fin rays 7; gill rakers on first arch (outer/inner) 0/10–13, on second arch 9–11/10–12; longitudinal scales 28–35; transverse scale rows below first dorsal-fin origin 5.5–9, below first dorsal-fin midbase 5–7, below second dorsal-fin origin 6–8, above anal-fin origin 21–27.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 30–36 (40–52); orbit diameter 25–33 (36–47); postorbital length 38–46 (55–64); postrostral length 66–74; orbit–preopercle distance 43–49 (62–70); suborbital width 11–13 (15–19); upper-jaw length 34–43 (50–60); length of rictus 27–34 (39–47); length of premaxillary tooth band 22–27 (31–37); preoral length 19–28 (26–41); length of terminal snout scute 2 (3); length of lateral nasal ridge 18–24 (24–36); length of suborbital ridge 88–93 (122–135); snout width 22–29 (31–42); internasal width 23–28 (32–41); interorbital width 28–33 (39–49); occipital width 9–11 (12–16); body width over pectoral-fin bases 33–57 (47–81); body depth at first dorsal-fin origin 64–77 (92–112); body depth at anal-fin origin 50–70 (71–99); prepelvic length 108–124 (154–174); preanus length 132–146 (186–208); preanal length 147–164 (209–235); isthmus–pelvic distance 34–45 (51–65); isthmus–anus distance 56–72 (82–101); isthmus–anal distance 68–90 (101–130); pelvic–anal distance 32–50 (44–72); anus–anal distance 13–23 (20–34); pelvic-fin length 33–44 (45–61); pectoral-fin length 61–74 (88–104); predorsal length 102–112 (143–169); height of first dorsal fin 41–57 (59–83); length of first dorsal-fin base 21–31 (30–44); interdorsal length 39–60 (58–83); length of gill slit 15–22 (23–30); length of posterior nostril 7–9 (10–13); barbel length 4–6 (6–9); length of pyloric caecum 9 (13).

Size. Attains about 96 cm TL (NSMT-P 463, 963+ mm TL, probably Japan).

Distribution. Restricted to the northwestern Pacific around Japan, including Taiwan and Korea. Known from the Pacific off Japan from Kushiro (144.38°E) southward

FIGURE 67. Fresh specimens of *Coelorinchus macrochir*. (A–C) BSKU 110140, 30.5 mm HL, 100+ mm TL, Suruga Bay, Shizuoka Pref., Pacific, 200–450 m depth; (D) HUMZ 206465, 128 mm HL, 643 mm TL, south of Tomakomai, Hokkaido, Pacific, 725 m depth; (E) KPM-NI 34639, 139 mm HL, 689+ mm TL, Kanagawa Pref., Pacific. (A, D–E) Lateral views; (B) dorsal and (C) ventral views of the head and trunk. [Photos: (A–C) N. Nakayama; (D) HUMZ; (E) H. Masuda]

to Bungo Channel (132.06°E), Okinawa Trough, Sea of Japan off Niigata and the Korean Peninsula, and southern Okhotsk Sea off Iturup Island, at depths of 158–830 m (Balanov 2003; Shao *et al.* 2008b; Kim *et al.* 2009b; this study; Appendix 3-4B). The only species of *Coelorinchus* occurring in the Okhotsk Sea. Common off the Pacific coasts of northern Japan north of Choshi (35.74°N; Chiba Pref.), but rare in other parts of its range.

Remarks. *Coelorinchus macrochir* was originally described as a species of *Macrurus* (= *Macrourus* Bloch, 1786) based on a large *Challenger* specimen collected from Sagami Bay off Eno-shima (Fig. 68). For a full description see Okamura (1970a; as *Abyssicola macrochir*). This is

the only species of Japanese grenadiers that is widely distributed around the archipelago, including off the Pacific coasts of Japan, East China Sea, Sea of Japan, and Okhotsk Sea. Recently, Shao *et al.* (2008b) reported the species from the Da-xi fish market in northeastern Taiwan, and their specimens was briefly described and illustrated by Iwamoto *et al.* (2015:61, fig. 7). It was also recorded from the east of Korea by Kim *et al.* (2009b).

Relationships and comparisons. *Coelorinchus macrochir* is peculiar within the genus due to its unusual position of the anus. In most species of the genus, the anus abuts the anal-fin origin or is slightly separated from it by a few scale rows, whereas that of *C. macrochir* is located

FIGURE 68. Holotype of *Macrurus macrochir* (= *Coelorinchus macrochir*). BMNH 1887.12.7.123, 142 mm HL, 657+ mm TL, Sagami Bay, Pacific, 631 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

FIGURE 69. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of *Coelorinchus macrochir*. (A–B) BSKU 19160, 59.0 mm HL. (C–D) BSKU 22773, 97.7 mm HL. (A, C) Views from above; (B, D) oblique views. [Photos: NSMT]

about midway along the pelvic–anal distance. It further differs from most congeners in its unique dentition: teeth arranged in narrow band in the premaxillary and two rows in the dentary (vs. moderately wide band in both jaws). Based on these features, previous authors often placed the species in a monotypic genus, *Abyssicola* Goode & Bean, 1896. However, Iwamoto (1990:172) synonymized the genus with *Coelorinchus* (based on O. Okamura's unpublished data). Also, recent molecular studies (Roa-Varón & Orti 2009; Fukui *et al.* 2010) have shown that the species is deeply nested with a clade of *Coelorinchus*, further supporting their synonymy.

Coelorinchus macrochir is unlikely to belong to any groups of the genus mentioned in this study. It is most similar in general appearance to *C. matamua* (McCann & McKnight, 1980) known from temperate waters in the Southern Hemisphere, but differs notably from that species in having a small dermal fossa of the light organ anterior to the anus (vs. no external evidence of light organ) and the underside of the head completely scaled (vs. broadly naked anteriorly), and in lacking a broad dark band encircling the trunk (vs. present).

***Coelorinchus matsubarae* Okamura, 1982**

[Japanese name: Mato-hige]

(Figs. 47B, 70–71; Appendix 3–4C)

Coelorhynchus matsubarae Okamura, 1982:167, fig. 101 (original description; holotype: BSKU 29440, from Kyushu-Palau Ridge, 28°06.7'N, 134°39.9'E, in 550–600 m; 46 paratypes from Kyushu-Palau Ridge and Emperor Seamounts; new Japanese name: “Matō-hige”).

Coelorinchus matsubarae: Okamura 1984b:96, pl. 82, fig. M (compiled); Okamura 1988:96, pl. 82, fig. M (compiled); Okamura 1997:126, fig. 7 (compiled); Nakabo & Kai 2013:507 (in key; Japan); Motomura 2020:38 (listed; Japan).

Coelorinchus matsubarae: Nakabo 1993:367 (in key; Japan); Nakabo 2000:431 (in key; Japan); Nakabo 2002:431 (in key; Japan).

Diagnosis. Ventral light organ externally represented by long dark line, terminated with 1 fossa at each end, one immediately anterior to anus, the other on chest just posterior to isthmus (duct connecting each fossa often difficult to distinguish). Underside of head completely naked except along lateral nasal ridges; dorsal surface of snout almost fully scaled, naked clefts only narrowly developed along each side of median rostral ridge. Snout long, sharply pointed, length 70–88% PRL; dorsal profile of snout slightly concave in lateral view; terminal scute long, slender, spearhead-shaped, length 18–31% PRL; distal portion of snout narrowly attenuated when viewed dorsally; lateral nasal ridge incompletely supported by nasal bone. Interdorsal space less than first dorsal-fin base length. Anal fin originating posterior to vertical through

second dorsal-fin origin. First dorsal fin high, but second spinous ray not especially elongate, height of fin 87–104% PRL. Second dorsal fin well developed, its rays as long as opposites of anal fin. Anus immediately anterior to anal-fin origin. Premaxillary teeth small, slender, conical in short, uniformly wide band, none especially enlarged; posterior end of tooth band falling far short of lateral corner of mouth. Body scales covered with short, reclined, needle-like spinules in subparallel to slightly divergent rows; spinules in each row almost equal in height, free from one another, with last spinule extending slightly beyond posterior scale margin; buttresses scarcely developed. Orbit diameter 44–50% PRL; postorbital length 49–56% PRL; upper-jaw length 38–46% PRL; preoral length 61–85% PRL; barbel length 8–18% PRL. Transverse scale rows below first dorsal-fin midbase 4–6.5. Spinules on scales anterior to light organ not rudimentary. Prominent large dark blotch between lateral line and pectoral-fin base; about 3 faint dark saddles on tail; fine dark vermiculations on dorsal surface of head (uniformly dark brown in preserved specimens); scale pockets on dorsal half of body outlined by dark brown; underside of head dusky, heavily peppered; lips pale; oral cavity, and gular and branchiostegal membranes blackish; first dorsal fin pale with blackish margin; pelvic fin dark, but outermost ray paler distally.

Material examined. 60 specimens. **Holotype of *Coelorhynchus matsubarae*:** BSKU 29440 (86.6 mm HL, 308+ mm TL), Kyushu-Palau Ridge, Japan, 28.1117°N, 134.6650°E, 550–600 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, tr. 9, bottom trawl, coll. O. Okamura *et al.*, 19 Dec. 1979. **Paratypes of *C. matsubarae*:** BSKU 32226 (1, 73+ mm HL, 286+ mm TL), *BSKU 32229 (1, 68+ mm HL, 277 mm TL), *BSKU 32230 (1, 69+ mm HL, 211+ mm TL), BSKU 32231 (1, 66+ mm HL, 259 mm TL), BSKU 32232 (1, 71.7 mm HL, 227+ mm TL), *BSKU 32233 (1, 65.6 mm HL, 234+ mm TL), BSKU 32234 (1, 67.5 mm HL, 245+ mm TL), BSKU 32235 (1, 66.6 mm HL, 266 mm TL), BSKU 32236 (1, 62.5 mm HL, 239 mm TL), *BSKU 32237 (1, 60.8 mm HL, 233+ mm TL), BSKU 32238 (1, 64.0 mm HL, 242 mm TL), BSKU 32239 (1, 64.1 mm HL, 244+ mm TL), BSKU 32240 (1, 55.1 mm HL, 210 mm TL), BSKU 32241 (1, 57.9 mm HL, 225+ mm TL), BSKU 32242 (1, 53.5 mm HL, 214+ mm TL), BSKU 32243 (1, 62.9 mm HL, 239+ mm TL), *BSKU 32244 (1, 57.6 mm HL, 206+ mm TL), BSKU 32245 (1, 52.8 mm HL, 201+ mm TL), BSKU 32246 (1, 57.3 mm HL, 215+ mm TL), BSKU 32248 (1, 55.2 mm HL, 189+ mm TL), BSKU 32249 (1, 51.2 mm HL, 204 mm TL), BSKU 32251 (1, 53.2 mm HL, 211 mm TL), *BSKU 32252 (1, 56.2 mm HL, 195+ mm TL), *BSKU 32253 (1, 50+ mm HL, 185+ mm TL), *BSKU 32254 (1, 51.1 mm HL, 197+ mm TL), *BSKU 32255 (1, 53.3 mm HL, 208 mm TL), BSKU 32256 (1, 43.0 mm HL, 166+ mm TL), BSKU 32257 (1, 41.7 mm HL, 146+ mm TL), collected with holotype; *ZMUT KP448

FIGURE 70. Fresh specimens of *Coelorinchus matsubarai*. (A) ZMUT unknown, Kyushu-Palau Ridge, Pacific; (B) BSKU 29440, holotype, 86.6 mm HL, 308+ mm TL, Kyushu-Palau Ridge, Pacific, 550–600 m depth. Lateral views. [Photos: (A) ZMUT; (B) BSKU]

(1, 62.5 mm HL, 197+ mm TL), *ZMUT KP450 (1, 43.0 mm HL, 161+ mm TL), *ZMUT KP451 (1, 68.6 mm HL, 253+ mm TL), *ZMUT KP456 (1, 66.1 mm HL, 252+ mm TL), *ZMUT KP459 (1, 64.8 mm HL, 244+ mm TL), *ZMUT KP463 (1, 53.4 mm HL, 203+ mm TL), *ZMUT KP468 (1, 54.2 mm HL, 214 mm TL), *ZMUT KP470 (1, 53.6 mm HL, 197+ mm TL), *ZMUT KP475 (1, 56.9 mm HL, 210 mm TL), *ZMUT KP478 (1, 74.6 mm HL, 285 mm TL), *ZMUT KP479 (1, 77.1 mm HL, 295 mm TL), *ZMUT KP481 (1, 58.2 mm HL, 241+ mm TL), *ZMUT KP489 (1, 64.3 mm HL, 246 mm TL), *ZMUT KP493 (1, 63.7 mm HL, 254+ mm TL), *ZMUT KP495 (1, 73.8 mm HL, 285+ mm TL), *ZMUT KP498 (1, 72.0 mm HL, 283 mm TL), *ZMUT KP596 (1, 55.2 mm HL, 209 mm TL), *ZMUT KP620 (1, 61.9 mm HL, 231+ mm TL), *ZMUT KP626 (1, 65.7 mm HL, 249 mm TL), *ZMUT KP640 (1, 59.2 mm HL, 215 mm TL), Kyushu-Palau Ridge, 28.0833°N, 134.7000°E, 550 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, bottom trawl, coll. K. Mochizuki *et al.*, 17 Jan. 1980; BSKU 29527 (1, 60.8 mm HL, 231 mm TL), BSKU 30600 (1, 54.2 mm HL, 208+ mm TL), *BSKU 30601 (1, 60.3 mm HL, 240+ mm TL), BSKU 30603 (1, 75.5 mm HL, 287+ mm TL), Kyushu-Palau Ridge, 28.0667°N, 134.6833°E, 523 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, tr. 14, bottom trawl, coll. O. Okamura *et al.*, 18 Dec. 1978.

Non-types: Japan: ZMUT KP490 (1, 61.0 mm HL, 226+ mm TL), ZMUT KP616 (1, 42.7 mm HL, 156+ mm TL), ZMUT KP621 (1, 52.8 mm HL, 189+ mm TL), ZMUT KP622 (1, 62.7 mm HL, 239+ mm TL), ZMUT KP641 (1, 48.8 mm HL, 187+ mm TL), Kyushu-Palau Ridge, date unknown; BSKU 32237 (1, 60.4 mm HL, 234+ mm TL), BSKU 32254 (1, 50.8 mm HL, 197+ mm TL), collected with holotype.

Counts and measurements. Based on 30 specimens (41.3–86.6 mm HL, 145+–308+ mm TL). Counts: first dorsal-fin rays II, 8–10; pectoral-fin rays i16–i20; pelvic-fin rays 6–7; gill rakers on first arch (outer/inner) 0/8–10, on second arch 6–7/9; longitudinal scales 36–43; transverse scale rows below first dorsal-fin origin 5.5–7.5, below first dorsal-fin midbase 4–6.5, below second dorsal-fin origin 5.5–7, above anal-fin origin 16.5–22.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 41–47 (70–88); orbit diameter 25–28 (44–50); postorbital length 27–33 (49–56); postrostral length 54–60; orbit–preopercle distance 27–34 (50–58); suborbital width 12–14 (21–24); upper-jaw length 21–27 (38–46); length of rictus 17–22 (30–37); length of premaxillary tooth band 10–12 (17–21); preoral length 36–46 (61–85); length of terminal snout scute 10–17 (18–31); length of lateral nasal ridge 28–35 (48–65); length of suborbital

FIGURE 71. Holotype of *Coelorhynchus matsubarai* (= *Coelorinchus matsubarai*). BSKU 29440, 86.6 mm HL, 308+ mm TL, Kyushu-Palau Ridge, Pacific, 550–600 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

ridge 91–97 (153–175); snout width 24–27 (42–51); internasal width 17–19 (29–35); interorbital width 20–23 (34–40); occipital width 9–11 (15–20); body width over pectoral-fin bases 30–45 (57–76); body depth at first dorsal-fin origin 43–53 (76–89); body depth at anal-fin origin 33–38 (58–68); prepelvic length 101–111 (177–203); preanus length 140–156 (254–275); preanal length 144–163 (263–280); isthmus–pelvic distance 25–30 (46–53); isthmus–anus distance 59–80 (107–136); isthmus–anal distance 63–87 (114–149); pelvic–anal distance 40–61 (73–105); anus–anal distance 2–8 (3–13); pelvic-fin length 33–40 (57–74); pectoral-fin length 36–45 (64–79); predorsal length 106–111 (185–202); height of first dorsal fin 48–58 (87–104); length of first dorsal-fin base 17–22 (31–38); interdorsal length 9–14 (16–24); length of gill slit 10–13 (19–23); length of posterior nostril 7–10 (11–18); barbel length 4–10 (8–18).

Size. To at least 31 cm TL (BSKU 29440, holotype, 308+ mm TL).

Distribution. So far known only from the Kyushu-Palau Ridge (Appendix 3-4C) and the southern part of the Emperor Seamounts at depths of 315–600 m (Okamura 1982; Sazonov 1994; Iwamoto & Okamoto 2015). Common in its range.

Remarks. *Coelorinchus matsubarai* was described from 57 specimens collected from the Kyushu-Palau Ridge and the Emperor Seamounts (Figs. 70B, 71). The reader can refer to the original description (Okamura 1982) for further morphological details. This species is abundant in its range, but it does not occur on the continental slope along the Pacific coasts of southern Japan.

Relationships and comparisons. *Coelorinchus matsubarai* belongs to the *C. hubbsi* group (see the Relationships of *C. brevirostris*), and is most similar to *C. gladius* Gilbert & Cramer, 1897 confined to Hawaii and *C. spilonotus* Sazonov & Iwamoto, 1992 known

from the Sala y Gomez Ridge and Hawaii. They are diagnosed within the group by having a prominent dark blotch between the lateral line and the pelvic-fin base, and having about 2–3 faint dark saddles on the body.

Coelorinchus matsubarai differs from *C. gladius* and *C. spilonotus* in that the spinules on the body scales are much shorter, with the last spinule in each row extending only slightly beyond the posterior scale margin (Fig. 47B). In contrast, the spinules of the other two species are much longer, with the tip of the last spinules extending well beyond the scale margin. The mottled color pattern of the dorsal surface of the head appears to be unique to *C. matsubarai*, although it is completely or almost completely faded in preserved specimens. Unfortunately, it is unknown whether the pattern is also found in *C. gladius* and *C. spilonotus*, as their fresh coloration has not been reported. *Coelorinchus matsubarai* is further distinguished from *C. gladius* by having a shorter scute at the snout tip (18–31% PRL vs. 45–52%). Due to the different size of the scute, the two species are readily distinguished by several morphometric characters, including the snout length (70–88% PRL in *C. matsubarai* vs. 111% in *C. gladius*), preoral length (61–85% PRL vs. 104–106%), pre-dorsal length (185–202% PRL vs. 218–222%), pre-pelvic length (117–203% PRL vs. 235%), and pre-anal length (263–280% PRL vs. 316%). *Coelorinchus matsubarai* also differs from *C. spilonotus* in that the dorsal surface of the snout is entirely scaled except for narrow naked clefts along the median rostral ridge (vs. prominent naked areas dorsally posterior to lateral nasal ridges). It is further distinguished from *C. spilonotus* by having a longer terminal scute (18–31% PRL vs. 15–16%) and more gill rakers [inner rakers on the first arch 8–10 vs. 5–6 in the two *C. spilonotus* specimens examined (6–8 in Sazonov & Iwamoto 1992:50); inner rakers on the second arch 9 vs. 6 (6–8)]. The position of the saddles is

also different between the two species. In *C. matsubarae*, the first saddle originates one first dorsal-fin base length posterior to the second dorsal-fin origin, and the second saddle is located well posterior to it. In *C. spilonotus*, the first saddle is situated anterior to the first dorsal-fin origin, and the position of the second saddle almost corresponds with that of the first of *C. matsubarae* (see Sazonov & Iwamoto 1992: fig. 18).

***Coelorhynchus multispinulosus* Katayama, 1942**

[Japanese name: Yari-hige]

(Figs. 34D, 72–74; Appendix 3-4D)

[?] *Macrourus japonicus* (not Temminck & Schlegel 1846): Temminck & Schlegel 1846:256, pl. CXII, figs. II, Ila, Ilb [original description; in part, 2 paralectotypes, RMNH D1405 and RMNH D1406; “bays in provinces Oomura and Shimabara” (Nagasaki Pref.)]; Boeseman 1947:184 (re-examination of type specimens; in part, 2 paralectotypes, RMNH D1405 and RMNH D1406).

Coelorhynchus anagirostris [sic] (not *Coelorhynchus anatirostris* Jordan & Gilbert in Jordan and Starks 1904): Jordan & Thompson 1914:306 [1 spec. from Misaki; FMNH 57216 (ex. CM 6155a)].

[?] *Coelorhynchus japonicus* (not Temminck & Schlegel 1846): Kikuchi 1931:9 (listed; Toyama Bay); Katayama 1940:26 (listed; Toyama Bay); Mori 1952:72 (listed; east of Korean Peninsula); Mori 1956:10 (listed; Kasumi and Hamada, San'in, Sea of Japan); Uchida *et al.* 1957:28 (listed; Tsuyazaki, Fukuoka Pref.); Yamashita 1959:22 (listed; Genkai-nada off Saga Pref.); Lindberg & Legeza 1969:243, fig. 239 (compiled; brief description; Sea of Japan); Tsuda 1990:566, unnumbered figs. (compiled; Sea of Japan); Shinohara *et al.* 2014:241 (listed; Sea of Japan).

Coelorhynchus multispinulosus Katayama, 1942:332, fig. 1 [original description; 1 spec. from “Tsuiyama Market, Hyogo-ken”, catalog number of holotype not indicated (now known as NSMT-P 18224); new Japanese name: “Yari-hige”]; Matsubara 1955:1310, pl. 133, fig. 454 (in key; Japan; *C. vermicularis* a junior synonym of *C. multispinulosus*); Mori 1956:10 (listed; Tsuiyama, San'in, Sea of Japan); Kamohara 1958:73 (listed; Kochi Pref.); Okada *et al.* 1959:83 (listed; Kumano-nada); Chyung & Kim 1959:7, pl. II, figs. 15, 17 [first record from Korea; brief description; 6 spec. from Korea (Pusan, Masan, Tongyung, and Yeosu)]; Kamohara 1961b:68, pl. 68, fig. 1 [compiled; Japan; erroneously noted Japanese name as “Yaridara” (Inase-hige)]; Chu *et al.* 1963:172, fig. 138 [description; 4 spec. from China (Wenzhou, Zhoushan, Jiangsu, and northern part of East China Sea)]; Kamohara 1964:95 (listed; Kochi Pref.); Tsuda 1964:6 (listed; Toyama Bay); Tataru *et al.* 1965:121 (listed; off Shikoku); Matsubara 1965:505 (compiled; Japan); Lindberg & Legeza 1969:243, figs. 233, 234A, 237, 238A (compiled; brief description; Sea of Japan); Okamura 1970a:156, pl. XXXIII, text-fig. 63 (description; biological notes; 89 spec. from Pacific off

southern Japan from Heta to Shibushi and East China Sea); Takegawa & Morino 1970:388 (listed; Wakasa Bay, Sea of Japan); Honma *et al.* 1972:56, fig. 6 (brief description; 1 spec. from Niigata Pref.); Shiohaki & Dotsu 1973:30 (listed; Nomozaki, Nagasaki Pref.); Dotsu & Nakamura 1976:285 (listed; Tsushima Island); Takata *et al.* 1979: table 1 (listed; Sea of Japan off Tsushima Island); Kataoka & Tomida 1981:78 (listed; Mie Pref.); Tominaga & Uyeno 1981:488 (listed; Japan); Ohta 1983: table A (listed; Suruga Bay); Tsuda 1990:569, unnumbered figs. (compiled; Sea of Japan); Shinohara *et al.* 2011:41 [listed (based on holotype); Sea of Japan]; Shinohara *et al.* 2014:241 (listed; Sea of Japan).

Coelorhynchus vermicularis Matsubara, 1943:133, figs. 1A, 2A, 3 [original description; holotype: FAKU 2495 (ex SKFC 3), from Kumano-nada; 3 paratypes from Kumano-nada]; Kuroda 1947:31 (brief description; 1 spec. from Suruga Bay); Kuroda 1951:391 (listed; Suruga Bay); Kamohara 1952:99 (spec. from Kochi Pref.).

[?] *Coelorhynchus parallelus* (not Günther 1877): Yanai 1950:22 (listed; San'in; Sea of Japan); Lindberg & Legeza 1969:244, figs. 236, 240 (compiled; brief description; Sea of Japan); Tsuda 1990:569, unnumbered figs. (compiled; Sea of Japan).

Coelorhynchus intermedius Chu & Lo in Chu, Chan & Chen, 1963:173, fig. 139 [original description; holotype: SFU 1773, from East China Sea off Zhejiang, China; 5 paratypes from Zhejiang, China).

Coelorhynchus abbreviatus Chu & Lo in Chu, Chan & Chen, 1963:174, fig. 140 (original description; 4 spec. from East China Sea off Zhejiang, China).

[?] *Coelorhynchus longissimus* (not Matsubara 1943): Tsuda 1964:6 (listed; Toyama Bay); Tsuda 1990:569, unnumbered figs. (compiled; Sea of Japan).

[?] *Coelorhynchus kamoharai* (not Matsubara 1943): Tsuda 1964:6 (listed; Toyama Bay); Takata *et al.* 1979: table 1 (listed; Sea of Japan off Tsushima Island); Tsuda 1990:566, unnumbered figs. (compiled; Sea of Japan).

Coelorhynchus (Quincuncia) multispinulosus: Okamura 1970b: table 1 (listed; Japan).

Coelorhynchus multispinulosus: Yatou 1984:223, 365, fig. 158 [brief description; 7 spec. from Okinawa Trough; photo based on BSKU spec. (catalog no. unknown)]; Okamura 1984b:96, pl. 82, fig. H (compiled); Kamohara & Okamura 1985:32, pl. 31, fig. 153 (compiled; Japan); Kishida 1986:100 (brief description; East China and Yellow Seas); Okamura 1988:96, pl. 82, fig. H (compiled); Iwamoto 1990:174, fig. 398 (synopsis); Fujioka 1991:119 [listed; Yamaguchi Pref. (Murozumi, Seto Inland Sea)]; Suzuki & Kataoka 1997:82, pl. 33, fig. 178 (brief description; 1 spec. from Kumano-nada); Okamura 1997:126, fig. 8 (compiled); Youn 2002:538 (listed; Korea); Shao *et al.* 2008b: table 2 (11 spec. listed from northeastern Taiwan); Tamai *et al.* 2012:34, fig. 2F (1 spec. from Enshu-nada); Nakabo & Kai 2013:506 (in key; Japan); Ikeda & Nakabo 2015:320, pl. 66, figs. 8–11 (brief description; spec. from Wakayama Pref.); Iwamoto *et al.* 2015:63 (brief description; 46 spec. from northeastern and northern Taiwan); Iwatsubo & Motomura 2017:59, unnumbered fig. (listed;

Kagoshima Bay); Fujiwara *et al.* 2018:51, fig. 5E, table 1 (3 spec. listed from Yamaguchi Pref.); Sonoyama *et al.* 2020:29 (4 spec. listed from Yamaguchi Pref., Sea of Japan); Motomura 2020:38 (listed; Japan).

Coelorhynchus multispinulosus [sic]: Akazaki 1984:265 (listed; Aoshima, Miyazaki Pref.).

Caelorinchus multispinulosus: Nakabo 1993:366 (in key; Japan); Shao 1993:170, fig. 37-10 (compiled; Taiwan); Yamada *et al.* 1995:96, fig. 23 (compiled; East China and Yellow Seas); Shinohara & Matsuura 1997:291 (listed; Suruga Bay); Shinohara *et al.* 1998:111 (listed; Tachibana Bay, Nagasaki); Nakabo 2000:430 (in key; Japan); Shinohara *et al.* 2001:304 (15 spec. listed from Tosa Bay); Nakabo 2002:430 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Nakajima 2003:53, pl. 15, fig. 86 (brief description; 1 spec. from Enshu-nada); Matsumoto 2005: table 1 (listed; Shimane Pref.); Shinohara *et al.* 2005:417 (42 spec. listed from Ryukyu Islands); Kim *et al.* 2005:173, fig. 294 (compiled; Korea); Yamada *et al.* 2007:312, pl. 14, fig. 5 (brief description; 1 spec.; East China and Yellow Seas; biological notes); Kim *et al.* 2009a:13 (listed; Jeju Island, Korea); Furuhashi *et al.* 2010: table 2 (27 spec. listed from northern Okinawa Trough); Kawano *et al.* 2014: appendix table (listed; Sea of Japan); Iwatsuki *et al.* 2017:32 (listed; Hyuga-nada).

Caelorinchus (Quincuncia) multispinulosus: Chiou *et al.* 2004a: table 1 (listed; Taiwan).

Caelorinchus mulispinulosus [sic]: Nishida *et al.* 2004: table 1 (listed; Tsuyazaki, Fukuoka Pref.).

[?] *Caelorinchus japonicus* (not Temminck & Schlegel 1846): Nishida *et al.* 2004: table 1 (listed; Tsuyazaki, Fukuoka Pref.); Kawano *et al.* 2014: appendix table (listed; Sea of Japan).

Caelorinchus sp.: Matsuura *et al.* 2009:24 (in part, NSMT-P 73336; 1 spec. from Fukushima Pref.).

[?] *Coelorhynchus kamoharai* (not Matsubara 1943): Kawano *et al.* 2014: appendix table (listed; Sea of Japan).

[?] *Coelorhynchus longissimus* (not Matsubara 1943): Kawano *et al.* 2014: appendix table (listed; Sea of Japan).

[?] *Coelorhynchus parallelus* (not Günther 1877): Kawano *et al.* 2014: appendix table (listed; Sea of Japan); Shinohara *et al.* 2014:241 (listed; Sea of Japan).

Diagnosis. Light organ tubular, completely covered with scales, externally represented by long black streak extending from immediately anterior to anus to chest just posterior to isthmus. Underside of head completely naked except for overlapping scales posterior to lateral nasal ridges; dorsal surface of snout broadly naked along each side of median rostral ridge. Snout long, sharply pointed, length 68–81% PRL, its dorsal profile straight in lateral view; terminal scute short, arrowhead-shaped, dorsoventrally flattened, length 8–20% PRL. Lateral nasal ridge incompletely supported by nasal bone. Anus abutting anal-fin origin. Premaxillary teeth small, slender, conical in moderately wide, long, tapered band, with outer series slightly enlarged; posterior margin of tooth band almost

reaching lateral corner of mouth. Body scales covered with short, erect, needle-like spinules in quincunx order; tips of spinules recurved posteriorly; base of each spinule lacking buttresses. Second dorsal fin distinctly lower than anal fin. Upper-jaw length 45–52% PRL; pectoral-fin length 56–72% PRL; height of first dorsal fin 81–100% PRL; interdorsal length 25–41% PRL; length of gill slit 21–27% PRL; barbel length 16–24% PRL. Nape and dorsal half of body overlain with dark vermiculations or small irregular spots; faint dark blotches serially arranged along lower 1/3 of body; ventral half of head and body silvery white when viewed laterally; median nasal bone blackish; underside of head generally pale, without short, black, hair-like papillae; lips pale to dusky; oral cavity immaculate; gular and branchiostegal membranes heavily peppered; chest dark; first dorsal fin dusky, but second spinous ray prominently blackish; pelvic fin pale to dusky.

Material examined. 35 specimens. **Holotype of *Coelorhynchus multispinulosus*:** NSMT-P 18224 (52.1 mm HL, 190+ mm TL), Tsuiyama market, Hyogo Pref., Japan, Sea of Japan, Apr. 1942. **Holotype of *Coelorhynchus intermedius*:** SFU 1773 (51.7 mm HL, 190+ mm TL), northern part of East China Sea, off Zhejiang, China, Feb. 1958. **Non-types: Japan:** BSKU 27774 (1, 68.3 mm HL, 258+ mm TL), west of Amami-oshima Island, Okinawa Trough, 28.4167°N, 127.0000°E, 200 m, F/V *Ryoan-maru*, No. 28, tr. 53, bottom trawl, coll. Y. Kinoshita and S. Hagino, 12 Mar. 1978; BSKU 28671 (1, 62.8 mm HL, 270 mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.9167°N, 127.4833°E, 130–160 m, F/V *Ryoan-maru*, No. 28, tr. 81, bottom trawl, coll. Y. Kinoshita and S. Hagino, 22 Mar. 1978; BSKU 91328 (1, 66.1 mm HL, 278+ mm TL), Saga fish market, otter trawl, 27 Jun. 2007; BSKU 99273 (1, 69.9 mm HL, 290+ mm TL), BSKU 99274 (1, 63.5 mm HL, 254+ mm TL), BSKU 99275 (1, 53.0 mm HL, 208+ mm TL), Saga fish market, otter trawl, 27 May 2007; BSKU 99390 (1, 67.2 mm HL, 257+ mm TL), BSKU 99391 (1, 61.9 mm HL, 219+ mm TL), BSKU 99392 (1, 57.3 mm HL, 244+ mm TL), BSKU 99393 (1, 68.1 mm HL, 254+ mm TL), BSKU 99394 (1, 69.7 mm HL, 265+ mm TL), BSKU 99398 (1, 61.4 mm HL, 224+ mm TL), Saga fish market, otter trawl, 8 Jun. 2007; BSKU 99263 (1, 46.6 mm HL, 195+ mm TL), Mimase fish market, bottom trawl, 1 Mar. 2006; BSKU 99362 (1, 59.8 mm HL, 243+ mm TL), BSKU 99364 (1, 60.2 mm HL, 275+ mm TL), BSKU 99365 (1, 37.2 mm HL, 143+ mm TL), BSKU 99370 (1, 70.1 mm HL, 295+ mm TL), Mimase fish market, bottom trawl, 11 Jan. 2007; BSKU 9672 (1, 78.7 mm HL, 332+ mm TL), Mimase fish market, coll. T. Kamohara, 24 Dec. 1950; KSHS 19279 (1, 80.7 mm HL, 346+ mm TL), Mimase fish market, coll. T. Yamakawa, 3 Mar. 1981; BSKU 95364 (1, 57.3 mm HL, 227+ mm TL), Mimase fish market, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama, 12 Mar. 2008; BSKU 92369

(1, 55.4 mm HL, 249 mm TL), Mimase fish market, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama, 16 Nov. 2007; BSKU 108900 (1, 78.8 mm HL, 332+ mm TL), off Kannoura, Mimase fish market, bottom trawl, 2007; BSKU 11562 (1, 79.3 mm HL, 288+ mm TL), Mimase or Urado fish markets, bottom trawl, coll. A. Kashiwagi and S. Furuno, 11 Oct. 1965; BSKU 112976 (1, 35.8 mm HL, 166 mm TL), Etomo fish market, Matsue, Sea of Japan, F/V *Ichiwa-maru*, bottom trawl, coll. Y. Misawa, 17 Mar. 2014; BSKU 112572 (1, 52.4 mm HL, 230+ mm TL), BSKU 112574 (1, 54.1 mm HL, 218+ mm TL), *BSKU 112578 (1, 64.3 mm HL, 292+ mm TL), Etomo fish market, Matsue, Sea of Japan, F/V *Myojin-maru*, bottom trawl, coll. Y. Misawa, 17 Mar. 2014; KPM-NI 23523 (1, 60.0 mm HL, 281+ mm TL), west of Akita, Sea of Japan, coll. H. Sugiyama, 2008; KPM-NI 20135 (1, 42.3 mm HL, 198 mm TL), west of Akita, Sea of Japan, 120 m, F/V *Kane-maru*, coll. H. Sugiyama, 28 Mar. 2007. **South China Sea:** BSKU 17171 (1, 50.4 mm HL, 209+ mm TL), BSKU 17172 (1, 45.8 mm HL, 169+ mm TL), southeast of Ho Chi Minh, 6.8600°N, 108.7867°E, 135–137 m, R/V *Hakuho-maru*, cr. KH-72-01, sta. 50, beam trawl, coll. O. Okamura, 10–11 Jul. 1972; SFU SH1607 (1, 52.1 mm HL, 191+ mm TL), SFU SH1614 (1, 40.6 mm HL, 170+ mm TL), off Paracel Islands, Dec. 1972.

Counts and measurements. Based on 34 specimens (35.8–80.7 mm HL, 143+–346+ mm TL). Counts: first dorsal-fin rays II,8–10; pectoral-fin rays i12–i18; pelvic-fin rays 7; gill rakers on first arch (outer/inner) 0/7–9, on second arch 5–6/7–9; longitudinal scales 39–50; transverse scale rows below first dorsal-fin origin 5.5–8.5, below first dorsal-fin midbase 4–6, below second dorsal-fin origin 4.5–7, above anal-fin origin 14–20.5.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 41–45 (68–81); orbit diameter 23–28 (40–47); postorbital length 30–36 (53–61); postrostral length 56–60; orbit–preopercle distance 34–39 (58–66); suborbital width 11–14 (20–25); upper-jaw length 26–30 (45–52); length of rictus 20–25 (34–41); length of premaxillary tooth band 17–22 (30–37); preoral length 31–38 (52–68); length of terminal snout scute 5–11 (8–20); length of lateral nasal ridge 25–31 (42–56); length of suborbital ridge 93–99 (158–175); snout width 22–26 (38–45); internasal width 19–22 (33–39); interorbital width 22–25 (38–43); occipital width 4–8 (7–14); body width over pectoral-fin bases 32–44 (56–75); body depth at first dorsal-fin origin 47–60 (81–104); body depth at anal-fin origin 42–53 (73–92); prepelvic length 103–113 (177–196); preanus length 138–156 (241–264); preanal length 141–160 (245–272); isthmus–pelvic distance 33–44 (58–75); isthmus–anus distance 74–85 (125–144); isthmus–anal distance 67–94 (119–160); pelvic–anal distance 38–54 (67–92); anus–anal distance 2–4 (4–7); pelvic-fin length 25–36 (42–62); pectoral-fin length 32–42 (56–72); predorsal length 103–113 (178–197); height of first dorsal fin 48–57 (81–100);

length of first dorsal-fin base 18–26 (31–41); interdorsal length 15–24 (25–41); length of gill slit 12–16 (21–27); length of posterior nostril 5–8 (8–15); barbel length 9–14 (16–24).

Size. To about 38 cm TL (Okamura 1970a).

Distribution. Distributed in the northwestern Pacific from Japan to China. Known from off the Pacific coasts of Japan northward to Iwaki (37.05°N; Fukushima Pref.), the Sea of Japan south of Akita (ca. 40°N), and the East and South China Seas (Chyung & Kim 1959; Shao *et al.* 2008a, 2008b; this study; Appendix 3–4D). Depth range 78–400 m, but usually less than 200 m. Common in Pacific Japan west of the Kii Channel (134.92°E) and the Sea of Japan west of the Oki Islands (133.16°E), as well as on the continental shelf of the East China Sea.

Nomenclatural discussion. *Coelorinchus multispinulosus* was originally described by Katayama (1942) based on a single specimen obtained at the Tsuiyama fish market in Hyogo Pref., Japan (collected from the Sea of Japan; Fig. 73). A year later, Matsubara (1943) described *C. vermicularis* based on four specimens collected from the Kumano-nada, Japan, and then (1955:1310) subsequently synonymized the species with *C. multispinulosus*. Unfortunately, the holotype of *C. vermicularis* was lost during WWII (see also the Comments on holotype of *C. hige*), and no paratypes were rediscovered during this study. Matsubara's (1943) detailed description of *C. vermicularis* agrees well with the holotype and additional specimens of *C. multispinulosus*, and thus these two names are considered synonymous in this study.

Chu & Lo (*in Chu et al.* 1963) described *C. abbreviatus* and *C. intermedius* from the northwestern part of the East China Sea off China, based on four and six specimens respectively. Subsequently, Okamura (1970a) synonymized these two species with *C. formosanus* Okamura, 1963 without explanation of type specimens of *C. abbreviatus* and *C. intermedius*, and all subsequent authors have followed his treatment. During a visit to Shanghai Ocean University in 2018, the present author had an opportunity to examine one of the type specimens labeled as the holotype of *C. intermedius* (SFU 1773, 51.7 mm HL, 190+ mm TL; Fig. 74). Unfortunately, no other types of *C. intermedius* and *C. abbreviatus* were found among the SFU fish collection. While there were two specimens labeled as “paratypes of *C. intermedius*” (SFU SH1607, 52.1 mm HL, 191+ mm TL; SFU SH1614, 40.6 mm HL, 170+ mm TL), these specimens were collected in December 1978 (after the original description), and apparently not included in the type series of *C. intermedius*.

An examination of the holotypes of *C. multispinulosus* and *C. intermedius*, along with 32 additional specimens from Japan, China, and the South China Sea, revealed that these two species are not distinguished morphologically. All measurements of the holotype of *C. intermedius* lie within the range

FIGURE 72. Fresh specimens of *Coelorinchus multispinulosus*. (A–C) BSKU 112572, 52.4 mm HL, 230+ mm TL, Etomo fish market, Shimane Pref., Sea of Japan; (D–F) BSKU 95364, 57.3 mm HL, 227+ mm HL, Mimase fish market, Kochi Pref., Pacific; (G–I) BSKU 112578, 64.3 mm HL, 292+ mm TL, collected with BSKU 112572. (A, D, G) Lateral views; (B, E, H) dorsal and (C, F, I) ventral views of the head and trunk. [Photos: N. Nakayama]

of variation for the *C. multispinulosus* specimens examined. Unfortunately, the coloration of the holotype of *C. intermedius* is completely faded due to its long preservation in formalin, and the presence of the unique vermiculations characterizing *C. multispinulosus*

could not be confirmed in this specimen. However, no significant differences were found in other features including dentition, squamation, and spinulation on the body scales; and thus there is no doubt about their conspecificity.

Although type specimens of *C. abbreviatus* were not available for this study, the original description strongly suggests that the species simply represents an occasional variant of *C. multispinulosus*. Chu & Lo (in Chu *et al.* 1963) distinguished *C. abbreviatus* from *C. multispinulosus* by the development of the “caudal fin” [sic] (prominent vs. rudimentary) and the numbers of second dorsal- (47–55 vs. 93–104) and anal-fin rays (44–52 vs. 87–97). However, a true caudal fin is absent in macrourids (Iwamoto 1970; Okamura 1970b). As noted by Iwamoto (1970:322) and Okamura (1970b:110–111, fig. 58), the tip of the tail is frequently damaged during ontogeny, and a pseudocaudal often replaces the broken end. Most likely, Chu & Lo’s (in Chu *et al.* 1963) “caudal fin” refers to a developed pseudocaudal. Because the size of the pseudocaudal reflects how much the tail tip is damaged, it is an unreliable character in grenadier taxonomy. The numbers of second dorsal- and anal-fin rays are also affected by damage to the tail tip; therefore,

they cannot be used to separate species of grenadiers. The illustration of *C. abbreviatus* provided by Chu & Lo (in Chu *et al.* 1963: fig. 140) clearly depicts that the tail of the species is largely damaged at the distal end, where a prominent pseudocaudal is developed. Furthermore, Chu & Lo (in Chu *et al.* 1963) distinguished *C. abbreviatus* from *C. intermedius*, the latter of which is also regarded here as a junior synonym of *C. multispinulosus* (see previous paragraphs), by the length from the anus to the “caudal-fin base” [sic, probably the base of the pseudocaudal] (about equal to vs. 1.5–1.6 in preanal length), and the numbers of second dorsal- (47–55 vs. 61–70) and anal-fin rays (44–52 vs. 60–68). However, these differences can also be attributed to the damaged tail tip in the type specimens of *C. abbreviatus*. Consequently, *C. abbreviatus* is herein synonymized with *C. multispinulosus*.

While *C. abbreviatus* and *C. intermedius* were previously regarded as junior synonyms of *C. formosanus* Okamura, 1963, their senior synonym *C. multispinulosus*

FIGURE 73. Holotype of *Coelorhynchus multispinulosus* (= *Coelorinchus multispinulosus*). NSMT-P 18224, 52.1 mm HL, 190+ mm TL, Tsuiyama market, Hyogo Pref., Sea of Japan. (A) Lateral, (B) dorsal, and (C) ventral views. Preserved condition. [Photos: N. Nakayama]

FIGURE 74. Preserved specimen of *Coelorinchus multispinulosus*. SFC 1773, holotype of *Coelorhynchus intermedius*, 51.7 mm HL, 190+ mm TL, off Zhejiang, China, East China Sea. (A) Lateral, (B) dorsal, and (C) ventral views. [Photos: N. Nakayama]

is readily differentiated from *C. formosanus* in that the underside of the head is completely naked, except for characteristic overlapping scales posterior to the lateral nasal ridges; these features were described in the original descriptions of the former two species, and also confirmed by the examination of the holotype of *C. intermedius* in this study. By contrast, the underside of the head of *C. formosanus* is broadly scaled on the ventral surfaces of the preopercles and posterior portions of the mandibular rami.

Remarks, relationships, and comparisons. For further morphological details see Okamura (1970a). *Coelorinchus multispinulosus* belongs to the *C. argentatus* group (see the Relationships for *C. formosanus*) and is most similar to *C. pardus* Iwamoto & Williams, 1999 known from the Arafura Sea, but the two species differ most notably from each other in the body markings. A thorough comparison of the two species was provided in the original description of *C. pardus*.

Because of their close resemblance, in Japanese waters *C. multispinulosus* may be confused with *C. kamoharai* Matsubara, 1943 or *C. longissimus* Matsubara, 1943. However, it is readily distinguished from *C. kamoharai* by lacking short, dark, hair-like papillae on the underside of the head and the top of the snout (vs. present and densely scattered), and from *C. longissimus* by having spinules on the body scales arranged in quincunx order (vs. subparallel to narrowly divergent rows).

Coelorinchus nox sp. nov.

[New Japanese name: Yami-hige]

(Figs. 75–77, 78A; Table 3; Appendix 3–4E)

Diagnosis. Light organ externally represented by short, narrow, naked fossa immediately anterior to anus, its anterior margin falling far short of line connecting inner pelvic-fin bases. Underside of head mostly scaled except for narrow to moderately broad naked area above upper lip, and anterior portions of mandibular rami; dorsal surfaces of snout fully scaled. Snout long, sharply pointed, length 62–81% PRL; dorsal profile of snout slightly concave in lateral view; terminal scute moderately short, diamond-shaped, dorsoventrally flattened, length 12–17% PRL; lateral nasal ridge completely supported by nasal bone, moderately convex in dorsal view, giving spade-shaped appearance to snout. Anus slightly separated from anal-fin origin. Premaxillary teeth small, conical, arranged in short, uniformly wide band, none distinctly enlarged (outer series slightly larger); posterior margin of tooth band falling far short of lateral corner of mouth. Body scales covered with short, reclined, keel-like spinules in widely divergent, coarsely saw-tooth ridges; every spinule row complete, extending to posterior scale margin; spinules in each row greatly overlapping, increasing in height posteriorly; buttresses narrowly developed. Occipital scales covered with short, erect, needle-like spinules in 4 or more widely divergent, comb-like rows. Orbit diameter 45–52% PRL; suborbital width 18–23% PRL; internasal width 27–31% PRL; interorbital width 39–41% PRL; pectoral-fin length 57–69% PRL; pectoral-fin rays i16–i17; transverse scale rows below first dorsal-fin midbase 3.5–4.5. Body uniformly dark brown, without either dark saddles or blotches; lips pale to dusky; oral cavity blackish; gular and branchiostegal membranes heavily peppered with small melanophores; first dorsal, pectoral, and pelvic fins uniformly blackish.

Material examined. 14 specimens. **Holotype:** BSKU 109205 (99.2 mm HL, 346+ mm TL), west of Amami-oshima Island, Okinawa Trough, 28.5679°N, 127.0421°E, 607 m, T/V *Nagasaki-maru*, cr. N365, sta. J6, tr. 1, 3-m ORE beam trawl, coll. N. Nakayama, 16 Nov. 2012. **Paratypes:** BSKU 26321 (1, 90.6 mm HL, 313+ mm TL), BSKU 26322 (1, 127 mm HL, 437+ mm TL), BSKU 26323 (1, 98.3 mm HL, 339+ mm TL), off Senkaku Islands, Okinawa Trough, 25.6283°N, 122.8933°E, 560–692 m, F/V *Yuryo-maru*, No. 8, tr. 3, bottom trawl, coll. T. Kitajima *et al.*, 14 Jan. 1978; BSKU 28498 (1, 99.9 mm HL, 338+ mm TL), BSKU 28499 (1, 105 mm HL, 335+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.7333°N, 128.0500°E, 815 m, F/V *Ryoan-maru*, No. 28, tr. 74, bottom trawl, coll. Y. Kinoshita and S. Hagino, 20 Mar. 1978; BSKU 27985 (1, 116 mm HL, 354+ mm TL), BSKU 27988 (1, 111 mm HL, 379+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 28.8333°N, 127.2333°E, 700–740 m, F/V *Ryoan-maru*, No. 28, tr. 59, bottom trawl, coll. Y. Kinoshita and S. Hagino, 16 Mar. 1978; BSKU 27582 (1, 91.7 mm HL, 293+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 28.7167°N, 127.2333°E, 750–755 m, F/V *Ryoan-maru*, No. 28, tr. 48, bottom trawl, coll. Y. Kinoshita and S. Hagino, 11 Mar. 1978; BSKU 109016 (1, 79.8 mm HL, 271+ mm TL), west of Amami-oshima Island, Okinawa Trough, 28.5686°N, 127.0419°E, 604 m, T/V *Nagasaki-maru*, cr. N365, sta. J6, tr. 4, 3-m ORE beam trawl, coll. N. Nakayama, 16 Nov. 2012; BSKU 109011 (1, 87.3 mm HL, 297+ mm TL), west of Amami-oshima Island, Okinawa Trough, 28.5513°N, 127.0375°E, 621 m, T/V *Nagasaki-maru*, cr. N365, sta. J6, tr. 3, 3-m ORE beam trawl, coll. N. Nakayama, 16 Nov. 2012; BSKU 109206 (1, 81.8 mm HL, 270+ mm TL), collected with holotype; BSKU 27673 (1, 112 mm HL, 375+ mm TL), BSKU 27675 (1, 111 mm HL, 378+ mm TL), west of Okinawa-jima Island, Okinawa Trough, 28.7333°N, 127.0167°E, 610–640 m, F/V *Ryoan-maru*, No. 28, tr. 49, bottom trawl, coll. Y. Kinoshita and S. Hagino, 11 Mar. 1978.

Description. General features are shown in Figs. 75–76. Counts and measurements are given in Table 3. Body deepest at first dorsal-fin origin, progressively tapering to long, laterally compressed tail. Trunk short, moderately compressed, width over pectoral-fin bases 1.1–1.3 in depth below first dorsal-fin origin. Head large, HL about 3.1–3.5 in TL. Snout long, sharply pointed, protruding well beyond upper jaw, length 1.2–1.8 times as long as orbit diameter; dorsal contour of snout slightly concave in lateral view; lateral nasal ridge completely supported by nasal bone; anterolateral margins of snout moderately convex when viewed dorsally, giving spade-shaped appearance to snout. Orbit large, its greatest diameter 0.9–1.2 in postorbital length. Interorbital space broad, flat, width 1.1–1.3 in orbit diameter. Mouth moderately large, inferior, protrusible, upper-jaw length 0.9–1.3 in orbit

diameter; posterior margin of maxilla extending beyond vertical through midorbit; lateral corner of mouth barely restricted by skin folds; lips thin, slightly papillose near tooth bands. Suborbital region divided by longitudinal bony ridge passing from tip of snout to posteroventral angle of opercle; its upper half almost vertical, lower half sharply inclined mesially. Preopercle large, posterior margin inclined, forming moderately angular lobe at posteroventral corner. Subopercle prolonged ventrally as slender flap; posterior margin deeply concave. Gill membranes broadly connected across, and attached mesially to isthmus, with free posterior fold. Outer gill slit moderately restricted by skin folds, length 2.0–2.5 in orbit diameter. Gill rakers small, low, tubercular, armed with short, fine spines; no rakers on outer side of first arch and inner side of fourth arch; gill filaments moderately long. Chin barbel short, slender, length 2.3–3.6 in orbit diameter.

Anus only slightly separated from anal-fin origin by 1–2 scale rows. Light organ externally represented by short, narrow, naked streak anterior to anus; its anterior margin not extending to about posterior 1/3 of distance from outer pelvic-fin bases to anus, and falling far short of line connecting inner pelvic-fin bases.

Teeth small, sharp, conical, gently incurved, in narrow bands in both jaws. Premaxillary teeth in short, uniformly wide band, with about 5 tooth rows near symphysis; posterior margin of tooth band falling far short of lateral corner of mouth; teeth becoming progressively smaller inwardly, with outer series only slightly larger. Mandibular teeth uniformly small, in narrow tapered band, with about 4 tooth rows across widest point near symphysis; posterior margin of tooth band extending beyond lateral corner of mouth. All teeth deeply embedded in thick layer of gum papillae.

Scales on body large, thin, moderately deciduous, covered with short, reclined, keel-like spinules in widely divergent rows (Fig. 77); in 86.6 mm HL specimen (BSKU 109011), those on dorsum below interdorsal space with 6–8 rows of spinules; middle row usually longest with about 8 spinules, but not especially high or enlarged compared with adjacent rows; every spinule row complete throughout; spinules forming angle of about 40° to scale surface; spinules greatly overlapping, closely adjoined to one another, but their distal tips usually free, forming saw-toothed ridges that give spiny appearance to body surfaces; height of spinules increasing posteriorly, with tip of last in each row extending moderately beyond posterior scale margin; buttresses of spinules narrowly developed; no reticulate structures on unexposed portion. Body fully scaled except for fins and ventral light organ.

Scales on head ridges thickened and coarsely modified; those on median rostral ridge covered with radiating rows of spinules. Terminal snout scute moderately short, diamond-shaped, dorsoventrally flattened, its length 2.7–4.3 in orbit diameter. Supraoccipital scute

stout, prominent, armed with single row of spinules; posttemporal scutes present, but not especially marked. Occipital scales covered with short, erect, needle-like spinules in widely divergent rows (Fig. 78A); spinules in each row free from one another, forming comb-like ridges, and their height gradually increasing posteriorly; scales on dorsal surfaces of snout posterior to leading edges armed with divergent rows or small cluster of short, erect, needle-like spinules. Other scales on head variable in size, generally similar to those on body, but spinules more erect; those on opercle, preopercle, and postorbital and supratemporal canals largest. Nasal fossa and dorsal surfaces of snout fully scaled; naked clefts along each side of median rostral ridge rudimentary. Underside of head completely scaled, except for narrow to moderately broad area above upper lip, anterior portions of mandibular rami, and posteroventral margins of preopercles; scales on ventral surfaces of head small, non-imbricate, covered with small cluster of short, erect, needle-like spinules.

No open pores along cephalic sensory canals. Free neuromasts scattered over head surfaces, especially on underside of snout, but not prominently marked. Anterior nostril small and circular; posterior large, bean-shaped; septum between nostrils forming anteriorly opened hood. Lateral line complete, not interrupted throughout.

Origins of first dorsal and pelvic fins on about same vertical; first dorsal fin moderately high, its height 2.1–3.0 times as long as its base length; second spinous ray not elongated, smooth along its leading edge; its tip usually extending to second dorsal-fin origin or beyond when laid back. Interdorsal space 1.4–2.2 times as long as first dorsal-fin base length. Second dorsal fin originating posterior to anal-fin origin (above bases of 2nd–7th anal-fin rays). Pectoral fin inserted slightly anterior to vertical through pelvic-fin base. Outer pelvic-fin ray prolonged, with fine distal tip.

Color when fresh (Fig. 75). Head and body uniformly dark brown; gill cover slightly iridescent; all fins blackish.

Color in alcohol (Fig. 76). Head and body uniformly dark brown; abdomen dark blue; lips pale to dusky; oral and gill cavities blackish; gill rakers and filaments light brown, arches pallid; gular and branchiostegal membranes heavily peppered with small melanophores; all fins uniformly blackish.

Size. To about 44 cm TL (BSKU 26322, paratype, 437+ mm TL).

Distribution. So far known from the Okinawa Trough at depths of 560–815 m (Appendix 3-4E). Uncommon.

Etymology. The specific epithet is derived from the Latin noun *nox*, meaning “darkness”, in reference to the characteristic dark coloration of the species. The Japanese name, Yami-hige, is derived from a combination of a word meaning darkness (= yami) and a Japanese name for grenadiers (= hige).

Relationships and comparisons. *Coelorinchus nox* **sp. nov.** belongs to the *C. japonicus* group (sensu stricto; see the Relationships of *C. japonicus*), and is most closely similar to *C. smithi* Gilbert & Hubbs, 1920, with which it has been confused in the East China Sea. The latter species includes at least two morphologically distinct populations, of which one is known from Japan and the other from the East Indies (the type locality of *C. smithi* is Indonesia; see also the Remarks of the latter species). *Coelorinchus nox* readily differs from the both populations of *C. smithi* in having the occipital scales covered with short, erect, needle-like spinules arranged in comb-like ridges (vs. moderately reclined, keel-like to knife-like spinules in saw-toothed ridges; Fig. 78A vs. B), while young specimens of Japanese *C. smithi* occasionally have a similar spinulation pattern. It is further distinguished from *C. smithi* from the East Indies by having a relatively

FIGURE 75. Holotype of *Coelorinchus nox* **sp. nov.** BSKU 109205, 99.2 mm HL, 346+ mm TL, Okinawa Trough, East China Sea, 607 m depth. Lateral view. Fresh condition. [Photo: N. Nakayama]

TABLE 3. Morphometric measurements and meristic counts* for *Coelorinchus commutabilis*, *C. nox* sp. nov., and *C. smithi*.

Species	<i>C. commutabilis</i>	<i>C. nox</i>		<i>C. smithi</i>		
Type status	HT	HT	PT	HT	NT	
Locality	Philippines	Japan	Japan	Indonesia	Philippines	Japan
Number of specimens	<i>n</i> = 1	<i>n</i> = 1	<i>n</i> = 13	<i>n</i> = 1	<i>n</i> = 5	<i>n</i> = 45
Measurements (mm)						
TL	314+	346+	270+–437+	261+	158+–322+	136+–528+
HL	93.7	99.2	79.8–127	80.7	54.2–95.6	42.9–135
% of HL (% of PRL)						
Snout length	42 (70)	40 (66)	39–45 (62–81)	45 (80)	45–46 (81–86)	41–49 (69–96)
Orbit diameter	27 (44)	30 (49)	25–33 (45–52)	28 (49)	25–28 (45–51)	25–28 (43–52)
Postorbital length	34 (57)	32 (52)	30–34 (49–56)	29 (52)	27–31 (50–56)	26–35 (47–58)
Postrostral length	60 (–)	61 (–)	55–63 (–)	56 (–)	54–55 (–)	51–60 (–)
Orbit–preopercle distance	35 (59)	35 (58)	32–37 (52–61)	31 (56)	30–35 (55–63)	28–37 (52–62)
Suborbital width	12 (20)	14 (23)	11–13 (18–22)	12 (22)	12–13 (22–23)	11–13 (19–23)
Upper-jaw length	27 (45)	26 (43)	23–32 (40–53)	25 (44)	21–26 (39–47)	19–27 (36–45)
Length of rictus	21 (34)	21 (34)	18–22 (30–36)	18 (31)	16–20 (30–36)	15–21 (28–36)
Length of PMX tooth band	12 (20)	12 (20)	11–13 (19–22)	10 (18)	10–12 (19–21)	9–13 (17–22)
Preoral length	35 (59)	36 (60)	31–40 (52–71)	40 (71)	39–43 (71–80)	33–47 (55–91)
Length of terminal snout scute	– (–)	– (–)	7–10 (12–17)	9 (17)	9–11 (16–20)	7–13 (11–24)
Length of lateral nasal ridge	28 (46)	26 (43)	26–32 (43–57)	32 (56)	30–34 (55–63)	26–37 (44–73)
Length of suborbital ridge	93 (155)	94 (156)	93–96 (148–169)	94 (167)	94–97 (172–178)	92–99 (154–190)
Snout width	25 (41)	24 (39)	25–28 (41–49)	26 (46)	25–27 (46–49)	24–28 (42–54)
Internasal width	18 (30)	16 (27)	17–18 (28–31)	18 (32)	15–16 (28–30)	16–19 (30–37)
Interorbital width	24 (40)	23 (39)	22–25 (39–41)	23 (42)	18–20 (33–36)	19–23 (36–42)
Occipital width	9 (16)	10 (16)	9–11 (15–18)	9 (16)	8 (14–16)	8–11 (14–19)
Body width over P. bases	33 (55)	46 (76)	35–45 (58–78)	38 (67)	37–41 (69–74)	29–50 (54–84)
Body depth at 1D. origin	– (–)	54 (89)	48–51 (83–85)	47 (83)	47–49 (85–91)	41–63 (77–107)
Body depth at A. origin	41 (68)	49 (80)	39–47 (66–78)	44 (77)	38–45 (69–81)	35–58 (64–98)
Pre-V. length	105 (175)	111 (183)	103–115 (170–200)	106 (188)	111–112 (206–208)	103–120 (185–214)
Pre-anus length	137 (229)	136 (225)	133–141 (220–246)	126 (224)	129–130 (238–241)	120–148 (218–260)
Pre-A. length	145 (242)	143 (235)	140–146 (231–260)	133 (236)	136–137 (252–255)	126–155 (227–276)

...Continued on the next page

TABLE 3. (Continued)

Species	<i>C. commutabilis</i>	<i>C. nox</i>		<i>C. smithi</i>		
Type status	HT	HT	PT	HT	NT	
Locality	Philippines	Japan	Japan	Indonesia	Philippines	Japan
Number of specimens	<i>n</i> = 1	<i>n</i> = 1	<i>n</i> = 13	<i>n</i> = 1	<i>n</i> = 5	<i>n</i> = 45
Isthmus–V. distance	30 (50)	30 (49)	26–32 (45–58)	27 (47)	25–30 (47–54)	20–34 (34–58)
Isthmus–anus distance	62 (104)	51 (85)	47–53 (82–88)	47 (84)	– (–)	43–65 (77–110)
Isthmus–A. distance	70 (116)	58 (95)	54–69 (94–124)	53 (95)	48–53 (89–98)	48–74 (87–125)
V.–A. distance	41 (69)	34 (56)	31–40 (54–70)	31 (55)	26–31 (48–57)	25–45 (46–77)
Anus–A. distance	7 (12)	7 (12)	6–10 (10–17)	6 (11)	4–9 (8–16)	4–10 (7–18)
V. length	39 (65)	50 (83)	35–53 (61–87)	39 (70)	33–40 (60–74)	30–44 (53–74)
P. length	44 (73)	39 (64)	35–42 (57–69)	37 (65)	35–37 (64–68)	33–45 (60–76)
Pre-1D. length	108 (180)	110 (181)	106–114 (178–202)	109 (193)	101–106 (183–196)	104–111 (181–209)
Height of 1D.	49 (82)	– (–)	34–43 (61–71)	38 (67)	38–39 (68–73)	37–51 (66–89)
Length of 1D. base	17 (28)	15 (25)	13–17 (22–29)	15 (27)	13–15 (24–27)	13–19 (24–32)
Interdorsal length	27 (45)	28 (46)	21–35 (36–58)	25 (45)	17–20 (31–35)	20–34 (37–58)
Length of gill slit	11 (19)	12 (20)	11–14 (19–23)	12 (22)	12 (22)	8–17 (15–29)
Length of posterior nostril	9 (15)	7 (12)	7–8 (11–14)	10 (17)	7–9 (14–16)	7–10 (13–18)
Barbel length	9 (14)	10 (17)	8–13 (14–22)	6 (10)	6–8 (11–14)	6–10 (11–17)
Counts						
1D. rays	II,9	II,8	II,8–9	II,9	II,8–9	II,8–10
P. rays	i19–i20	i16–i17	i16–i17	i17	i14–i18	i16–i20
V. rays	7	7	7	7	7	7
Outer GR on first arch	0	0	0	0	0	0
Inner GR on first arch	8	8	7–9	7	8	7–8
Outer GR on second arch	6	6	5–6	6	6	6–7
Inner GR on second arch	7	8	7–9	9	7–9	7–9
Longitudinal scales	39	35	32–41	42	37–43	31–45
Scales below 1D. origin	5.5	4.5	4–6.5	4.5	5.5–6	4.5–7
Scales below 1D. midbase	4.5	3.5	3.5–4.5	4	4–4.5	3.5–5.5
Scales below 2D. origin	5.5	4	4–6	4.5	4.5–6	4–5.5
Scales above A. origin	19	16	15–18	18.5	17–18	15–21

*Abbreviations: 1D.—first dorsal fin; 2D.—second dorsal fin; A.—anal fin; GR—gill rakers; HT—holotype; NT—non-type; P.—pectoral fin; PMX—premaxillary; PT—paratype; V.—pelvic fin

short snout (snout length 62–81% PRL vs. 80–86%; preoral length 52–71% PRL vs. 71–80%) and a longer chin barbel (14–22% PRL vs. 10–14%) (see also Table 3), and from Japanese *C. smithi* by its uniformly dark brown

coloration (Fig. 75 vs. distinctly paler ventrally, Fig. 85A–F). The body of Japanese *C. smithi* is occasionally darker in larger specimens than smaller specimens (Fig. 85G), but it is distinctly paler than that of the new species.

FIGURE 76. Holotype of *Coelorinchus nox* **sp. nov.** BSKU 109205, 99.2 mm HL, 346+ mm TL, Okinawa Trough, East China Sea, 607 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

FIGURE 77. Scanning electron micrographs showing a body scale (from the dorsum below the interdorsal space) of *Coelorinchus nox* **sp. nov.** BSKU 109205, holotype, 99.2 mm HL. (A) View from above; (B) oblique view. [Photos: NSMT]

Coelorinchus nox also superficially resembles *C. commutabilis* Smith & Radcliffe in Radcliffe, 1912 from the East Indies, *C. charius* Iwamoto & Williams, 1999 from off Western Australia, and *C. sereti* Iwamoto & Merrett, 1997 from the tropical southwestern Pacific [the holotype of *C. commutabilis* (USNM 72945, 93.7 mm HL) and type specimens of *C. charius* (19 spec., 51.3–98.2 mm HL; including holotype: AMS I.31166-005, 80.6 mm HL) and *C. sereti* (7 spec., 62.1–124 mm HL; including the holotype: MNHN 1994-925, 119 mm HL) were examined in the course of this study]. However, the new species differs most notably from *C. commutabilis*

and *C. charius* in that the occipital scales are covered with needle-like spinules in comb-like ridges (vs. keel-like to knife-like spinules in saw-toothed ridges). In contrast, it differs primarily from *C. sereti* in that the dorsal contour of the snout is moderately concave in lateral view (Fig. 76B vs. slightly convex; Iwamoto & Merrett 1997: fig. 15a), and the snout is progressively attenuated toward its tip when viewed dorsally, giving a sharply pointed appearance to the snout (vs. broadly convex anteriorly so as to give a much broader appearance to the snout; Iwamoto & Merrett 1997: fig. 15 a'). It further differs from *C. commutabilis* in having fewer pectoral-fin rays

FIGURE 78. Photographs showing spinulation of the occipital scales in two species of *Coelorinchus*. (A) *C. nox* **sp. nov.**, BSKU 109205, holotype, 99.2 mm HL; (B) *C. smithi*, BSKU 112398, 94.5 mm HL. Preserved specimens. [Photos: N. Nakayama]

(i16–i17 vs. i19–i20), a slightly larger orbit (45–52% PRL vs. 44%), and a shorter postorbital length (49–56% PRL vs. 57%); from *C. charius* in having a uniformly dark body (distinctly paler ventrally; Iwamoto & Williams 1999: fig. 9); and from *C. charius* and *C. sereti* in having a moderately short terminal snout scute (12–17% PRL vs. 19–24% and 9–13% in *C. charius* and *C. sereti* respectively) and a narrower internasal width (27–31% PRL vs. 31–35% and 33–37% PRL).

***Coelorinchus parallelus* (Günther, 1877)**

[Japanese name: Soroi-hige]

(Figs. 31C–F, 79–81; Table 4; Appendix 3–4F)

Macrurus parallelus Günther, 1877:439 [original description; in part; lectotype: BMNH 1887.12.7.65 (designated herein), from “off Inoshima” (= Eno-shima Island), *Challenger* sta. 232, in 345 fth (631 m); paralectotypes. in BMNH, MNHN, NMI; see the Comments on type specimens]; Günther 1887:125, pl. XXIX, figs. A, A', A", a, a', a", a''' (description; in part, BMNH 1887.12.7.65–66, 2 spec., from Sagami Bay off Eno-shima Island); Wheeler & O’Riordan 1969:95 (comments on type specimens).

Coelorhynchus parallelus: Jordan & Gilbert in Jordan & Starks 1904:618 (description; 1 spec. from Misaki); Franz 1910:26 (3 spec. from Aburatsubo); Jordan *et al.* 1913:418 (listed; Japan; new Japanese name: “Soroi-hige”); Gilbert & Hubbs 1916:181 (brief description; 4 spec. from East China Sea); Kamohara 1931a:95 (questionable; listed; Kochi); Kamohara 1934a:54 (questionable; listed; Kochi); Okada & Matsubara 1938:451 (in key; Japan); Kamohara 1938:73 (questionable; spec. from Kochi Pref.); Kamohara 1950:276 (questionable; listed; Kochi and Wakayama Pref.); Uchinomi 1951:76 (questionable; listed; southern part of Kii Peninsula); Kamohara 1952:99 (questionable; spec. from Kochi Pref.); Matsubara 1955:1313 (in key; Japan); Kamohara 1958:73 (questionable; listed; Kochi Pref.); Kamohara 1964:96 (questionable; listed; Kochi Pref.); Okamura 1970a:198, pl.

XLIII, text-fig. 86 (description; 2 spec. from Sagami Bay and East China Sea; erroneously considered type locality New Zealand); Tominaga & Uyeno 1981:488 (listed; Japan).

Coelorhynchus (Oxymacrus) parallelus: Okamura 1970b: table 1 (listed; Japan).

Coelorinchus sparsilepis Okamura & Yatou, 1984:241, fig. 170 (original description; holotype: BSKU 28441, from Okinawa Trough, 29°46.0'N, 127°59.0'E, in 710–713 m; new Japanese name: “Mabara-hige”); Okamura 1988:453, pl. 373, fig. K (compiled); Nakabo & Kai 2013:509 (in key; Japan); Motomura 2020:38 (listed; Japan).

Coelorinchus parallelus: Yatou 1984:245, 370, fig. 171 (brief description; 9 spec. from Okinawa Trough; photo based on BSKU 29896); Okamura 1984b:97, pl. 83, fig. I (compiled); Okamura 1988:97, pl. 83, fig. I (compiled); Iwamoto 1990:178, fig. 404 (synopsis); Okamura 1997:127, fig. 20 (compiled); Shao *et al.* 2008b: table 2 (3 spec. listed from southwestern Taiwan); Nakabo & Kai 2013:509 (in key; Japan); Iwamoto *et al.* 2015:65 (brief description; 6 spec. from northeastern and southwestern Taiwan, and South China Sea); Motomura 2020:38 (listed; Japan).

Caelorinchus parallelus: Nakabo 1993:369 (in key; Japan); Shao 1993:170 (compiled; Taiwan); Shinohara & Matsuura 1997:291 (listed; Suruga Bay); Nakabo 2000:433 (in key; Japan); Shinohara *et al.* 2001:304 (2 spec. listed from Tosa Bay); Nakabo 2002:433 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Shinohara *et al.* 2005:417 (listed; Ryukyu Islands); Senou *et al.* 2006:420 (listed; Sagami Sea).

Caelorinchus sparsilepis: Nakabo 1993:369 (in key; Japan); Nakabo 2000:433 (in key; Japan); Nakabo 2002:433 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Shinohara *et al.* 2005:417 (listed; Ryukyu Islands).

Caelorinchus (Oxygadus) parallelus: Chiou *et al.* 2004a: table 1 (listed; Taiwan).

Coelorinchus commutabilis (not Smith & Radcliffe in Radcliffe 1912): Shao *et al.* 2008b: table 2 [2 spec. listed from South China Sea (as first record from Taiwan)].

Diagnosis. No external evidence of light organ anterior to anus. Underside of head, including ventral surfaces of preopercles, completely and uniformly scaled except for anterior portions of mandibular rami; dorsal surface of snout fully scaled; nasal fossa usually heavily scaled anteroventrally. Snout long, sharply pointed, length 67–89% PRL; dorsal profile of snout almost straight to slightly concave in lateral view; terminal scute moderately long, diamond-shaped, dorsoventrally flattened; lateral nasal ridge incompletely supported by nasal bone. Anus abutting anal-fin origin. Premaxillary teeth small, conical, arranged in short, uniformly wide band, none especially enlarged; posterior margin of tooth band falling far short of lateral corner of mouth. Body scales covered with stout, broadly trihedral, shield-like spinules in parallel ridge-like rows; middle row notably enlarged and higher, giving striated appearance to body surface; lateral rows often incomplete, not extending to posterior scale margin; lateral buttresses broadly developed. Transverse scale rows below first dorsal-fin origin 4.5–7. Occipital scales covered with single crest-like row of short, keel-like spinules; no prominent naked area atop head between parietal ridges. Modified scales on head ridges prominently spiny and thickened. Free neuromasts on underside of head not especially enlarged. Height of first dorsal fin 48–77% PRL. Head and body uniformly dark brown; orbit narrowly edged in black; all fins blackish.

Material examined. 48 specimens. **Lectotype of *Macrurus parallelus*** (designated herein): BMNH 1887.12.7.65 (93.2 mm HL, 325+ mm TL), off Eno-shima Island, Kanagawa Pref., Sagami Bay, Japan, 35.1833°N, 139.4667°E, 345 fms (631 m), *Challenger* sta. 232, trawl and dredge, 12 Mar. 1875. **Paralectotype of *M. parallelus*** (designated herein): BMNH 1887.12.7.66 (102 mm HL, 335+ mm TL), collected with lectotype. **Holotype of *Coelorinchus sparsilepis***: BSKU 28441 (73.3 mm HL, 227+ mm TL), northwest of Amami-oshima Island, Kagoshima Pref., Okinawa Trough, Japan, East China Sea, 29.7667°N, 127.9833°E, 713–770 m, F/V *Ryoan-maru*, No. 28, tr. 73, bottom trawl, coll. Y. Kinoshita and S. Hagino, 20 Mar. 1978. **Non-types: Japan:** BSKU 26327 (1, 52.8 mm HL, 277+ mm TL), off Senkaku Islands, Okinawa Trough, 25.6283°N, 122.8933°E, 560–692 m, F/V *Yuryo-maru*, No. 8, tr. 3, bottom trawl, coll. T. Kitajima *et al.*, 14 Jan. 1978; BSKU 57860 (1, 102 mm HL, 394+ mm TL), north of Miyako-jima Island, Okinawa Trough, 26.2500°N, 125.2833°E, 955–991 m, R/V *Tansei-maru*, cr. KT-02-03, sta. E-2, trawl, coll. H. Endo, 26 Apr. 2002; BSKU 26557 (1, 61.8 mm HL, 221 mm TL), BSKU 26559 (1, 70.6 mm HL, 251+ mm TL), off Miyako-jima Island, Okinawa Trough, 26.4633°N, 124.8483°E, 680–770 m, F/V *Yuryo-maru*, No. 8, tr. 10, bottom trawl, coll. S. Kishida and Y. Kinoshita, 23 Jan. 1978; BSKU 26427 (1, 52.3 mm HL, 185+ mm TL), BSKU 26428 (1, 71.2 mm HL, 273 mm TL), north of Miyako-jima Island, Okinawa Trough, 26.3350°N, 124.8450°E, 1000–1140 m,

F/V *Yuryo-maru*, No. 8, tr. 7, bottom trawl, coll. S. Kishida and Y. Kinoshita, 22 Jan. 1978; BSKU 26498 (1, 137 mm HL, 502+ mm TL), BSKU 26499 (1, 90.2 mm HL, 330+ mm TL), BSKU 26500 (1, 103 mm HL, 357+ mm TL), northwest of Miyako-jima Island, Okinawa Trough, 25.2850°N, 124.7800°E, 910–990 m, F/V *Yuryo-maru*, No. 8, tr. 8, bottom trawl, coll. S. Kishida and Y. Kinoshita, 22 Jan. 1978; BSKU 29896 (1, 96.9 mm HL, 321+ mm TL), west of Yaku-shima Island, Okinawa Trough, 30.7067°N, 128.4683°E, 790 m, F/V *Yuryo-maru*, No. 8, sta. 6-T100, bottom trawl, coll. S. Tawara, 16 Jan. 1980; BSKU 26670 (1, 119 mm HL, 457+ mm TL), west of Yaku-shima Island, Okinawa Trough, 30.0300°N, 128.3650°E, 900 m, F/V *Yuryo-maru*, No. 8, tr. 16-1, bottom trawl, coll. Y. Kinoshita and S. Hagino, 3 Feb. 1978; BSKU 29883 (1, 87.0 mm HL, 298+ mm TL), west of Yaku-shima Island, Okinawa Trough, 30.2067°N, 128.2833°E, 780–810 m, F/V *Yuryo-maru*, No. 8, sta. 3-T1, bottom trawl, coll. H. Maeda, 26 Oct. 1979; BSKU 33463 (1, 103 mm HL, 377+ mm TL), BSKU 33466 (1, 99.3 mm HL, 360+ mm TL), BSKU 33467 (1, 92.3 mm HL, 311+ mm TL), BSKU 33468 (1, 85.5 mm HL, 284+ mm TL), BSKU 33470 (1, 93.1 mm HL, 345+ mm TL), BSKU 33471 (1, 79.8 mm HL, 285+ mm TL), BSKU 33472 (1, 79.8 mm HL, 275+ mm TL), BSKU 33474 (1, 75.3 mm HL, 271+ mm TL), BSKU 33476 (1, 89.7 mm HL, 288+ mm TL), BSKU 33478 (1, 71.0 mm HL, 249+ mm TL), BSKU 33480 (1, 59.3 mm HL, 210+ mm TL), west of Yaku-shima Island, Okinawa Trough, 30.2067°N, 128.2833°E, 780–810 m, F/V *Yuryo-maru*, No. 8, sta. 3, tr. T1, bottom trawl, coll. H. Maeda, 26 Oct. 1979; BSKU 28548 (1, 115 mm HL, 385+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.8167°N, 128.1000°E, 915–932 m, F/V *Ryoan-maru*, No. 28, tr. 75, bottom trawl, coll. Y. Kinoshita and S. Hagino, 20 Mar. 1978; BSKU 28507 (1, 75.6 mm HL, 247+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.7333°N, 128.0500°E, 815 m, F/V *Ryoan-maru*, No. 28, tr. 74, bottom trawl, coll. Y. Kinoshita and S. Hagino, 20 Mar. 1978; BSKU 28440 (1, 51.0 mm HL, 165+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.7667°N, 127.9833°E, 713–770 m, F/V *Ryoan-maru*, No. 28, tr. 73, bottom trawl, coll. Y. Kinoshita and S. Hagino, 20 Mar. 1978; BSKU 26989 (1, 113 mm HL, 432+ mm TL), BSKU 26991 (1, 116 mm HL, 431+ mm TL), BSKU 26993 (1, 90.3 mm HL, 296+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.6467°N, 127.9733°E, 830 m, F/V *Yuryo-maru*, No. 8, tr. 19, bottom trawl, coll. Y. Kinoshita and S. Hagino, 4 Feb. 1978; BSKU 26949 (1, 98.9 mm HL, 334+ mm TL), BSKU 26950 (1, 67.9 mm HL, 223+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.6483°N, 127.9333°E, 750–760 m, F/V *Yuryo-maru*, No. 8, tr. 18, bottom trawl, coll. Y. Kinoshita and S. Hagino, 4 Feb. 1978; BSKU 27115 (1, 81.6 mm HL, 283+ mm TL), BSKU 27116 (1, 91.0 mm HL, 312+ mm TL), BSKU 27117 (1, 69.7 mm HL, 251+ mm TL),

northwest of Amami-oshima Island, Okinawa Trough, 29.3833°N, 127.5150°E, 650 m, F/V *Yuryo-maru*, No. 8, tr. 21, bottom trawl, coll. Y. Kinoshita and S. Hagino, 5 Feb. 1978; SNFR 15177 (1, 60.2 mm HL, 205+ mm TL), East China Sea, 29.3050°N, 127.4933°E, 649–663 m, FRV *Yoko-maru*, sta. H-2, tr. 13, coll. T. Irie *et al.*, 11 Sept. 1993; BSKU 28066 (1, 80.6 mm HL, 279+ mm TL), BSKU 28067 (1, 87.1 mm HL, 272+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 28.8833°N, 127.3000°E, 820–830 m, F/V *Ryoan-maru*, No. 28, tr. 60, bottom trawl, coll. Y. Kinoshita and S. Hagino, 16 Mar. 1978; BSKU 23132 (1, 69.9 mm HL, 250+ mm TL), Tosa Bay, 33.1500°N, 133.6600°E, 615 m, FRV *Kaiyo-maru*, cr. SK-75-2, sta. 10, tr. 1, 3-m beam trawl, 25 Jan. 1975; BSKU 69277 (1, 70.6 mm HL, 235+ mm TL), Tosa Bay, 33.1417°N, 133.6433°E, 582–597 m, FRV *Kotaka-maru*, otter trawl, 16 Jun. 1999; BSKU 46137 (1, 123 mm HL, 430+ mm TL), Tosa Bay, 33.0400°N, 133.6098°E, 675–708 m, FRV *Kotaka-maru*, otter trawl, 22 May 1989; BSKU 44784 (1, 108 mm HL, 413+ mm TL), Tosa Bay, 700 m, FRV *Kotaka-maru*, otter trawl, 24 May 1988; MSM 13-11 (1, 118 mm HL, 455+ mm TL), off Miho, Suruga Bay, 35.0032°N, 138.5430°E, 1282 m, 1.3-m larval net, 21 Feb. 2012; BSKU 14176 (1, 102 mm HL, 398+ mm TL), Sagami Bay, coll. O. Okamura, date unknown; NSMT-P 101339 (1, 117 mm HL, 393+ mm TL), off Yoron-jima Island, East China Sea, 27.1133°N, 128.6983°E, 870–945 m, R/V *Hakuho-maru*, cr. KH-73-02, sta. 4, 3-m SA beam trawl, 24 Feb. 1973. **South China Sea:** *ASIZP 66785 (1, 60.7 mm HL, 212+ mm TL), South China Sea, 20.8333°N, 117.4500°E, 731 m, otter trawl, coll. H.-M. Yeh *et al.*, 17 Aug. 2005.

Redescription based on Japanese specimens.

General features are shown in Figs. 79–81. Counts and measurements are given in Table 4. Body deepest at first dorsal-fin origin, gradually tapering to long, laterally compressed tail. Trunk short, moderately compressed, width over pectoral-fin bases 1.0–1.3 in depth below first dorsal-fin origin. Head large, HL about 3.1–3.9 in TL (5.1 in BSKU 26327, 52.8 mm HL). Snout long, acutely pointed, protruding well beyond upper jaw, length 1.4–2.0 times as long as orbit diameter; dorsal contour of snout almost straight to slightly concave in lateral view; lateral nasal ridge incompletely supported by nasal bone (often difficult to determine without radiographs or dissection, especially in large specimens); anterolateral margins of snout slightly convex when viewed dorsally. Orbit large, its greatest diameter 1.1–1.4 in postorbital length. Interorbital space broad, almost flat, width 1.0–1.3 in orbit diameter. Mouth moderately small, inferior, protrusible, upper-jaw length 0.9–1.3 in orbit diameter; posterior margin of maxilla extending to vertical through hind 1/3 of orbit or beyond (but not reaching hind rim of orbit); lateral corner of mouth slightly restricted by skin folds; lips thin, slightly papillose near tooth bands. Suborbital region divided by stout, longitudinal bony ridge passing

from tip of snout to posteroventral angle of opercle; its upper half almost vertical, lower half sharply inclined mesially. Preopercle large, posterior margin inclined, forming moderately angular lobe at posteroventral corner. Subopercle produced posteroventrally into narrowly pointed flap; posterior margin deeply concave. Gill membranes broadly connected across, and attached mesially to isthmus, with free posterior fold. Outer gill slit moderately restricted by skin folds, length 1.9–2.6 in orbit diameter. Gill rakers small, low, tubercular, armed with short, fine spines; no rakers on outer side of first arch and inner side of fourth arch; gill filaments moderately long. Chin barbel short, slender; length 2.3–4.4 in orbit diameter.

Anus abutting anal-fin origin. Periproct poorly developed. No external evidence of light organ anterior to anus. Pyloric caeca 12 (based on BMNH 1887.12.7.66, paralectotype).

Teeth small, conical, sharp, gently incurved, in narrow bands in both jaws. Premaxillary teeth in short, uniformly wide band, with about 5 tooth rows anteriorly; no teeth significantly enlarged; posterior margin of tooth band falling far short of lateral corner of mouth; teeth becoming progressively smaller inwardly. Mandibular teeth uniformly small, in narrow tapered band, with about 4 tooth rows across widest point near symphysis; posterior margin of tooth band reaching lateral corner of mouth. All teeth deeply embedded in thick layer of gum papillae.

Body scales large, thin, adherent, covered with stout, broadly trihedral, shield-like spinules in parallel to subparallel rows (Fig. 31C–F); in 122 mm HL specimen (BSKU 46137), those on dorsum below interdorsal space with 3–5 rows of spinules; middle row with about 5 spinules, notably higher and enlarged compared with adjacent rows, giving striated appearance to body surfaces; lateral rows often incomplete, not extending to posterior scale margin; spinules forming angle of about 45° to scale surface, greatly overlapping and closely adjoined to one another; height of spinules increasing posteriorly, with last spinule extending moderately beyond posterior scale margin (when rows complete); transverse buttresses well developed on each side of basal portion of each spinule; reticulate structures absent on unexposed portion. Body fully scaled except for fins.

Scales on head ridges prominently spiny and thickened; those on median rostral ridge usually covered with radiating rows of spinules. Terminal snout scute prominent, moderately long, diamond-shaped, dorsoventrally flattened, its length 2.1–3.7 in orbit diameter. Supraoccipital and posttemporal scutes stout, prominent, armed with single row of spinules. Scales on dorsal and lateral surfaces of head (including those between parietal ridges) covered with uniserial row of short, erect, keel-like spinules; spinules increasing in height posteriorly, broadly connected basally to one another, forming crest-like ridges; scales on opercle,

FIGURE 79. Fresh specimens of *Coelorinchus parallelus*. (A) BSKU 46137, 123 mm HL, 430+ mm TL, Tosa Bay, Kochi Pref., Pacific, 675–708 m depth; (B) BSKU 44784, 108 mm HL, 413+ mm TL, Tosa Bay, Pacific, 700 m depth. Lateral views. [Photos: BSKU]

FIGURE 80. Lectotype of *Macrurus parallelus* (= *Coelorinchus parallelus*). BMNH 1887.12.7.65, 93.2 mm HL, 325+ mm TL, Sagami Bay, Pacific, 631 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

TABLE 4. Morphometric measurements and meristic counts* for *Coelorinchus parallelus*.

Nominal species	<i>C. parallelus</i>			<i>C. sparsilepis</i>
Type status	LT	PLT	NT	HT
Number of specimens	<i>n</i> = 1	<i>n</i> = 1	<i>n</i> = 44	<i>n</i> = 1
Measurements (mm)				
TL	325+	335+	165+–502+	227+
HL	93.2	102	51.0–137	73.3
% of HL (% of PRL)				
Snout length	44 (80)	47 (87)	41–47 (67–89)	47 (87)
Orbit diameter	26 (46)	25 (46)	24–28 (41–50)	25 (46)
Postorbital length	31 (55)	29 (55)	28–35 (50–58)	29 (55)
Postrostral length	56 (–)	54 (–)	53–61 (–)	54 (–)
Orbit–preopercle distance	30 (54)	30 (56)	30–61 (53–62)	30 (56)
Suborbital width	14 (26)	14 (27)	12–15 (21–27)	13 (24)
Upper-jaw length	21 (38)	22 (41)	20–27 (36–46)	21 (39)
Length of rictus	14 (25)	17 (32)	15–26 (28–39)	16 (30)
Length of PMX tooth band	8 (14)	8 (15)	7–13 (14–22)	9 (17)
Preoral length	43 (78)	43 (80)	35–45 (59–84)	45 (85)
Internasal width	20 (35)	18 (34)	18–21 (31–35)	18 (34)
Length of terminal snout scute	11 (20)	– (–)	7–11 (12–22)	– (–)
Length of lateral nasal ridge	34 (60)	36 (67)	27–36 (44–67)	35 (64)
Length of suborbital ridge	93 (168)	96 (179)	92–97 (152–177)	97 (180)
Snout width	30 (54)	30 (56)	26–31 (44–57)	29 (54)
Interorbital width	22 (40)	23 (42)	21–26 (37–44)	20 (38)
Occipital width	– (–)	– (–)	7–11 (13–18)	– (–)
Body width over P. bases	44 (80)	45 (84)	34–52 (61–89)	41 (76)
Body depth at 1D. origin	49 (87)	44 (82)	42–58 (75–95)	42 (78)
Body depth at A. origin	46 (82)	– (–)	31–48 (56–84)	34 (64)
Pre-V. length	117 (210)	112 (209)	101–117 (170–208)	101 (188)
Pre-anus length	– (–)	143 (268)	130–161 (222–274)	– (–)
Pre-A. length	153 (273)	148 (277)	133–168 (228–282)	137 (255)
Isthmus–V. distance	28 (50)	27 (50)	22–33 (38–56)	22 (40)
Isthmus–anus distance	– (–)	– (–)	48–83 (84–140)	– (–)
Isthmus–A. distance	– (–)	– (–)	52–90 (90–151)	– (–)
V.–A. distance	43 (77)	40 (74)	31–60 (55–101)	37 (69)
Anus–A. distance	– (–)	5 (9)	2–7 (3–12)	– (–)
V. length	31 (55)	29 (54)	25–42 (44–71)	33 (62)
P. length	31 (56)	– (–)	31–40 (54–69)	35 (65)
Pre-1D. length	107 (192)	111 (207)	105–114 (179–204)	108 (201)
Height of 1D.	– (–)	34 (64)	27–46 (48–77)	36 (66)
Length of 1D. base	15 (26)	13 (24)	12–17 (21–29)	13 (25)
Interdorsal length	17 (31)	17 (32)	10–24 (18–39)	10 (19)
Length of gill slit	12 (21)	12 (23)	10–14 (18–23)	11 (20)
Length of posterior nostril	7 (13)	7 (13)	5–9 (10–15)	6 (11)
Barbel length	8 (14)	7 (14)	6–11 (10–18)	7 (13)
Counts				
1D. rays	II,8	II,8	II,7–9	II,8
P. rays	i15–i16	i16	i15–i19	i16–i19
V. rays	7	7	7	7
Outer GR on first arch	0	0	0	0
Inner GR on first arch	8	8	7–10	9
Outer GR on second arch	6	6	6–8	6
Inner GR on second arch	7	7	8–10	8
Longitudinal scales	35	37	30–45	40
Scales below 1D. origin	6	6.5	4.5–7	6

...Continued on the next page

TABLE 4. (Continued)

Nominal species	<i>C. parallelus</i>			<i>C. sparsilepis</i>
Type status	LT	PLT	NT	HT
Number of specimens	<i>n</i> = 1	<i>n</i> = 1	<i>n</i> = 44	<i>n</i> = 1
Scales below 1D. midbase	4	5.5	3–6.5	4.5
Scales below 2D. origin	5	5	4.5–6	5
Scales above A. origin	16	19	14–21	–

*Abbreviations: 1D.—first dorsal fin; 2D.—second dorsal fin; A.—anal fin; GR—gill rakers; HT—holotype; LT—lectotype; NT—non-type; P.—pectoral fin; PLT—paralectotype; PMX—premaxillary; V.—pelvic fin

preopercle, and postorbital and supratemporal canals largest. Scales on ventral surfaces of head similar to those on dorsum, but much smaller and not imbricate. Nasal fossa usually heavily scaled anteroventrally (scales rarely absent in some specimens). Dorsal surfaces of snout fully scaled, without naked clefts along each side of median rostral ridge. Underside of head completely and uniformly scaled except for narrow naked band above upper lip and anterior portion of mandibular rami. Posteroventral margins of preopercles only narrowly naked.

No open pores along cephalic sensory canals. Free neuromasts prominent, serially arranged along head canals and lateral line, also sparsely distributed on dorsal surface of body; those on underside of head not especially enlarged; short hair-like papillae scattered over head surfaces (especially on underside of snout). Anterior nostril small and circular; posterior large, bean-shaped; septum between nostrils forming anteriorly opened hood. Lateral line complete, not interrupted throughout.

Origins of first dorsal and pelvic fins posterior to vertical through pectoral-fin base; first dorsal fin moderately high, its height 2.3–3.3 times as long as its base length (1.8 in BSKU 44784, 108 mm HL); second spinous ray not elongate, smooth along its leading edge (small denticle present distally in BSKU 33467, 89.7 mm HL); its tip usually extending beyond second dorsal-fin origin (not reaching this level in BSKU 44784). Interdorsal space narrow, 0.8–1.5 times as long as first dorsal-fin base length. Anal fin originating below or slightly posterior to second dorsal-fin origin (below bases of 1st–5th rays). Outer pelvic-fin ray prolonged, with fine distal tip.

Color when fresh (Fig. 79). Head, body, and fins uniformly dark brown (Fig. 79A) to swarthy (Fig. 79B).

Color in alcohol (Figs. 80–81). Head and body uniformly dark brown; orbit narrowly edged in black; lips dark; oral and gill cavities blackish; gill rakers and filaments light brown, arches darker; all fins blackish.

Size. Attains about 50 cm TL (BSKU 26498, 502 mm TL, Okinawa Trough, Japan).

Distribution. Known from off the Pacific coasts of southern Japan northward to the Miura Peninsula (139.62°E), Okinawa Trough, and northern South China Sea, at depths of 360–1034 m (Gilbert & Hubbs 1916; Shao *et al.* 2008a, 2008b; this study; Appendix 3-4F). Elsewhere reported from the western Coral Sea, and the

Chesterfield and Bellona Plateaus, at depths of 412–1013 m (Weber 1913; Weber & de Beaufort 1929; Iwamoto & Merrett 1997; Merrett & Iwamoto 2000; Last *et al.* 2014). Moderately common in the East China Sea, but rare in other parts of its range.

Comments on type specimens. *Coelorinchus parallelus* was originally described by Günther (1877) from “off Inoshima” (= Enoshima Island, Kanagawa Pref.), Sagami Bay, Japan. Unfortunately, Günther’s (1877) original description is brief, without information on how many specimens he examined. However, in the “*Challenger*” Report, Günther (1887:125–127) provided a more detailed description of the species, noting that eight specimens were collected from “*Hyalonema*-ground, off Inoshima” (*Challenger* sta. 232). Four of these were examined in the course of this study at the Natural History Museum, London (BMNH 1887.12.7.65–68, 221+–355+ mm TL; Fig. 80), two of which (BMNH 1887.12.7.67–68, 221+–355+ mm TL) were re-identified as *C. gilberti* Jordan & Hubbs, 1925. According to Wheeler & O’Riordan (1969), a 314 mm TL specimen is now deposited in the National History Museum of Ireland, Dublin (NMI block reg. no. 114.1899, fide Fricke *et al.* 2020). Another 280 mm TL specimen has also been transferred to the Muséum national d’Histoire naturelle, Paris (MNHN 1890-0125; Bailly *et al.* 1999; Fricke *et al.* 2020). The whereabouts of the remaining two specimens are unknown. To fix the name *C. parallelus*, BMNH 1887.12.7.65 (which represents what many authors have called *C. parallelus*) is herein designated as a lectotype of this species (Fig. 80), and the other specimens used in the original description automatically become paralectotypes.

Günther (1887) reported additional *Challenger* specimens of *C. parallelus* from off New Zealand (1 spec. from sta. 169), off the Kermadec Islands (3 and 2 spec. from sta. 170A and 171 respectively), and the south of Japan (1 spec. from sta. 235), but these specimens are not included in the type series of this species. Five of them are present in the BMNH collection; none represents the true *C. parallelus*, but rather *C. gilberti* (BMNH 1887.12.7.69, 1 spec., 136+ mm TL, from sta. 235; Günther 1887: pl. XXIX, fig. A’), *C. kermadecus* Jordan & Gilbert in Jordan & Starks, 1904 (BMNH 1887.12.7.60–62, 3 spec., 176+–549+ mm TL, from sta. 170A; Günther 1887: pl. XXIX, fig. A), and *C. trachycarus* Iwamoto, McMillan &

FIGURE 81. Preserved specimen of *Coelorinchus parallelus*. BSKU 28441, holotype of *C. sparsilepis*, 73.3 mm TL, 227+ mm TL, Okinawa Trough, East China Sea, 770–713 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

Shcherbachev, 1999 (BMNH 1887.12.7.59, 1 spec., 448+ mm TL, from sta. 169).

Nomenclatural discussion. Okamura & Yatou (1984) described *C. sparsilepis* based on a single specimen (Fig. 81) collected from the Okinawa Trough, Japan. Despite its close similarity to *C. parallelus* and their sympatric distribution in the area, Okamura & Yatou (1984) did not compare the two species. Yatou (1984:224–225, 370) also reported *C. parallelus* in the same book without comments on *C. sparsilepis*. The close similarity between *C. sparsilepis* and *C. parallelus* was first recognized by Iwamoto & Merrett (1997:502), who suggested that the two species are not separable except for the suborbital width (38% of orbit diameter in *C. sparsilepis* vs. 50–62.5% in *C. parallelus*) and orbit diameter (2.1 in snout length vs. 1.7–1.8).

A re-examination of the holotype of *C. sparsilepis* confirmed that the suborbital width is 52% of the orbit diameter, and the orbit is 1.90 in the snout length. Furthermore, these data fall within the ranges of variation for the additional specimens of *C. parallelus* examined (suborbital width 46–59% of orbit diameter; orbit diameter 1.44–1.95 snout length). Additionally, no significant differences were found in other characters, including counts and measurements (Table 4), dentition, and squamation. Consequently, *C. sparsilepis* is herein regarded as a junior synonym of *C. parallelus*.

Remarks. Yanai (1950) listed *C. parallelus* from the Sea of Japan off San'in, noting that the species was commonly found in Matsue and Hamada in Shimane Pref. On the basis of Yanai's list, many subsequent authors (e.g., Lindberg & Legeza 1969; Shinohara *et al.* 2014) believed the species occurred in the Sea of Japan.

However, considering its deeper vertical distribution (usually >600 m depth), Yanai's *C. parallelus* appears to represent misidentified *C. multispinulosus* Katayama, 1942, the only species of grenadiers commonly found in the southern part of the Sea of Japan. In fact, no voucher specimens of *C. parallelus* collected from the area were found in museum collections.

Ueno (1965, 1971) reported *C. parallelus* from off the Pacific coast of Hokkaido, but it is most likely to represent misidentified *C. gilberti*, a species occasionally found in the Pacific off northern Japan. Maeda & Maruyama (1991), Maeda & Tsutsui (2003), and Nakabo & Kai (2013:1874) also doubted Ueno's records of *C. parallelus*.

According to Okamura (1970a:198), other records of *C. parallelus* by Japanese ichthyologists before the 1970s should be considered suspect, and probably represent misidentifications of *C. kamoharai* Matsubara, 1943 or *C. multispinulosus*.

Günther (1887) reported *C. parallelus* from Kermadec Island and New Zealand, but Jordan & Gilbert (*in* Jordan & Starks 1904:618–619) considered Günther's (1887) additional specimens specifically distinct from the Japanese population, providing a new scientific name, *C. kermadecus*, for the former. Subsequently, in their synonymy of *C. parallelus*, Gilbert & Hubbs (1920:515) suggested that the New Zealand specimen represents another undescribed species. The species is now known as *C. trachycarus* widely distributed in Australasian waters.

Weber (1913) reported *C. parallelus* from Indonesia, including the Savu, Flores, Ceram, and Halmahera Seas. However, Gilbert & Hubbs (1920:151) suggested that Weber's (1913) specimens have a much shorter snout than

the Japanese material they examined. Weber (1913: pl. IV, fig. 3) also illustrated his *C. parallelus* as lacking scales on the nasal fossa. However, the fossa of Japanese specimens is heavily scaled ventrally, which further implies that Weber's specimens represent a different species.

A Hawaiian record of *C. parallelus* by Gilbert & Cramer (1897) was re-identified as *C. aratum* Gilbert, 1905 in the original description of the latter species.

Smith's (1953) *C. parallelus* from the western Indian Ocean is attributed to a misidentification of *C. braueri* Barnard, 1925. His illustration (fig. 232) clearly shows important diagnostic features of that species, including an elongate spinous ray of the first dorsal fin, a long, attenuated snout, and the dorsal contour of the snout greatly convex in lateral view.

Kyushin *et al.* (1973) recorded *C. parallelus* from the Andaman Sea, but his specimen (HUMZ 90186) is re-identified here as *Coelorinchus* sp. cf. *kermadecus* in this study.

The following records of *C. parallelus* from the Atlantic and Indian Oceans seem to represent other species of the subgenus *Oxygadus* Gilbert & Hubbs, 1920 (see also Iwamoto 1990:179): Alcock (1889, 1899) from the Gulf of Manner; Brauer (1906) from the South Atlantic; Zugmayer (1911) from the Mid-Atlantic Ridge in the North Atlantic; and Fowler (1936) from the eastern Atlantic.

Relationships and comparisons. *Coelorinchus parallelus* is a distinctive species of the *C. parallelus* group (see the Relationships of *C. divergens*), and differs most notably from other members of that group in that the occipital scales are covered with a single crest-like row of spinules (vs. multiple divergent rows of spinules). In addition, it differs notably from *C. acanthiger* Barnard, 1925 in having fewer scales below the first dorsal-fin origin (4.5–7 vs. 7–10) and lacking a prominent naked area atop the head between the parietal ridges; from *C. amirantensis* Iwamoto, Golani, Baranes & Goren, 2006 and *C. yurii* Iwamoto, Golani, Baranes & Goren, 2006 in that the underside of the head is densely scaled (vs. sparsely scaled); from *C. braueri* and *C. yurii* in having a lower first dorsal fin (height 48–77% PRL vs. >98%); from *C. mycterismus* McMillan & Paulin, 1993 in that the dorsal contour of the snout is almost straight to slightly concave in lateral view (vs. distinctly concave); and from *C. trachycarus* in that the ventral surfaces of the preopercles are completely scaled (vs. broadly naked). *Coelorinchus parallelus* also differs from all except *C. divergens* Okamura & Yatou, 1984, *C. kermadecus*, *C. mycterismus*, and *C. spinifer* Gilbert & Hubbs, 1920 in that the nasal fossa is usually heavily scaled (vs. completely or generally naked); and from *C. amirantensis* and *C. spinifer* in having a shorter snout (67–89% PRL vs. >93%).

***Coelorinchus sheni* Chiou, Shao & Iwamoto, 2004**

[Japanese name: Tora-hige]

(Figs. 42C–D, 82–84; Appendix 3-5A)

Coelorinchus sheni Chiou, Shao & Iwamoto, 2004a:37, figs. 1–4 [original description; holotype: ASIZP 61348, from off Tashi (= Da-xi, northeastern Taiwan); 2 paratypes from Taiwan; “*Coelorinchus (Oxygadus) sheni*” in table 1].

Coelorinchus sheni: Shao *et al.* 2008b: table 2 (3 spec. listed from northeastern and southwestern Taiwan); Iwamoto *et al.* 2015:67 (brief description; 13 spec. from northeastern and southeastern Taiwan); Ho & Shao 2011:36 (type catalog); Nakayama *et al.* 2018:122, figs. 1–2 (description; 1 spec. from Japan; comments on type specimens; new Japanese name: “Tora-hige”); Motomura 2020:38 (listed; Japan).

Diagnosis. Light organ externally represented by short, narrow, naked streak immediately anterior to anus (difficult to distinguish in some specimens), its anterior margin falling far short of line connecting inner pelvic-fin bases. Underside of head completely and uniformly covered with scales, except for anterior portions of mandibular rami; top of snout completely scaled; nasal fossa scaled ventrally. Snout long, sharply pointed, length 57–73% PRL, its dorsal profile straight or slightly convex in lateral view; terminal scute pointed distally, arrowhead-shaped; lateral nasal ridge incompletely supported by nasal bone. Anus immediately anterior to anal-fin origin, but they slightly separated by a few scale rows. Premaxillary teeth in long, tapered band, with outer series prominently enlarged; posterior margin of tooth band reaching near lateral corner of mouth. Body scales covered with short, reclined, keel-like spinules in widely divergent, coarsely saw-toothed ridges; every spinule row complete, extending to posterior scale margin; spinules in each row overlapping but free at tips, gradually increasing in height posteriorly; buttresses poorly developed. Occipital scales covered with 2–4 divergent rows of spinules. Preoral length 37–59% PRL; barbel length 13–21% PRL. About 6 dark saddles on body (faint in large individuals); ventral margins of saddles extending beyond mid-lateral axis of body; lips paler; oral cavity blackish; gular and branchiostegal membranes dusky.

Material examined. 11 specimens. **Holotype of *Coelorinchus sheni*:** ASIZP 61348 (115 mm HL, 412+ mm TL), off Da-xi, Yilan, northeastern Taiwan, East China Sea, 24.91°N, 122.06°E, 400–600 m depth, F/V *Gin Ton Long*, bottom trawl, coll. by M.-L. Qiu, 30 May 2001. **Paratypes of *C. sheni*:** ASIZP 61232 (1, 249 mm HL, 868+ mm TL), off Lauto, southeastern Taiwan, ca. 22.6667°N, 121.6667°E, rod and reel, 1 Mar. 1999; CAS 215541 (1, 112 mm HL, 424+ mm TL), Da-xi fish market, Yilan, northeastern Taiwan, East China Sea, coll. M.-L. Chiou and H.-C. Ho, date unknown. **Non-types: Japan:** OCF-P 3492 (1, 196 mm HL, 692+ mm TL), Izena Bank, northwest of Ie-jima Island, 27.1343°N, 127.2403°E, 650

m, coll. T. Arakaki, 15 Sept. 2016. **Taiwan:** BSKU 116417 (1, 154 mm HL, 558+ mm TL), Da-xi fish market, Yilan, coll. H.-C. Ho, 18 Jun. 2014; ASIZP 70210 (6, 41.3–98.8 mm HL, 86+–380+ mm TL), Da-xi fish market, Yilan, coll. M.-L. Qiu, 25 Jun. 2005.

Counts and measurements. Based on 11 specimens (41.3–249 mm HL, 86+–868+ mm TL). Counts: first dorsal-fin rays II,8–9; pectoral-fin rays i17–i20; pelvic-fin rays 7; gill rakers on first arch (outer/inner) 0–3/8–9, on second arch 6–8/9–10; longitudinal scales 33–43; transverse scale rows below first dorsal-fin origin 5.5–6.5, below first dorsal-fin midbase 4–5.5, below second dorsal-fin origin 5–6, above anal-fin origin 20.5–24.5; pyloric caeca 46–58.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 36–43 (57–73); orbit diameter 21–27 (33–44); postorbital length 35–42 (58–66); postrostral length 58–63; orbit–preopercle distance 36–46 (59–76); suborbital width 12–15 (21–25); upper-jaw length 27–37 (45–59); length of rictus 21–31 (36–49); length of premaxillary tooth band 16–22 (28–35); preoral length 24–35 (37–59); length of terminal snout scute 3–6 (5–10); length of lateral nasal ridge 19–27 (30–44); length of suborbital ridge 88–97 (139–163); snout width 23–29 (37–47); internasal width 17–21 (26–34); interorbital width 23–26 (37–43); occipital width 9–11 (14–18); body width over pectoral-fin bases 36–50 (59–79); body depth at first dorsal-fin origin 54–65 (90–107); body depth at anal-fin origin 43–54 (71–89); prepelvic length 102–114 (168–192); preanus length 129–148 (214–244); preanal length 133–154 (222–252); isthmus–pelvic distance 28–39 (46–64); isthmus–anus distance 52–74 (86–122); isthmus–anal distance 58–80 (96–131); pelvic–anal distance 32–46 (52–75); anus–anal distance 5–46 (7–72); pelvic-fin length 33–56 (54–93); pectoral-fin length 45–52 (77–86); predorsal length 105–

112 (173–183); height of first dorsal fin 42–50 (69–82); length of first dorsal-fin base 17–21 (29–34); interdorsal length 12–28 (20–46); length of gill slit 13–19 (22–32); length of posterior nostril 7–9 (12–15); barbel length 8–13 (13–21).

Size. Attains about 87 cm TL (ASIZP 61232, paratype, 868+ mm TL, southwestern Taiwan).

Distribution. Known from the southern East China Sea off Taiwan and Japan, and also southeastern Taiwan (Appendix 3-5A). Depth range 400–650 m (based on specimens associated with reliable depth records; see Nakayama *et al.* 2018). Very rare.

Remarks. *Coelorinchus sheni* has been well described by previous authors, and thus does not need repeating here. This species was originally described by Chiou *et al.* (2004a) based on three specimens collected from Taiwan (Figs. 83–84). Recently, Iwamoto *et al.* (2015) reported additional specimens from the type locality (Da-xi fish market), and Nakayama *et al.* (2018) newly recorded this species from Japanese waters based on a single specimen collected from the East China Sea off Ie-jima Island, Okinawa Pref. Nakayama *et al.* (2018) also emended localities of type specimens of the species. One of the paratypes (ASIZP 61232), which was previously not found among the collection of the Academia Sinica, Taipei (Nakayama *et al.* 2018), was rediscovered during the author's visit in November 2017 (Fig. 84).

Relationships and comparisons. *Coelorinchus sheni* is a member of the *C. tokiensis* group (see the Relationships of *C. hexafasciatus*), and most closely related to *C. quadricristatus* (Alcock in Wood-Mason & Alcock, 1891), which is distributed in the Indian Ocean. The two species are unique within the group in that the terminal snout scute is sharply pointed. They also differ from all but *C. celanostoma* McMillan & Paulin, 1993 and *C. osipullus* McMillan & Iwamoto, 2009 in that the

FIGURE 82. Fresh specimen of *Coelorinchus sheni*. OCF-P03492, 196 mm HL, 692+ mm TL, northwest of Ie-jima Island, Okinawa Pref., East China Sea, 650 m depth. Lateral view. [Photo: reproduced from Nakayama *et al.* (2018: fig. 1) with permission from the Japanese Society of Systematic Zoology]

FIGURE 83. Holotype of *Caelorinchus sheni* (= *Coelorinchus sheni*). ASIZP 61348, 115 mm HL, 412+ mm TL, Da-xi fish market, northeastern Taiwan, East China Sea. (A) Lateral, (B) dorsal, and (C) ventral views. Preserved condition. [Photos: N. Nakayama]

FIGURE 84. Paratype of *Caelorinchus sheni* (= *Coelorinchus sheni*). ASIZP 61232, 249 mm HL, 868+ mm TL, off Lauto, southeastern Taiwan. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

underside of the head is completely scaled. *Coelorinchus sheni* readily differs from *C. quadricristatus* in having a shorter snout [36–43% HL vs. as long as a half HL (fide Alcock in Wood-Mason & Alcock 1891:119)] and a deeper body [depth below first dorsal-fin origin 1.3–1.6 times as long as snout length vs. “greatest height of the body barely equal to length of the snout” (Alcock 1899:106)]. Furthermore, saddles of *C. sheni* extend ventrally beyond the mid-lateral axis of the body (Fig. 83), whereas those

of *C. quadricristatus* reach only this level (see Alcock 1892: pl. III, fig. 1).

***Coelorinchus smithi* Gilbert & Hubbs, 1920**

[Japanese name: Kishu-hige]

(Figs. 52B, 53E–F, 54B, 85–87; Tables 3, 5; Appendix 3-5B)

Coelorhynchus commutabilis (not Smith & Radcliffe in Radcliffe

1912); Smith & Radcliffe in Radcliffe 1912:128, pl. 29, fig. 2 (original description; part of “typical examples”, Indonesia and Philippines).

Coelorhynchus smithi Gilbert & Hubbs, 1920:493, fig. 20 [original description; holotype: USNM 78212, from “between Gillolo and Makyan Islands”, *Albatross* sta. 5621, 0°15'00"N, 127°24'35"E, in 298 ftm (545 m); 18 paratypes]; Okamura 1963a:27, fig. 5 (first record from Japan; description; 40 spec. from Aichi, Mie, and Kochi Pref.; new Japanese name: “Kishû-hige”); Kamohara 1964:96 (listed; Kochi Pref.); Okamura 1970a:179, pl. XXXIX, text-figs. 75–77 (description; biological notes; 63 spec. from Pacific off southern Japan from Heta to Totoro); Kuroda 1971:56 [listed from Suruga Bay after Okamura (1970a)]; Tominaga & Uyeno 1981:488 (listed; Japan); Yatou 1982:171, 353, fig. 103 (description; 8 spec. from Tosa Bay; photo based on BSKU 29521); Ozawa 1983:13 (listed; off Makurazaki, Kagoshima Pref., East China Sea).

Coelorhynchus (Oxymacrus) smithi: Okamura 1970b: table 1 (listed; Japan).

Coelorrinchus smithi: Yatou 1984:231, 368, fig. 164 (brief description; 10 spec. from Okinawa Trough; photo based on BSKU 29588); Okamura 1984b:97, pl. 83, fig. C (compiled); Okamura 1988:97, pl. 83, fig. C (compiled); Okamura 1997:127, fig. 15 (compiled); Shao *et al.* 2008b: table 2 (8 spec. from northeastern and southwestern Taiwan, and South China Sea); Nakabo & Kai 2013:510 (in key; Japan); Iwamoto *et al.* 2015:67 (brief description; 7 spec. from northeastern and southwestern Taiwan); Motomura 2020:38 (listed; Japan).

Caelorinchus smithi: Nakabo 1993:369 (in key; Japan); Shinohara & Matsuura 1997:291 (listed; Suruga Bay); Nakabo 2000:433 (in key; Japan); Shinohara *et al.* 2001:305 (18 spec. listed from Tosa Bay); Nakabo 2002:433 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Shinohara *et al.* 2005:417 (listed; Ryukyu Islands); Suetsugu & Ohta 2005: table 3 (listed; Enshu-nada); Furuhashi *et al.* 2010: table 2 (38 spec. listed from northern Okinawa Trough); Iwatsuki *et al.* 2017:32 (listed; Hyuga-nada).

Caelorinchus (Oxymacrus) smithi: Chiou *et al.* 2004a: table 1 (listed; Taiwan).

Diagnosis. Light organ externally represented by short naked fossa immediately anterior to anus, its anterior margin falling far short of line connecting inner pelvic-fin bases. Underside of head almost completely scaled except for narrow naked area above upper lip and anterior portions of mandibular rami; dorsal surface of snout fully scaled (narrow naked clefts rarely present along each side of mid rostral ridge). Snout long, sharply pointed, length 69–96% PRL; dorsal profile of snout moderately concave in lateral view; terminal scute moderately short, diamond-shaped, dorsoventrally flattened, length 11–24% PRL; lateral nasal ridge completely supported by nasal bone, only slightly convex in dorsal view, giving attenuated appearance to snout. Anus slightly separated from anal-fin origin. Premaxillary teeth small, conical, arranged in short, uniformly wide band, none distinctly enlarged

(outer series slightly larger); posterior margin of tooth band falling far short of lateral corner of mouth. Body scales covered with short, reclined, keel-like spinules in widely divergent, coarsely saw-toothed ridges; every spinule row complete, extending to posterior scale margin; spinules in each row greatly overlapping, increasing in height posteriorly; buttresses narrowly developed. Occipital scales covered with short, moderately reclined, keel-like (rarely knife-like) spinules in 4 or more widely divergent, saw-toothed rows. Orbit diameter 43–52% PRL; suborbital width 19–23% PRL; internasal width 28–37% PRL; interorbital width 33–42% PRL; pelvic-fin length 53–74% PRL; pectoral-fin length 60–76% PRL; transverse scale rows below first dorsal-fin midbase 3.5–5.5, below second dorsal-fin origin 4–6. Body generally darker dorsally, paler ventrally; about 2 faint saddles dorsally on tail (much prominent in smaller specimens); lips white; oral cavity blackish; gular and branchiostegal membranes pale to dusky; first dorsal, pectoral, and pelvic fins generally dusky to dark.

Material examined. 55 specimens. **Holotype of *Coelorhynchus smithi*:** USNM 78212 (80.7 mm HL, 261+ mm TL), Molucca Sea, between Gilolo and Makyan Islands, Indonesia, western Pacific, 0.2500°N, 127.4097°E, 298 ftm (545 m), *Albatross* sta. 5621, 12-ft Agassiz beam trawl, 28 Nov. 1909. **Non-types: Japan:** KPM-NI 24972 (1, 110 mm HL, 407+ mm TL), off Okinawa, 600 m, long line, coll. K. Yunokawa, 2009; KPM-NI 28333 (1, 135 mm HL, 528+ mm TL), ca. 25 km southeast of Aguni-jima Island, East China Sea, 26.4120°N, 127.3947°E, 774–780 m, R/V *Tansei-maru*, cr. KT-08-33, sta. AG-1, 3-m beam trawl, coll. T. Sato, 17 Dec. 2008; KPM-NI 28341 (1, 71.1 mm HL, 256+ mm TL), KPM-NI 28342 (1, 58.7 mm HL, 207+ mm TL), ca. 9 km east of Aguni-jima Island, East China Sea, 26.5700°N, 127.3307°E, 583–589 m, R/V *Tansei-maru*, cr. KT-08-33, sta. AG-6, 3-m beam trawl, coll. T. Sato, 17 Dec. 2008; BSKU 29588 (1, 102 mm HL, 382+ mm TL), northwest of Miyako-jima Island, Okinawa Trough, 25.7917°N, 124.5117°E, 500–550 m, F/V *Yuryo-maru*, No. 8, sta. 1-T9, bottom trawl, coll. T. Kitajima, 15 Sept. 1979; BSKU 32512 (1, 102 mm HL, 308+ mm TL), northwest of Miyako-jima Island, Okinawa Trough, 25.7917°N, 124.5117°E, 500–550 m, F/V *Yuryo-maru*, No. 8, sta. 1-T9, bottom trawl, coll. M. Hirose and T. Kume, 10 Oct. 1979; BSKU 32583 (1, 84.0 mm HL, 313+ mm TL), northwest of Miyako-jima Island, Okinawa Trough, 25.7900°N, 124.3900°E, 600 m, F/V *Yuryo-maru*, No. 8, sta. 1-T13, bottom trawl, coll. T. Kitajima, 20 Sept. 1979; BSKU 29637 (1, 93.2 mm HL, 332+ mm TL), northwest of Miyako-jima Island, Okinawa Trough, 25.7900°N, 124.3333°E, 600 m, F/V *Yuryo-maru*, No. 8, sta. 1-T13, bottom trawl, coll. T. Kitajima, 20 Sept. 1979; BSKU 29844 (1, 68.1 mm HL, 236+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 30.0115°N, 127.8387°E, 400–420 m, F/V *Yuryo-maru*, No. 8, sta. 3-T4, bottom trawl, coll. H. Maeda, 26 Oct. 1979; BSKU

97991 (1, 45.3 mm HL, 152+ mm TL), south of Muroto, 33.0790°N, 134.1645°E, 414–513 m, R/V *Tansei-maru*, cr. KT-05-29, sta. K-500, beam trawl, coll. H. Endo *et al.*, 17 Nov. 2005; BSKU 77899 (1, 67.6 mm HL, 217+ mm TL), Tosa Bay, 33.2083°N, 133.6433°E, 364–382 m, FRV *Kotaka-maru*, otter trawl, coll. H. Endo, 27 Sept. 1999; BSKU 112076 (1, 82.9 mm HL, 311 mm TL), Tosa Bay, 33.1428°N, 133.6248°E, 450–490 m, FRV *Kotaka-maru*, 1 Aug. 1988; BSKU 69308 (1, 80.0 mm HL, 274+ mm TL), Tosa Bay, 33.1800°N, 133.6217°E, 412–423 m, FRV *Kotaka-maru*, otter trawl, 8 Sept. 1998; BSKU 43829 (1, 56.0 mm HL, 211+ mm TL), Tosa Bay, 400 m, FRV *Kotaka-maru*, otter trawl, 11 Jun. 1987; BSKU 40217 (1, 80.9 mm HL, 239+ mm TL), BSKU 40218 (1, 67.4 mm HL, 210+ mm TL), Tosa Bay, 400 m, FRV *Kotaka-maru*, otter trawl, 25 Jul. 1984; BSKU 51398 (1, 42.9 mm HL, 136+ mm TL), Tosa Bay, 400 m, FRV *Kotaka-maru*, otter trawl, 27 Sept. 1999; BSKU 65947 (1, 97.2 mm HL, 346+ mm TL), BSKU 97954 (1, 74.7 mm HL, 275+ mm TL), Tosa Bay, 400 m, FRV *Kotaka-maru*, otter trawl, 9 May 2000; BSKU 40335 (1, 77.8 mm HL, 271+ mm TL), Tosa Bay, 400 m, FRV *Kotaka-maru*, otter trawl, 9 Oct. 1984; BSKU 10041 (1, 64.3 mm HL, 201+ mm TL), BSKU 10045 (1, 66.9 mm HL, 239+ mm TL), BSKU 10046 (1, 59.3 mm HL, 214 mm TL), Mimase fish market, coll. T. Kamohara, 10 Feb. 1961; BSKU 4382 (1, 49.4 mm HL, 179+ mm TL), BSKU 4383 (1, 75.7 mm HL, 264+ mm TL), Mimase fish market, coll. T. Kamohara, 19 Dec. 1954; BSKU 8834 (1, 84.9 mm HL, 304+ mm TL), Mimase fish market, coll. T. Kamohara, 30 Jan. 1951; BSKU 12747 (1, 65.5 mm HL, 226+ mm TL), Mimase fish market, coll. Y. Mishima (= Y. Machida), 15 Feb. 1967; BSKU 13782 (1, 64.2 mm HL, 246 mm TL), Mimase fish market, coll. Y. Mishima (= Y. Machida), Jan.–Mar. 1968; BSKU 106852 (1, 77.0 mm HL, 281 mm TL), Mimase fish market, F/V *Kosei-maru*, 15 Apr. 2012; BSKU 102699 (1, 51.6 mm HL, 180+ mm TL), BSKU 102700 (1, 51.5 mm HL, 157+ mm TL), BSKU 102702 (1, 59.8 mm HL, 192+ mm TL), BSKU 102703 (1, 56.4 mm HL, 199+ mm TL), BSKU 102705 (1, 60.0 mm HL, 207+ mm TL), Mimase fish market, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama, 21 Feb. 2010; BSKU 102690 (1, 72.9 mm HL, 267+ mm TL), Mimase fish market, F/V *Kosei-maru*, tr. 3, bottom trawl, coll. N. Nakayama, 21 Feb. 2010; BSKU 13918 (1, 64.4 mm HL, 232+ mm TL), Mimase fish market, Jan. 1968; BSKU 13881 (1, 66.1 mm HL, 229+ mm TL), Mimase fish market, Mar. 1968; BSKU 44652 (1, 58.7 mm HL, 216 mm TL), off Okitsu, Tosa Bay, 200–230 m, 20 Apr. 1988; BSKU 29521 (1, 79.0 mm HL, 271+ mm TL), Tosa Bay, 32.9833°N, 133.4833°E, 453 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, cr. 4-Ky, sta. T18, bottom trawl, coll. O. Okamura *et al.*, 21 Dec. 1979; BSKU 30481 (1, 74.2 mm HL, 254+ mm TL), Tosa Bay, 32.9833°N, 133.4833°E, 453 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, tr. 17, beam trawl, coll. O. Okamura *et al.*, 21 Dec. 1979; BSKU 13107 (1,

72.1 mm HL, 282 mm TL), BSKU 13108 (1, 67.2 mm HL, 249 mm TL), ca. 15 km east of Cape Ashizurimisaki, Tosa Bay, 420–555 m, 2–6 Jun. 1968; *BSKU 71585 (1, 34.5 mm HL, 123+ mm TL), Tosa Bay, 400 m, FRV *Kotaka-maru*, otter trawl, 9 May 2000; *BSKU 112398 (1, 94.5 mm HL), *BSKU 112402 (1, 88.0 mm HL, 302+ mm TL), *BSKU 112403 (1, 55.5 mm HL, 213 mm TL), off Susaki, Tosa Bay, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama, 4 Mar. 2014; MSM 13-10 (1, 93.9 mm HL, 333+ mm TL), southeast of Fuji River, Suruga Bay, 35.1013°N, 138.6823°E, 502–529 m, 1.3-m larval net, 26 Jan. 2010; FRLM 3736 (1, 83.6 mm HL, 322+ mm TL), off Goza, Kumano-nada, long line, coll. Fisheries Research Institute of Mie Pref., 13 Oct. 1982; BSKU 19190 (1, 62.9 mm HL, 226+ mm TL), south of Miura Peninsula, Sagami Bay, 35.1250°N, 139.5100°E, 700 m, FRV *Soyo-maru*, sta. 181, beam trawl, 10 Mar. 1961. **Philippines:** BSKU 15694 (1, 95.6 mm HL, 322+ mm TL), Sulu Sea, 8.3467°N, 118.3300°E, 738 m, R/V *Hakuho-maru*, cr. KH-72-1, sta. 13, 3-m beam trawl, coll. O. Okamura, 27 May 1972; BSKU 15665 (1, 59.3 mm HL, 169+ mm TL), BSKU 15666 (1, 57.8 mm HL, 195+ mm TL), BSKU 15667 (1, 65.6 mm HL, 199+ mm TL), BSKU 15668 (1, 54.2 mm HL, 158+ mm TL), Sulu Sea, 8.3167°N, 118.1517°E, 495–500 m, R/V *Hakuho-maru*, cr. KH-72-1, sta. 12, 3-m beam trawl, coll. O. Okamura, 27 May 1972.

Redescription based on Japanese specimens.

General features are shown in Figs. 54B and 85. Counts and measurements are given in Tables 3 and 5. Body deepest at first dorsal-fin origin, gradually tapering from this level to long, laterally compressed tail. Trunk short, moderately compressed, width over pectoral-fin bases 1.1–1.4 in depth below first dorsal-fin origin. Head large, HL about 3.1–3.9 in TL. Snout long, sharply pointed, protruding well beyond upper jaw, length 1.4–2.0 times as long as orbit diameter; dorsal contour of snout moderately concave in lateral view; lateral nasal ridge completely supported by nasal bone; anterolateral margins of snout only slightly convex when viewed dorsally, giving attenuated appearance to snout. Orbit large, its greatest diameter 0.9–1.2 in postorbital length (1.4 in largest specimen; KPM-NI 28333, 135 mm HL). Interorbital space broad, flat, width 1.2–1.4 in orbit diameter. Mouth moderately large, inferior, protrusible, upper-jaw length 1.1–1.4 in orbit diameter (0.9 in largest specimen); posterior margin of maxilla extending to vertical through hind 1/3 of orbit or beyond; lateral corner of mouth barely restricted by skin folds; lips thin, slightly papillose near tooth bands. Suborbital region divided by longitudinal bony ridge passing from tip of snout to posteroventral angle of opercle; its upper half almost vertical, lower half sharply inclined mesially. Preopercle large, posterior margin inclined, forming moderately angular lobe at posteroventral corner. Subopercle produced into narrowly pointed tab posteroventrally; posterior margin deeply

FIGURE 85. Fresh specimens of *Coelorinchus smithi*. (A–C) BSKU 112403, 55.5 mm HL, 213 mm TL, Tosa Bay off Susaki, Kochi Pref., Pacific, ca. 400 m depth; (D–F) BSKU 112402, 88.0 mm HL, 302+ mm TL, collected with BSKU 112403; (G) KPM-NI 24972, 110 mm HL, 407+ mm TL, off Okinawa-jima Island, Okinawa Pref., 600 m depth. (A, D, G) Lateral, (B, E) dorsal, and (C, F) ventral views. [Photos: (A–F) N. Nakayama; (G) H. Senou]

TABLE 5. Frequency distribution of the number of pectoral-fin rays for *Coelorinchus smithi*.

Locality	<i>n</i>	Counts*						
		14	15	16	17	18	19	20
Indonesia	2				2^H			
Philippines	10	1		5	3	1		
Japan	89			1	10	34	41	3

*Excluding the uppermost rudimentary ray; numbers in bold and that denoted by superscript “H” indicate modal counts and value for the holotype, respectively

concave. Gill membranes broadly connected across, and attached mesially to isthmus, with free posterior fold. Outer gill slit moderately restricted by skin folds, length 2.1–3.3 in orbit diameter (1.5 in largest specimen). Gill rakers small, low, tubercular, armed with short, fine spines; no rakers on outer side of first arch and inner side of fourth arch; gill filaments moderately long. Chin barbel short, slender; length 2.8–4.9 in orbit diameter (2.4 in largest specimen).

Anus only slightly separated from anal-fin origin by a few scale rows. Light organ externally represented by short naked fossa anterior to anus; its anterior margin extending to about posterior 1/3 of distance from outer pelvic-fin bases to anus, and falling far short of line connecting inner pelvic-fin bases.

Teeth small, sharp, conical, gently incurved, in narrow bands in both jaws. Premaxillary teeth in short, uniformly wide band, with about 6–7 tooth rows near symphysis; posterior margin of tooth band falling far short of lateral corner of mouth; teeth becoming progressively smaller inwardly, with outer series only slightly larger. Mandibular teeth uniformly small, in narrow tapered band, with about 5 tooth rows near symphysis; posterior margin of tooth band extending beyond lateral corner of mouth. All teeth deeply embedded in thick layer of gum papillae.

Scales on body large, thin, moderately deciduous, covered with short, reclined, keel-like spinules in widely divergent rows (Fig. 53E–F); in 92.5 mm HL specimen (BSKU 112399), those on dorsum below interdorsal space with 6 rows of spinules; middle row usually with about 6–7 spinules, but not especially high or enlarged compared with adjacent rows; every spinule row complete throughout; spinules forming angle of about 45° to scale surface; spinules greatly overlapping, closely adjoined to one another, but their distal tips occasionally free, forming saw-toothed ridges that give spiny appearance to body surfaces; height of spinules increasing posteriorly, with last spinules in each row extending moderately beyond posterior scale margin; lateral buttresses of spinules narrowly developed; reticulate structures absent on unexposed portion. Body fully scaled except for fins and ventral light organ.

Scales on head ridges thickened and coarsely spinulated; those on median rostral ridge covered with radiating rows of spinules. Terminal snout scute moderately short, diamond-shaped, dorsoventrally flattened, its

length 2.1–4.0 in orbit diameter. Supraoccipital scute stout, prominent, armed with divergent rows of spinules; posttemporal scutes thickened, but not especially enlarged, armed with single row of spinules. Occipital scales covered with short, moderately reclined, keel-like (rarely knife-like) spinules in widely divergent rows (Fig. 52B); spinules in each row connected basally to one another, forming saw-toothed ridges, and their height gradually increasing posteriorly; scales on dorsal surfaces of snout posterior to lateral nasal ridges armed with divergent rows or cluster of short, erect, needle-like or knife-like spinules. Other scales on head variable in size, generally similar to those on body, but spinules much shorter and more erect; those on opercle, preopercle, and postorbital and supratemporal canals largest. Nasal fossa usually heavily scaled anteroventrally. Dorsal surfaces of snout fully scaled, without prominent naked clefts along each side of median rostral ridge. Underside of head completely scaled except for narrow naked area above upper lip, anterior portions of mandibular rami, and posteroventral margins of preopercles; scales on ventral surfaces of head small, non-imbricate, covered with small cluster of short, erect, knife-like spinules.

No open pores along cephalic sensory canals. Free neuromasts on head not prominently marked. Anterior nostril small and circular; posterior large, bean-shaped; septum between nostrils forming anteriorly opened hood. Lateral line complete, not interrupted throughout.

Origins of first dorsal and pelvic fins on about same vertical; first dorsal fin moderately high, its height 2.3–3.2 times as long as its base length; second spinous ray not elongated, smooth along its leading edge (a few rudimentary denticles rarely present distally); its tip usually extending to second dorsal-fin origin or beyond when laid back. Interdorsal space 1.2–2.2 times as long as first dorsal-fin base length. Second dorsal fin originating posterior to anal-fin origin (above base of 3rd–9th anal fin rays). Pectoral-fin base slightly anterior to vertical through pelvic-fin base. Outer pelvic-fin ray prolonged, with fine distal tip.

Color when fresh (Fig. 85). Head and body dark brown dorsally, whitish ventrally (especially lower 1/3 of body), but ventral parts moderately darker in large specimens; about 2 faint, dark saddles on tail in smaller specimens (Fig. 85A), first below about second dorsal-fin origin, second further posterior on tail, separated from first

FIGURE 86. Holotype of *Coelorhynchus smithi* (= *Coelorinchus smithi*). USNM 78212, 80.7 mm HL, 261+ mm TL, between Gilolo and Makyan Islands, Indonesia, Molucca Sea. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

FIGURE 87. Fresh specimens of *Coelorhynchus smithi*. (A–B) BSKU 15665, 59.3 mm HL, 169+ mm TL, Sulu Sea, 495–500 m depth; (C–D) BSKU 15694, 95.6 mm HL, 322+ mm TL, Sulu Sea, 738 m depth. (A, C) Lateral and (B, D) ventral views of the head and trunk. [Photos: BSKU]

by distance about equal to postorbital length; gill cover slightly shiny, except for large specimens; lips white, gill and branchiostegal membranes pale to dusky; light organ and periproct dull black; first dorsal fin uniformly dark, but whitish apically; pectoral and second dorsal fins generally dusky; pelvic fin dark, but outermost ray whitish; second dorsal and anal fins dusky to dark, much darker posteriorly.

Color in alcohol. Head and body generally dark brown dorsally; lower 1/3 of body distinctly paler; abdomen slightly dark in ventral view; 2 faint, dark saddles on tail

(especially marked in small specimens, and often difficult to distinguish in large specimens); lips white; oral and gill cavities blackish; gill rakers and filaments light tan, arches dark dusky; gular and branchiostegal membranes pale to dusky; first dorsal, pectoral, and pelvic fins generally uniformly dark, but outer pelvic-fin rays paler distally; second dorsal and anal fin heavily peppered; anal fin prominently blackish posteriorly.

Size. To at least 53 cm TL (KPM-NI 28333, 528+ mm TL, Okinawa Trough, Japan).

Distribution. Widely distributed from Japan

to Australia, including Taiwan, the Philippines, and Indonesia, at depths of 196–1110 m (Gilbert & Hubbs 1920; Iwamoto & Williams 1999; Iwamoto & Graham 2001). In Japan and adjacent waters, the species is known from the Pacific off southern Japan northward to the Boso Peninsula (139.89°E), Okinawa Trough, and northern South China Sea, at depths of 196–780 m (Shao *et al.* 2008b; this study; Appendix 3-5B). Common.

Remarks. *Coelorinchus smithi* was originally described by Gilbert & Hubbs (1920) based on 19 specimens collected from Indonesia and the Philippines, with the type locality in the Molucca Sea (Fig. 86). These specimens were previously included in the type series of *C. commutabilis* Smith & Radcliffe in Radcliffe, 1912. *Coelorinchus smithi* was first reported from Japanese waters by Okamura (1963a).

Comparisons of specimens collected from Japan with those from the East Indies (holotype and 5 Philippine specimens) revealed subtle differences in fresh coloration and pectoral-fin ray counts. In the Japanese specimens, the ventral half of the body is whitish (Fig. 85A–F) except for one large specimen examined (KPM-NI 24972, 407+ mm TL; Fig. 85G), whereas the whole body is uniformly and intensely dark in similar-sized specimens from the Philippines (Fig. 87). Unfortunately, the color of the 261+ mm TL holotype is somewhat faded (Fig. 86), but the dark coloration of East Indies' specimens was also depicted in the original description, where Gilbert & Hubbs (1920:497) stated that the overall coloration of the type specimens was “brown, usually darker than in *C. commutabilis*; it is darker above than below” (for *C. commutabilis* see the Relationships and comparisons given below). Iwamoto & Merrett (1997:489) pointed out that “color can be quite variable within a species, especially from one region to another, and it appears to be influenced considerably by the substratum”. However, the difference between the two populations of *C. smithi* is too distinctive to be ascribed to intraspecific variation. In addition, the Japanese specimens commonly have two faint saddles on the tail (indistinct in large individuals), of which one is situated below the second dorsal-fin origin, and the other is separated from the first saddle by a distance about equal to the postorbital length (Fig. 85A). These saddles are absent in the Philippine specimens examined (unknown in the holotype due to discoloration), but Gilbert & Hubbs (1920:497) also noted the presence of similar markings in the original description. Moreover, counts of pectoral-fin rays are slightly higher in the Japanese specimens than in the holotype and the Philippine specimens examined [i16–i20 (usually i18 or i19, modally i19) vs. i14–i18 (modally i16 or i17); Table 5], although their ranges overlap. The modal difference in the fin-ray counts further suggests that the Japanese population represents a distinct species or subspecies with allopatric distribution from the true *C. smithi*. A redescription based on the Japanese specimens examined is provided here for future investigations, and

a molecular study will help to elucidate their taxonomic status.

Relationships and comparisons. *Coelorinchus smithi* belongs to the *C. japonicus* group (*sensu stricto*; see the Relationships of *C. japonicus*), and closely resembles *C. commutabilis* known from the East Indies. When similar-sized specimens are compared, the Japanese *C. smithi* examined readily differ from the holotype of *C. commutabilis* (USNM 72945, 93.7 mm HL; examined here) in that the ventral half of the body is distinctly paler (Fig. 85 vs. body uniformly dark), but otherwise they are similar to each other. According to Gilbert & Hubbs (1920:497) *C. smithi* (from the East Indies) is distinguished from *C. commutabilis* by having “constantly larger scales” and “several carinae instead of a median keel of the scales of occipital ridges”. However, examinations of their holotypes along with additional specimens of *C. smithi* revealed no significant differences in the numbers of transverse scale rows (below first dorsal-fin origin 4.5–6 vs. 5.5 in *C. smithi* and *C. commutabilis* respectively; below first dorsal-fin midbase 4–4.5 vs. 4.5; and below second dorsal-fin origin 4.5–6 vs. 5.5). The holotype of *C. commutabilis* is also differentiated in that most scales on the parietal ridges (“occipital ridges” of Gilbert & Hubbs 1920) are covered with only a few rows of stout spinules, whereas those of *C. smithi* are generally armed with multiple rows so as to give a highly spinulated appearance to the scales. The difference in spinulation of the modified scales may warrant the recognition of separate species, but further verification based on more specimens of *C. commutabilis* is required.

The Japanese population of *C. smithi* is also similar to *C. charius* Iwamoto & Williams, 1999 known from off Western Australia, both of which are characterized by their paler body color and the presence of faint saddles on the tail when young. An examination of the holotype and 18 paratypes of *C. charius* (51.3–98.2 mm HL) revealed that these species are difficult to distinguish from each other by counts and measurements, squamation, and scale spinulation. However, a subtle difference was found in the saddle pattern of small specimens. In the young *C. smithi* examined, only two saddles were present dorsally on the tail (Figs. 54B, 85A), whereas similar-sized *C. charius* specimens had more than five saddles (Iwamoto & Williams 1999: fig. 9b). Although this difference is pronounced only in small specimens, it indicates the specific separation between Japanese *C. smithi* and *C. charius*.

Coelorinchus sereti Iwamoto & Merrett, 1997 known from the tropical southwestern Pacific is another species closely related to *C. smithi*. They are quite similar in overall appearance, with no pronounced differences in counts and measurements, except for the pelvic-fin length (53–74% PRL in *C. smithi* vs. 72–90% in *C. sereti*). However, the shape of the snout is substantially different between the two species. In *C. smithi*, the dorsal contour of the snout is moderately concave in lateral view (Figs.

85A, D, G, 86A, 87A, C), whereas that of *C. sereti* is gently convex (Iwamoto & Merrett 1997: fig. 15a). Also, when viewed dorsally (Figs. 85B, E, 86B), the snout of *C. smithi* is progressively attenuated toward its tip, giving a sharply pointed appearance, whereas the leading edges of *C. sereti* are broadly convex anteriorly, giving a much wider appearance to the snout (Iwamoto & Merrett 1997: fig. 15a').

In Japanese waters, *C. smithi* is likely to be confused with *C. nox* **sp. nov.**, but they are distinguished by the combination of squamation and coloration (see the Relationships and comparisons of the latter). *Coelorinchus smithi* also resembles *C. japonicus* (Temminck & Schlegel, 1846), with which it has often been confused in museum collections. However, it differs notably from the latter species in that the occipital scales are covered with four or more widely divergent rows of spinules (vs. uniserial, rarely 2–3 narrowly divergent rows). As noted by Okamura (1963a:31), the ventral light organ of *C. smithi* is much broader and more prominent compared with that of *C. japonicus* (Fig. 85C, F vs. Fig. 51C). In addition, *C. smithi* tends to have relatively fewer scales below the second dorsal-fin origin (4–6, modally 5) than *C. japonicus* (5–7.5, modally 6).

***Coelorinchus tokiensis* (Steindachner & Döderlein, 1887)**

[Japanese name: Miyako-hige]

(Figs. 42E–F, 88–90; Appendix 3-5C)

Macrurus tokiensis Steindachner & Döderlein, 1887:283 [original description; syntype: MZS 1804, from “Tokio”, but probably from Eno-shima Island, Sagami Bay; 1 of 2 syntypes missing].

Coelorhynchus tokiensis: Jordan & Snyder 1901:120 (listed; Japan); Jordan & Gilbert in Jordan & Starks 1904:620 [after Steindachner & Döderlein (1887)]; Jordan *et al.* 1913:418 (listed; Japan; new Japanese name: “Miyako-hige”); Gilbert & Hubbs 1916:179 (description; 2 spec. from East China Sea); Okada & Matsubara 1938:450 (in key; Japan); Kamohara 1938:73 (spec. from Kochi Pref.); Kamohara 1950:277 (listed; Kochi and Wakayama Pref.); Uchinomi 1951:76 (listed; southern part of Kii Peninsula); Kamohara 1952:100 (spec. from Kochi Pref.); Matsubara 1955:1313 (in key; Japan); Kamohara 1958:73 (listed; Kochi Pref.); Kamohara 1964:96 (listed; Kochi Pref.); Okamura 1970a:194, pl. XLII, text-fig. 83 (description; 5 spec. from Pacific off southern Japan from Sagami Bay to Mimase); Tominaga & Uyeno 1981:488 (listed; Japan).

Coelorhynchus (Oxymacrus) tokiensis: Okamura 1970b: table 1 (listed; Japan).

Coelorhynchus longicephalus Okamura, 1982:177, fig. 105 (original description; holotype: BSKU 26144, from Kyushu-Palau Ridge, 27°54.2'N, 134°39.5'E, in 700 m; 16 paratypes from Kyushu-Palau Ridge, in 336–700 m; new Japanese name:

“Zunaga-sokodara”; in part, BSKU 26102 is *C. gilberti*).

Coelorinchus tokiensis: Yatou 1984:235, 370, fig. 168 (brief description; 5 spec. from Okinawa Trough; photo based on BSKU 29872); Okamura 1984b:97, pl. 83, fig. H (compiled); Okamura 1988:97, pl. 83, fig. H (compiled); Suzuki & Kataoka 1997:84, pl. 33, fig. 183 (brief description; 1 spec. from Kumano-nada off Namikiri); Okamura 1997:127, fig. 17 (compiled); Amaoka 2007: fig. 2-B (photograph of type specimen); Nakabo & Kai 2013:511 (in key; Japan); Nakayama *et al.* 2015c:249, figs. 1–7 (redescription; 59 spec. from Japan; *C. hexafasciatus* a junior synonym of *C. tokiensis*); Miyazaki *et al.* 2019: fig. 3D, table 1 (1 spec. listed from Tokyo Submarine Canyon); Motomura 2020:38 (listed; Japan).

Coelorinchus longicephalus: Okamura 1984b:97, pl. 83, fig. I (compiled); Okamura 1988:97, pl. 83, fig. I (compiled); Iwamoto 1990:169, fig. 390 (synopsis); Okamura 1997:127, fig. 18 (compiled); Nakabo & Kai 2013:512 (in key; Japan); Ikeda & Nakabo 2015:322, pl. 69, figs. 6–9 (brief description; 2 spec. from Wakayama Pref.).

Caelorinchus tokiensis: Nakabo 1993:370 (in key; Japan); Miya *et al.* 1995: table 2 (questionable; 1 spec. from off Boso Peninsula; midwater capture); Nakabo 2000:434 (in key; Japan); Shinohara *et al.* 2001:305 (listed; Tosa Bay); Nakabo 2002:434 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Shinohara *et al.* 2005:417 (listed; Ryukyu Islands); Senou *et al.* 2006:420 (listed; Sagami Sea).

Caelorinchus longicephalus: Nakabo 1993:371 (in key; Japan); Nakabo 2000:435 (in key; Japan); Nakabo 2002:435 (in key; Japan); Furuhashi *et al.* 2010: table 2 (6 spec. listed from northern Okinawa Trough).

Diagnosis. Light organ externally represented by short, narrow, naked streak immediately anterior to anus, its anterior margin falling far short of line connecting inner pelvic-fin bases. Underside of head completely naked except for oval scaly patches on ventral surfaces of preopercles; longitudinal naked clefts along each side of mid rostral ridge scarcely developed. Snout long with bluntly pointed tip, length 70–85% PRL, its dorsal profile straight to gently convex in lateral view; snout tipped with 3 bluntly pointed scutes; lateral nasal ridge incompletely supported by nasal bone. Anus slightly separated from anal-fin origin. Premaxillary teeth in long, tapered band, with outer series prominently enlarged; posterior margin of tooth band almost reaching lateral corner of mouth. Body scales covered with short, reclined, keel-like spinules in widely divergent, fine saw-toothed ridges; every spinule row complete, extending to posterior scale margin; spinules in each row overlapping but free at tips, gradually increasing in height posteriorly; buttresses poorly developed. Occipital scales usually covered with single row of spinules. Preoral length 47–69% PRL; barbel length 9–19% PRL. About 6–7 dark saddles on body (faint in large individuals); underside of head dusky; lips white; oral cavity blackish; gular and branchiostegal membranes dusky.

Material examined. 57 specimens. **Syntype of *Macrurus tokiensis*:** MZS Pis1804 (1, 173 mm HL, 589+ mm TL), Tokyo, Japan, coll. L. Döderlein, date unknown (examined by Nakayama *et al.* 2015c). **Holotype of *Coelorhynchus longicephalus*:** BSKU 26144 (252 mm HL, 875+ mm TL), Kyushu-Palau Ridge, Japan, 27.9167°N, 134.6500°E, 700 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. 2-3, bottom trawl, coll. Y. Kobayashi *et al.*, 20 Feb. 1978. **Paratypes of *C. longicephalus*:** BSKU 29535 (1, 122 mm HL, 453 mm TL), BSKU 29536 (1, 208 mm HL, 680+ mm TL), Kyushu-Palau Ridge, 27.7500°N, 135.3167°E, 336 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. T19, bottom trawl, coll. Y. Kobayashi *et al.*, 17 Dec. 1979; BSKU 30438 (1, 127 mm HL, 459 mm TL), BSKU 30439 (1, 252 mm HL, 870+ mm TL), BSKU 30444 (1, 197 mm HL, 685+ mm TL), Kyushu-Palau Ridge, 26.7500°N, 135.3167°E, 336 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. T8, bottom trawl, coll. Y. Kobayashi *et al.*, 17 Dec. 1979; BSKU 32223 (1, 113 mm HL, 363+ mm TL), Kyushu-Palau Ridge, 28.1117°N, 134.6656°E, 550–600 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, tr. 9, bottom trawl, coll. O. Okamura *et al.*, 19 Dec. 1979; BSKU 29435 (1, 127 mm HL, 363+ mm TL), Kyushu-Palau Ridge, 28.1070°N, 134.6593°E, 550–600 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. 4-T9, bottom trawl, coll. O. Okamura *et al.*, 19 Dec. 1979; BSKU 26222 (1, 227 mm HL, 755+ mm TL), BSKU 26223 (1, 197 mm HL, 634+ mm TL), Kyushu-Palau Ridge, 27.9000°N, 134.6550°E, 695 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. 2-2, bottom trawl, coll. Y. Kobayashi *et al.*, 15 Feb. 1978; BSKU 26100 (1, 198 mm HL, 659+ mm TL), BSKU 26101 (1, 227 mm HL, 779+ mm TL), BSKU 26103 (1, 211 mm HL, 742+ mm TL), BSKU 26104 (1, 202 mm HL, 672+ mm TL), BSKU 26145 (1, 229 mm HL, 784+ mm TL), BSKU 26146 (1, 167 mm HL, 514+ mm TL), collected with holotype. **Non-types: Japan:** KPM-NI 24973 (1, 240 mm HL, 878 mm TL), near Okinawa, 600 m, long line, coll. K. Yunokawa, 2009; BSKU 106833 (1, 55.3 mm HL, 190+ mm TL), BSKU 106834 (1, 73.8 mm HL, 266 mm TL), southeast of Shimokoshiki-jima Island, East China Sea, 31.5675°N, 129.8930°E, 393 m, F/V *Maruko-maru*, tr. 2, bottom trawl, coll. N. Nakayama *et al.*, 24 Apr. 2012; BSKU 106825 (1, 53.3 mm HL, 175+ mm TL), southeast of Shimokoshiki-jima Island, East China Sea, 31.5658°N, 129.8915°E, 380 m, F/V *Maruko-maru*, tr. 1, bottom trawl, coll. N. Nakayama *et al.*, 24 Apr. 2012; BSKU 106815 (1, 60.8 mm HL, 215 mm TL), southeast of Shimokoshiki-jima Island, East China Sea, 31.5652°N, 129.8878°E, 407 m, F/V *Maruko-maru*, tr. 3, bottom trawl, coll. N. Nakayama *et al.*, 24 Apr. 2012; BSKU 106773 (1, 69.6 mm HL, 228+ mm TL), BSKU 106774 (1, 59.8 mm HL, 204+ mm TL), BSKU 106781 (1, 80.5 mm HL, 289+ mm TL), BSKU 106782 (1, 75.5 mm HL, 268+ mm TL), BSKU 106783 (1, 70.1 mm HL, 250

mm TL), BSKU 106784 (1, 71.5 mm HL, 262 mm TL), BSKU 106785 (1, 74.6 mm HL, 263+ mm TL), BSKU 106786 (1, 67.1 mm HL, 238+ mm TL), BSKU 106787 (1, 67.1 mm HL, 236+ mm TL), BSKU 106788 (1, 74.6 mm HL, 263 mm TL), southeast of Shimokoshiki-jima Island, East China Sea, 31.5560°N, 129.8847°E, 415 m, F/V *Maruko-maru*, tr. 4, bottom trawl, coll. N. Nakayama *et al.*, 24 Apr. 2012; FFNU P-00630 (1, 73.1 mm HL, 249 mm TL), west of Amakusa Knoll, East China Sea, 32.1465°N, 129.4870°E, 511 m, T/V *Nagasaki-maru*, cr. N211, sta. B5-4, 3-m ORE beam trawl, coll. J. Hashimoto, 10 May 2006; FFNU P-00642 (1, 84.8 mm HL, 300 mm TL), east of Hirazi Bank, East China Sea, 32.2440°N, 129.4802°E, 421 m, T/V *Nagasaki-maru*, cr. N224, sta. B4, 3-m ORE beam trawl, coll. J. Hashimoto, 28 Oct. 2006; BSKU 29872 (1, 129 mm HL, 472+ mm TL), east of Danjo Islands, Okinawa Trough, 32.0033°N, 128.7050°E, 360–400 m, F/V *Yuryo-maru*, No. 8, sta. 3-L18, long line, coll. H. Maeda, 2 Nov. 1979; FFNU P-00672 (1, 140 mm HL, 521 mm TL), northwest of Daiichi-Danjo Knoll, East China Sea, 31.7013°N, 128.5800°E, 500 m, T/V *Nagasaki-maru*, cr. N237, sta. E5-2, 3-m ORE beam trawl, coll. J. Hashimoto, 9 May 2007; HUMZ 95252 (1, 241 mm HL, 877+ mm TL), northwest of Amami-oshima, East China Sea, 28.9817°N, 128.0233°E, 760–790 m, hook and line, 14 Nov. 1981; BSKU 28710 (1, 205 mm HL, 760 mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.7333°N, 127.8000°E, 495–510 m, F/V *Ryoan-maru*, No. 28, tr. 72, bottom trawl, coll. Y. Kinoshita and S. Hagino, 19 Mar. 1978; BSKU 33258 (1, 150 mm HL, 549+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.7167°N, 127.7000°E, 520 m, F/V *Yuryo-maru*, No. 8, sta. 2-T12, bottom trawl, coll. M. Hirose and T. Kume, 14 Oct. 1979; BSKU 27223 (1, 165 mm HL, 600 mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.4333°N, 127.5167°E, 570 m, F/V *Yuryo-maru*, No. 8, tr. 22, bottom trawl, coll. Y. Kinoshita and S. Hagino, 5 Feb. 1978; BSKU 27140 (1, 82.8 mm HL, 290 mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.3833°N, 127.5150°E, 650 m, F/V *Yuryo-maru*, No. 8, tr. 21, bottom trawl, coll. Y. Kinoshita and S. Hagino, 5 Feb. 1978; BSKU 27584 (1, 55.2 mm HL, 198+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 28.7167°N, 127.2333°E, 750–755 m, F/V *Ryoan-maru*, No. 28, tr. 48, bottom trawl, coll. Y. Kinoshita and S. Hagino, 11 Mar. 1978; BSKU 44426 (1, 194 mm HL, 746+ mm TL), BSKU 44427 (1, 152 mm HL, 514+ mm TL), off Nahari, Tosa Bay, 300 m, 13 Jan. 1988; BSKU 44361 (1, 231 mm HL, 820+ mm TL), BSKU 44362 (1, 190 mm HL, 641+ mm TL), off Yasuda, Tosa Bay, 400 m, 9 Dec. 1987; BSKU 43855 (1, 245 mm HL, 824+ mm TL), off Kochi, 11 Jul. 1987; BSKU 81498 (1, 53.4 mm HL, 197 mm TL), Tosa Bay, 600 m, FRV *Kotaka-maru*, otter trawl, 16 May 1988; BSKU 45506 (1, 194 mm HL, 690+ mm TL), off Tosa-Shimizu, long line, 24 Aug. 1988; BSKU 45500 (1, 269 mm HL, 879+ mm

FIGURE 88. Fresh specimens of *Coelorinchus tokienensis*. (A–C) BSKU 106781, 80.5 mm HL, 289+ mm TL, off Shimokoshiki-jima Island, Kagoshima Pref., East China Sea; (D) BSKU 44427, 152 mm HL, 514+ mm TL, off Nahari, Kochi Pref., Tosa Bay, Pacific, 300 m depth; (E) BSKU 29536, paratype of *C. longicephalus*, 208 mm HL, 680+ mm TL, Kyushu-Palau Ridge, Pacific, 336 m depth. (A, D–E) Lateral views; (B) dorsal and (C) ventral views of the head and trunk. [Photos: (A–C) N. Nakayama; (D–E) reproduced from Nakayama *et al.* (2015c: fig. 7d, f) with permission from the Ichthyological Society of Japan]

TL), BSKU 45501 (1, 251 mm HL, 901+ mm TL), BSKU 45503 (1, 245 mm HL, 898 mm TL), off Tosa-Shimizu, long line, Oct. 1988; MSM 10-8 (1, 217 mm HL, 833+ mm TL), Numazu fish market, date unknown; FRLM 3578 (1, 165 mm HL, 573 mm TL), off Mie, Kumano-nada, long line, coll. Fisheries Research Institute of Mie Pref., 25 May 1982; NSMT-P 100178 (1, 67.7 mm HL, 226+ mm TL), northeast of Miyake-jima Island, Shichito-Iojima Ridge, 34.1287°N, 139.6413°E, 484–565 m, R/V *Tansei-maru*, cr. KT-89-06, sta. ME10, 3-m ORE beam

trawl, 19 May 1989; FAKU 34265 (1, 139 mm HL, 478+ mm TL), off Oiso, Sagami Bay, Apr. 1969; FAKU 35862 (1, 111 mm HL, 366+ mm TL), Sagami Bay, Apr. 1962.

Counts and measurements. Based on 57 specimens (53.3–269 mm HL, 175+–901+ mm TL). Counts: first dorsal-fin rays II,8–10; pectoral-fin rays i16–i19; pelvic-fin rays 6–7; gill rakers on first arch (outer/inner) 0–2/8–11, on second arch 7–9/9–11; longitudinal scales 37–53; transverse scale rows below first dorsal-fin origin 5.5–9, below first dorsal-fin midbase 4.5–7, below second

FIGURE 89. Syntype of *Macrurus tokiensis* (= *Coelorinchus tokiensis*). MZS Pis1804, 173 mm HL, 589+ mm TL, Tokyo, Pacific. (A) Dorsal and (B) ventral views of the head and trunk; (C) lateral view. Preserved condition. [Photos: reproduced from Nakayama *et al.* (2015c: fig. 1) with permission from the Ichthyological Society of Japan]

dorsal-fin origin 5.5–8, above anal-fin origin 18.5–28.5; pyloric caeca 51–61.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 42–46 (70–85); orbit diameter 20–27 (36–46); postorbital length 31–36 (54–65); postrostral length 54–59; orbit–preopercle distance 33–40 (58–72); suborbital width 10–13 (18–22); upper-jaw length 25–31 (44–57); length of rictus 19–25 (34–45); length of premaxillary tooth band 16–22 (29–38); preoral length 26–39 (47–69); length of terminal snout scute 18 (33); length of lateral nasal ridge 24–28 (40–51); length of suborbital ridge 92–98 (162–178); snout width 20–27 (35–48); internasal width 16–20 (28–34); interorbital width 18–23 (32–39); occipital width 6–9 (11–16); body width over pectoral-fin bases 32–46 (56–82); body depth at first dorsal-fin origin 42–57 (74–102); body depth at anal-fin origin 35–50 (60–90); prepelvic length 99–115 (172–205); preanus length 122–147 (209–265); preanal length 128–155 (220–277); isthmus–pelvic distance 22–33 (39–59); isthmus–anus distance 49–63 (85–112); isthmus–anal distance 51–74 (88–131); pelvic–anal distance 26–44 (46–78); anus–anal distance 4–10 (8–17); pelvic-fin length 25–45 (44–78); pectoral-fin length 32–46 (54–78); predorsal length 102–

109 (178–196); height of first dorsal fin 35–43 (62–74); length of first dorsal-fin base 15–20 (26–35); interdorsal length 10–24 (17–43); length of gill slit 14–18 (24–31); length of posterior nostril 6–9 (11–16); barbel length 5–11 (9–19); length of pyloric caecum 3–5 (6–8).

Size. Attains about 90 cm TL (BSKU 45501, 901+ mm TL, off Tosa-Shimizu, Kochi, Japan).

Distribution. In Japan, known from the Pacific off southern Japan northward to the Boso Peninsula (139.89°E), off the Izu Islands, the Okinawa Trough, and the Kyushu-Palau Ridge, at depths of 300–790 m (Appendix 3-5C). Elsewhere, known only from the South China Sea (Iwamoto *in* Randall & Lim 2000). Common in the northern Okinawa Trough and Kyushu-Palau Ridge, but rare in other parts of its range.

Remarks. *Coelorinchus tokiensis* was originally described as a species of *Macrurus* (= *Macrourus* Bloch, 1786) on the basis of two specimens collected from Tokyo, Japan (probably originally from off Eno-shima Island, Sagami Bay; see Amaoka 2007:28). One of the two syntypes was recently found among the ichthyological collection of the Musée Zoologique de l'Université Louis Pasteur, Strasbourg (MZS Pis1804; Amaoka 2007: fig. 2b; Fig. 89). Nakayama *et al.* (2015c) redescribed *C.*

FIGURE 90. Preserved specimen of *Coelorinchus tokiensis*. BSKU 26144, holotype of *Coelorhynchus longicephalus*, 252 mm HL, 875+ mm TL, Kyushu-Palau Ridge, Pacific, 700 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: reproduced from Nakayama *et al.* (2015c: fig. 2) with permission from the Ichthyological Society of Japan]

tokiensis with an examination of the MZS syntype, and recognized *C. longicephalus* Okamura, 1982 (Fig. 90) as a junior synonym of the former.

Relationships and comparisons. *Coelorinchus tokiensis* is one of six species of the *C. tokiensis* group as redefined by Nakayama *et al.* (2015c; see the Relationships of *C. hexafasciatus*), and is readily distinguished from its similar congeners by having the combination of a long snout (70–85% PRL) and the underside of the head completely naked except for oval scaly patches on the ventral surfaces of the preopercles.

Comments on “*Caelorinchus* (*Quincuncia*) *logispinosus*”

Chiou *et al.* (2004a: table 1) introduced the name “*Caelorinchus* (*Quincuncia*) *logispinosus*” in their list of

Taiwanese grenadiers. While the specific epithet is new to macrourid taxonomy, they provided neither a description nor definition of this ambiguous species. Because the name fails to implement Art. 13 of the ICZN (1999), it is considered here as a *nomen nudum*.

Genus *Coryphaenoides* Gunnerus, 1765

[Japanese name: Hokakedara-zoku]

Coryphaenoides Gunnerus, 1765:50 (type species: *Coryphaenoides rupestris* Gunnerus, 1765, by monotypy).

Lepturus Gronow *in* Gray, 1854:165 [type species: *Lepturus brevirostris* Gronow *in* Gray, 1854, by monotypy; objectively invalid, preoccupied by *Lepturus* Moehring, 1758 (see also Moehring 1752) and *Lepturus* Brisson, 1760 *in* Aves (see also Grant & Mackworth-Præd 1956)].

Chalinura Goode & Bean, 1883:198 [type species: *Chalinura*

simula Goode & Bean, 1883, by original designation; objectively invalid, preoccupied by *Chalinura* Dalman, 1826 in Arachnida (see also Dunlop & Jekel 2008), replaced by *Fuyangia* Whitley, 1931].

Lionurus Günther, 1887:141 [as subgenus of *Macrurus*; type species: *Coryphaenoides filicauda* Günther, 1878, by subsequent designation by Jordan & Evermann (1898)].

Nematonurus Günther, 1887:150 [as subgenus of *Macrurus*; type species: *Macrurus armatus* Hector, 1875, by subsequent designation by Jordan & Evermann (1898)].

Moseleya Goode & Bean, 1896:417 (type species: *Coryphaenoides longifilis* Günther, 1877, by original designation; objectively invalid, preoccupied by *Moseleya* Quelch, 1884 in Coelenterata, replaced by *Dolloa* Jordan, 1900).

Albatrossia Jordan & Everman, 1898:2573 (type species: *Macrourus pectoralis* Gilbert, 1891, by original designation).

Bogoslovius Jordan & Evermann, 1898:2574 (type species: *Bogoslovius clarki* Jordan & Gilbert in Jordan & Evermann, 1898, by original designation).

Dolloa Jordan, 1900:897 (type species: *Coryphaenoides longifilis* Günther, 1877, by original designation; replacement name for *Moseleya* Goode & Bean, 1896).

Ateleobranchium Gilbert & Burke, 1912:94 (type species: *Ateleobranchium pterotum* Gilbert & Burke, 1912, by original designation).

Fuyangia Whitley, 1931:334 (type species: *Chalinura simula* Goode & Bean, 1883, by original designation; replacement name for *Chalinura* Goode & Bean, 1883).

Hemimacrurus Fraser-Brunner, 1935:322 (type species: *Macrurus acrolepis* Bean, 1884, by original designation).

Cariburus Parr, 1946:57 (type species: *Macrurus zaniophorus* Vaillant, 1888, by original designation).

Diagnosis. Anus immediately anterior to anal-fin origin; periproct absent or rudimentary. Ventral light organ absent. Infraorbital ridge not connected with preopercular ridge, separated by distinct gap. Second spinous ray of first dorsal fin serrated along its leading edge (serrations rarely rudimentary or lost). Tip and lateral angles of snout armed with stout tubercles in most species. Chin barbel present. Pelvic fin inserted below or posterior to (rarely slightly anterior to) pectoral-fin base. Anal-fin origin well posterior to hind margin of first dorsal-fin base. Body scales lacking reticulate structure; buttresses of spinules generally absent. Scales along second dorsal and anal fins not enlarged. Grooved lateral line complete (rarely interrupted anteriorly). Cephalic sensory pores present or absent. Rete-gas gland complexes 4–7. Branchiostegal rays 6. Color variable, without prominent silvery reflection when fresh. [Modified from Iwamoto & Sazonov (1988).]

Remarks. *Coryphaenoides* is the second largest genus of the family, with about 65 species currently recognized (Iwamoto *et al.* 2015; Nakayama & Endo 2016a; Fricke *et al.* 2020), 15 of which are herein recorded from the study area. Species of the genus are widely known from

the deep demersal habitat of the world's oceans, with a few occurring at hadal depths [i.e., *C. armatus* (Hector, 1875), *C. leptolepis* Günther, 1877, and *C. yaquinae* Iwamoto & Stein, 1974; see the Remarks and Range extension of each species]. *Coryphaenoides* is seemingly abundant in mid- and high-latitude regions, but appears to be rare in tropical waters. The genus also includes some commercially exploited species (Iwamoto 1990; Iwamoto *et al.* 2015), viz., *C. rupestris* Gunnerus, 1765 (northeastern Atlantic), *C. acrolepis* (Bean, 1884) (North Pacific), and *C. pectoralis* (Gilbert, 1891) (North Pacific).

Key to species of *Coryphaenoides* from Japan and adjacent waters

- 1a Mandibular teeth arranged in 1 or 2 rows (especially evident laterally and posteriorly) 2
- 1b Mandibular teeth arranged in distinct band 6
- 2a Interdorsal space shorter than (0.3–0.8 times as long as) first dorsal-fin base length; barbel rudimentary or short, length 1–8% HL 3
- 2b Interdorsal space longer than (1.2–3.0 times as long as) first dorsal-fin base length; barbel well developed, length 13–25% HL 4
- 3a Pelvic-fin rays 5–8; first dorsal-fin rays II, 8–10; mandibular teeth arranged in uniserial row; snout conical in lateral view, moderately protruding beyond upper jaw; outer pelvic-fin ray not especially prolonged *C. pectoralis*
- 3b Pelvic-fin rays 9–10; first dorsal-fin rays II, 12–15; mandibular teeth arranged in 2 rows, with inner row distinctly enlarged; snout bluntly rounded in lateral view, barely protruding beyond upper jaw; outer pelvic-fin ray greatly prolonged *C. longifilis*
- 4a Premaxillary teeth in wide tapered band, with outer row of enlarged canines and inner band of villiform teeth; body generally pale overall *C. leptolepis*
- 4b Premaxillary teeth arranged in 1–2 rows or narrow band; body uniformly dark brown 5
- 5a Inner premaxillary teeth, if present, arranged in uniserial row; mandibular teeth arranged in 1 distinct row near symphysis; scaly patches present on lower half of the suborbital region and posterior portion of mandibular rami *C. armatus*
- 5b Inner premaxillary teeth arranged in 2–3 irregular rows or narrow band; mandibular teeth arranged in 2 irregular rows near symphysis; lower half of suborbital region and mandibular rami completely naked *C. yaquinae*
- 6a Mouth small, posterior margin of maxilla not reaching vertical through midorbit (upper-jaw length 28–32% HL), lateral corner of mouth moderately restricted by lip folds 7
- 6b Mouth large, posterior margin of maxilla extending to vertical through midorbit or beyond (upper-jaw length 33–44% HL); lateral corner of mouth not restricted by lip folds 9
- 7a Spinules on body scales arranged in tightly packed convergent rows; leading edges of snout narrowly naked; interpelvic width distinctly greater than (or rarely about equal to) pelvic-

- fin base length *C. nasutus*
- 7b** Spinules on body scales arranged in tightly packed parallel to subparallel rows; leading edges of snout fully scaled; interpelvic width less than pelvic-fin base length **8**
- 8a** Last spinules on body scales scarcely overlapping posterior scale margin; underside of snout with broad naked band above upper lip; preopercle naked along ventral margin *C. marginatus*
- 8b** Last spinules on body scales greatly overlapping posterior scale margin; underside of snout only narrowly naked above upper lip; preopercle almost completely scaled *C. microps*
- 9a** Outer gill slit greatly restricted by skin folds, length less than half orbit diameter (5–10% HL) **10**
- 9b** Outer gill slit moderately wide, length greater than half orbit diameter (15–26% HL) **12**
- 10a** Underside of snout fully scaled; spinules on body scales arranged in tightly packed convergent rows or quincunx order; pelvic-fin rays 9–10 *C. rudis*
- 10b** Underside of snout with prominent naked area; spinules on body scales arranged in slightly divergent rows; pelvic-fin rays 11–12 **11**
- 11a** Middle row of body-scale spinules distinctly higher than adjacent rows; dorsal contour of head smoothly curved from snout tip to first dorsal-fin origin; orbit diameter 23% HL; pectoral-fin rays i24 *C. asper*
- 11b** Middle row of body-scale spinules not especially higher than adjacent rows; dorsal contour of head prominently humped over nape, with distinct depression above orbits; orbit diameter 20–21% HL; pectoral-fin rays i19–i22 *C. soyooae*
- 12a** Scales on head and body adherent, those along leading edge of snout and infraorbital ridge slightly thickened and enlarged; barbel well developed, length 11–19% HL; pelvic-fin rays usually 8 (rarely 7 or 9); body uniformly black in adults *C. acrolepis*
- 12b** Scales on head and body deciduous, those along leading edge of snout and infraorbital ridge not modified; barbel rudimentary to short, length 3–11% HL; pelvic-fin rays usually 9 (rarely 8 or 10); body pale to dusky **13**
- 13a** No prominent modified scales on snout; pectoral fin not reaching vertical through anal-fin origin *C. altipinnis*
- 13b** Snout tipped with prominent enlarged tubercles; pectoral fin extending to or beyond vertical through anal-fin origin **14**
- 14a** Snout relatively broad, its width, 24–30% HL, internasal width 21–27% HL; spinules on body scales arranged in parallel to subparallel rows *C. cinereus*
- 14b** Snout relatively narrow, its width, 19–22% HL, internasal width 16–18% HL; spinules on body scales arranged in narrowly divergent rows *C. filifer*

Coryphaenoides acrolepis (Bean, 1884)

[Japanese name: Ibara-hige]

(Figs. 91, 92A, 93–94, 95A–C; Table 6; Appendix 3-5D)

Macrurus [sic] *acrolepis* Bean, 1884:362 [original description; holotype: USNM 32496, from off Cape Flattery, Washington (from stomach of seal)].

Macrurus firmisquamis Gill & Townsend, 1897:234 (original description; holotype: USNM 48772, from southwest of Pribilof Islands, Bering Sea).

Chalinura spinulosa Gilbert & Burke, 1912:92, figs. 36, 36a (original description; holotype: USNM 74397, from off Avatcha Bay, Albatross sta. 4797).

Coryphaenoides bona-nox Jordan & Thompson, 1914:305, pl. XXXVIII, figs. 1, 1a [original description; holotype: FMNH 57520 (ex. CM 6462), from Sagami Bay].

Nematonurus bona-nox: Gilbert & Hubbs 1916:162 (new combination; brief description; 2 spec. from Eno-shima Island).

Coryphaenoides altipinnis (not Günther 1877): Schmidt 1931:154 (description 1 spec. from Sagami Bay) [re-identified here from description].

Coryphaenoides (Nematonurus) acrolepis: Okada *et al.* 1935:258, pl. 154, fig. 3 (first record from Japan; description; 1 spec. probably from Choshi; new Japanese name: “Ibara-hige”).

Coryphaenoides spinulosus: Taranetz 1937:169 (new combination; listed; Soviet Far East); Ueno 1971:101 (listed; northern Kuril Islands; new Japanese name: “Toge-sokodara”).

Coryphaenoides acrolepis: Taranetz 1937:170 (listed; Soviet Far East); Okamura 1970a:125, pl. XXVII, text-fig. 51 (description; biological notes; 11 spec. from Pacific off Japan from Hokkaido to Kamogawa); Ueno 1971:101 (listed; Pacific off Hokkaido); Tominaga & Uyeno 1981:488 (listed; Japan); Shiogaki 1982:33 (listed; Aomori Pref.); Sawada 1983:111, 248, fig. 63 (brief description; 5 spec. from Pacific off Tohoku and Okhotsk Sea; photos based on HUMZ 78122 and HUMZ 79709); Amaoka & Nakaya 1983:44, fig. 59 (compiled; northern Japan); Okamura 1984b:95, pl. 82, fig. C (compiled); Okamura 1988:95, pl. 82, fig. C (compiled); Iwamoto 1990:202, fig. 472 (synopsis); Maeda & Maruyama 1991:366 (listed; Hokkaido); Nakabo 1993:362 (in key; Japan); Masuda & Kobayashi 1994:51, fig. 1 (in situ observation; off Kamaishi); Amaoka *et al.* 1995:101, fig. 150 (compiled; northern Japan); Hori 1996:28 (listed; Ibaraki Pref.); Shinohara *et al.* 1996:169 (2 spec. listed from Pacific off Tohoku); Okamura 1997:124, fig. (compiled); Funabashi 1998:85 (listed; Ibaraki Pref.); Nakabo 2000:427 (in key; Japan); Zama 2001:40 (listed; Miyagi Pref.); Nakabo 2002:427 (in key; Japan); Maeda & Tsutsui 2003:488 (listed; Hokkaido); Shiogaki *et al.* 2004:55 (listed; Pacific off Aomori Pref.); Senou *et al.* 2006:420 (listed; Sagami Sea); Senou 2008:367, fig. 31.34A–B (in situ observation; east of Hokkaido and Sagami Bay); Kitagawa *et al.* 2008:40, unnumbered fig. (brief description; spec. from Pacific off Tohoku); Shinohara *et al.* 2009:708 (1 spec. listed from Pacific off Tohoku); Amaoka *et al.* 2011:129, fig. 162 (compiled; Hokkaido); Nakabo &

Kai 2013:503 (in key; Japan); Ikeda & Nakabo 2015:320, pl. 66, figs. 1–7 (brief description; 3 spec. from Kii Channel off Wakayama Pref.); Shinohara *et al.* 2020:6 (description; 7 spec. from Kurile-Kamchatka Trench, at midwater depths of 883–1350 m; stomach contents); Amaoka *et al.* 2020:161, fig. 216 (listed; Hokkaido); Motomura 2020:39 (listed; Japan).

Nematonurus acrolepis: Okada & Matsubara 1938:448, pl. 110, fig. 1 (in key; Japan; *C. bonanox* a junior synonym of *C. acrolepis*); Okada 1955:424, fig. 383; Matsubara 1955:1308 (in key; Japan); Ueno 1965:19 (listed; Pacific off Hokkaido); Matsubara 1965:504 (compiled; Japan); Nakajima 1975:55 (listed; Enshu-nada); Hikita 1981:78, fig. 1 (spec. from Okhotsk Sea off Shiretoko Peninsula).

[?] *Coryphaenoides (Chalinura) angustifrons* Rass, 1955:331, fig. 4 (original description; holotype: ZIN 44885, from Okhotsk Sea, 52°20'N, 150°20'E, 1030 m depth; 9 paratypes from Kuril-Kamchatka Trench and Okhotsk Sea).

Coryphaenoides (Coryphaenoides) acrolepis: Okamura 1970b: table 1 (listed; Japan).

Nematonurus [sic] *acrolepis*: Maruyama 1971:33 (listed; Iwate Pref.).

[?] *Coryphaenoides angustifrons* [sic]: Ueno 1971:101 (listed; northern Kuril Islands and southern Okhotsk Sea; new Japanese name: “Chishima-sokodara”).

Ventrifossa sp. 1: Masuda & Kobayashi 1994:50, fig. 4 (in situ observation; Sagami Bay) [re-identified here from figure].

Ventrifossa sp. 2: Masuda & Kobayashi 1994:50, fig. 5 (in situ observation; off Kamaishi) [re-identified here from figure].

Ventrifossa sp. 3: Masuda & Kobayashi 1994:50, fig. 6 (in situ observation; off Kamaishi) [re-identified here from figure].

Diagnosis. Pelvic-fin rays usually 8 (rarely 7 or 9). Snout conical, moderately protruding beyond upper jaw. Tip and lateral angles of snout with stout tubercles; scales along head ridges slightly thickened and enlarged. Mouth large, posterior margin of upper jaw extending to below posterior 1/3 of orbit or beyond; upper-jaw length 33–40% HL; lateral corner of mouth not restricted by skin folds. Outermost gill slit moderately wide, length 15–21% HL. Barbel well developed, length 11–19% HL. Teeth small, bluntly pointed, in broad bands in both jaws, outer premaxillary series slightly enlarged. Body scales covered with short, reclined, conical spinules in widely divergent rows; tip of last spinule in each row extending well beyond posterior scale margin. Transverse scale rows below first dorsal-fin midbase 7–10. Snout fully scaled dorsally, but variously naked ventrally (underside of snout fully scaled in some larger specimens). Interdorsal space less than first dorsal-fin base length. Height of first dorsal fin less than HL (60–94% HL); second spinous ray not prolonged, finely serrated along its leading edge; first dorsal-fin rays II, 8–11. Body uniformly blackish in adults.

Material examined. 39 specimens. **Holotype of *Macrurus* [sic] *acrolepis*:** *USNM 32496 (badly deteriorated; originally 365 mm TL, fide Bean 1884:362), off Cape Flattery, Washington (from stomach of seal), USA,

northeastern Pacific, coll. J.G. Swan, 1 Feb. 1883. **Holotype of *Coryphaenoides bonanox*:** FMNH 57520 (ex. CM 6462, 141 mm HL, 581+ mm TL), Sagami Bay, Kanagawa Pref., Japan, date unknown. **Holotype of *Chalinura spinulosa*:** USNM 74397 (badly deteriorated; originally 41 cm TL, fide Gilbert & Burke 1912:92), off Avacha Bay, east coast of Kamchatka, northwestern Pacific, 52.6250°N, 158.8333°E, 682 ftm (1248 m), *Albatross* sta. 4797, 9ft *Albatross-Blake* beam trawl, 20 Jun. 1906. **Non-types: Japan:** BSKU 49891 (1, 92.1 mm HL, 462+ mm TL), BSKU 49893 (1, 166 mm HL, 808+ mm TL), BSKU 49894 (1, 130 mm HL, 655+ mm TL), BSKU 49895 (1, 142 mm HL, 710+ mm TL), BSKU 49896 (1, 103 mm HL, 512+ mm TL), BSKU 49899 (1, 148 mm HL, 776+ mm TL), 1500 m, R/V *Hakuho-maru*, cr. KH-91-6, sta. 12, 16 Nov. 1991; BSKU 83765 (1, 78.1 mm HL, 316+ mm TL), off Tohoku, 450 m, FRV *Wakataka-maru*, sta. E, otter trawl, Oct. 1997; BSKU 19556 (1, 107 mm HL, 565 mm TL), off Ito, Sagami-nada, 34.9467°N, 139.3500°E, 1470–1500 m, FRV *Soyo-maru*, sta. T21, beam trawl, 15 Dec. 1964; BSKU 103028 (1, 138 mm HL, 714+ mm TL), off Manazuru, Sagami Bay, 450 m, rod and reel, coll. T. Okamoto, date unknown; BSKU 14188 (1, 104 mm HL, 478+ mm TL), BSKU 14192 (1, 91.2 mm HL, 430+ mm TL), off Odawara, Sagami Bay, 35.2014°N, 139.2003°E, 670–750 m, 14 Apr. 1969; BSKU 97738 (1, 77.5 mm HL, 385+ mm TL), southeast of Cape Erimomisaki, Japan Trench, 41.2353°N, 144.0357°E, 2977–3014 m, R/V *Tansei-maru*, cr. KT-93-15, sta. M7, 3-m beam trawl, coll. Y. Machida, 17 Nov. 1993; BSKU 35672 (1, 168 mm HL, 867+ mm TL), off Kuji, 40.3683°N, 143.3800°E, 1970–2000 m, R/V *Hakuho-maru*, cr. KH-81-4, sta. 3, 10 Jul. 1981; BSKU 35681 (1, 83.9 mm HL, 422+ mm TL), BSKU 35682 (1, 108 mm HL, 520+ mm TL), BSKU 35684 (1, 138 mm HL, 683+ mm TL), BSKU 35685 (1, 147 mm HL, 717+ mm TL), BSKU 35688 (1, 141 mm HL, 734+ mm TL), off Kinkazan, 38.7333°N, 143.1850°E, 1950–1960 m, R/V *Hakuho-maru*, cr. KH-81-4, sta. 8, 25 Jul. 1981; BSKU 96504 (1, 147 mm HL, 705+ mm TL), off Kinkazan, 38.4640°N, 143.0518°E, 1853–1906 m, R/V *Tansei-maru*, cr. KT-08-27, sta. K-2, 3-m ORE beam trawl, coll. H. Endo and N. Nakayama, 23 Oct. 2008; BSKU 96421 (1, 80.4 mm HL, 398 mm TL), *BSKU 96425 (1, 62.1 mm HL, 323+ mm TL), off Miyako, 39.5668°N, 142.9008°E, 1586–1617 m, R/V *Tansei-maru*, cr. KT-08-27, sta. M-2, 3-m ORE beam trawl, coll. H. Endo and N. Nakayama, 21 Oct. 2008; BSKU 97783 (1, 67.2 mm HL, 348 mm TL), off Miyako, 39.5747°N, 142.6852°E, 1132–1170 m, R/V *Tansei-maru*, cr. KT-93-15, sta. O4, 3-m ORE beam trawl, coll. Y. Machida, 13 Nov. 1993; BSKU 97743 (1, 85.4 mm HL, 403+ mm TL), off Kuji, 40.0613°N, 142.6825°E, 1005–1016 m, R/V *Tansei-maru*, cr. KT-93-15, sta. M3, 3-m beam trawl, coll. Y. Machida, 18 Nov. 1993; BSKU 97841 (1, 78.2 mm HL, 365+ mm TL), BSKU 97846 (1, 67.9 mm HL, 332+ mm TL), off Otsuchi, 39.0538°N, 142.6342°E, 1195–1202 m, R/V *Tansei-maru*, cr. KT-89-03, sta. SR91, 10 Mar. 1989; BSKU 97820 (1, 90.8 mm

FIGURE 91. Fresh specimens of *Coryphaenoides acrolepis*. (A) BSKU 45528, 45.0 mm HL, 206+ mm TL, KT-89-03, sta. SR89; (B) BSKU 96425, 62.1 mm HL, 323+ mm TL, off Miyako, Iwate Pref., Pacific, 1586–1617 m depth; (C) BSKU 96421, 80.4 mm HL, 398 mm TL, collected with BSKU 96425; (D) BSKU 96393, 98.0 mm HL, 476+ mm TL, south of Cape Erimomisaki, Hokkaido, Pacific, 2022–2027 m depth; (E–G) BSKU 103028, 138 mm HL, 714+ mm TL, Sagami Bay, Pacific, 450 m depth. (A–E) Lateral views; (F) dorsal and (G) ventral views of the head and trunk. [Photos: (A) BSKU; (B–D) H. Endo and N. Nakayama; (E–G) N. Nakayama]

HL, 408+ mm TL), off Otsuchi, 39.1444°N, 142.3748°E, 847–935 m, R/V *Tansei-maru*, cr. KT-89-03, sta. SR89, 10 Mar. 1989; *BSKU 45528 (1, 45.0 mm HL, 206+ mm TL), BSKU 45529 (1, 71.9 mm HL, 363 mm TL), off Otsuchi, 39.1444°N, 142.3748°E, 847–935 m, R/V *Tansei-maru*, cr. KT-89-03, sta. SR89, 3-m ORE beam trawl, 10 Mar. 1989; BSKU 97728 (1, 52.2 mm HL, 277 mm TL), BSKU 97729 (1, 59.1 mm HL, 297+ mm TL), BSKU 97784 (1, 90.2 mm HL, 440+ mm TL), BSKU 97785 (1, 72.1 mm HL, 360+ mm TL), BSKU 97786 (1, 78.6 mm HL, 382+ mm TL), south-southeast of Akkeshi, 42.6052°N, 144.9938°E, 1113–1161 m, R/V *Tansei-maru*, cr. KT-93-15, sta. T8, 3-m ORE beam trawl, coll. Y. Machida, 16 Nov. 1993; BSKU 96393 (1, 98.0 mm HL, 476+ mm TL), BSKU 96394 (1,

146 mm HL, 727+ mm TL), BSKU 96395 (1, 129 mm HL, 663+ mm TL), south of Cape Erimomisaki, 41.6675°N, 143.3268°E, 2022–2027 m, R/V *Tansei-maru*, cr. KT-08-27, sta. S-2, 3-m ORE beam trawl, coll. H. Endo and N. Nakayama, 20 Oct. 2008.

Counts and measurements. See Table 6.

Size. Attains about 97 cm TL (NSMT-P 102193, 973+ mm TL, off Boso Peninsula, Japan).

Development. Larval morphology was described by Stein (1980).

Distribution. Widely distributed in the North Pacific from Japan to North America including the Okhotsk and Bering Seas as well as the Emperor Seamounts (Iwamoto & Stein 1974; Borets 1986; Iwamoto 1990; Mecklenburg

TABLE 6. Morphometric measurements and meristic counts* for *Coryphaenoides acrolepis*.

Nominal species	<i>C. acrolepis</i>		<i>C. bonanox</i>	<i>C. spinulosus</i>	
Type status	HT	NT	HT	HT	
Source	Ori. descr.	This study	This study	Ori. descr.	This study
Number of specimens	<i>n</i> = 1	<i>n</i> = 34	<i>n</i> = 1	<i>n</i> = 1	<i>n</i> = 1
Measurements (mm)					
TL	635+	277–867+	581+	410	–
HL	139	52.2–168	141	80.5	–
% of HL (% of PRL)					
Snout length	29	27–32 (37–46)	28 (37)	26	–
Orbit diameter	29	23–32 (31–45)	27 (36)	30	–
Postorbital length	–	42–51 (59–70)	49 (65)	–	–
Postrostral length	–	69–76 (–)	75 (–)	–	–
Orbit–preopercle distance	–	35–43 (48–59)	42 (56)	–	–
Suborbital width	–	10–13 (14–19)	12 (16)	–	–
Upper-jaw length	36	33–40 (46–55)	38 (50)	39	–
Length of rictus	–	28–35 (39–48)	32 (43)	–	–
Length of PMX tooth band	–	23–29 (32–39)	28 (37)	–	–
Preoral length	–	13–21 (18–30)	15 (20)	–	–
Dist. tip and lat. angle of snout	–	13–17 (17–23)	12 (16)	–	–
Snout width	–	19–25 (25–35)	19 (25)	–	–
Internasal width	–	14–18 (19–25)	16 (21)	–	–
Interorbital width	22	19–26 (26–36)	23 (30)	21	–
Body width over P. bases	–	45–67 (62–92)	53 (70)	–	–
Body depth at 1D. origin	–	65–94 (89–129)	– (–)	80	–
Body depth at A. origin	49	51–73 (70–101)	72 (96)	–	–
Pre-V. length	91	102–122 (140–173)	101 (135)	–	–
Pre-anus length	–	139–173 (190–242)	151 (201)	–	–
Pre-A. length	180	143–178 (196–249)	160 (214)	–	–
Isthmus–V. distance	–	37–47 (52–65)	43 (58)	–	–
Isthmus–anus distance	–	75–107 (104–148)	– (–)	–	–
Isthmus–A. distance	–	81–115 (111–157)	– (–)	–	–
V.–A. distance	–	44–69 (60–96)	62 (82)	–	–
Anus–A. distance	–	4–8 (5–12)	9 (12)	–	–
V. length	56	46–73 (64–101)	73 (97)	50	–
P. length	47	45–61 (63–82)	58 (77)	53	–
Pre-1D. length	107	108–118 (146–159)	113 (151)	111	–
Height of 1D.	45+	60–94 (81–132)	60 (80)	80**	–
Length of 1D. base	29	22–29 (29–40)	24 (32)	25	–
Interdorsal length	–	8–21 (11–29)	18 (24)	13	–
Length of gill slit	–	15–21 (21–28)	17 (23)	–	–
Length of posterior nostril	–	6–10 (9–14)	7 (10)	–	–
Barbel length	–	11–19 (15–27)	13 (18)	17	–
Counts					
1D. rays	II,11	II,8–11	II,9	II,9	–
P. rays	i20	i15–i22	i20–i21	19***	i18
V. rays	8	7–9	8	8	8
Outer GR on first arch	–	3–7	5–7	6	–
Inner GR on first arch	–	12–15	13	–	–
Outer GR on second arch	–	11–14	12–13	–	–
Inner GR on second arch	–	12–15	14–15	–	–
Longitudinal scales	–	39–48	41	–	–
Scales below 1D. origin	ca. 7 or 8	8.5–14	–	ca. 8	–
Scales below 1D. midbase	–	7–10	8.5	–	8.5
Scales below 2D. origin	–	6.5–11	8.5	–	9.5 or 10.5
Scales above A. origin	–	24.5–35.5	–	–	–

...See the next page for the table footer

*Abbreviations: 1D.—first dorsal fin; 2D.—second dorsal fin; A.—anal fin; GR—gill rakers; HT—holotype; NT—non-type; P.—pectoral fin; PMX—premaxillary; V.—pelvic fin

**Length of the second spinous ray

***Probably i18

et al. 2002; this study). A single specimen also recorded from the southwestern Atlantic off the Falkland Islands (Laptikhovsky *et al.* 2013). Depth range about 300–3000 m. In Japanese waters, known from off the Pacific coasts southward to Cape Omaezaki (138.22°E) and the southern Okhotsk Sea at depths of 450–3014 m (Appendix 3-5D). Common, especially in the north of Choshi (35.74°N; Chiba Pref.).

FIGURE 92. Type specimens of two species of *Coryphaenoides*. (A) Holotype of *Macrurus* [sic] *acrolepis*, USNM 32496, badly deteriorated, off Cape Flattery, Washington, USA, northeastern Pacific; (B) syntype of *Chalinura filifera*, USNM 48626, disintegrated, off Queen Charlotte Island, British Columbia, Canada, northeastern Pacific, 2905 m depth. [Photos: N. Nakayama]

Remarks. *Coryphaenoides acrolepis* was originally described by Bean (1884) based on a single specimen obtained from the stomach of a seal found in the northeastern Pacific off Cape Flattery, Washington, United States. As mentioned by Bean (1884:362), the holotype is “mutilated” and “a small portion of the tail is wanting, and the belly is almost wholly digested” (Fig. 92A).

Coryphaenoides acrolepis has been well described by previous authors; for a full description see Okamura (1970a) and Iwamoto & Stein (1974). This species is one of the most common grenadiers in the North Pacific (Iwamoto 1990:202).

Nomenclatural discussion. Jordan & Thompson

(1914) described *Coryphaenoides bonanox* (originally spelled as “*bona-nox*”) from a single specimen collected in Sagami Bay (Fig. 93), and many authors synonymized the species with *C. acrolepis* without explanation (e.g., Okada & Matsubara 1938; Okamura 1970a; Iwamoto & Stein 1974; Iwamoto 1990; Mecklenburg *et al.* 2002). The holotype of *C. bonanox* was examined here to investigate this synonymy. In the original description of *C. bonanox*, Jordan & Thompson (1914:305) wrote that “the dentition and scales of this species distinguish it from previously known forms”. Examination of the holotype confirmed that teeth of the specimen are small, sharp, slightly incurved, arranged in long tapered bands in the both jaws, with an outer row of premaxillary teeth enlarged. Also, its body scales are covered with short, reclined, conical spinules in 7–9 widely divergent rows. These features are typically found in the *C. acrolepis* specimens examined, and thus the two species cannot be distinguished by these characters. Counts and measurements also show no significant differences between the two species (Table 6). This study reconfirms that *C. bonanox* is a junior synonym of *C. acrolepis*.

Gilbert & Burke (1912) described *Coryphaenoides spinulosus* (as *Chalinura spinulosa*) from a single specimen collected from off Avacha Bay, southern Kamchatka (Fig. 94). This is a poorly known grenadier, and no additional specimens have been found since the original description. According to Fricke *et al.* (2020), the species is a junior synonym of *C. acrolepis*, but no published data exist to support this synonymy. Although *C. spinulosus* has not been recorded from Japanese waters, the holotype of this species was examined here to determine its taxonomic status. Unfortunately, the specimen is badly deteriorated (Fig. 94), and it was impossible to obtain meristic and morphometric characters except for fin ray and scale row counts. However, adequate information is available from its original description, with excellent illustrations of the holotype and a body scale from the dorsum of the body (Gilbert & Burke 1912: figs. 36, 36a). According to Gilbert & Burke’s (1912) description, *C. spinulosus* can be distinguished from *C. acrolepis* by having a somewhat shorter snout (26% HL vs. 27–32%; Table 6), but otherwise the two species are very similar to each other. Their figure also shows that the snout of *C. spinulosus* is relatively high and only slightly protrudes beyond the upper jaw (fig. 36). In *C. acrolepis*, the snout is usually much longer, giving a more conical appearance to the anterior portion of the head (Fig. 91). However, these subtle differences most likely reflect individual variation or the poor condition of the holotype of *C. spinulosus*.

FIGURE 93. Preserved specimen of *Coryphaenoides acrolepis*. FMNH 57520, holotype of *C. bonanox*, 141 mm HL, 581+ mm TL, Sagami Bay, Pacific. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

FIGURE 94. Preserved specimen of *Coryphaenoides acrolepis*. USNM 74397, holotype of *Chalinura spinulosa*, not measured due to its poor condition, off Avacha Bay, east of Kamchatka, Pacific, 1247 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

Consequently, *C. spinulosus* is herein regarded as a junior synonym of *C. acrolepis*.

Relationships and comparisons. *Coryphaenoides acrolepis* belongs to the subgenus *Coryphaenoides* (sensu Iwamoto 1990). According to Iwamoto (1990:203), the species is closely related to *C. filifer* (Gilbert, 1896) with which it occurs sympatrically in the North Pacific. In meristic and morphometric features, *C. acrolepis* differs from *C. filifer* in having lower counts of first dorsal-fin rays (II,8–11 vs. II,12–15) and a slightly longer chin barbel (11–19% HL vs. 8–11%). Spinulation

of body scales further distinguishes the two species, with *C. acrolepis* having much stouter spinules than *C. filifer* (Fig. 95A–C vs. Fig. 109). Furthermore, small but conspicuous pores are present along each canal of the cephalic sensory system in *C. acrolepis*, whereas the pores of *C. filifer* are too tiny to observe without a microscope, and none appear to be present along the postorbital and supratemporal canals.

FIGURE 95. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of two species of *Coryphaenoides*. (A–B) *Co. acrolepis*, BSKU 103028, 138 mm HL; (C) *Co. acrolepis*, USNM 74397, holotype of *Chalinura spinulosa*, not measured due to its poor condition; (D) *Co. altipinnis*, BMNH 1887.12.7.90, holotype, 102 mm HL. (A, C–D) Views from above; (B) oblique view. [Photos: (A–B) NSMT; (C–D) N. Nakayama]

Coryphaenoides altipinnis Günther, 1877

[Japanese name: Hokake-dara]

(Figs. 95D, 96; Appendix 3-5E)

Coryphaenoides altipinnis: Günther, 1877:439 [original description; holotype: BMNH 1887.12.7.90, from “south of Japan” (southeast of Boso Peninsula), *Challenger* sta. 237, in 1875 fth (3429 m)]; Jordan & Snyder 1901:120 (listed; Japan); Jordan & Gilbert in Jordan & Starks 1904:608 [after Günther (1877, 1887)]; Jordan *et al.* 1913:416 (listed; Japan; new Japanese name: “Hokake-dara”); Okada & Matsubara 1938:449 (in key; Japan); Kuroda 1951:391 (questionable; listed; Suruga Bay); Kuroda 1952:176 (questionable; 1 spec. from Suruga Bay); Matsubara 1955:1309 (in key; Japan); Okamura 1970a:129 [brief description after Günther (1877, 1887)]; Tominaga & Uyeno 1981:488 (listed; Japan); Okamura 1984b:95, pl. 345, fig. G (compiled); Okamura 1988:95, pl. 345, fig. G (compiled); Nakabo 1993:364 (in key; Japan); Nakabo 2000:428 (in key; Japan); Nakabo 2002:428 (in key;

Japan); Motomura 2020:39 (listed; Japan).

Macrurus altipinnis: Günther 1887:138, pl. XXXIX, fig. A [description; in part, holotype, BMNH 1887.12.7.90; other spec. represent *C. nasutus* and *Nezumia proxima*].

Coryphaenoides (Coryphaenoides) altipinnis: Okamura 1970b: table 1 (listed; Japan).

Coryphaenoides altipennis [sic]: Eschmeyer 1998:79 (listed); Nakabo & Kai 2013:504 (in key; Japan).

Diagnosis. Pelvic-fin rays 9. Snout short, slightly protruding beyond upper jaw. No prominent modified scales on head ridges and tip and lateral angles of snout. Mouth large, posterior margin of upper jaw almost reaching vertical through hind rim of orbit; upper-jaw length 41% HL; lateral corner of mouth not restricted by skin folds. Outermost gill slit moderately wide, length 24% HL. Barbel short, length 9% HL. Teeth small, conical, in broad tapered bands in both jaws, outer premaxillary series enlarged. Body scales covered with short, greatly

reclined, needle-like spinules in narrowly divergent rows; tip of last spinule in each row barely extending beyond posterior scale margin. Transverse scale rows below first dorsal-fin midbase 11. Snout almost fully scaled. Interdorsal space less than first dorsal-fin base length. Height of first dorsal fin 93% HL; second spinous ray slightly elongate, finely serrated along its leading edge; first dorsal-fin rays II,13.

Material examined. 1 specimen. **Holotype of *Coryphaenoides altipinnis*:** BMNH 1887.12.7.90 (1, 102 mm HL, 473+ mm TL), southeast of Boso Peninsula, Chiba, Pref., Japan, 34.6167°N, 140.5333°E, 1875 ftn (3429 m), *Challenger* sta. 237, trawl, 17 Jun. 1875.

Counts and measurements. Counts: first dorsal-fin rays II,13; pectoral-fin rays i21; pelvic-fin rays 9; gill rakers on first arch (outer/inner) 8–10/14, on second arch 12–13/14; longitudinal scales 38; transverse scale rows below first dorsal-fin origin 15, below first dorsal-fin midbase 11, below second dorsal-fin origin 12.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 25 (32); orbit diameter 25 (32); postorbital length 54 (69); postrostral length 77; orbit–preopercle distance 48 (62); suborbital width 11 (14); upper-jaw length 41 (54); length of rictus 35 (45); length of premaxillary tooth band 33 (42); preoral length 10 (13); distance between tip and lateral angle of snout 13 (16); snout width 21 (27); internasal width 17 (22); interorbital width 25 (32); body width over pectoral-fin bases 49 (63); body depth at first dorsal-fin origin 77 (99); body depth at anal-fin origin 56 (72); prepelvic length 100 (130); preanus length 163 (211); preanal length 168 (217); isthmus–pelvic distance 46 (59); pectoral-fin length 56 (72); predorsal length 106 (137); height of first dorsal fin 93 (120); length of first dorsal-fin base 31 (40); interdorsal length 27 (35); length of gill slit 24 (31); length of posterior nostril 8 (10); barbel length 9 (12).

Redescription of holotype. General features are shown in Fig. 96. Body deepest at first dorsal-fin origin, progressively tapering to long tail. Trunk short, deep, compressed, width over pectoral-fin bases 1.6 in depth at first dorsal-fin origin. Head large, HL about 4.7 in TL; head bones moderately stout; flesh firm. Dorsal contour of head steeply elevated from snout tip to above nasal fossa, then straightened from this point to first dorsal-fin origin. Snout short, conical in preserved specimen, but probably more or less inflated when fresh; its tip low, slightly protruding beyond upper jaw, ventral contour oblique; snout length equal to orbit diameter. Orbit moderately large, circular, greatest diameter 2.2 in postorbital length. Interorbital space broad, width equal to orbit diameter. Mouth large, oblique, subinferior, upper-jaw length 0.6 in orbit diameter; posterior margin of maxillary almost reaching vertical through hind rim of orbit; lateral corner of mouth not restricted by lip folds; lips thin. Suborbital region moderately flat, only weakly ridged. Preopercle

large, hind margin inclined from vertical, forming broad lobe posteroventrally; preopercular ridge poorly marked. Interopercle mostly hidden behind preopercle. Subopercle and opercle smoothly adjoined, forming posterior margin of gill cover. Gill membranes narrowly connected across isthmus; gill opening wide, extending ventrally to below posterior margin of lower jaw. Outermost gill slit moderately wide, length equal to orbit diameter. Gill rakers developed into small tubercles, tipped with short, slender spines. Barbel short, slender, length 2.7 in orbit diameter.

Anus immediately anterior to anal-fin origin; periproct poorly developed. Ventral light organ absent.

Teeth short, conical, slightly incurved, in long tapered band in both jaws. Premaxillary teeth arranged in about 5 rows at widest point near symphysis, in about 2 rows posteriorly; outermost series enlarged; posterior margin of tooth band almost reaching lateral corner of mouth. Dentition in lower jaw similar to that in premaxillary, but no teeth especially enlarged; mandibular band much shorter than premaxillary band.

Body scales small, thin, highly deciduous, covered with short, greatly reclined, needle-like spinules in narrowly divergent, discrete rows (Fig. 95D); those on dorsum below interdorsal space with about 5 spinule rows; length of spinules gradually increasing posteriorly, but last spinule in each row barely extending beyond posterior scale margin; lateral buttresses not developed; reticulate structure absent. Body fully and uniformly scaled except for fins.

Neither head ridges nor tip and lateral angles of snout armed with coarsely modified tubercles. Head almost fully scaled; naked areas confined to nasal fossa, anterior portion of mandibular rami, lips, chin barbel, and gular and branchiostegal membranes; ventral surface of snout completely scaled, except for narrow naked area above upper lip.

Cephalic sensory canals broad; infraorbital and mandibular canals probably with small open pores. Lateral line complete, not interrupted throughout.

First dorsal-fin origin slightly posterior to pectoral-fin base; second spinous ray of first dorsal fin slightly prolonged, its tip extending to base of 9th second dorsal-fin ray when laid back; height of first dorsal fin 3.0 times as long as its base length; leading edge of second spinous ray finely serrated. Second dorsal-fin origin more or less above anal-fin origin; interdorsal space 0.9 times as long as first dorsal-fin base length. Pectoral fin long, but its tip not reaching vertical through anal-fin origin. Pelvic fin inserted slightly anterior to pectoral-fin base, outermost elongate (its tip slightly damaged). Anal fin well developed, much deeper than second dorsal fin.

Color when fresh. Unknown.

Color in alcohol (Fig. 96). Head and body uniformly pale brown; fins darker; gular and branchiostegal membranes mostly pale, posterior margins of gill membranes prominently blackish.

FIGURE 96. Holotype of *Coryphaenoides altipinnis*. BMNH 1887.12.7.90, 102 mm HL, 473+ mm TL, southeast of Boso Peninsula, Chiba Pref., Pacific, 3429 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

Size. To at least 47 cm TL.

Distribution. So far known only from the holotype collected from off the Boso Peninsula at a depth of 3429 m (Appendix 3-5E). Very rare.

Remarks. *Coryphaenoides altipinnis* was originally described by Günther (1877) from the “south of Yeddo” (= southeast of the Boso Peninsula), Japan, at a depth of 3429 m (*Challenger* sta. 237). According to Günther (1887), only a single specimen was collected from the type locality (BMNH 1887.12.7.90, holotype; Fig. 96). He (1887) also reported two additional specimens from off Japan at a depth of 1034 m (*Challenger* sta. 235), both of which were re-identified in this study, one as a different species of *Coryphaenoides* (probably *C. nasutus*; BMNH 1887.12.7.91, 27.0 mm HL, 128+ mm TL) and the other as *Nezumia proxima* (BMNH 1887.12.7.92, 24.0 mm HL, 149+ mm TL). Unfortunately, no additional specimens were discovered during the course of this study. The paucity of available materials could be attributable to its apparent preference for very deepwater habitats. Because Günther’s (1877, 1887) description is brief, the holotype is redescribed here for future studies. It should be noted that the original spelling of the specific epithet is “*altipinnis*”, instead of “*altipennis*”.

Based on a 386 mm TL specimen, Kuroda (1952) reported *C. altipinnis* from Suruga Bay (see also Kuroda 1951). Citing Kuroda’s record, Shinohara & Matsuura (1997) included this species in their list of deep-sea fishes occurring in this area. Although Kuroda’s specimen is presumably lost, it was likely a misidentification, considering that the specimen was obviously collected by local fisheries confined to upper slope depths.

Relationships and comparisons. *Coryphaenoides*

altipinnis belongs to the subgenus *Coryphaenoides* (sensu Iwamoto 1990), and is closely similar to *C. cinereus* (Gilbert, 1896) and *C. filifer* (Gilbert, 1896). The absence of prominent modified scales on the snout readily distinguishes the species from the latter two (vs. snout tipped with large prominent tubercles). It further differs from these two species in that the tip of the pectoral fin does not reach a vertical through the anal-fin origin (vs. extending to or beyond this level), and in having more scale rows below the first dorsal-fin origin (15 vs. 10–14 in *C. cinereus* and 8.5–13.5 in *C. filifer*). Regarding morphometric characters, it differs from *C. cinereus* and *C. filifer* in the preoral length (10% HL vs. 13–19% and 11–13%), postorbital length (54% HL vs. 43–52% and 48–53%), and pectoral-fin length (56% HL vs. 60–68% and 56–70%); from *C. cinereus* in the snout length (25% HL vs. 26–29%) and snout width (21% HL vs. 34–30%); and from *C. filifer* in the orbit–preopercle distance (48% HL vs. 42–46%).

Okamura (1970a:136) suggested a close similarity between *C. altipinnis* and *C. spinulosus* (Gilbert & Burke, 1912), the latter of which is regarded here as a junior synonym of *C. acrolepis* (Bean, 1884). *Coryphaenoides altipinnis* is readily distinguished from *C. acrolepis* by several meristic characters, including first dorsal-fin rays (II,13 in *C. altipinnis* vs. II,8–11 in *C. acrolepis*), outer gill rakers on the first arch (8–10 vs. 3–7), and transverse scale rows (below first dorsal-fin origin 15 vs. ≤14; below first dorsal-fin midbase 11 vs. 7–10; below second dorsal-fin origin 12 vs. 6.5–11). It further differs from *C. acrolepis* in having a shorter snout (snout length 25% HL vs. 27–32%; preoral length 10% HL vs. 13–21%), a smaller barbel (9% HL vs. 11–19%), a longer postorbital (54% HL vs.

42–51%) and orbit–preopercle distance (48% HL vs. 35–43%), a broader interdorsal space (27% HL vs. 8–21%), and a wider gill slit (24% HL vs. 15–21%). Body-scale structure is also different between the two species, with *C. altipinnis* having much weaker spinulation than *C. acrolepis* (Fig. 95D vs. A–C). In addition, cephalic sensory canals are completely closed in *C. altipinnis* except for a few tiny pores on the mandibular and infraorbital canals, whereas in *C. acrolepis*, small pores are serially present along all canals.

***Coryphaenoides armatus* (Hector, 1875)**

[Japanese name: Yoroidara]

(Figs. 97–98, 99A, 100A–B; Appendix 3-5F)

Macrurus armatus Hector, 1875:81 [original description; holotype: BMNH 1982.6.8.1, from ca. 200 miles west of Cape Farewell, New Zealand, in 400 ftm (732 m)].

Coryphaenoides variabilis Günther, 1878:27 [original description; lectotype: BMNH 1889.12.7.130, from Mid-Pacific, *Challenger* sta. 246, in 2050 ftm (3749 m); paralectotypes from sta. 146, 157, 246, 271, and 300; in part, see the Comments on type specimens of *C. leptolepis*; lectotype designated by Wilson & Waples (1983:1136)].

Macrurus asper Goode & Bean, 1883:196 [original description; 6 syntypes (MCZ and USNM), from northwestern Atlantic, *Blake* sta. 308 and 309, in 304–1242 ftm (556–2271 m)].

Macrurus goodii Günther, 1887:136 [original description based on type specimens of *M. asper* Goode & Bean, 1883; replacement name for the latter species preoccupied by *C. asper* Günther, 1877).

Coryphaenoides gigas Vaillant, 1888:232, pl. XX, figs. 2, 2a, 2b, 2c [original description; lectotype: MNHN 1886-0117, from northeastern Atlantic, in 4255 m; 1 paralectotype from northeastern Atlantic; lectotype designated by Eschmeyer (1998:645)].

Nematonurus cyclolepis Gilbert, 1896:458 [original description; 2 syntypes (CAS-SU and USNM), from off Queen Charlotte Island, British Columbia, northeastern Pacific, *Albatross* sta. 3342, in 1588 ftm (2904 m)].

Macrurus (Nematonurus) suborbitalis Gill & Townsend, 1897:234 [original description; holotype: USNM 48773, from southwest of Pribilof Islands, Bering Sea, *Albatross* sta. 3603, 1771 ftm (3129 m)].

Nematonurus abyssorum Gilbert, 1915:374, pl. 21, fig. 23 [original description; holotype: USNM 75827, from off Santa Catalina Island, northeastern Pacific, *Albatross* sta. 4390, in 1350–2182 ftm (2469–3990 m)].

Coryphaenoides armatus: Iwamoto 1990:205, fig. 478 (synopsis); Endo & Okamura 1992:433, fig. 1 (first record from Japan; brief description; 5 spec. from northwestern Pacific; new Japanese name: “Yoroidara”); Nakabo 1993:363 (in key; Japan); Okamura 1997:125, fig. 6 (compiled); Endo *et al.* 2000:16, photo C (in situ observation; Nankai Trough); Nakabo 2000:427 (in key; Japan); Nakabo 2002:427 (in

key; Japan); Shinohara *et al.* 2009:708 (listed; Pacific off Tohoku); Nakabo & Kai 2013:503 (in key; Japan); Amaoka *et al.* 2020:161, fig. 217 (listed; Hokkaido); Motomura 2020:39 (listed; Japan).

[?] *Coryphaenoides* sp.: Senou 2008:367, fig. 31.36 (in situ observation; Chishima Trench).

Diagnosis. Pelvic-fin rays 9–12 (usually 11). Snout rounded in adults (conical in young specimens), slightly protruding beyond upper jaw. Tip and lateral angles of snout armed with small scute-like scales; scales along head ridges slightly thickened but not especially enlarged. Mouth large, posterior margin of upper jaw extending to about vertical through hind rim of orbit; upper-jaw length 35–41% HL; lateral corner of mouth not restricted by skin folds. Outermost gill slit moderately wide, length 15–20% HL. Barbel long, stout at base, length 13–20% HL. Teeth slender, bluntly pointed; premaxillary teeth in 2 distinct rows, with outer series enlarged (inner row usually absent in larger specimens); mandibular teeth in uniserial row. Body scales covered with short, reclined, needle-like to knife-like spinules in subparallel to slightly divergent rows; tip of last spinule in each row extending to posterior scale margin or slightly beyond. Transverse scale rows below first dorsal-fin midbase 6–8. Top of snout narrowly naked posterior to leading edges; underside of head mostly naked, with narrow scaly patches below infraorbital ridges and on posterior parts of mandibular rami. Interdorsal space greater than first dorsal-fin base length. Height of first dorsal fin less than HL (64–90% HL); second spinous ray not prolonged, weakly serrated along its leading edge; first dorsal-fin rays II,9–10. Body uniformly dark brown.

Material examined. 18 specimens. **Japan:** BSKU 96414 (1, 44.5 mm HL, 249 mm TL), southeast of Cape Erimomisaki, Japan Trench, 41.2005°N, 143.9330°E, 2889–2995 m, R/V *Tansei-maru*, cr. KT-08-27, sta. S-4, 3-m ORE beam trawl, coll. H. Endo and N. Nakayama, 21 Oct. 2008; BSKU 96459 (1, 48.8 mm HL, 242+ mm TL), off Miyako, 39.5342°N, 143.6835°E, 3150–3201 m, R/V *Tansei-maru*, cr. KT-08-27, sta. M-4, 3-m ORE beam trawl, coll. H. Endo and N. Nakayama, 22 Oct. 2008; BSKU 96478 (1, 37.1 mm HL, 202+ mm TL), off Kinkazan, 38.4748°N, 143.3470°E, 2698–2814 m, R/V *Tansei-maru*, cr. KT-08-27, sta. K-3, 3-m ORE beam trawl, coll. H. Endo and N. Nakayama, 23 Oct. 2008; NSMT-P 30199 (1, 92.1 mm HL, 527+ mm TL), east of Tori-shima Island, western Pacific, 30.1687°N, 146.8190°E, 6273 m, R/V *Tokyo-daigaku-maru*, fish trap, 11 Oct. 1985; NSMT-P 97647 (1, 146 mm HL, 848+ mm TL), NSMT-P 97648 (1, 150 mm HL, 805+ mm TL), NSMT-P 97649 (1, 158 mm HL, 902+ mm TL), NSMT-P 97651 (1, 89.7 mm HL, 515 mm TL), NSMT-P 97669 (1, 103 mm HL, 576+ mm TL), NSMT-P 98008 (1, 110 mm HL, 587+ mm TL), off Miyako, 38.7983°N, 144.1345°E, 7340–7433 m, R/V *Hakuho-maru*, cr. KH-01-02, sta. TD-7, 30 Sept. 2001;

BSKU 35722 (1, 91.7 mm HL, 542+ mm TL), BSKU 35723 (1, 103 mm HL, 626+ mm TL), off Kuji, Japan Trench, 40.1133°N, 143.9700°E, 4200–4220 m, R/V *Hakuho-maru*, cr. KH-84-1, sta. 7-1, 15 Jul. 1981; BSKU 37297 (1, 141 mm HL, 753+ mm TL), BSKU 37299 (1, 107 mm HL, 642+ mm TL), east of Choshi, Japan Trench, 35.8367°N, 142.2083°E, 4100 m, T/V *Bosei-maru*, fish trap, 4 Nov. 1981; NSMT-P 30197 (1, 144 mm HL, 664 mm TL), NSMT-P 30200 (1, 118 mm HL, 633+ mm TL), east of Cape Inbozaki, 35.8333°N, 142.2000°E, 4100 m, T/V *Bosei-maru*, fish trap, 2 Nov. 1981. **Northeastern Pacific:** BSKU 44504 (1, 99.2 mm HL, 524+ mm TL), off New Port, 44.9500°N, 126.6167°W, 2850 m, 19 Feb. 1971; BSKU 44505 (1, 89.7 mm HL, 469+ mm TL), 2700 m, Feb. 1971.

Counts and measurements. Based on 18 specimens (37.1–158 mm HL, 202+–902+ mm TL). Counts: first dorsal-fin rays II,9–10; pectoral-fin rays i17–i21; pelvic-fin rays 9–12; gill rakers on first arch (outer/inner) 7–10/12–15, on second arch 11–14/11–14; longitudinal scales 29–44; transverse scale rows below first dorsal-fin origin 7.5–13, below first dorsal-fin midbase 6–8, below second dorsal-fin origin 7.5–10, above anal-fin origin 25.5–35.5; pyloric caeca 11.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 23–29 (28–39); orbit diameter 18–25 (24–33); postorbital length 51–61 (67–77); postrostral length 74–81; orbit–preopercle distance 44–49 (57–63); suborbital width 9–13 (12–18); upper-jaw length 35–41 (46–55);

length of rictus 29–36 (39–49); length of premaxillary tooth band 22–30 (29–40); preoral length 8–14 (10–19); distance between tip and lateral angle of snout 10–15 (12–20); snout width 17–24 (21–32); internasal width 13–20 (17–27); interorbital width 21–31 (28–41); occipital width 2–6 (3–8); body width over pectoral-fin bases 45–77 (60–96); body depth at first dorsal-fin origin 70–127 (93–160); body depth at anal-fin origin 57–105 (75–132); prepelvic length 102–118 (129–154); preanus length 164–200 (209–252); preanal length 170–207 (216–262); isthmus–pelvic distance 44–57 (58–72); isthmus–anus distance 99–154 (131–194); isthmus–anal distance 103–161 (135–202); pelvic–anal distance 61–106 (80–134); anus–anal distance 3–10 (4–13); pelvic-fin length 52–81 (68–109); pectoral-fin length 52–76 (69–102); predorsal length 114–136 (153–172); height of first dorsal fin 64–90 (81–118); length of first dorsal-fin base 27–37 (34–48); interdorsal length 37–88 (49–111); length of gill slit 15–20 (20–26); length of posterior nostril 3–7 (4–9); barbel length 13–20 (17–25); length of pyloric caecum 60 (78).

Size. Attains about 102 cm TL (Mecklenburg *et al.* 2002).

Development. Larval morphology was briefly described by Merrett (2006) based on an illustration given by Fahay & Markle (1984: fig. 140d; as *Coryphaenoides* sp.).

Distribution. Known from the continental slope to hadal depths of the world's oceans except the Arctic (Iwamoto & Stein 1974; Wilson & Waples 1983; Iwamoto

FIGURE 97. Fresh specimens of *Coryphaenoides armatus*. (A) BSKU 96478, 37.1 mm HL, 202+ mm TL, off Kinkazan, Miyagi Pref., Pacific, 2698–2814 m depth; (B–C) BSKU 96459, 48.8 mm HL, 242+ mm TL, off Miyako, Iwate Pref., Pacific, 3150–3201 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: H. Endo and N. Nakayama]

1990; Endo & Okamura 1992; Mecklenburg *et al.* 2002; this study). In Japan, known only from off the Pacific coasts of Honshu, Hokkaido, and Shikoku at depths of 2698–7433 m (Appendix 3-5F). Moderately common in its bathymetric range.

FIGURE 98. Scatter diagrams showing relationships of the (A) interorbital width and (B) interdorsal length (as % HL) to HL (mm) in *Coryphaenoides armatus*.

Remarks. *Coryphaenoides armatus* has been well described by previous authors; for a full description see Nybelin (1957; as *Nematonurus armatus*), Iwamoto & Stein (1974), and Iwamoto & Sazonov (1988). This species has a cosmopolitan distribution, and was first recorded from Japan by Endo & Okamura (1992). Based on morphological and electrophoretic analyses, Wilson & Waples (1984) recognized two subspecies of *C. armatus*, namely, *C. a. variabilis* (Günther, 1878) confined to the North Pacific and *C. a. armatus* widely distributed in the other areas of the world's oceans. According to their study, the North Pacific subspecies is characterized morphologically by having a narrower interorbital width (usually <24% HL in *C. a. variabilis* vs. usually ≥24% in *C. a. armatus*), a shorter interdorsal length (usually ≤60% HL vs. usually >60%), and higher counts of first dorsal-fin rays [II,9–10 vs. II,7–10 (usually II,8–9)] and pelvic-fin rays [20–23 total of both sides (mode 22) vs. 17–21 (20)]. The Japanese specimens examined agree well with *C.*

a. variabilis in the meristic characters, with first dorsal-fin rays modally II,9 (range II,9–10) and total pelvic-fin rays usually 22 (19–24). However, a great variation exists in both the interorbital and interdorsal spaces, and their values (relative to HL) become greater with growth (Fig. 98).

Iwamoto & Sazonov (1998) found *C. a. variabilis* from northeastern Pacific to have much darker color than in *C. a. armatus* from the southeastern Pacific. They also noted the northeastern Pacific *C. a. variabilis* as having less-developed serrations along the first dorsal fin. The Japanese specimens examined are uniformly dark in color (Fig. 97), and their dorsal-fin serrations are only weakly developed except in smaller specimens. These features are consistent with Iwamoto & Sazonov's (1998) description of *C. a. variabilis*.

Many authors believed *C. armatus* to be a deep-slope/upper-rise species, with its vertical distribution generally being restricted to about 2000–4300 m (Iwamoto & Stein 1974; Wilson & Waples 1983, 1984; Endo & Okamura 1992; King & Priede 2008; Jamieson *et al.* 2012). However, the specimens examined here includes seven specimens collected from two hadal stations: NSMT-P 30199 (1 spec., 91.6 mm HL, 515+ mm TL) from the east of Tori-shima Island (30°10.12'N, 146°49.14'E) by a fish trap deployed at a depth of 6273 m; and NSMT-P 97647–97649, 97651, 97669 and 98008 (6 spec., 89.4–158 mm HL, 515–902+ mm TL) from the Japan Trench off Miyako (38.7983°N, 144.1345°E) at a depth of 7340–7433 m. These specimens confirm the occurrence of *C. armatus* in the hadal zone in the northwestern Pacific, representing the deepest record for the species. *Coryphaenoides armatus* was considered to be segregated bathymetrically from its sister species, *C. yaquinae* Iwamoto & Stein, 1974 (ca. 2000–4300 m in *C. armatus* vs. ca. 3400–6500 m in *C. yaquinae*; Wilson & Waples 1983; Endo & Okamura 1992; Jamieson *et al.* 2012). However, the above finding suggests that *C. armatus* occurs, albeit rarely, in much deeper waters than previously thought and where *C. yaquinae* is predominantly distributed.

Relationships and comparisons. *Coryphaenoides armatus* belongs to the subgenus *Nematonurus* (sensu Iwamoto 1990), and is most similar to *C. yaquinae*. The two species are readily distinguished by differences in dentition. In *C. armatus*, the mandibular teeth are arranged in one distinct row near the symphysis, whereas those of *C. yaquinae* are in two irregular series. It further differs from *C. yaquinae* in that inner premaxillary teeth, if present, are arranged in a single row (vs. 2–3 irregular rows or narrow band). Also, in large specimens of *C. armatus*, outer teeth are compressed distally, whereas those of *C. yaquinae* are conical with a slender tip (Fig. 99A vs. B). In the northwestern Pacific off Japan, *C. armatus* is distinguished from *C. yaquinae* by the presence of scaly patches on the lower half of the suborbital area and the posterior portion of the mandibular rami. Spinulation of body scales is much

FIGURE 99. Dentition of two species of *Coryphaenoides*. (A) *C. armatus*, BSKU 37299, 107 mm HL; (B) *C. yaquinae*, BSKU 35737, 88.7 mm HL. Anterior views of the mouth. Preserved condition. [Photos: N. Nakayama]

FIGURE 100. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of two species of *Coryphaenoides*. (A–B) *C. armatus*, BSKU 37299, 107 mm HL; (C–D) *C. yaquinae*, BSKU 35734, 107 mm HL. (A, C) Views from above; (B, D) oblique views. [Photos: NSMT]

coarser in *C. armatus* than in similar-sized specimens of *C. yaquinae* (Fig. 100A–B vs. C–D), although a considerable variation exists within each species.

***Coryphaenoides asper* Günther, 1877**

[Japanese name: Zara-dara]

(Figs. 101–102; Appendix 3-6A)

Coryphaenoides asper Günther, 1877:440 [original description; lectotype (designated herein): BMNH 1887.12.7.88, from southeast of Boso Peninsula, *Challenger* sta. 237, in 1875 ftm (3429 m); in part (see Remarks)]; Okada & Matsubara 1938:449 (in key; Japan); Matsubara 1955:1309 (in key; Japan); Okamura 1970a:136 [brief description after Günther (1877, 1887)]; Tominaga & Uyeno 1981:488 (listed; Japan); Okamura 1984b:95, pl. 345, fig. I (compiled); Okamura 1988:95, pl. 345, fig. I (compiled); Nakabo 1993:364 (in key; Japan); Nakabo 2000:428 (in key; Japan); Nakabo 2002:428 (in key; Japan); Nakabo & Kai 2013:504 (in key; Japan); Motomura 2020:39 (listed; Japan).

Macrurus asper: Günther 1887:137, pl. XXXVI, fig. A [brief description; 1 spec. from “south of Japan” (= off Boso Peninsula), *Challenger* sta. 237].

Macrourus asper: Jordan & Snyder 1901:120 (listed; Japan); Jordan & Gilbert in Jordan & Starks 1904:617 [after Günther (1877, 1887)]; Jordan *et al.* 1913:417 (listed; Japan; new Japanese name: “Zara-dara”).

Coryphaenoides (Coryphaenoides) asper: Okamura 1970b: table 1 (listed; Japan).

Diagnosis. Pelvic-fin rays 11–12. Snout bluntly pointed, only slightly protruding beyond upper jaw. Lateral angles of snout armed with small, stout, modified scales. Scales on anterior half of infraorbital ridge slightly thickened. Dorsal contour of head smoothly curved from snout tip to first dorsal-fin origin. Orbit diameter 23% HL. Mouth large, posterior margin of upper jaw extending to below midorbit; upper-jaw length 35% HL; lateral corner of mouth not restricted by skin folds. Outermost gill slit greatly restricted, length 10% HL. Barbel well developed, slender, length 12% HL. Teeth in tapered bands in both jaws, with outer premaxillary series enlarged. Body scales covered with narrowly divergent rows of long, reclined, slender, needle-like spinules; tip of last spinule in each row extending well beyond posterior scale margin; middle row distinctly higher than those of adjacent rows. Transverse scale rows below first dorsal-fin midbase 5. Underside of snout anterior to lateral angles broadly naked; top of snout narrowly naked posterior to leading edges. Height of first dorsal fin 109% HL; second spinous ray serrated along its leading edge; first dorsal-fin rays II,11. Pectoral-fin rays i24. Outermost pelvic-fin ray moderately prolonged, its tip extending well beyond anal-fin origin. Body uniformly dark.

Material examined. 1 specimen. **Lectotype of**

Coryphaenoides asper (designated herein): BMNH 1887.12.7.88 (65.7 mm HL, 316+ mm TL), southeast of Boso Peninsula, Chiba Pref., Japan, 34.6167°N, 140.5333°E, 1875 ftm (3429 m), *Challenger* sta. 237, trawl, 17 Jun. 1875.

Counts and measurements. Counts: first dorsal-fin rays II,11; pectoral-fin rays i24; pelvic-fin rays 11–12; gill rakers on first arch (outer/inner) 5–6/10–11, on second arch 8–9/9–11; longitudinal scales 36; transverse scale rows below first dorsal-fin origin 7–8, below first dorsal-fin midbase 5, below second dorsal-fin origin 6–7.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 28 (36); orbit diameter 23 (31); postorbital length 54 (71); postrostral length 76; orbit–preopercle distance 51 (67); suborbital width 15 (20); upper-jaw length 35 (46); length of rictus 29 (38); length of premaxillary tooth band 24 (31); preoral length 13 (17); interorbital width 28 (37); body width over pectoral-fin bases 54 (71); body depth at first dorsal-fin origin 93 (122); body depth at anal-fin origin 77 (101); isthmus–pelvic distance 50 (65); isthmus–anal distance 56 (74); pelvic-fin length 85 (111); pectoral-fin length 62 (81); predorsal length 112 (147); height of first dorsal fin 109 (143); length of first dorsal-fin base 27 (36); interdorsal length 22 (29); length of gill slit 10 (13); length of posterior nostril 7 (9); barbel length 12 (16).

Redescription of lectotype. General features are shown in Fig. 101. Body deepest at first dorsal-fin origin, gradually tapering to long tail. Trunk short, moderately compressed, width over pectoral-fin bases 1.7 in depth at first dorsal-fin origin. Head large, HL about 4.8 in TL; head bones and flesh moderately firm. Predorsal contour smoothly curved, but slightly elevated anterior to first dorsal-fin origin. Snout short, only slightly protruding beyond upper jaw, length 1.2 times as long as orbit diameter; ventral contour oblique in lateral view; snout low, its tip on horizontal through midorbit. Orbit moderate in size, circular, greatest diameter 2.3 in postorbital length. Interorbital space broad, flat, width 0.8 in orbit diameter. Mouth large, oblique, subinferior, upper-jaw length 0.7 in orbit diameter; posterior margin of maxilla reaching vertical through midorbit; lateral corner of mouth not restricted by lip folds; lips somewhat thin. Suborbital region weakly ridged; upper half almost vertical, lower half moderately inclined mesially. Preopercle large, hind margin slightly inclined from vertical, forming broad lobe posteroventrally; preopercular ridge low, stout, with distinct backward extension at angle. Interopercle mostly hidden behind preopercle. Subopercle and opercle smoothly adjoined, forming posterior margin of gill cover. Gill membranes narrowly connected across, and attached mesially to isthmus, with posterior free fold; gill opening moderately wide, extending forward to below posterior margin of lower jaw. Outermost gill slit greatly restricted by skin folds, length 2.3 in orbit diameter. Lower gill rakers developed into small, low tubercles; upper rakes somewhat

FIGURE 101. Lectotype of *Coryphaenoides asper*. BMNH 1887.12.7.88, 65.7 mm HL, 316+ mm TL, southeast of Boso Peninsula, Chiba Pref., Pacific, 3429 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

FIGURE 102. Scanning electron micrographs showing a body scale (from the dorsum below the interdorsal space) of *Coryphaenoides asper*. BMNH 1887.12.7.88, lectotype, 65.7 mm HL. (A) View from above; (B) oblique view. [Photos: N. Nakayama]

pad-like; outer rakers of first arch much smaller than others; all rakers armed with short, slender spines. Barbel well developed, slender, length 1.9 in orbit diameter.

Anus immediately anterior to anal-fin origin; periproct poorly developed. Ventral light organ absent.

Teeth short, slender, conical, in long tapered band in both jaws. Premaxillary teeth with about 7 tooth rows at widest point near symphysis, with about 4 tooth rows posteriorly; outermost series enlarged, incurved; posterior margin of tooth band almost reaching lateral corner of mouth. Mandibular band much narrower than premaxillary band, but teeth much larger than those in premaxillary, with about 5 tooth rows anteriorly, about 2 rows posteriorly;

posterior margin of tooth band falling short of lateral corner of mouth; no teeth significantly enlarged.

Body scales small, thin, deciduous, covered with narrowly divergent rows of long, slender, reclined, needle-like spinules (Fig. 102); those on dorsum below interdorsal space with 3–6 spinule rows; middle row with about 4–5 spinules, distinctly higher than other rows; height of spinules increasing posteriorly, with last spinules extending well beyond posterior scale margin; spinules of each row greatly overlapping, but free from one another; lateral buttresses barely developed; no reticulate structure on scale surfaces. Body completely and uniformly scaled except for fins.

Scales on anterior half of infraorbital ridge slightly thickened. Lateral angles of snout armed with small, but stout, modified scales; presence/absence of terminal snout scute unknown due to poor condition of specimen. Head scales small except for those on opercle and postorbital canal, similar to those on body, but spinules more erect, giving spiny appearance to head surfaces. Head mostly covered with scales. Underside of snout broadly naked; naked area extending posteriorly to below lateral angles of snout. Dorsal surface of head fully scaled, but top of snout narrowly naked posterior to its leading edges. Other naked areas confined to anterior 1/3 of mandibular rami, gular and branchiostegal membranes, and nasal fossa.

No open pores on cephalic sensory canals. Naked area on underside of snout heavily covered with small free neuromasts and short hair-like papillae. Similar neuromasts scattered on snout, interorbital space, suborbital area, mandibular rami, preopercle, and postorbital canal. Lateral line not interrupted throughout.

Origin of first dorsal fin slightly posterior to pelvic-fin base; first dorsal fin well developed, height 4.0 times as long as its base length; second spinous ray moderately long, but not especially prolonged, armed along its leading edge with 23 short, sharp, slender denticles; denticles becoming much slender and erect distally; tip of second spinous ray extending to base of 15th second dorsal-fin ray when laid back. Second dorsal fin moderately separated from first dorsal, interdorsal space 0.8 times as long as first dorsal-fin base length. Pectoral fin moderately short, its tip extending to vertical through anal-fin origin; pectoral-fin base slightly anterior to pelvic-fin base. Outermost pelvic-fin ray moderately elongate, its tip reaching base of 8th anal-fin ray when laid back. Anal fin well developed, originating slightly anterior to second dorsal-fin origin.

Color when fresh. Unknown.

Color in alcohol (Fig. 101). Head and body dusky brown overall; abdomen and gill cover much darker than surroundings; first dorsal fin mostly blackish (about distal 2/3), but paler basally; second dorsal fin probably blackish originally; pectoral fin uniformly dark; pelvic fin generally pale, but distal margin darker; free portion of outer pelvic-fin ray blackish; basal half of anal fin pale, distal half blackish; oral cavity grayish brown; gill cavity black; gill rakers, arches, and filaments pale brown; barbel and gular membrane pale, but branchiostegal membranes blackish posteriorly; anterior margin of posterior nostril narrowly edged with black; periproct intensely black.

Size. To about 32 cm TL.

Distribution. So far known only from the lectotype collected from off the Boso Peninsula at a depth of 3429 m (Appendix 3-6A). Very rare.

Comments on type specimens. According to the original description, Günther (1877) described *Coryphaenoides asper* based on specimens collected from the “south of the Philippines and Japan”. However, in the “*Challenger*” Report, Günther (1887) listed only

a single specimen for this species (BMNH 1887.12.7.88, 65.7 mm HL, 316+ mm TL; Fig. 101). Although it is uncertain how many specimens were involved in the original description, it is most likely that the material from the Philippines (originally reported as *C. asper*) was re-identified as other species in the “*Challenger*” Report. Among the specimens examined by Günther (1887) was only a single specimen of macrourid grenadiers collected from the Philippines (*Challenger* sta. 214; see also Günther 1887:143, pl. XXVI, fig. B), viz., *Trachonurus* sp., BMNH 1887.12.7.106, 45 mm HL, 246 mm TL (see the Comments on type specimen of *T. villosus*). Species of *Trachonurus* are more or less similar to *C. asper* in general appearance, and thus the above *Trachonurus* specimen could have been misidentified as *C. asper* in the original description of the latter species. In order to avoid further confusion, BMNH 1887.12.7.88 is herein designated as the lectotype of *C. asper*. Consequently, other specimens (if any exist) used in the original description automatically become paralectotypes of the species.

Remarks. Jordan & Thompson (1914) reported *C. asper* based on two specimens collected from Misaki (Sagami Bay), Japan. Although their specimens were not found among the ichthyological collection of the Field Museum of Natural History (FMNH) during the author’s visit in 2012, their figure (pl. 38, fig. 2) depicts a typical form of *Ventrifossa misakia* (Jordan & Gilbert in Jordan & Starks, 1904). Kuroda (1965) also reported *C. asper* from Suruga Bay based on three specimens collected by local fishers. Because he referred to a description and figure of “*C. asper*” given by Jordan & Thompson (1914), his materials are likely a misidentification of *V. misakia*.

The lectotype was the only representative of this rare species until Shao *et al.* (2008a, 2008b) reported the second specimen from the northern part of the South China Sea off Taiwan. Iwamoto *et al.* (2009) provided a description of this additional specimen, noting a difference in body-scale spinulation between the lectotype and their material (spinules in narrowly divergent rows in the lectotype vs. subparallel to slightly convergent rows in the Taiwanese specimen). They also noted that “we cannot evaluate the significance of this difference without additional specimens for comparisons”. Subsequently, Iwamoto *et al.* (2015) re-identified the Taiwanese specimen as *Coryphaenoides* sp. cf. *asper*.

As suggested by previous authors, the additional specimen from Taiwan (ASIZP 66107, 92.3 mm HL) differs notably from the lectotype in the morphology of the body scales (see Iwamoto *et al.* 2009: fig. 3C). In the Taiwanese specimen, the spinules on the scales are arranged in about 15 convergent rows, with the last spinule in each row barely extending beyond the posterior scale margin, whereas those in the lectotype are in five divergent rows and the last spinules extend well beyond the scale margin. The Taiwanese specimen also has much shorter spinules on the scales, which give a smooth

texture to the body (vs. spinules distinctly longer in the lectotype, giving a spiny appearance to body surfaces). These differences are notable even considering the Taiwanese specimen is much larger than the lectotype (65.7 mm HL). It further differs from the lectotype in having lower counts of pectoral- (i20–i21 vs. i24) and pelvic-fin rays (10 vs. 11–12) and inner gill rakers on the first arch (7 vs. 10–11). In addition, the head is completely scaled in the Taiwanese specimen, but the lectotype has a narrow naked area on the underside of the snout. These differences further support their specific separation, and the Taiwanese specimen will be officially described by the author in a separate publication as a representative of a new species.

Consequently, *C. asper* is currently represented by only the lectotype from Japan. Because Günther's (1877, 1887) description of the species is brief, the specimen is redescribed here for future studies.

Relationships and comparisons. *Coryphaenoides asper* belongs to the subgenus *Coryphaenoides* (sensu Iwamoto 1990). This species is most similar to *C. soyoe* Nakayama & Endo, 2016, with which it shares most of the diagnostic characters distinguishing the latter species from its congeners. However, the two species are readily distinguished from each other by scale spinulation: the spinules of *C. asper* are much longer as compared with *C. soyoe*, with the middle row enlarged, giving a spiny appearance to the head and body (Fig. 102), whereas those of *C. soyoe* are much shorter, so as to give a smooth texture to the head and body surfaces (Fig. 133). In addition, the predorsal profile of *C. asper* is smoothly curved from the snout tip to the first dorsal-fin origin (Fig. 101A), whereas in *C. soyoe*, there is a pronounced hump over the nape, with a distinct depression above the orbits (Figs. 131, 132A). Other differences include the number of pectoral-fin rays (i24 in *C. asper* vs. i19–i22 in *C. soyoe*), orbit diameter (23% HL vs. 20–21%), orbit–preopercle distance (51% HL vs. 49%), height of first dorsal fin (109% HL vs. 87%), pectoral-fin length (62% HL vs. 49%), and length of outer gill slit (10% HL vs. 7–8%), although Nakayama & Endo (2016a) noted the necessity to confirm the meristic and morphometric differences when more specimens become available. *Coryphaenoides asper* superficially resembles *C. asprellus* (Smith & Radcliffe in Radcliffe, 1912), *C. hextii* (Alcock, 1890), and *C. woodmasoni* (Alcock, 1899), all known from the Indian Ocean (*C. asprellus* is also distributed in the East Indies). However, it chiefly differs from these species in having higher counts of pelvic-fin rays [11–12 vs. 7–10; data for the latter three species are from Shcherbachev & Iwamoto (1995)].

Coryphaenoides cinereus (Gilbert, 1896)

[Japanese name: Karafuto-sokodara]

(Figs. 103–106; Table 7; Appendix 3–6B)

Macrurus cinereus Gilbert, 1896:457 [original description; numerous syntypes; type locality: North Pacific off Unalaska and Ookamok Islands, *Albatross* sta. 3307, 3329 and 3340, in 399–1033 ftm (730–1889 m)].

Coryphaenoides cinereus: Gilbert & Hubbs 1916:167 (new combination; first record from Japan; brief description; 2 spec. from Okhotsk Sea); Taranetz 1937:169 (listed; Soviet Far East); Okada & Matsubara 1938:449 (in key; Japan; new Japanese name: “Karafuto-sokodara”); Matsubara 1955:1309 (in key; Japan); Lindberg 1959 (listed; southern Okhotsk Sea); Ueno 1965:19 (listed; Hokkaido); Okamura 1970a:129, pl. XXVIII, text-fig. 52 (description; biological notes; 10 spec. from Pacific off Tohoku and Choshi); Ueno 1971:101 (listed; Hokkaido); Tominaga & Uyeno 1981:489 (listed; Japan); Hikita 1981:78, fig. 1 (spec. from Okhotsk Sea off Shiretoko Peninsula); Shiogaki 1982:33 (listed; Aomori Pref.); Sawada 1983:111, 194, 249, figs. 62, 139 (brief description; 8 spec. from Pacific off Tohoku and Okhotsk Sea); Ohta 1983: tables A–B (listed with a question mark; Suruga Bay; in situ observation); Amaoka & Nakaya 1983:45, fig. 61 (compiled; northern Japan); Okamura 1984b:95, pl. 82, fig. D (compiled); Okamura 1988:95, pl. 82, fig. D (compiled); Iwamoto 1990:207, fig. 482 (synopsis); Maeda & Maruyama 1991:366 (listed; Hokkaido); Endo *et al.* 1993:222, fig. 3 (larval description; 9 spec. from North Pacific); Nakabo 1993:364 (in key); Amaoka *et al.* 1995:102, fig. 152 (compiled; northern Japan); Miya *et al.* 1995: table 2 (1 spec. from off Boso Peninsula; midwater capture); Shinohara *et al.* 1996:169 (2 spec. listed from Pacific off Tohoku); Okamura 1997:126, fig. 1 (compiled); Shinohara & Matsuura 1997:291 (listed; Suruga Bay); Goto 2000:6 (1 spec. from Iwate Pref.); Nakabo 2000:428 (in key; Japan); Nakabo 2002:428 (in key; Japan); Maeda & Tsutsui 2003:488 (listed; Hokkaido); Shiogaki *et al.* 2004:55 (listed; Pacific off Aomori Pref.); Senou 2008:368, fig. 31.37 (in situ observation; east of Hokkaido); Kitagawa *et al.* 2008:40, unnumbered fig. (brief description; spec. from Pacific off Tohoku); Shinohara *et al.* 2009:708 (1 spec. listed from Pacific off Tohoku); Amaoka *et al.* 2011:130, fig. 164 (compiled; Hokkaido); Nakabo & Kai 2013:504 (in key; Japan); Shinohara *et al.* 2020:8 (description; 1 spec. from Kurile-Kamchatka Trench); Amaoka *et al.* 2020:163, fig. 220 (listed; Hokkaido); Motomura 2020:39 (listed; Japan).

Coryphaenoides filamentosus Okamura, 1970a:133, pl. XXIX, text-fig. 53 (original description; holotype: BSKU 14208, from “Tohoku District”; 1 paratype collected with holotype; new Japanese name: “Hokuyo-sokodara”); Tominaga & Uyeno 1981:489 (listed; Japan); Okamura 1984b:95, pl. 345, fig. H (compiled); Okamura 1988:95, pl. 345, fig. H (compiled); Nakabo 1993:364 (in key); Nakabo 2000:428 (in key; Japan); Nakabo 2002:428 (in key; Japan); Shinohara *et al.* 2009:708 (listed; Pacific off Tohoku); Nakabo & Kai 2013:504 (in key; Japan); Motomura 2020:39 (listed; Japan).

Coryphaenoides (Coryphaenoides) cinereus: Okamura 1970b: table 1 (listed; Japan).

Coryphaenoides (Coryphaenoides) filamentosus: Okamura 1970b: table 1 (listed; Japan).

Diagnosis. Pelvic-fin rays usually 9 (rarely 8 or 10–11). Snout short, slightly protruding beyond upper jaw, tipped with prominent enlarged tubercles; lateral angles of snout armed with small scute-like scales (highly deciduous). Snout width 24–30% HL; internasal width 21–27% HL. No modified scales along head ridges. Mouth large, posterior margin of upper jaw extending to below posterior 1/3 of orbit of beyond (but not reaching hind rim or orbit); upper-jaw length 34–40% HL; lateral corner of mouth not restricted by skin folds. Outermost gill slit moderately wide, length 18–26% HL. Barbel short, length 3–9% HL. Teeth small, conical, in narrow bands in both jaws, none especially enlarged. Body scales covered with short, reclined, needle-like spinules in parallel to subparallel rows; tip of last spinule in each row extending slightly beyond posterior scale margin. Transverse scale rows below first dorsal-fin midbase 6.5–10.5. Dorsal surface of snout narrowly naked posterior to leading edges; underside of head broadly naked from snout tip to vertical under posterior nostril. Height of first dorsal fin 79–106% HL; second spinous ray slightly elongate, finely serrated along its leading edge; first dorsal-fin rays II, 10–13. Outer pelvic-fin ray greatly prolonged, its tip extending well beyond anal-fin origin. Body pale to dusky.

Material examined. 37 specimens. **Syntypes of *Macrurus cinereus*:** CAS-SU 3084 (3, 40.9–69.4 mm HL, 230+–382+ mm TL), south of Chirikof Island, Alaska, North Pacific, 55.4333°N, 155.4333°W, 695 ftn (1272 m), *Albatross* sta. 3340, 29 Aug. 1890. **Holotype of *Coryphaenoides filamentosus*:** BSKU 14208 (67.9 mm HL, 426+ mm TL), off east coast of Japan, no further information. **Paratype of *C. filamentosus*:** BSKU 14209 (61.5 mm HL, 350+ mm TL), collected with holotype. **Non-types: Japan:** BSKU 86816 (1, 81.1 mm HL, 466+ mm TL), southwest of Okino-shima Island, 32.4033°N, 132.2671°E, 1501–1516 m, R/V *Tansei-maru*, cr. KT-99-18, sta. BT-4, coll. H. Endo and S. Nagatomo, 16 Dec. 1999; BSKU 86850 (1, 90.8 mm HL, 555+ mm TL), Kumano-nada, 33.7332°N, 136.5788°E, 2035–2051 m, R/V *Tansei-maru*, cr. KT-99-18, sta. BT-7, beam trawl, coll. H. Endo and S. Nagatomo, 20 Dec. 1999; BSKU 97789 (1, 89.5 mm HL, 502+ mm TL), off Kuji, 40.3727°N, 143.3827°E, 1923–1995 m, R/V *Tansei-maru*, cr. KT-93-15, sta. M5, 3-m beam trawl, coll. Y. Machida, 18 Nov. 1993; BSKU 97757 (1, 81.8 mm HL, 471+ mm TL), BSKU 97763 (1, 79.1 mm HL, 480+ mm TL), BSKU 97765 (1, 88.2 mm HL, 502+ mm TL), south of Cape Erimomisaki, 41.0675°N, 143.3268°E, 2022–2027 m, R/V *Tansei-maru*, cr. KT-08-27, sta. S-2, 3-m ORE beam trawl, coll. H. Endo and N. Nakayama, 20 Oct. 2008; BSKU 35706 (1, 73.8 mm HL, 418+ mm TL), BSKU

35708 (1, 81.1 mm HL, 462 mm TL), BSKU 35710 (1, 85.3 mm HL, 472+ mm TL), off Kesennuma, 38.7333°N, 143.1850°E, 1950–1960 m, R/V *Hakuho-maru*, cr. KH-81-4, sta. 8, 25 Jul. 1981; BSKU 96497 (1, 92.5 mm HL, 543+ mm TL), off Kinkazan, 38.4640°N, 143.0518°E, 1853–1906 m, R/V *Tansei-maru*, cr. KT-08-27, sta. K-2, 3-m ORE beam trawl, coll. H. Endo and N. Nakayama, 23 Oct. 2008; *BSKU 96471 (1, ca. 31 mm HL, 140+ mm TL), off Kinkazan, 38.4647°N, 143.5283°E, 3137–3223 m, R/V *Tansei-maru*, cr. KT-08-27, sta. K-4, 3-m ORE beam trawl, coll. H. Endo and N. Nakayama, 23 Oct. 2008; BSKU 97748 (1, 31.9 mm HL, 180+ mm TL), off Kuji, 40.0613°N, 142.6825°E, 1005–1016 m, R/V *Tansei-maru*, cr. KT-93-15, sta. M3, 3-m beam trawl, coll. Y. Machida, 18 Nov. 1993; BSKU 97852 (1, 49.2 mm HL, 291+ mm TL), BSKU 97861 (1, 36.8 mm HL, 164+ mm TL), off Rikuzen-Takata, 39.0538°N, 142.6342°E, 1195–1202 m, R/V *Tansei-maru*, cr. KT-89-3, sta. SR91, 3-m beam trawl, 10 Mar. 1989; BSKU 97833 (1, 36.5 mm HL, 214+ mm TL), off Kamaishi, 39.1444°N, 142.3748°E, 847–935 m, R/V *Tansei-maru*, cr. KT-89-3, sta. SR89, 3-m beam trawl, 10 Mar. 1989; BSKU 58779 (1, 82.7 mm HL, 492+ mm TL), off Kuril Islands, 43.0630°N, 147.1152°E, 451–661 m, FRV *Wakataka-maru*, sta. G06, coll. D. Kitagawa, 21 Jun. 1996; BSKU 97733 (1, 63.6 mm HL, 362+ mm TL), BSKU 97771 (1, 59.5 mm HL, 328+ mm TL), BSKU 97773 (1, 48.2 mm HL, 290+ mm TL), BSKU 97774 (1, 58.1 mm HL, 338+ mm TL), BSKU 97776 (1, 57.8 mm HL, 344 mm TL), BSKU 97788 (1, 51.8 mm HL, 290+ mm TL), south-southeast of Akkeshi, 42.6052°N, 144.9938°E, 1113–1161 m, R/V *Tansei-maru*, cr. KT-93-15, sta. T8, 3-m ORE beam trawl, coll. Y. Machida, 16 Nov. 1993; BSKU 96392 (1, 75.7 mm HL, 440+ mm TL), BSKU 96401 (1, 89.3 mm HL, 487+ mm TL), BSKU 96405 (1, 74.0 mm HL, 406+ mm TL), south of Cape Erimomisaki, 41.6675°N, 143.3268°E, 2022–2027 m, R/V *Tansei-maru*, cr. KT-08-27, sta. S-2, 3-m ORE beam trawl, coll. H. Endo and N. Nakayama, 20 Oct. 2008; BSKU 97780 (1, 65.4 mm HL, 391+ mm TL), BSKU 97792 (1, 60.1 mm HL, 348+ mm TL), Pacific off northern Japan, R/V *Tansei-maru*, cr. KT-93-15, 3-m beam trawl, coll. Y. Machida, date unknown; *BSKU 49888 (1, 56.4 mm HL, 296+ mm TL), Japan, 1500 m, R/V *Hakuho-maru*, cr. KH-91-6, sta. 12, 16 Nov. 1991; NSMT-P 90682 (4, 64.6–74.1 mm HL, 276+–437+ mm TL), Kitami-Yamato Bank, Okhotsk Sea, coll. H. Matsubara, 2 Jun. 2008.

Counts and measurements. See Table 7.

Size. Attains about 57 cm TL (HUMZ 88325, 565 mm TL, Bering Sea).

Sexual dimorphism. The pelvic-fin length is sexually dimorphic, with males having a longer outer ray than those of similar-sized females (Gilbert & Hubbs 1920; Iwamoto & Stein 1974; Iwamoto 1990).

Development. Larval morphology was described by Endo *et al.* (1993).

Distribution. Widely distributed in the North Pacific

FIGURE 103. Fresh specimens of *Coryphaenoides cinereus*. (A) BSKU 96471, ca. 31 mm HL, 140+ mm TL, off Kinkazan, Miyagi Pref., Pacific, 3137–3223 m depth; (B) BSKU 49888, 56.4 mm HL, 296+ mm TL, KH-91-6, sta. 12; (C) BSKU 96392, 75.7 mm HL, 440+ mm TL, off Cape Erimomisaki, Hokkaido, Pacific, 2022–2027 m depth. Lateral views. [Photos: (A, C) H. Endo and N. Nakayama; (B) BSKU]

TABLE 7. Morphometric measurements and meristic counts* for *Coryphaenoides cinereus*.

Nominal species	<i>C. cinereus</i>		<i>C. filamentosus</i>	
Type status	ST	NT	HT	PT
Number of specimens	<i>n</i> = 3	<i>n</i> = 30	<i>n</i> = 1	<i>n</i> = 1
Measurements (mm)				
TL	230+–382+	164+–555+	426+	350+
HL	40.9–69.4	31.9–92.5	67.9	61.5
% of HL (% of PRL)				
Snout length	26–27 (35)	26–29 (34–39)	26 (35)	28 (37)
Orbit diameter	31 (40–41)	25–33 (32–45)	28 (36)	28 (37)
Postorbital length	45–47 (60–62)	43–52 (58–68)	51 (67)	48 (64)
Postrostral length	74–77 (–)	74–78 (–)	77 (–)	75 (–)
Orbit–preopercle distance	44–45 (58–59)	43–49 (56–66)	45 (58)	45 (59)
Suborbital width	13–14 (17–19)	12–15 (15–19)	13 (17)	12 (16)
Upper-jaw length	35–38 (47–49)	34–40 (44–53)	36 (47)	35 (47)
Length of rictus	31–32 (41–43)	29–35 (37–46)	32 (41)	30 (40)
Length of PMX tooth band	27–28 (35–37)	24–30 (30–39)	27 (35)	27 (35)
Preoral length	13–16 (17–21)	14–19 (18–26)	16 (21)	18 (24)
Dist. tip and lat. angle of snout	14–17 (18–22)	14–18 (19–25)	17 (22)	16 (22)
Snout width	26 (34)	24–30 (31–40)	– (–)	26 (34)
Internasal width	22–23 (29–31)	21–27 (27–35)	– (–)	23 (31)
Interorbital width	26–28 (35–36)	25–30 (33–40)	26 (34)	27 (36)
Body width over P. bases	40–49 (54–65)	40–64 (54–83)	61 (80)	54 (72)
Body depth at 1D. origin	73–81 (99–105)	66–87 (88–113)	86 (112)	82 (109)
Body depth at A. origin	52–61 (69–78)	47–69 (62–89)	69 (91)	66 (87)
Pre-V. length	101–107 (133–139)	102–125 (132–162)	115 (150)	108 (144)
Pre-anus length	132–142 (177–184)	137–164 (176–215)	160 (209)	151 (200)
Pre-A. length	138–146 (185–188)	140–169 (182–222)	166 (217)	156 (207)
Isthmus–V. distance	30–32 (40–42)	35–45 (48–59)	38 (50)	38 (50)
Isthmus–anus distance	– (–)	75–98 (99–131)	85 (110)	79 (105)
Isthmus–A. distance	69–74 (93–96)	79–103 (105–136)	92 (120)	85 (113)
V.–A. distance	38–42 (51–54)	40–62 (53–81)	55 (72)	49 (66)
Anus–A. distance	– (–)	3–9 (3–11)	7 (9)	6 (8)
V. length	66–70 (86–90)	68–113 (86–147)	92 (120)	84 (112)
P. length	60 (78)	64–86 (83–112)	71 (92)	69 (92)
Pre-1D. length	109–114 (141–150)	105–116 (135–155)	114 (148)	109 (145)
Height of 1D.	88 (115)	79–106 (106–140)	– (–)	102 (136)
Length of 1D. base	26–31 (36–41)	26–36 (35–47)	35 (46)	31 (41)
Interdorsal length	12–15 (16–21)	12–48 (17–64)	13 (17)	23 (31)
Length of gill slit	18–21 (24–28)	18–26 (24–34)	20 (27)	19 (26)
Length of posterior nostril	8–9 (10–12)	8–11 (10–15)	10 (14)	9 (12)
Barbel length	6 (7–8)	3–9 (4–11)	7 (9)	7 (9)
Counts				
1D. rays	II,10–11	II,10–13	II,11	II,12
P. rays	i19–i23	i17–i23	i20	i18–i19
V. rays	9–11	8–10	9	9
Outer GR on first arch	9–11	6–13	10	9
Inner GR on first arch	14–15	13–16	14	15
Outer GR on second arch	13–14	12–16	14	13
Inner GR on second arch	13–14	13–16	14	13
Longitudinal scales	43	31–45	38	37
Scales below 1D. origin	10–12	10–14	12	11
Scales below 1D. midbase	9.5–10.5	6.5–10	8	7
Scales below 2D. origin	8	7–12	10	9
Scales above A. origin	–	23–39	29	–

...See the next page for the table footer

*Abbreviations: 1D.—first dorsal fin; 2D.—second dorsal fin; A.—anal fin; GR—gill rakers; HT—holotype; NT—non-type; P.—pectoral fin; PMX—premaxillary; PT—paratype; ST—syntype; V.—pelvic fin

FIGURE 104. Syntype of *Macrurus cinereus* (= *Coryphaenoides cinereus*). CAS-SU 3084 (1 of 3 spec.), 59.8 mm HL, 365 mm TL, south of Chirikof Island, Alaska, North Pacific, 1272 m depth. (A) Lateral, (B) dorsal, and (C) ventral views. Preserved condition. [Photos: N. Nakayama]

from Japan to the United States, including the Okhotsk and Bering Seas as well as the Emperor Seamounts, at depths of 135–3223 m (Iwamoto & Stein 1974; Borets 1986; Iwamoto 1990; Mecklenburg *et al.* 2002; Orlov & Tokranov 2008; this study). In Japanese waters, known from off the Pacific coasts southward to Miyazaki (31.90°N) and the southern Okhotsk Sea at depths of 350–3223 m (Appendix 3-6B). Abundant in northern Japan, but rare in the west of the Boso Peninsula (139.89°E).

Remarks. *Coryphaenoides cinereus* was originally described by Gilbert (1896) based on “numerous

specimens” collected from the North Pacific off Unalaska and Ookamok Islands (Fig. 104). This species has been well described by previous authors, and thus there is no need for repetition here. For a full description see Okamura (1970a) and Iwamoto & Stein (1974). This species is one of the most abundant continental slope fishes found in the North Pacific (Orlov & Tokranov 2008), with the highest abundance in the Bering Sea, Kamchatka, and the Okhotsk Sea (Iwamoto & Stein 1974). Orlov & Tokranov (2008) provided detailed information on the biology of this species off the Kuril and Kamchatka Islands.

FIGURE 105. Preserved specimen of *Coryphaenoides cinereus*. BSKU 14208, holotype of *C. filamentosus*, 67.9 mm HL, 426+ mm TL, “off east coast of Japan”, Pacific. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

Geographical variations between the Japanese and Bering Sea populations of *C. cinereus* have been noted by several authors (Gilbert & Hubbs 1916, 1920; Okamura 1970a; Iwamoto & Stein 1974; Iwamoto 1990), and some were reconfirmed in this study. The Japanese specimens examined have a slightly longer abdomen than three syntypes collected from Alaska, with a longer preanus length (137–164% HL vs. 132–142%), isthmus–pelvic distance (35–45% HL vs. 30–32%), isthmus–anal distance (79–103% HL vs. 69–74%), and pelvic–anal distance (40–62% HL vs. 38–42%). Paired fins of the Japanese specimens examined are also longer than the syntypes, viz., pelvic-fin length (68–113% HL vs. 66–70%) and pectoral-fin length (64–86% HL vs. 60%). These differences are tentatively considered to be geographical variation within a single species, although more Alaskan specimens should be examined to warrant this conclusion.

Nomenclatural discussion. Okamura (1970a) described *C. filamentosus* based on two specimens collected from off the “east coast of Japan” (Fig. 105). Unfortunately, no further information is available on the type locality. Okamura (1970a) considered *C. filamentosus* to be closely similar to *C. cinereus*, one of the most common grenadiers found in the area. In his key

to species of Japanese *Coryphaenoides*, *C. filamentosus* is distinguished from *C. cinereus* by the following combination of features: head bones firm (vs. soft in *C. cinereus*); body depth greater than postrostral length (vs. about equal to postrostral); spinules on the body scales arranged in divergent rows (vs. parallel rows); interorbital width greater than orbit diameter (vs. less than orbit diameter); upper-jaw length equal to 1/3 HL (vs. distinctly longer than 1/3 HL); orbit diameter 1.8 in postrostral length (vs. 1.4–1.6); barbel length 3.1–3.3 in orbit diameter (vs. 4.5–6.0).

An examination of the types of *C. filamentosus*, along with 33 specimens of *C. cinereus* including three syntypes, revealed no significant differences between the two species. They could not be distinguished from each other by meristic and morphometric characters, including those Okamura (1970a) considered as diagnostic characters for separating the two species, viz., body depth, interorbital width, upper-jaw length, orbit diameter, and barbel length (Table 7). The head bones of *C. filamentosus* are not firm (distinctly soft), but this is probably due to lengthy preservation in formalin. Furthermore, in the additional specimens of *C. cinereus* examined, the spinules are often arranged in weakly divergent rows (Fig. 106E); the condition is almost consistent with Okamura’s (1970a)

FIGURE 106. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of *Coryphaenoides cinereus*. (A–B) BSKU 14209, paratype of *C. filamentosus*, 61.5 mm HL; (C–D) NSMT-P 90682, 72.2 mm HL; (E–F) BSKU 97789, 89.5 mm HL. (A, C, E) Views from above; (B, D, F) oblique views. [Photos: NSMT]

illustration for *C. filamentosus* (text-fig. 53). The close similarity of the two species is amply sufficient to warrant their conspecificity, and *C. filamentosus* is therefore regarded as a junior synonym of *C. cinereus*.

Relationships and comparisons. *Coryphaenoides*

cinereus belongs to the subgenus *Coryphaenoides* (sensu Iwamoto 1990), and is most similar to *C. filifer* (Gilbert, 1896) widely distributed in the North Pacific. It differs from *C. filifer* in having a much broader snout (snout width 24–30% HL vs. 19–22%; internasal width 21–27% HL vs.

16–18%), a longer preoral (13–19% vs. 11–13%), and a shorter chin barbel (3–9% HL vs. 8–11%) In *C. cinereus*, the body scales are covered with short, reclined, needle-like spinules in parallel to subparallel rows (Fig. 106A, C, E), whereas in *C. filifer*, the spinules are arranged in narrowly divergent rows (Fig. 109). Furthermore, they are distinguished from each other in the presence/absence of open pores along the cephalic sensory canals: the canals are completely closed in *C. cinereus*, whereas in *C. filifer* small and inconspicuous pores are present at least along the supraorbital, infraorbital, and preopercular canals.

***Coryphaenoides filifer* (Gilbert, 1896)**

[Japanese name: Kitano-sokodara]

(Figs. 92B, 107–109; Appendix 3-6C)

Chalinura filifera Gilbert, 1896:458 [original description; 3 syntypes, from off Queen Charlotte Island, *Albatross* sta. 3342, in 1588 ftm (2904 m)].

Macrurus dorsalis Gill & Townsend, 1897:233 [original description; holotype: USNM 48770, from southwest of Pribilof Islands, Bering Sea, *Albatross* sta. 3640, in 1401 ftm (2562 m)].

Macrurus lepturus Gill & Townsend, 1897:233 [original description; holotype: USNM 48770, from southwest of Pribilof Islands, Bering Sea, *Albatross* sta. 3640, in 1401 ftm (2562 m)].

Coryphaenoides filifer: Gilbert & Hubbs 1916:143 (listed); Endo *et al.* 1994:330, fig. 2 (first record from Japan; brief description; 2 spec. from Okhotsk Sea; new Japanese name: “Kitano-sokodara”); Amaoka *et al.* 1995:102, fig. 151 (compiled; northern Japan); Okamura 1997:125, fig. 7 (compiled); Nakabo 2000:426 (in key; Japan); Nakabo 2002:426 (in key; Japan); Maeda & Tsutsui 2003:488 (listed; Hokkaido); Amaoka *et al.* 2011:129, fig. 163 (compiled; Hokkaido); Nakabo & Kai 2013:502 (in key; Japan); Amaoka *et al.* 2020:162, fig. 219 (listed; Hokkaido).

Macrourus albatrossus Townsend & Nichols, 1925:17, pl. IV, fig. 2 [original description; holotype: USNM 87555 (ex. AMNH 8345), from off northern Baja California, northeastern Pacific, *Albatross* sta. 1692, in 1076 ftm (1968 m)].

Coryphaenoides filiferus: Motomura 2020:39 (listed; Japan).

Diagnosis. Pelvic-fin rays 9–10. Snout short, slightly protruding beyond upper jaw. Tip and lateral angles of snout armed with enlarged tubercles; no scute-like scales along head ridges. Snout width 19–22% HL; internasal width 16–18% HL. Mouth large, posterior margin of upper jaw extending to about vertical through hind rim of orbit; upper-jaw length 38–43% HL; lateral corner of mouth not restricted by skin folds. Outermost gill slit moderately wide, length 19–23% HL. Barbel short, slender, length 8–11% HL. Teeth small, conical, in narrow bands in both jaws; outer premaxillary series slightly enlarged. Body scales covered with short, reclined, needle-like spinules in narrowly divergent rows; tip of last spinule in each row extending distinctly beyond posterior scale margin.

Transverse scale rows below first dorsal-fin midbase 6–11.5. Snout fully scaled dorsally, narrowly naked ventrally. Height of first dorsal fin 91–105% HL; second spinous ray slightly elongate, weakly serrated along its leading edge; first dorsal-fin rays II, 12–15. Outer pelvic-fin ray greatly prolonged, its tip extending well beyond anal-fin origin. Body dark overall.

Material examined. 15 specimens. **Syntype of *Chalinura filifera*:** *USNM48626 (1, badly disintegrated), off Queen Charlotte Island, British Columbia, Canada, northeastern Pacific, 52.6608°N, 132.6333°W, 1588 ftm (2905 m), *Albatross* sta. 3342, large beam trawl, 3 Sept. 1890. **Non-types: Japan:** HUMZ 126365 (1, 92.5 mm HL, 506+ mm TL), northwest of Shiretoko Peninsula, Okhotsk Sea, 44.5450°N, 145.0867°E, 1348 m, bottom trawl, coll. D. Tsutsui, 26 May 1993; HUMZ 126444 (1, 116 mm HL, 626+ mm TL), northwest of Shiretoko Peninsula, Okhotsk Sea, 44.5067°N, 145.0767°E, 1285 m, bottom trawl, coll. D. Tsutsui, 27 May 1993. **Northeastern Pacific:** FMNH 117113 (5, 83.6–104 mm HL, 412+–557+ mm TL), off Newport, 44.7617°N, 125.5214°W, 2800 m, R/V *Wecoma*, cr. W9704A, sta. 9, 40-ft otter trawl, coll. J. Voightt, 19 Apr. 1997; FMNH 117112 (5, 78.8–102 mm HL, 400–511+ mm TL), off Newport, 44.7658°N, 125.5289°W, 2850 m, R/V *Wecoma*, cr. W9704A, sta. 3, 40-ft otter trawl, coll. J. Voightt, 17 Apr. 1997; FMNH 117111 (1, 99.1 mm HL, 487+ mm TL), off Newport, 44.7594°N, 125.5311°W, 2850 m, R/V *Wecoma*, cr. W9704A, sta. 2, 40-ft otter trawl, coll. J. Voightt, 17 Apr. 1997; FMNH 119802 (1, 105 mm HL, 557+ mm TL), off Newport, 44.7594°N, 125.5311°W, 2850 m, R/V *Wecoma*, sta. 2, otter trawl, coll. J. Voightt, 17 Apr. 1997.

Counts and measurements. Based on 14 specimens (78.8–116 mm HL, 400–626+ mm TL). Counts: first dorsal-fin rays II, 12–15; pectoral-fin rays i20–i24; pelvic-fin rays 9–10; gill rakers on first arch (outer/inner) 6–12/12–16, on second arch 12–16/12–15; longitudinal scales 40–50; transverse scale rows below first dorsal-fin origin 8.5–13.5, below first dorsal-fin midbase 6–11.5, below second dorsal-fin origin 8–12, above anal-fin origin 36.5–37.5.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 25–27 (33–37); orbit diameter 24–29 (32–38); postorbital length 48–53 (64–69); postrostral length 74–77; orbit–preopercle distance 42–46 (56–61); suborbital width 10–13 (13–17); upper-jaw length 38–43 (50–57); length of rictus 33–37 (43–48); length of premaxillary tooth band 29–33 (37–43); preoral length 11–13 (15–18); distance between tip and lateral angle of snout 12–16 (16–20); snout width 19–22 (25–29); internasal width 16–18 (21–24); interorbital width 24–28 (32–37); body width over pectoral-fin bases 45–60 (60–78); body depth at first dorsal-fin origin 70–83 (93–111); body depth at anal-fin origin 54–72 (71–97); prepelvic length 98–108 (129–144); preanus length 141–163 (182–218); preanal length 146–

FIGURE 107. Fresh specimen of *Coryphaenoides filifer*. HUMZ 126365, 92.5 mm HL, 506+ mm TL, northwest of Shiretoko Peninsula, Hokkaido, Okhotsk Sea, 1348 m depth. Lateral view. [Photo: HUMZ]

FIGURE 108. Preserved specimen of *Coryphaenoides filifer*. HUMZ 126365, 92.5 mm HL, 506+ mm TL, northwest of Shiretoko Peninsula, Hokkaido, Okhotsk Sea, 1348 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

FIGURE 109. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of *Coryphaenoides filifer*. (A) HUMZ 126365, 92.5 mm HL; (B) HUMZ 126444, 116 mm HL. Views from above. [Photos: N. Nakayama]

169 (189–226); isthmus–pelvic distance 38–47 (51–62); isthmus–anus distance 108 (145); isthmus–anal distance 87–115 (116–154); pelvic–anal distance 46–72 (61–96); anus–anal distance 7 (10); pelvic-fin length 76–103 (99–140); pectoral-fin length 56–70 (74–91); predorsal length 98–116 (133–152); height of first dorsal fin 91–105 (121–139); length of first dorsal-fin base 28–36 (37–48); interdorsal length 11–27 (15–36); length of gill slit 19–23 (25–31); length of posterior nostril 8–10 (10–14); barbel length 8–11 (11–15).

Size. Attains about 66 cm TL (Mecklenburg *et al.* 2002).

Development. Descriptions of larvae and juveniles were given by Stein (1980).

Distribution. Widely distributed in the North Pacific from the Okhotsk Sea off Hokkaido (Appendix 3-6C), north to Alaska, and east to southern California (Iwamoto & Stein 1974; Endo *et al.* 1994; Mecklenburg *et al.* 2002; this study). Depth range 1285–1348 m in Japan, but much greater in other parts of its range (2065–2904 m; Iwamoto & Stein 1974). Very rare in Japanese waters.

Remarks. *Coryphaenoides filifer* was originally described by Gilbert (1896) based on three specimens collected from the northeastern Pacific off Queen Charlotte Island, Canada. One of these specimens deposited at the Smithsonian Institution (USNM 48626) is in poor condition, with the head and body badly deteriorated into many parts (Fig. 92B), which made it impossible to reconfirm the species' diagnostic features. Unfortunately, the other two types were not found in the course of this study. This species was first recorded from Japan by Endo *et al.* (1994) based on two specimens collected from the Kitami-Yamato Bank in the southern margin of the Okhotsk Sea. For further morphological information see Iwamoto & Stein (1974) and Endo *et al.* (1994).

Nomenclatural discussion. The spelling of the specific epithet of this species is open to discussion.

Although most previous authors used “*filifer*” (e.g., Gilbert & Hubbs 1916; Iwamoto & Stein 1974; Iwamoto 1990; Nakabo 1993, 2000, 2002; Eschmeyer 1998; Mecklenburg *et al.* 2002; Nakabo & Kai 2013; Nakabo & Hirashima 2015), some recent authors considered “*filiferus*” to be grammatically correct (Fricke *et al.* 2020; Motomura 2020). As a species of the feminine nominal genus *Chalinura* Goode & Bean, 1883 (= *Fuyangia* Whitley, 1931; see the Relationships of *C. leptolepis*), Gilbert (1896) originally described the species as *Ch. filifera*. This specific epithet is considered a feminine form of “*filifer*”, which is derived from the combination of the Latin noun *filum*, meaning a thread, and the Latin adjectival suffix *-fer* (*-fera*, *-ferum*), denoting carrying (probably in allusion to the species' elongate outer pelvic-fin ray). However, *filifera* can also be regarded as a feminine form of “*filiferus*”, because the suffix *-fer* is occasionally modified to *-ferus* (*-fera*, *-ferum*; Glare 1968:685). Because Gilbert (1896) did not mention the etymology of his *filifera*, the masculine form of this Latin adjective is currently unclear. However, the spelling “*filifer*” is considered to have priority in this case for the following three reasons: 1) Gilbert & Hubbs (1916), who were the first to recognize the species under a masculine genus (*Coryphaenoides*), inflected the spelling of the word as “*filifer*”; 2) the spelling “*filifer*” has been consistently used for the species for more than a century (with *Coryphaenoides*); and 3) the suffix *-ferus* can be regarded as a supplementary modification of *-fer* [in the *Oxford Latin Dictionary* (Glare 1968:685) the explanation of *-ferus* appears within the heading of *-fer*], and there are no grammatical reasons to choose *-ferus*, rather than *-fer*, for the species.

Relationships and comparisons. *Coryphaenoides filifer* belongs to the subgenus *Coryphaenoides* (sensu Iwamoto 1990). It is superficially similar to two Japanese congeners, *C. altipinnis* Günther, 1877 and *C. cinereus* (Gilbert, 1896). For comparisons with the latter two

species see the Relationships and comparisons of *C. altipinnis* and *C. cinereus*.

***Coryphaenoides leptolepis* Günther, 1877**

[Japanese name: Hiragashira-dara]

(Figs. 110–114; Table 8; Appendix 3-6D)

Coryphaenoides leptolepis Günther, 1877:441 (original description; 4 spec., from “off the coasts of Brazil, Japan, and Mid-Pacific”, *Challenger* sta. 122, 237, and 246); Gilbert & Hubbs 1916:143 (listed); Nybelin 1957:268, tables V–VI (synonymy; description; North Atlantic; comparison with “*Co. leptolepis*”); Iwamoto & Stein 1974:43, fig. 23 (synonymy; description); Geistdoerfer 1986:652 (North Atlantic); Geistdoerfer 1990:546 (eastern tropical Atlantic; compiled; synonymy); Iwamoto 1990:187 (in key); McAllister 1990:122 (listed; Canada); Mecklenburg *et al.* 2002:274 (brief description; Alaska; compiled; distribution); Moor *et al.* 2003:209 (New England); Mundy 2005:250 [record from Hess Seamount (not Hawaii), after Wilson & Waples (1983)].

Coryphaenoides variabilis (not Günther 1878): Günther 1878:27 (original description; in part, 1 paralectotype, BMNH 1887.12.7.132, Mid-Pacific, *Challenger* sta. 246).

Chalinura simula Goode & Bean, 1883:199 [original description; 1 spec. from south of Nova Scotia, northwestern Atlantic, *Blake* sta. 308, in 1242 ftn (2271 m); 3 other specimens, juveniles, conditionally referred]; Goode & Bean 1896:412, fig. 345 (description; northwestern Atlantic); Garman 1899:395 (listed); Parr 1946:65, fig. 20 (description; illustration; western Atlantic); Nybelin 1957:264, tables V–VI, pl. III, figs. 3–4 (synonymy; description; North Atlantic; comparison with “*Ch. leptolepis*”).

Macrurus leptolepis: Günther 1887:144, pl. XXXI (new combination; redescription based on lectotype, BMNH 1887.12.7.109, from “off Pernambuco”, *Challenger* sta. 122; other spec. of original type series designated as types for new species, *M. liocephalus*).

Macrurus simulus: Günther 1887:145 [new combination; description after Goode & Bean (1883)]; Koefoed 1927:100 (synonymy; description; comparison with similar congeners).

Macrurus liocephalus Günther, 1887:145, pl. XXXVIII, fig. A. [original description; 3 spec., including BMNH 1887.12.7.111 and BMNH 1887.12.7.112 (also paralectotypes of *Cor. leptolepis*), from “Japan and Mid-Pacific”, *Challenger* sta. 237 and 246, in 1875–2025 ftn (3429–2750 m)].

Macrurus armatus (not Hector 1875): Günther 1887:150 (description; in part, 1 paralectotype of *Cor. variabilis*, BMNH 1887.12.7.132, Mid-Pacific, *Challenger* sta. 246).

Chalinura serrula Bean, 1890:37 [original description; holotype: USNM 199612, from off Prince of Wales Island, *Albatross* sta. 2859, 55°20'N, 136°20'W, in 1569 ftn (2870 m); 2 paratypes collected with holotype]; Garman 1899:395 (listed).

Chalinura leptolepis: Goode & Bean 1896:414 (new combination; synonymy; description); Garman 1899:395 (listed); Marshall 1973:590, fig. 26 (lectotype designation; synonymy;

description; western North Atlantic); Headrich & Merrett 1988:1332, pl. 39 (listed; North Atlantic Basin); Arruda 1997:66 (listed; Azores); Santos *et al.* 1997 (listed; Azores).

Chalinura liocephala: Jordan & Snyder 1901:119 (listed; Japan); Jordan & Gilbert in Jordan & Starks 1904:608 [new combination; after Günther (1877, 1887)]; Jordan *et al.* 1913:415 (listed; Japan; new Japanese name: “Hira-gashira-dara”).

Coryphaenoides liocephalus: Gilbert & Hubbs 1916:143 (new combination; listed); Okada & Matsubara 1938:449 (in key; Japan); Matsubara 1955:1309 (in key; Japan); Okamura 1970a:125 [brief description after Günther (1887)]; Hart 1973:225 (description; illustration; Canada); Tominaga & Uyeno 1981:489 (listed; Japan); Okamura 1984b:95, pl. 345, fig. F (compiled); Okamura 1988:95, pl. 345, fig. F (compiled); Iwamoto 1990:187 (in key); McAllister 1990:122 (listed; Canada); Nakabo 1993:362 (in key; Japan); Nakabo 2000:427 (in key; Japan); Nakabo 2002:427 (in key; Japan); Nakabo & Kai 2013:503 (in key; Japan); Motomura 2020:39 (listed; Japan).

Coryphaenoides simulus: Gilbert & Hubbs 1916:143 (new combination; listed); Fowler 1936:457 (synonymy; description; Congo).

Coryphaenoides serrulus: Gilbert & Hubbs 1916:144 (new combination; listed).

Coryphaenoides (Coryphaenoides) liocephalus: Okamura 1970b: table 1 (listed; Japan).

Coryphaenoides sp.: Endo *et al.* 2000:16, photo D (in situ observation; Nankai Trough).

Diagnosis. Pelvic-fin rays usually 9 or 10, rarely 8. Snout low, blunt, scarcely protruding beyond upper jaw. No stout tubercular scales on snout and head ridges. Uniserial row of small embedded scales along leading edges of snout, behind which a pair of naked lunate areas narrowly developed. Underside of head completely naked. Premaxillary teeth in wide tapered band, with outer series of enlarged canines and inner band of villiform teeth; mandibular teeth in uniserial row. Body scales covered with short, greatly reclined, needle-like spinules in widely spaced parallel rows; tip of last spinule in each row scarcely extending beyond posterior scale margin. Transverse scale rows below first dorsal-fin midbase 5–8. Orbit small, greatest diameter 17–21% HL. Mouth large, posterior margin of upper jaw extending to about vertical through hind rim of orbit or slightly beyond; upper-jaw length 39–46% HL; lateral corner of mouth not restricted by lip folds. Outer gill slit moderately wide, length 22–28% HL. Barbel long, slender, length 16–25% HL. Interdorsal space greater than first dorsal-fin base length. Height of first dorsal fin less than HL (64–75% HL); second spinous ray serrated along its leading edge; first dorsal-fin rays II, 8–10. Origin of second dorsal fin posterior to that of anal fin. Outer pelvic-fin ray greatly prolonged, its tip extending well beyond anal-fin origin. Body generally pale, but posterior portion of tail much darker.

Material examined. 45 specimens. **Lectotype of *Coryphaenoides leptolepis*:** BMNH 1887.12.7.109 (91.6 mm HL, 440+ mm TL), off Pernambuco, Brazil, southwestern Pacific, 9.0833°S, 34.8333°W, 350 ftm (641 m), *Challenger* sta. 122, trawl, 10 Sept. 1873. **Paralectotypes of *Co. leptolepis*:** BMNH 1887.12.7.111 (1, 84.5 mm HL, 406+ mm TL; also lectotype of *Macrurus liocephalus*, designated herein), off Boso Peninsula, Chiba Pref., Japan, 34.6167°N, 140.5333°E, 1875 ftm (3429 m), *Challenger* sta. 237, trawl, 17 Jun. 1875. BMNH 1887.12.7.112 (1, 36.0 mm HL, 198 mm TL; also paralectotype of *M. liocephalus*, designated herein), east of Emperor Seamounts, Mid-Pacific, 36.1667°N, 178.0000°E, 2050 ftm (3750 m), *Challenger* sta. 246, trawl, 2 Jul. 1875. **Holotype of *Chalinura serrula*:** USNM 45358 (57.5 mm HL, 283+ mm TL), off Prince of Wales Island, Canada, Gulf of Alaska, northeastern Pacific, 55.3667°N, 136.3333°W, 1569 ftm (2870 m), *Albatross* sta. 2859, large beam trawl, 28 Aug. 1888. **Paratypes of *Ch. serrula*:** USNM 199612 (2, 34.2–38.3 mm HL, 146+–198+ mm TL), collected with holotype. **Holotype of *Chalinura simula*:** *MCZ 25824 (458 mm TL, fide Goode & Bean 1883:199; examination based on only photographs; Fig. 111F), south of Nova Scotia, off northeastern extremity of George's Bank, northwestern Atlantic, 41.4117°N, 65.5917°W, 1242 ftm (2272 m), R/V *Blake*, sta. 308, 29 Jun. 1880 (examination based on only photographs). **Paralectotype of *Coryphaenoides variabilis*:** BMNH 1887.12.7.132 (1, 36.1 mm HL, 179+ mm TL), east of Emperor Seamounts, Mid-Pacific, 36.1667°N, 178.0000°E, 2050 ftm (3750 m), *Challenger* sta. 246, trawl, 2 Jul. 1875. **Non-types:** **Japan:** BSKU 96940 (1, 63.2 mm HL, 330+ mm TL), BSKU 96941 (1, 71.2 mm HL, 292+ mm TL), BSKU 96942 (1, 66.0 mm HL, 355+ mm TL), BSKU 96943 (1, 71.2 mm HL, 377 mm TL), BSKU 96944 (1, 60.3 mm HL, 313 mm TL), BSKU 96945 (1, 56.9 mm HL, 292 mm TL), BSKU 96946 (1, 56.8 mm HL, 297+ mm TL), southeast of Cape Erimomisaki, Japan Trench, 41.2353°N, 144.0357°E, 2977–3014 m, R/V *Tansei-maru*, cr. KT-93-15, sta. M7, 3-m beam trawl, coll. Y. Machida, 17 Nov. 1993; BSKU 96473 (1, 95.1 mm HL, 509 mm TL), off Kinkazan, 38.4748°N, 143.3470°E, 2698–2814 m, R/V *Tansei-maru*, cr. KT-08-27, sta. K-3, 3-m ORE beam trawl, coll. H. Endo and N. Nakayama, 23 Oct. 2008; NSMT-P 97650 (3, 60.8–84.9 mm HL, 306+–444 mm TL), off Kesennuma, Japan Trench, 38.7983°N, 144.1345°E, 7340–7433 m, R/V *Hakuho-maru*, cr. KH-01-02, sta. TD-7, 30 Sept. 2001. **Australia:** FMNH 5777 (1, 39.3 mm HL, 137+ mm TL), south of Western Australia, 50.0167°S, 123.0667°E, 1800 ftm (3292 m), *Challenger* sta. 158, trawl, 7 Mar. 1874. **Northeastern Pacific:** FMNH 117120 (2, 28.9–47.7 mm HL, 141+–221+ mm TL), off Newport, 44.7658°N, 125.5294°W, 1750 m, R/V *Wecoma*, cr. W9704A, sta. 5, 40-ft otter trawl, coll. J. Voightt, 17 Apr. 1997; FMNH 117117 (1, 49.5 mm HL, 248+ mm TL), off Newport, 44.7581°N, 125.5361°W, 2850 m, R/V *Wecoma*, cr. W9704A, sta. 1,

40-ft otter trawl, coll. J. Voightt, 16 Apr. 1997; *CMNFI 1965-0321.3 (1, 84 mm HL, 342+ mm TL; examination based on only photographs provided by S. Tessier), south of Moresby Island, 51.0500°N, 132.2200°W, 2762 m, 8 Aug. 1965. **North Atlantic:** ZMUB 3384 (1 of 2, 85.8 mm HL, 315+ mm TL), 42.4333°N, 25.7500°W, 3120 m, R/V *Michael Sars*, sta. 88, 18 Jul. 1910; ZMUB 16803 (1, 62.4 mm HL, 326 mm TL), Mid-Atlantic Ridge, 42.9167°N, 30.3333°W, 2660–2670 m, R/V *G.O. Sars*, cr. MAR-ECO, sta. 1149, 7 Jul. 2004; ZMUB 16805 (1, 64.0 mm HL, 325+ mm TL), Mid-Atlantic Ridge, 53.0333°N, 33.6000°W, 2955–3071 m, R/V *G.O. Sars*, cr. MAR-ECO, sta. 1165, 24 Jul. 2004; *ZMUB 17836 (1, 71.0 mm HL, 352+ mm TL), Mid-Atlantic Ridge, 51.3297°N, 28.8690°W, 3505–3527 m, R/V *G.O. Sars*, cr. MAR-ECO, supersta. 54, local sta. 377, 16 Jul. 2004; ZMUB 17838 (1, 71.4 mm HL, 344+ mm TL), *ZMUB 17839 (1, 124+ mm TL, poor condition), *ZMUB 19319 (2, 74.7–75.3 mm HL, 378+–456 mm TL), Mid-Atlantic Ridge, 51.5493°N, 30.9683°W, 3452–3465 m, R/V *G.O. Sars*, cr. MAR-ECO, supersta. 64, local sta. 381, 21 Jul. 2004; ZMUB 17840 (1, 77.1 mm HL), ZMUB 19517 (5, 83.3–105 mm HL, 390+–578 mm TL), Mid-Atlantic Ridge, 51.0343°N, 33.6117°W, 2995–3071 m, R/V *G.O. Sars*, cr. MAR-ECO, supersta. 66, local sta. 383, 24 Jul. 2004; *ZMUB 19287 (1, 89.5 mm HL, 467+ mm TL), ZMUB 19288 (1, 107 mm HL, 567 mm TL), Mid-Atlantic Ridge, 42.9220°N, 28.1392°W, 2973–2979 m, R/V *G.O. Sars*, cr. MAR-ECO, supersta. 52, local sta. 374, 13 Jul. 2004; *ZMUB 19338 (1, 88.6 mm HL, 452+ mm TL), Mid-Atlantic Ridge, 53.1387°N, 34.7660°W, 2306–2374 m, R/V *G.O. Sars*, cr. MAR-ECO, supersta. 68, local sta. 384, 25 Jul. 2004; *ZMUB 17837 (1, 100+ mm TL, poor condition), Mid-Atlantic Ridge, 51.3297°N, 28.8690°W, 3505–3527 m, R/V *G.O. Sars*, cr. MAR-ECO, supersta. 54, local sta. 377, 16 Jul. 2004; ZMUC P372974 (1, 67.0 mm HL, 330+ mm TL), ZMUC P372975 (1, 91.3 mm HL, 416+ mm TL), northeastern Atlantic, 49.9167°N, 13.4333°W, *Challenger* sta. 51804, 26 Sept. 1983. **Uncertain variant (Japan):** BSKU 96947 (1, 80.9 mm HL, 384+ mm TL, uncertain variant), southeast of Cape Erimomisaki, Japan Trench, 40.7270°N, 143.7097°E, 2607–2609 m, R/V *Tansei-maru*, cr. KT-93-15, sta. M6, 3-m beam trawl, coll. Y. Machida, 18 Nov. 1993.

Redescription. General features are shown in Figs. 110–111. Counts and measurements are given in Table 8. Body deepest at first dorsal-fin origin, gradually tapering to long tail. Trunk short, moderately compressed, width over pectoral-fin bases 1.4–1.7 in depth at first dorsal-fin origin. Head large, HL about 4.8–5.2 in TL; head bones moderately stout; flesh firm. In the largest specimen (BSKU 96473, 80.9 mm HL), dorsal contour of nape moderately humped, forming distinct depression above orbit. Snout short, low, barely protruding beyond upper jaw, length 1.2–1.4 times as long as orbit diameter; ventral contour almost vertical in lateral view. Orbit small, greatest

FIGURE 110. Fresh specimens of *Coryphaenoides leptolepis*. (A) BSKU 96940, 63.2 mm HL, 330+ mm TL, southeast of Cape Erimomisaki, Hokkaido, Pacific, 2977–3014 m depth; (B–D) BSKU 96473, 95.1 mm HL, 509 mm TL, off Kinkazan, Miyagi Pref., Pacific, 2698–2814 m depth. (A–B) Lateral views; (C) dorsal and (D) ventral views of the head and trunk. [Photos: (A) BSKU; (B–D) H. Endo and N. Nakayama]

diameter 2.9–3.5 in postorbital length. Interorbital space slightly concave in preserved specimens, width 0.7–0.8 in orbit diameter. Mouth large, subterminal, upper-jaw length 0.4–0.5 in orbit diameter; posterior margin of maxilla reaching vertical through hind rim of orbit or slightly beyond; lateral corner of mouth not restricted by lip folds; lips thin, not papillose. Suborbital region almost flat, lacking stout bony ridge. Preopercle large, hind margin almost vertical, forming broad lobe posteroventrally; preopercular ridge with prominent backward extension at angle. Interopercle mostly covered by preopercle, but posterior portion exposed as narrow flap. Hind margin of subopercle concave, smoothly joined with opercle. Gill membranes narrowly joined across isthmus, with broad posterior free fold; gill opening wide, extending forward to below posterior margin of lower jaw. Outermost gill slit restricted by skin folds but moderately wide, length 0.7–0.9 in orbit diameter. Outer gill rakers on first arch small,

flat, lappet-like; other rakers moderately high, ridge-like; all rakers armed with short, slender spines distally; no rakers on inner side of fourth arch. Gill filaments moderately developed. Barbel long, slender throughout its length, 0.9–1.1 in orbit diameter.

Anus immediately anterior to anal-fin origin; periproct very narrow. Ventral light organ absent. Pyloric caeca 10 (based on lectotype of *C. liocephalus*).

Premaxillary teeth in tapered band, with wide villiform band of small conical teeth (about 8 rows near symphysis) and outer row of widely spaced, enlarged canines; posterior margin of tooth band falling short of lateral corner of mouth. Mandibular teeth slightly smaller than outer teeth of premaxillary, arranged in uniserial row, but rarely clustered near symphysis.

Body scales moderately small, highly deciduous, completely missing in most of the specimens examined. In 71.2 mm HL specimen (BSKU 96941), those on dorsum

TABLE 8. Morphometric measurements and meristic counts* for *Coryphaenoides leptolepis*.

Nominal species	<i>C. leptolepis</i>				<i>C. liocephalus</i>				<i>C. serrula</i>			
	Type status		NT		JPN		JPN-UV		CNP		NEP	
	LT	NA	NT	NA	JPN	JPN	JPN-UV	CNP	NEP	SEIO	JPN	PLT
Locality	Brazil	NA	NT	NA	JPN	JPN	JPN-UV	CNP	NEP	SEIO	JPN	PLT
Number of specimens	n = 1	n = 13	n = 13	n = 11	n = 11	n = 11	n = 1	n = 1**	n = 3	n = 1	n = 1	n = 1
Measurements (mm)												
TL	440+	315+–578		292+–509	384+	179+	141+–248+	137+	283+	198	406+	146+–198
HL	91.6	62.4–110		56.8–95.1	80.9	36.1	28.9–49.5	39.3	57.5	36.0	84.5	34.2–38.3
% of HL (% of PRL)												
Snout length	26	24–29		24–26	24	26	25–28	24	25	26	24	25–26
Orbit diameter	(35)	(30–38)		(31–35)	(31)	(35)	(33–36)	(32)	(33)	(36)	(31)	(33–34)
Postorbital length	18	17–21		18–20	20	21	19–21	21	20	21	17	20
Postrostral length	(24)	(22–27)		(23–26)	(25)	(28)	(25–27)	(27)	(26)	(29)	(22)	(26–27)
Orbit–preopercle distance	58	56–63		57–60	60	56	57–58	56	58	56	61	57
	(77)	(74–79)		(75–80)	(76)	(75)	(75–77)	(73)	(77)	(76)	(78)	(75–76)
	75	75–79		74–78	78	75	75–76	77	75	73	78	75–76
	(–)	(–)		(–)	(–)	(–)	(–)	(–)	(–)	(–)	(–)	(–)
	50	49–55		46–51	50	46	47–49	49	46	47	47	46
	(67)	(65–69)		(60–67)	(64)	(61)	(62–66)	(64)	(61)	(64)	(61)	(60)
Suborbital width	14	10–13		10–13	11	15	10–12	10	11	13	11	11
Upper-jaw length	(19)	(13–17)		(13–17)	(14)	(20)	(13–16)	(14)	(15)	(17)	(15)	(14–15)
Length of rictus	46	40–46		40–45	42	39	43–45	42	43	41	42	41
	(61)	(54–61)		(54–58)	(54)	(53)	(56–60)	(55)	(56)	(56)	(54)	(54)
	38	36–42		35–41	38	35	37–39	38	37	36	37	35
	(51)	(46–53)		(45–54)	(48)	(47)	(49–51)	(50)	(49)	(49)	(47)	(46–47)
Length of PMX tooth band	35	30–33		32–59	–	31	30–34	33	32	31	31	31–32
	(47)	(39–44)		(41–77)	(–)	(41)	(41–45)	(43)	(43)	(42)	(39)	(41–42)
Preoral length	–	9–12		10–13	11	11	10–12	9	9	11	9	10–12
	(–)	(12–16)		(12–17)	(15)	(15)	(13–16)	(12)	(12)	(16)	(12)	(13–16)
Dist. tip and lat. angle of snout	11	10–14		10–13	11	–	11–12	11	10	11	10	12
	(15)	(13–17)		(13–16)	(14)	(–)	(14–17)	(15)	(14)	(16)	(13)	(16)
Snout width	19	19–25		19–23	22	–	19–22	19	18	20	17	21
	(26)	(25–31)		(25–30)	(28)	(–)	(26–30)	(25)	(24)	(27)	(22)	(28)
Internasal width	19	17–19		16–19	17	19	19–20	19	17	19	17	19
	(25)	(22–26)		(20–25)	(22)	(25)	(24–27)	(25)	(22)	(27)	(22)	(25)
Interorbital width	23	21–25		22–26	24	23	24–27	23	22	23	24	24
	(31)	(28–33)		(29–34)	(31)	(31)	(31–36)	(29)	(30)	(32)	(30)	(32)

...Continued on the next page

TABLE 8. (Continued)

Nominal species	<i>C. leptolepis</i>						<i>C. liocephalus</i>				<i>C. serrula</i>							
	Type status		NT		LT		JPN		JPN-UV		CNP	NEP	SEIO	LT		PLT	HT	PT
	Brazil	NA	NA	JPN	JPN	LT	NA	JPN	JPN-UV	CNP	NEP	SEIO	JPN	CNP	PLT	HT	PT	
Locality	n = 1		n = 13	n = 11	n = 11	n = 1	n = 1	n = 1	n = 1	n = 1**	n = 3	n = 1	n = 1	n = 1	n = 1	n = 1	n = 1	n = 2
Number of specimens	46	(61)	39–55 (51–70)	33–57 (42–74)	33–57 (42–74)	47 (61)	47 (61)	47 (61)	47 (61)	44 (59)	35–39 (47–51)	33 (43)	46 (59)	46 (63)	46 (63)	42 (55)	42 (55)	37–43 (49–57)
Body width over P. bases	81	(108)	74–89 (95–112)	68–84 (89–111)	68–84 (89–111)	–	–	–	–	71 (96)	62–69 (82–91)	74 (96)	65 (83)	75 (103)	75 (103)	74 (99)	74 (99)	60 (79)
Body depth at A. origin	65	(86)	57–75 (74–94)	53–69 (68–89)	53–69 (68–89)	–	–	–	–	53 (71)	44–53 (59–69)	57 (74)	59 (75)	55 (75)	55 (75)	60 (80)	60 (80)	47–54 (63–71)
Pre-V. length	110	(146)	102–117 (135–150)	98–122 (127–160)	98–122 (127–160)	105 (135)	105 (135)	105 (135)	105 (135)	103 (137)	98–100 (130–131)	104 (135)	110 (141)	108 (148)	108 (148)	102 (136)	102 (136)	105–107 (139–142)
Pre-anus length	164	(218)	144–166 (187–217)	146–161 (191–211)	146–161 (191–211)	145 (186)	145 (186)	145 (186)	145 (186)	140 (188)	132–141 (176–185)	141 (184)	155 (199)	143 (196)	143 (196)	141 (187)	141 (187)	137–139 (181–185)
Pre-A. length	168	(224)	149–172 (193–224)	155–167 (202–218)	155–167 (202–218)	152 (194)	152 (194)	152 (194)	152 (194)	144 (193)	136–146 (182–193)	147 (191)	161 (206)	148 (202)	148 (202)	147 (195)	147 (195)	143 (189–190)
Isthmus–V. distance	–	(–)	38–49 (49–64)	35–44 (45–57)	35–44 (45–57)	39 (50)	39 (50)	39 (50)	39 (50)	36 (48)	33–36 (44–48)	41 (53)	–	–	37 (51)	–	–	–
Isthmus–anus distance	–	(–)	–	78–98 (103–127)	78–98 (103–127)	81 (104)	81 (104)	81 (104)	81 (104)	–	–	–	–	–	–	–	–	–
Isthmus–A. distance	–	(–)	83–87 (108–116)	84–103 (111–133)	84–103 (111–133)	88 (112)	88 (112)	88 (112)	88 (112)	–	71–85 (94–112)	86 (112)	–	–	–	–	–	–
V.–A. distance	66	(88)	44–70 (57–90)	53–65 (69–84)	53–65 (69–84)	50 (64)	50 (64)	50 (64)	50 (64)	49 (66)	46–52 (61–69)	46 (60)	58 (74)	49 (67)	49 (67)	47 (62)	47 (62)	44–46 (58–61)
Anus–A. distance	–	(–)	–	5–9 (7–11)	5–9 (7–11)	6 (8)	6 (8)	6 (8)	6 (8)	–	–	–	–	–	–	–	–	–
V. length	–	(–)	90–121 (118–153)	71–93 (92–123)	71–93 (92–123)	86 (111)	86 (111)	86 (111)	86 (111)	88 (117)	81–90 (108–119)	–	–	82 (113)	82 (113)	–	–	71 (94)
P. length	–	(–)	56–65 (74–86)	53–61 (69–79)	53–61 (69–79)	60 (77)	60 (77)	60 (77)	60 (77)	51 (69)	58–61 (77–82)	–	51 (65)	51 (70)	51 (70)	–	–	–
Pre-1D. length	122	(162)	114–124 (149–163)	112–118 (146–153)	112–118 (146–153)	116 (148)	116 (148)	116 (148)	116 (148)	107 (144)	113–115 (150–151)	114 (148)	104 (133)	106 (145)	106 (145)	115 (153)	115 (153)	107–110 (142–146)
Height of 1D.	–	(–)	64–75 (84–100)	68–70 (89–91)	68–70 (89–91)	71 (91)	71 (91)	71 (91)	71 (91)	–	70 (92)	–	–	–	–	–	–	–
Length of 1D. base	29	(39)	23–29 (30–38)	22–28 (29–36)	22–28 (29–36)	27 (35)	27 (35)	27 (35)	27 (35)	24 (33)	23–26 (31–35)	24 (31)	26 (33)	29 (40)	29 (40)	27 (36)	27 (36)	23–24 (30–32)

...Continued on the next page

TABLE 8. (Continued)

Nominal species	<i>C. leptolepis</i>				<i>C. liocephalus</i>				<i>C. serrula</i>	
	LT		NT		LT		PLT		HT	PT
Type status	Brazil	NA	NA	NT	JPN	JPN-UV	CNP	NEP	SEIO	
Locality	<i>n</i> = 1	<i>n</i> = 13	<i>n</i> = 11	<i>n</i> = 1	<i>n</i> = 1	<i>n</i> = 1	<i>n</i> = 1**	<i>n</i> = 3	<i>n</i> = 1	<i>n</i> = 1
Number of specimens										
Interdorsal length	43 (58)	37–54 (49–69)	33–50 (43–64)	41 (53)	30 (41)	32–37 (42–49)	45 (58)	36 (46)	42 (57)	37 (49)
Length of gill slit	28 (38)	24–28 (31–36)	23–28 (30–37)	22 (29)	25 (33)	24–28 (32–37)	24 (31)	26 (34)	26 (36)	26 (35)
Length of posterior nostril	6 (8)	5–7 (7–10)	6–7 (7–10)	7 (9)	8 (11)	6–7 (9–10)	– (–)	6 (7)	7 (10)	– (–)
Barbel length	– (–)	19–25 (25–33)	18–20 (23–27)	17 (22)	16 (21)	19–20 (25–27)	– (–)	16 (21)	18 (24)	– (–)
Counts										
1D. rays	II,8	II,8–9	II,8–9	II,8	II,8	II,9	II,8	II,9	II,10	II,9
P. rays	i17	i17–i20	i17–i19	i15– i18***	i17–i19	i18	i17–i19	i18	i19– i20	i18 i17–i18
V. rays	9	8–10	10	10	10	10	10	10	9–10	10
Outer GR on first arch	10	9–12	8–12	10	10–11	11	10	11–12	10–11	11
Inner GR on first arch	12–13	12–16	12–16	12–13	13	13	13	14	13–14	13
Outer GR on second arch	12	11–15	11–14	11–12	11–12	12	12	12	13–14	12
Inner GR on second arch	12	11–14	12–15	11–12	14	12	13	12–13	13–14	13
Longitudinal scales	35	35–42	35–41	36	39	41	39	38	38	–
Scales below 1D. origin	–	7.5–8.5	7.5–9.5	7.5	8.5	10	9	–	9.5	–
Scales below 1D. midbase	6	6–7.5	6–8	6.5	6	7	7	6	7	5
Scales below 2D. origin	9	9–11.5	8–11	7.5	8	9	9	9	9	10.5
Scales above A. origin	–	23–33	25–34	22	–	–	–	–	–	–

*Abbreviations: 1D.—first dorsal fin; 2D.—second dorsal fin; A.—anal fin; CNP—central North Pacific; GR—gill rakers; HT—holotype; JPN—Japan; NA—North Atlantic; NEP—northeastern Pacific; NT—non-type; P.—pectoral fin; PMX—premaxillary; PT—paratype; PLT—paralectotype; SEIO—southeastern Indian Ocean; UV—uncertain variant; V.—pelvic fin

**Paralectotype of *C. variabilis*

***i15 or i16 on the right side

FIGURE 111. Preserved specimens of *Coryphaenoides leptolepis*. (A) BMNH 1887.12.7.109, lectotype, 91.6 mm HL, 440+ mm TL, off Pernambuco, Brazil, southwestern Atlantic, 641 m depth; (B) BMNH 1887.12.7.111, paralectotype (also lectotype of *Macrurus liocephalus*), 84.5 mm HL, 406+ mm TL, off Boso Peninsula, Chiba Pref., Japan, 3429 m depth; (C) BMNH 1887.12.7.112, paralectotype (also paralectotype of *M. liocephalus*), 36.0 mm HL, 198 mm TL, east of Emperor Seamounts, North Pacific, 3750 m depth; (D) BMNH 1887.12.7.132, paralectotype of *Co. variabilis*, 36.1 mm HL, 179+ mm TL, east of Emperor Seamounts, Mid-Pacific, 3750 m; (E) USNM 45358, holotype of *Chalinura serrula*, 57.5 mm HL, 283+ mm TL, east of Prince of Wales Island, Gulf of Alaska, 2870 m depth; (F) MCZ 25824, holotype of *Ch. simula*, 458 mm TL (fide Goode & Bean 1883:199), south of Nova Scotia, northeastern Atlantic. Lateral views. [Photos: (A–E) N. Nakayama; (F) MCZ]

FIGURE 112. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of *Coryphaenoides leptolepis*. (A) BSKU 96941, typical specimen, 71.2 mm HL; (B) BMNH 1887.12.7.109, lectotype, 91.6 mm HL; (C–D) BSKU 96947, uncertain variant, 80.9 mm HL. Views from above. [Photos: N. Nakayama]

below interdorsal space covered with short, greatly reclined, needle-like spinules in 7 parallel rows (Fig. 112A); tip of last spinule in each row scarcely extending beyond posterior scale margin; no reticulate structures on unexposed portions. Body entirely and uniformly scaled except for fins.

No prominent enlarged scutes on tip and lateral angles of snout. Head ridges lacking stout modified scales. Single row of small embedded scales along leading edge of snout, behind which a pair of naked lunate areas narrowly developed. Ventral surfaces of head including snout, lower half of suborbital area, mandibular rami, lower preopercle, and exposed portion of interopercle completely naked. Nasal fossa also lacking scales.

Cephalic sensory canals well developed, broad, slightly inflated when fresh (Fig. 110). Open pores present and serially arranged along all canals (Fig. 113); those on infraorbital, preopercular, and mandibular canals enlarged, easily discernible even in preserved specimens.

Origin of first dorsal fin slightly posterior to vertical

through pectoral-fin base; first dorsal fin well developed, height 2.6–2.8 times as long as its base length; second spinous ray not especially elongate, armed along its leading edge with short, greatly reclined, needle-like denticles. Second dorsal fin well separated from first dorsal, interdorsal space 1.3–1.9 times as long as first dorsal-fin base length; the fin originating above base of 8th–10th anal-fin rays. Pelvic-fin base slightly anterior to vertical through pectoral-fin base; outermost ray greatly prolonged, its tip reaching bases of 5th–10th anal-fin rays, when laid back.

Color when fresh (Fig. 110). Ground color of head and body pale pinkish gray, gradually becoming darker posteriorly; abdomen bluish, especially when young; upper jaw and gill cover prominently darker; gular and branchiostegal membranes deep blue, but posterior margins of branchiostegal membranes narrowly edged with black; first dorsal, pectoral, and pelvic fins pale, but free portion of outer pelvic-fin ray blackish; anal fin uniformly dark dusky.

FIGURE 113. Diagrammatic illustrations showing the cephalic sensory system of *Coryphaenoides leptolepis*. (A) BSKU 96473, typical specimen, 95.1 mm HL; (B) BSKU 96947, uncertain variant, 80.9 mm HL. Arrows indicate occipital pores. [Illustrations: N. Nakayama]

Color in alcohol (Fig. 111). Dorsum trunk light beige, abdomen dark blue, tail gradually becoming darker posteriorly; head generally pallid, but opercle distinctly dark; orbit narrowly edged in black; lips blackish; septum between nostrils dark; oral and gill cavities dull black; gill rakers and arches dusky, filaments yellowish brown; chin dark, but gular membrane paler; branchiostegal membranes mostly pale, but abruptly blackish along their margins; chin barbel darker basally, paler distally; first dorsal fin pale to dusky overall, but tip of second spinous ray slightly blackish; second dorsal and anal fins mostly blackish, paler anteriorly; pectoral and pelvic fins generally pale, but uppermost pectoral-fin ray and outermost pelvic-fin ray prominently dark.

Size. The maximum size is about 58 cm TL (ZMUB 19517, 578 mm TL, Mid-Atlantic Ridge).

Variation. In juveniles, small but prominent black melanophores sparsely scattered over the body surface; this characteristic pigmentation disappears in large adults.

Development. Larval morphology was described by Stein (1980) and Merrett (1986).

Distribution. Widespread from the continental slope to hadal depths of the world's oceans in 640–7433 m (Marshall 1973; Iwamoto & Stein 1974; Mecklenburg *et al.* 2002; this study). In Japan, known only from the Pacific off Honshu and Hokkaido at depths of 2698–7433 m (Appendix 3-6D). Rare in its depth range.

Remarks. *Coryphaenoides leptolepis* was originally described by Günther (1877) based on specimens collected from Brazil, Japan, and the Mid-Pacific. A decade later, Günther (1887) provided a further description of this species (as *Macrurus leptolepis*) based on only the Brazilian syntype (from *Challenger* sta. 122; Fig. 111A), allocating the remaining type specimens from Japan (1 spec. from sta. 237; Fig. 111B) and the Mid-Pacific (2 spec. from sta. 246; Fig. 111C) to his new species *C. liocephalus* (described as *M. liocephalus* based on these three specimens). According to Günther (1887: annotation on p. 144), *C. liocephalus* differs from *C. leptolepis* by having a “shorter snout”, while he did not provide any other characters to separate the two species.

The synonymy between *C. leptolepis* and *C. liocephalus* has been uncertain over the past several decades (Iwamoto & Stein 1974; Iwamoto 1990; Mecklenburg *et al.* 2002). Although *C. leptolepis* was widely recorded from the Northern Hemisphere of the Atlantic and eastern Pacific, no additional specimens were collected from the type locality (off Brazil). By contrast, *C. liocephalus* was represented by only its type specimens, except for a record from off British Columbia by Hart (1973). Due to a lack of additional materials of *C. liocephalus*, previous authors did not attempt to confirm the validity of this species, and they tentatively considered *C. liocephalus* valid (see the Nomenclatural

FIGURE 114. Uncertain variant of *Coryphaenoides leptolepis*. BSKU 96947, 80.9 mm HL, 384+ mm TL, southeast of Cape Erimomisaki, Hokkaido, Pacific, 2607–2609 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of head ant trunk. Preserved condition. [Photos: N. Nakayama]

discussion). A close similarity between the two nominal species was first suggested by Iwamoto & Stein (1974), who wrote that “*C. liocephalus* appears extremely close to, and may be conspecific with, *C. leptolepis*”. The difficulty of separating these species was also indicated by Iwamoto (1990); in his key to species of the genus (Iwamoto 1990:187–188), *C. leptolepis* keys out together with *C. liocephalus* at 18a and 21b, but no further couplets are provided to distinguish the two species. Citing a personal communication from Iwamoto (31 Aug. 1999), Mecklenburg *et al.* (2002:270) indicated that the tissue samples from their type localities could help determine their taxonomic status, and if significant differences exist, specimens called *C. leptolepis* from the North Pacific correspond with *C. liocephalus*, and *C. leptolepis* would be restricted to the Atlantic.

Furthermore, *C. leptolepis* has two other junior synonyms, viz., *C. simulus* (Goode & Bean, 1883) and *C. serrula* (Bean, 1980). Six years after Günther (1877), Goode & Bean (1883) described *C. simulus* (as *Chalinura simula*) based on a single specimen collected from the south of Nova Scotia, northwestern Atlantic (Blake sta. 308) (Fig. 111F). Although this species was initially considered to be most similar to *C. affinis* Günther, 1878, Koefoed (1927:102) subsequently recognized its close similarity with *C. leptolepis*, suggesting their differences to be “extremely small and uncertain”. Nybelin (1957:268), who re-examined the type of *C. leptolepis*, distinguished these two species by several morphometric characters, including the isthmus–anus distance, preanal length, predorsal length, and barbel length. However, the suggested differences were quite subtle and based on a

limited number of specimens, viz., only two specimens for *C. simulus* combined with data given by Koefoed (1927; 7 spec.) and Parr (1946; 1 spec.). Marshall (1970:591) indicated that the differences in proportions are likely due to dehydration of the type of *C. leptolepis*, or simply attributable to intraspecific variation, and *C. simulus* was therefore regarded as a junior synonym of *C. leptolepis*. Iwamoto & Stein (1974) further confirmed the two nominal species to represent a single species, showing greater variation for the morphometric characters based on additional specimens from the northeastern Pacific (16 spec.) and northwestern Atlantic (5 spec.).

The second synonymized species, *C. serrula*, was originally described by Bean (1890; as *Chalinura serrula*) based on three specimens collected in the Gulf of Alaska (Albatross sta. 2859) (Fig. 111E). Koefoed (1927:103) distinguished this species from *C. simulus* (= *C. leptolepis*) by the combination of the orbit diameter and the number of first dorsal-fin rays. However, Iwamoto & Stein (1974) showed complete overlap in these characters based on specimens from the northeastern Pacific and northwestern Atlantic, and thus *C. simulus* was regarded as a junior synonym of *C. leptolepis*.

During the KT-08-27 cruise of the R/V *Tansei-maru*, the author collected a single specimen of *Coryphaenoides* from off Kinkazan, Miyagi Pref., Japan, at a depth of 2698–2814 m (Fig. 110B–D). This specimen was then identified as *C. liocephalus*, but a further examination revealed that it can also be referable to *C. leptolepis*. In the course of this study, 11 additional Japanese specimens of the same species were found among the collections of BSKU and NSMT. The type series of *C. leptolepis* and *C.*

liocephalus were examined to elucidate their taxonomic relationship, along with the new materials from Japan and non-type specimens of *C. leptolepis* from both the Pacific and Atlantic. The type series of *C. simulus* and *C. serrula* were also examined to reconfirm their conspecificity with *C. leptolepis*.

Comments on type specimens. As noted in the last section, *C. leptolepis* was originally described from four specimens collected from Brazil (1 spec., BMNH 1887.12.7.109, 91.6 mm HL, 440+ mm TL, Fig. 111A), Japan (1 spec., BMNH 1887.12.7.111, 84.5 mm HL, 406+ mm TL, Fig. 111B), and the Mid-Pacific (2 spec., including BMNH 1887.12.7.112, 36.0 mm HL, 198 mm TL, Fig. 111C). However, Günther (1887:144) subsequently restricted this species to Brazil, allocating the other three specimens from Japan and the Mid-Pacific to the type series of *C. liocephalus*. Günther's (1887) act is not considered to be a lectotype designation for *C. leptolepis*; the lectotype was designated by Marshall (1973:590), who recognized the Brazilian specimen as "the type" of this species [see also Eschmeyer (1998), Art. 74.5 of the ICZN (1999), and Fricke *et al.* (2020)]. In accordance with Marshall's statement, the type specimens of *C. liocephalus* are also regarded as the paralectotypes of *C. leptolepis*. To fix the scientific name of *C. liocephalus*, BMNH 1887.12.7.111 from Japan is herein designated as the lectotype of this species, and BMNH 1887.12.7.112 from the Mid-Pacific automatically becomes a paralectotype. Unfortunately, one of the paralectotypes of *C. liocephalus* (and also *C. leptolepis*) from the Mid-Pacific was not found in the course of this study.

According to Eschmeyer (1998) and Fricke *et al.* (2020), BMNH 1887.12.7.113 (not discovered during this study) is not only a paralectotype of *C. leptolepis*, but also a syntype of *C. murrayi* Günther, 1878. This specimen was collected from New Zealand (*Challenger* sta. 168), and is obviously not a type specimen of *C. leptolepis*. In the original description of *C. murrayi*, Günther (1878) did not indicate how many specimens were involved in his description, nor designate a holotype of this species. According to Günther (1878:146), five specimens were collected from the type locality, and these specimens constitute the type series of *C. murrayi* (see Eschmeyer 1998; Fricke *et al.* 2020). Iwamoto & Shcherbachev (1991) considered BMNH 1887.12.7.113 as the "putative holotype", although they recognized that the type series includes multiple specimens. Despite that the use of the term "holotype" was wrong, Iwamoto & Shcherbachev's statement would be regarded as a lectotype designation for *C. murrayi*, based on Art. 74.5 of the ICZN (1999).

According to Eschmeyer (1998) and Fricke *et al.* (2020), BMNH 1887.12.7.132 (36.1 mm HL, 179+ mm TL, Fig. 111D) has the following type status: paralectotype of *C. leptolepis*; syntype of *C. liocephalus*; and paralectotype of *C. variabilis*. This specimen is only a paralectotype of *C. variabilis*, and is not included in the

type series of either *C. leptolepis* or *C. liocephalus*. This specimen was re-identified as *C. leptolepis* by Wilson & Waples (1983:1136), and their identification was reconfirmed in this study.

Nomenclatural discussion. The present examination of the type series of *C. leptolepis* and *C. liocephalus*, along with additional specimens from the Pacific and Atlantic, revealed that the two species are not separable morphologically. In the "*Challenger*" Report, Günther (1887: annotation on p. 144) distinguished the two species only by the snout length, with *C. liocephalus* having a "shorter snout". However, it was revealed here that their type specimens cannot be distinguished by the snout length (26% HL in *C. leptolepis* and 24–26% in *C. liocephalus*; Table 8). According to Iwamoto & Stein (1974:45, table 1), modal counts of first dorsal- and pectoral-fin rays are slightly different between the Pacific and Atlantic populations of *C. leptolepis* (modally 19 vs. 21). However, the differences are negligible when the additional specimens examined are included in the analysis. Based on Günther's (1887) descriptions, Iwamoto & Stein (1974:45) implied that *C. leptolepis* might be distinguished from *C. liocephalus* by its paler color ("dirty whitish" vs. "blackish"). However, this difference appears to be attributable to intraspecific geographical variation. Other characters, including dentition, squamation, and fin position, are almost identical between the Atlantic and Pacific populations, and *C. liocephalus* is herein synonymized with *C. leptolepis*.

Similarly, *C. serrula* was reconfirmed as a junior synonym of *C. leptolepis* in this study. The holotype of *C. serrula* (Fig. 111E) agrees well with the lectotype of *C. leptolepis* (Fig. 111A), with no substantial differences in morphological characters. Counts and measurements of the type series of *C. serrula* also lie within the range of variation for *C. leptolepis* (Table 8). These two species cannot be distinguished morphologically, and therefore represent a single species.

Regarding *C. simulus*, this study follows Marshall (1973) and Iwamoto & Stein (1974) in regarding the species as a junior synonym of *C. leptolepis*. Although the holotype of *C. simulus* was not directly examined in this study, its photographs (Fig. 111F) were kindly provided by Karsten E. Hartel, the former collection manager of the Museum of Comparative Zoology of Harvard University (MCZ). This specimen is now in poor condition, with the head skin badly deteriorated, all fins heavily damaged, and the posterior half of the tail separated from the rest of the body. Unfortunately, the original description given by Goode & Bean (1883) is too brief to determine the taxonomic status of this species, without any illustrations of its morphology. However, the general appearance of the holotype of *C. simulus* agrees well with what we regard as *C. leptolepis*, and there seems little doubt of their conspecificity.

Relationships. *Coryphaenoides leptolepis* was

previously considered to belong to the subgenus *Chalinura* Goode & Bean, 1883, whose type species is *C. simulus* (junior synonym of *C. leptolepis*; see the Nomenclatural discussion). Although originally established as a distinct genus, most recent authors relegated *Chalinura* to a subgenus of *Coryphaenoides* (e.g., Iwamoto & Stein 1974; Iwamoto & Sazonov 1988; Iwamoto 1990; Iwamoto & Shcherbachev 1991). This group has been widely recognized as a valid taxon of grenadiers, but the name is preoccupied by *Chalinura* Dalman, 1826 of the spider family Hersiliidae (Arachnida) (see also Dunlop & Jekel 2008; Fricke *et al.* 2020). A replacement name, *Fuyangia*, has been provided by Whitley (1931) for the Pisces *Chalinura*.

According to Iwamoto (1990) and Iwamoto & Shcherbachev (1991), this group comprises the following 12 species: *C. brevibarbis* (Goode & Bean, 1896) restricted in the North Atlantic; *C. fernandezianus* (Günther, 1887) known only from off Juan Fernandez Island, southeastern Pacific; *C. grahami* Iwamoto & Shcherbachev, 1991 distributed in the Indian Ocean, the southeastern Atlantic off South Africa, and the southwestern Pacific off Australia; *C. leptolepis*; *C. liocephalus*; *C. mcmillani* Iwamoto & Shcherbachev, 1991 widespread in the Southern Hemisphere, including the southeastern Atlantic, western and central Indian Ocean, southern Australia, and New Zealand; *C. mediterraneus* (Giglioli, 1893) sporadically recorded from the North Atlantic; *C. murrayi* Günther, 1878 known from the western Indian Ocean, southern Australia, New Zealand, and Fiji; *C. profundicola* (Nybelin, 1957) known only from the northeastern Atlantic; *C. serrulatus* Günther, 1878 widely distributed in the Indian Ocean and southwestern Pacific; and *C. striatura* Barnard, 1925 and *C. subserrulatus* Makushok, 1976 widespread in the Southern Hemisphere. *Coryphaenoides liocephalus* is synonymized with *C. leptolepis* in this study, which reduces the total number of species to 11. On the other hand, Iwamoto & Shcherbachev (1991) recognized two subspecies of *C. serrulatus*: *C. s. oceanus* Iwamoto & Shcherbachev, 1991 from the central and western Indian Ocean; and *C. s. subserrulatus* from New Zealand and Australia. Iwamoto & Graham (2001) also reported two juveniles collected from off the east coast of Australia under the name *Coryphaenoides* sp. cf. *fernandezianus*.

Comparisons. *Coryphaenoides leptolepis* readily differs from other species of the subgenus *Fuyangia*, except *C. brevibarbis*, *C. fernandezianus*, and *C. profundicola*, in having the combination of 8–10 (usually 9 or 10) pelvic-fin rays and prominent pores along the cephalic sensory canals. *Coryphaenoides leptolepis* differs notably from the above three species in having a longer barbel (16–25% HL vs. <12%). It further differs from *C. brevibarbis* and *C. fernandezianus* in having a wider gill slit (22–28% HL vs. 15–19% in *C. brevibarbis* and 18% in *C. fernandezianus*); from *C. brevibarbis* in

having a narrower interorbital space (21–27% HL vs. 30–37%); from *C. fernandezianus* in having a narrower snout (internasal width 16–20% HL vs. 23%); and from *C. profundicola* in having a larger orbit (17–21% HL vs. 14%), smaller mouth (upper-jaw length 39–46% HL vs. 48%), and a shorter pectoral fin (51–65% HL vs. 73%).

Uncertain variants. A specimen collected from the south of Hokkaido (BSKU 96947, 80.9 mm HL, 384+ mm TL; Fig. 114) differs from typical specimens of *C. leptolepis* in that the posterior ends of the occipital canals abut along the dorsal midline of the head, and there is no interspace between the occipital pores (Fig. 113B). In contrast, these pores are always separated from each other in the other specimens of *C. leptolepis* examined (Fig. 113A), with their interspace being 4–7% HL. Because such variation has not been reported for any species of grenadiers, this unusual specimen was initially considered to represent an undescribed species. However, similar variation was also found in the *C. yaquinae* examined (NSMT-P 78022, 82.7 mm HL, 441+ mm TL), and the difference in the occipital pores could be attributable to intraspecific variation. The Hokkaido specimen is further characterized by having a narrower gill slit (22% HL vs. 23–28% in the other *C. leptolepis* specimens examined), and fewer scale rows below the second dorsal-fin origin 7.5 (vs. 8–11.5) and above the anal-fin origin 22 (vs. 23–34). However, these differences seem too subtle to warrant their specific separation, and the disjunct datasets may overlap if more specimens are included in the analysis.

Melo *et al.* (2010) reported *Coryphaenoides* sp. cf. *leptolepis* based on a small damaged specimen (MNRJ 26899, 302+ mm TL) collected from off Rio de Janeiro. This specimen differs notably from *C. leptolepis* in having lower counts of pelvic-fin rays [7 vs. usually 9–10 (ZMUB 3384 exceptionally has 8 rays on the left side)] and higher counts of first dorsal-fin rays [II,11 vs. II,8–10 (usually II,8–9; II,10 only in the paralectotype of *C. liocephalus*)]. Melo *et al.* (2010) also indicated that their specimen has a shorter pelvic fin as compared with published data for *C. leptolepis* (60% HL vs. 96%). This difference needs further confirmation, considering the poor condition of the additional Brazilian specimen and the fragile nature of the outer pelvic-fin ray. Its length also shows a positive allometry in *C. leptolepis*, with a wide range of variation (71–121% based on the present study). Due to the paucity of available materials, the taxonomic status of the additional Brazilian specimen is uncertain, but the differences in the fin-ray counts suggest it may be distinct from *C. leptolepis*. A definitive conclusion should be made when more specimens become available for comparisons.

Range extension. *Coryphaenoides leptolepis* was previously known only from the Atlantic and the North Pacific Oceans (see the synonymy of this species). During a visit to the Field Museum of Natural History (FMNH) in Chicago, a small nineteenth century specimen of *C.*

leptolepis collected from the Indian Antarctic Ridge in the southeastern Indian Ocean (FMNH 5777, 39.3 mm HL, 137+ mm TL) was found among the collection. Surprisingly, the specimen was originally collected during the “*Challenger*” Expedition (sta. 158) in 1874. According to W. Leo Smith, a former curator of the FMNH fish collection, this historically important specimen was probably gifted to Chicago City for the World’s Columbian Exposition (also known as the Chicago World’s Fair) held in 1893. It represents not only the first record of *C. leptolepis* from the Indian Ocean, but also the second confirmed record of this species from the Southern Hemisphere. For future studies, counts and measurements for the Indian Ocean specimen are provided in Table 8. The discovery of *C. leptolepis* from the Indian Ocean suggests that the species has a cosmopolitan distribution at bathyal depths of the world’s oceans.

In addition, three specimens collected from off Kesenuma, Miyagi Pref., at a depth of 7340–7433 m represent the deepest record for *C. leptolepis*, and confirm its occurrence in the hadal zone in the northwestern Pacific.

***Coryphaenoides longifilis* Günther, 1877**

[Japanese name: Himo-dara]

(Figs. 115–118; Appendix 3-6E)

Coryphaenoides longifilis Günther, 1877:439 [original description; holotype: BMNH 1887.12.7.136, from “south of Yeddo” (off Cape Omaezaki), *Challenger* sta. 235, in 565 fth (1034 m)]; Okamura 1970a:121, pl. XXVI, text-fig. 50 (description; 6 spec. from Pacific off southern Japan from Kashima-nada to Mimase); Kuroda 1971:56 [listed from Suruga Bay after Okamura (1970a)]; Tominaga & Uyeno 1981:489 (listed; Japan); Shiogaki 1982:33 (listed; Aomori Pref.); Sawada 1983:109, 194, fig. 61 (brief description; 2 spec. from Pacific off Tohoku; photo based on HUMZ 78121); Ohta 1983: table B (listed; Suruga); Okamura 1984b:95, pl. 82, fig. B (compiled); Okamura 1988:95, pl. 82, fig. B (compiled); Nakabo 1993:362 (in key; Japan); Amaoka *et al.* 1995:103, fig. 153 (compiled; northern Japan); Hori 1996:28 (listed; Ibaraki Pref.); Shinohara *et al.* 1996:169 (2 spec. listed from Pacific off Tohoku); Shinohara & Matsuura 1997:291 (listed; Suruga Bay); Iwamoto 1990:211, fig. 488 (synopsis); Okamura 1997:125, fig. 5 (compiled); Funabashi 1998:85 (listed; Ibaraki Pref.); Nakabo 2000:426 (in key; Japan); Shinohara *et al.* 2001:305 (listed; Tosa Bay); Zama 2001:40 (listed; Miyagi Pref.); Nakabo 2002:426 (in key; Japan); Maeda & Tsutsui 2003:488 (listed; Hokkaido); Shiogaki *et al.* 2004:55 (listed; Pacific off Aomori Pref.); Senou *et al.* 2006:420 (listed; Sagami Sea); Senou 2008:366, fig. 31.33 (in situ observation; off Sanriku); Kitagawa *et al.* 2008:40, unnumbered fig. (brief description; spec. from Pacific off Tohoku); Shinohara *et al.* 2009:708 (1 spec. listed from Pacific off Tohoku); Amaoka *et al.* 2011:128, fig. 161 (compiled; Hokkaido); Nakabo & Kai

2013:502 (in key; Japan); Amaoka *et al.* 2020:160, fig. 215 (listed; Hokkaido); Motomura 2020:39 (listed; Japan).

Macrurus longifilis: Günther 1887:151, pl. XXXV (further description of holotype).

Bogoslovius clarki Jordan & Gilbert in Jordan & Evermann, 1898:2575 [original description; 4 syntypes in CAS and USNM; type locality: Bering Sea off Bogoslof Island, 54°51'00"N, 167°27'00"W, in 644 fth (1178 m), *Albatross* sta. 3634; this publication predates Jordan & Gilbert (1899)]; Jordan & Gilbert 1899:487, pl. LXXXIII [description as a “new species”; preoccupied by Jordan & Gilbert in Jordan & Evermann (1898); 4 spec. from *Albatross* sta. 3634].

Dolloa longifilis: Jordan 1900:987 (new combination); Jordan & Gilbert in Jordan & Starks 1904:607 [after Günther (1877, 1887)]; Jordan *et al.* 1913:411 (listed; Japan; new Japanese name: “Himo-dara”); Kamohara 1931b:544 (questionable; listed; Kochi).

Moseleya longifilis: Jordan & Snyder 1901:119 (listed; Japan).

Nematonurus longifilis: Gilbert & Hubbs 1916:159 (new combination; description; 3 spec. from Hyuga-nada and Enshu-nada); Okada & Matsubara 1938:448 (in key; Japan); Kamohara 1950:276 (questionable; listed; Kochi Pref.); Kamohara 1952:96 (questionable; listed; Kochi Pref.); Matsubara 1955:1308 (in key; Japan); Kamohara 1958:73 (questionable; listed; Kochi Pref.); Kamohara 1964:95 (questionable; listed; Kochi Pref.); Tataru *et al.* 1965:121 (questionable; listed; off Shikoku).

Coryphaenoides (Nematonurus) longifilis: Tarantetz 1937:169 (new combination; listed; Soviet Far East); Okamura 1970b: table 1 (listed; Japan).

Diagnosis. Pelvic-fin rays 9–10. Snout low, bluntly rounded in lateral view, protruding scarcely beyond large subterminal mouth. No modified scales on head ridges and tip and lateral angles of snout. Posterior margin of upper jaw extending to vertical through posterior 1/3 of orbit or beyond; upper-jaw length 40–46% HL; lateral corner of mouth not restricted by skin folds. Outermost gill slit moderately wide, length 14–23% HL. Barbel tiny, length 1–5% HL. Premaxillary teeth in 2 distinct rows with outer series notably enlarged; mandibular teeth in 2 irregular rows with inner series greatly enlarged. Body scales covered with short, reclined, needle-like to knife-like spinules in narrowly divergent rows (spinules rudimentary in larger specimens); tip of last spinule in each row scarcely extending beyond posterior scale margin (except in small specimens). Transverse scale rows below first dorsal-fin midbase 10–14.5. Head fully scaled except for narrow naked area on underside of snout. Interdorsal space less than first dorsal-fin base length. Height of first dorsal fin less than HL (71–94% HL); second spinous ray weakly serrated along its leading edge; first dorsal-fin rays II, 12–15. Upper pectoral- and outer pelvic-fin rays greatly elongate, extending well beyond anal-fin origin. Body pale to blackish.

Material examined. 29 specimens. **Holotype** of

***Coryphaenoides longifilis*:** BMNH 1887.12.7.136 (113 mm HL, 667+ mm TL), off Cape Omaezaki, Enshunada, Japan, 34.1167°N, 138.0000°E, 565 ftn (1034 m), *Challenger* sta. 235, trawl, 4 Jun. 1875. **Syntypes of *Bogoslovius clarki*:** CAS-SU 5748 (2, 53.5–68.2 mm HL, 301+–389+ mm TL), north-northeast of Bogoslof Island, Bering Sea, 54.8500°N, 167.4500°W, 664 ftn (1215 m), *Albatross* sta. 3634, 7 Jul. 1896. **Non-types:** **Japan:** *BSKU 43477 (1, 130 mm HL, 767+ mm TL), off Miyazaki, Hyuga-nada, 32.0650°N, 132.2950°E, 1463–1514 m, R/V *Tansei-maru*, cr. KT-86-16, sta. E, 3-m beam trawl, 3 Nov. 1986; *BSKU 14194 (1, 68.0 mm HL, 367+ mm TL), BSKU 14196 (1, 135 mm HL, 826+ mm TL), off Hitachi, 36.3189°N, 141.6667°E, 1690 m, 3 Dec. 1967; HUMZ 72560 (1, 124 mm HL, 785+ mm TL), off Fukushima, 37.1567°N, 141.9333°E, 900–920 m, F/V *Hatsue-maru*, No. 62, bottom trawl, coll. M. Toyoshima, 19 Jan. 1978; HUMZ 72597 (1, 135 mm HL, 860+ mm TL), off Fukushima, 37.1833°N, 141.9500°E, 810–820 m, F/V *Hatsue-maru*, No. 62, bottom trawl, coll. M. Toyoshima, 18 Jan. 1978; HUMZ 72659 (1, 124 mm HL, 772 mm TL), off Miyagi, 38.0333°N, 142.4833°E, 1100–1150 m, F/V *Hatsue-maru*, No. 62, bottom trawl, coll. M. Toyoshima, 7 Feb. 1978; HUMZ 72674 (1, 124 mm HL, 749+ mm TL), HUMZ 72675 (1, 118 mm HL, 535+ mm TL), off Miyagi, 38.7250°N, 142.3917°E, 1000–1100 m, F/V *Hatsue-maru*, No. 62, bottom trawl, coll. M. Toyoshima, 4 Feb. 1978; NSMT-P 104608 (1, 122 mm HL, 728+ mm TL), off Otsuchi, R/V *Tansei-maru*, cr. KT-87-05, sta. SR35, 3-m ORE beam trawl, 21 May 1985; HUMZ 78121 (1, 137 mm HL, 803 mm TL), off Iwate, 39.1217°N, 142.7750°E, 1480–1485 m, bottom trawl, coll. K. Sakamoto, 20 Sept. 1978; NSMT-P 79350 (1, 89.9 mm HL, 500+ mm TL), off Kesennuma, 38.5367°N, 142.8000°E, 1502 m, FRV *Wakataka-maru*, sta. E-1500, bottom trawl, coll. Y. Takata, 26 Oct. 2007; BSKU 35689 (1, 88.7 mm HL, 497+ mm TL), BSKU 35690 (1, 98.1 mm HL, 597+ mm TL), BSKU 35692 (1, 102 mm HL, 584+ mm TL), BSKU 35694 (1, 115 mm HL, 683+ mm TL), off Kinkazan, 38.7333°N, 143.1850°E, 1950–1960 m, R/V *Hakuho-maru*, cr. KH-81-4, sta. 8, 25 Jul. 1981; BSKU 97807 (1, 129 mm HL, 777+ mm TL), off Kuji, 40.3727°N, 143.3827°E, 1923–1995 m, R/V *Tansei-maru*, cr. KT-93-15, sta. M5, 3-m beam trawl, coll. Y. Machida, 18 Nov. 1993; *NSMT-P 102089 (2, 21.8–44.7 mm HL, 96+–226+ mm TL), east-southeast of Oshika Peninsula, 38.0367°N, 142.2250°E, 829–831 m, FRV *Wakataka-maru*, otter trawl, coll. T. Fujita, 19 May 1995; NSMT-P 90551 (1, 78.0 mm HL, 457+ mm TL), off Kushiro, 1500 m, R/V *Tansei-maru*, cr. KT-07-29, coll. G. Shinohara, 7 Nov. 2007; *BSKU 114807 (1, 27.0 mm HL, 119+ mm TL), south of Kushiro, 42.6542°N, 144.3770°E, 780 m, FRV *Wakataka-maru*, sta. D-700, otter trawl, coll. R. Misawa, 7 Jun. 2014; *BSKU 49902 (1, 96.2 mm HL, 522+ mm TL), NSMT-P 105269 (1, ex. BSKU 49909, 115 mm HL, 682+ mm TL), NSMT-P 105270 (1, ex. BSKU

49906, 128 mm HL, 770+ mm TL), Japan, 1500 m, R/V *Hakuho-maru*, cr. KH-91-06, sta. 12, 16 Nov. 1991. **Emperor Seamounts:** HUMZ 71903 (1, 126 mm HL, 690+ mm TL), HUMZ 71905 (1, 140 mm HL, 824+ mm TL), Ojin Seamount, 38.0000°N, 170.3000°E, 920–1080 m, coll. Japan Marine Fishery Resources Research Center (JAMARC), 17 Jul. 1977; HUMZ 88055 (1, 126 mm HL, 649+ mm TL), north of Nintoku Seamount, 42.2833°N, 170.4200°E, 1085–1200 m, F/V *Hokusei-maru*, hook and line, coll. K. Nakaya, 23 Jul. 1980.

Counts and measurements. Based on 23 specimens (53.5–140 mm HL, 301+–860+ mm TL). Counts: first dorsal-fin rays II,12–15; pectoral-fin rays i15–i19; pelvic-fin rays 9–10; gill rakers on first arch (outer/inner) 6–9/13–16, on second arch 12–15/13–15; longitudinal scales 38–52; transverse scale rows below first dorsal-fin origin 13–17, below first dorsal-fin midbase 10–14.5, below second dorsal-fin origin 11–16.5, above anal-fin origin 48.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 24–28 (30–37); orbit diameter 19–25 (25–32); postorbital length 53–58 (69–75); postrostral length 74–79; orbit–preopercle distance 38–44 (48–58); suborbital width 9–13 (12–17); upper-jaw length 40–46 (52–62); length of rictus 35–40 (46–53); length of premaxillary tooth band 28–35 (36–46); preoral length 4–7 (6–9); distance between tip and lateral angle of snout 8–11 (11–14); snout width 14–19 (18–24); internasal width 16–18 (21–24); interorbital width 22–26 (29–35); occipital width 2–8 (3–11); body width over pectoral-fin bases 30–56 (38–73); body depth at first dorsal-fin origin 70–86 (88–114); body depth at anal-fin origin 54–89 (71–118); prepelvic length 103–120 (132–159); preanus length 146–179 (191–237); preanal length 137–188 (174–249); isthmus–pelvic distance 42–54 (55–72); isthmus–anus distance 87–115 (114–153); isthmus–anal distance 90–124 (116–164); pelvic–anal distance 43–74 (55–98); anus–anal distance 3–9 (4–12); pelvic-fin length 114–222 (151–291); pectoral-fin length 66–123 (84–162); predorsal length 104–116 (134–152); height of first dorsal fin 71–94 (90–120); length of first dorsal-fin base 31–42 (40–55); interdorsal length 10–22 (12–27); length of gill slit 14–23 (19–29); length of posterior nostril 4–8 (5–10); barbel length 1–5 (1–6).

Size. Attains about 97 cm TL (NSMT-P 102193, 973+ mm TL, off Kesennuma, Japan).

Variation. Spinulation on the body scales is well developed in young specimens, and it tends to become weaker with increasing body size (Fig. 118). In adults, the spinulation is usually rudimentary and difficult to distinguish (Fig. 118B–C, F–G), but some specimens have prominent spinules so as to give a spiny appearance to body surfaces (Fig. 118D, H).

Distribution. Distributed in Japan and the Bering Sea, as well as on the Emperor Seamounts, at depths of 700–2025 m (Iwamoto & Stein 1974; Borets 1986; Iwamoto

FIGURE 115. Fresh specimens of *Coryphaenoides longifilis*. (A) BSKU 114807, 27.0 mm HL, 119+ mm TL, south of Kushiro, Hokkaido, Pacific; (B) BSKU 49902, 96.2 mm HL, 522+ mm TL, KH-91-6, sta. 12, 1500 m depth; (C) HUMZ 88055, 126 mm HL, 649+ mm TL, Emperor Seamounts; (D) BSKU 43477, 130 mm HL, 767+ mm TL, Hyuga-nada off Miyazaki Pref., Pacific, 1514–1463 m depth. Lateral views. [Photos: (A) N. Nakayama; (B, D) BSKU; (C) HUMZ]

1990; Mecklenburg *et al.* 2002; this study). In Japanese waters, known from off the Pacific coasts southward to Miyazaki (31.90°N) and the Shichito-Iojima Ridge at depths of 780–2000 m (Appendix 3-6E). Moderately common in northern Japan, but rare in the west of the Boso Peninsula (139.89°E) and at the Shichito-Iojima Ridge.

Remarks. For a full description see Okamura (1970a) and Iwamoto & Stein (1994). *Coryphaenoides longifilis* was originally described by Günther (1877) based on a single specimen (Fig. 116) collected during the “Challenger” Expedition from the Enshu-nada off Cape Omaezaki, Japan, at a depth of 1034 m (*Challenger*

sta. 235). Subsequently, Jordan & Gilbert (*in* Jordan & Evermann 1898) described *Bogoslovius clarki* as a new species based on four specimens collected from the Bering Sea off Bogoslof Island at a depth of 1214 m (Fig. 117). However, Makushok (1964) synonymized this species with *C. longifilis* based on a detailed comparison of literature data.

Relationships. The subgeneric affiliation of this enigmatic species needs further discussion. Although it was originally included in the genus *Coryphaenoides*, Günther (1887) subsequently classified the species under the subgenus *Nematonurus* Günther, 1887 of the genus *Macrurus* (= *Macrourus* Bloch, 1786). Goode & Bean

FIGURE 116. Holotype of *Coryphaenoides longifilis*. BMNH 1887.12.7.136, 113 mm HL, 667+ mm TL, off Cape Omaezaki, Enshu-nada, Pacific, 1034 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

FIGURE 117. Preserved specimen of *Coryphaenoides longifilis*. CAS-SU 5748, syntype of *Bogoslovius clarki* (1 of 2 spec.), 68.2 mm HL, 389+ mm TL, north-northeast of Bogoslof Island, Bering Sea, 1215 m depth. (A) Lateral, (B) dorsal, and (C) ventral views. Preserved condition. [Photos: N. Nakayama]

FIGURE 118. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of *Coryphaenoides longifilis*. (A, E) NSMT-P 102089, 44.7 mm HL; (B, F) BSKU 14194, 68.0 mm HL; (C, G) NSMT-P 104608, 122 mm HL; (D, H) BSKU 97807, 129 mm HL. (A–D) Views from above; (E–H) oblique views. [Photos: NSMT]

(1896) established *Moseleya* as a monotypic genus for *C. longifilis*, but the name is preoccupied by the stony coral genus *Moseleya* Quelch, 1884 (Cnidaria: Scleractinia: Lobophylliidae). Jordan (1900) subsequently provided a replacement name, *Dolloa*, for Goode & Bean's (1896) *Moseleya*. However, Gilbert & Hubbs (1916) regarded *Dolloa* as a junior synonym of *Nematonurus*, to which they gave full generic status. Okamura (1970a) and Iwamoto & Stein (1974) classified the species under *Coryphaenoides*, and relegated *Nematonurus* to subgeneric rank. Iwamoto & Sazonov (1988) listed *C. longifilis* under the subgenus *Coryphaenoides*, but Iwamoto (1990) transferred the species to the subgenus *Bogoslovius* Jordan & Evermann, 1898.

According to a molecular phylogeny based on the 12S mitochondrial ribosomal gene from 12 *Coryphaenoides* species (Wilson & Attia 2003: fig. 2), *C. longifilis* belongs to the subgenus *Coryphaenoides*, and is sister to a clade that comprises *C. acrolepis* (Bean, 1884), *C. cinereus* (Gilbert, 1896), *C. filifer* (Gilbert, 1896), and *C. pectoralis* (Gilbert, 1891). However, it is evident that more investigations are necessary for a better understanding of the subgeneric classification of *Coryphaenoides*. Considering that the genus represents the second largest group of the family, a more comprehensive analysis with broader taxon sampling is required to make a definitive conclusion on the species' phylogenetic position. This study follows Iwamoto's (1990) classification, and *C. longifilis* is considered to belong the subgenus *Bogoslovius*, of which the species is the only representative.

Comparisons. *Coryphaenoides longifilis* is unlikely to be confused with other congeners in its diagnostic dentition: the dentary teeth are arranged in two distinct series, with those of the inner row modified into enlarged canines. In addition, *C. longifilis* is easily recognized due to its unique physiognomy: unlike most other species of the genus, the snout of *C. longifilis* is broadly rounded and barely protrudes beyond the upper jaw; this feature is shared only with *C. gypsochilus* Iwamoto & McCosker, 2001 known from the Galápagos Islands and some species of the subgenus *Fuyangia* Whitley, 1931 (see also the Relationships of *C. leptolepis*). *Coryphaenoides longifilis* is one of the few species of the genus in which the outer pelvic-fin ray is extremely prolonged (its length greater than 1.5 times HL). Among the species of *Coryphaenoides*, the first dorsal fin is best developed in *C. longifilis*, and its base length appears to be the longest (31–42% HL); higher counts of first dorsal-fin rays (II, 12–15) also diagnose the species. Furthermore, body scales are much smaller in *C. longifilis* (below first dorsal-fin midbase 10–14.5; below second dorsal-fin origin 11–16.5) than in any other congeners.

***Coryphaenoides marginatus* Steindachner & Döderlein, 1887**

[Japanese name: Heri-dara]

(Figs. 119–121, 122A–B; Table 9; Appendix 3–6F)

Coryphaenoides marginatus Steindachner & Döderlein, 1887:284 [original description; 1 spec. from “Tokio”, but probably from Eno-shima Island, Sagami Bay; see Amaoka (2007:28)]; Jordan & Snyder 1901:120 (listed; Japan); Jordan & Gilbert in Jordan & Starks 1904:610 [after Steindachner & Döderlein (1887)]; Jordan *et al.* 1913:415 (listed; Japan; new Japanese name: “Heri-dara”); Gilbert & Hubbs 1916:164 (description; 43 spec. from East China Sea and Suruga Bay); Okada & Matsubara 1938:449 (in key; Japan); Kuroda 1951:391 (listed; Suruga Bay); Kuroda 1952:176 (2 spec. from Suruga Bay); Matsubara 1955:1309 (in key; Japan); Matsubara 1965:504 (compiled; Japan); Okamura 1970a:136, pl. XXX, text-figs. 54–55 (description; biological notes; 43 spec. from Pacific off southern Japan from Choshi to Suruga Bay); Tominaga & Uyeno 1981:489 (listed; Japan); Okamura 1982:165, 352, fig. 100 [brief description; 1 spec. from Tosa Bay (BSKU 29492)]; Ohta 1983: photo. 27, tables A–B (listed; Suruga Bay; in situ observation); Yatou 1984:219, 363, fig. 154 [brief description; 10 spec. from Okinawa Trough; photo based on BSU 29492 (from Tosa Bay)]; Okamura 1984b:95, pl. 82, fig. E (compiled); Okamura 1988:95, pl. 82, fig. E (compiled); Iwamoto 1990:214, fig. 492 (synopsis); Nakabo 1993:365 (in key; Japan); Suzuki & Kataoka 1997:82, pl. 33, fig. 177 (brief description; 1 spec. from Kumano-nada); Okamura 1997:126, fig. 2 (compiled); Shinohara & Matsuura 1997:291 (listed; Suruga Bay); Nakabo 2000:429 (in key; Japan); Shinohara *et al.* 2001:305 (12 spec. listed from Tosa Bay); Nakabo 2002:429 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Shinohara *et al.* 2005:417 (2 spec. listed from Ryukyu Islands); Fukui *et al.* 2008:286, figs. 3–4 (descriptions of early life stages; Suruga Bay); Kitagawa *et al.* 2008:41, unnumbered fig. (brief description; spec. from Pacific off Tohoku); Shinohara *et al.* 2009:708 (listed; Pacific off Tohoku); Kim *et al.* 2009b:107, fig. 2 (brief description; 10 spec. from Korea, Taiwan, and Japan; new Korean record); Furuhashi *et al.* 2010: table 2 (160 spec. listed from northern Okinawa Trough); Nakabo & Kai 2013:505 (in key; Japan); Miyazaki *et al.* 2019: fig. 3B, table 1 (4 spec. listed from Tokyo Submarine Canyon); Motomura 2020:39 (listed; Japan).

Coryphaenoides awae Jordan & Gilbert in Jordan & Starks, 1904:608, unnumbered fig. (original description; holotype: CAS-SU 8547, from “off Nanaura in Awa at the entrance of by of Tokyo”); Jordan *et al.* 1913:416, fig. 387 (listed; Japan; new Japanese name: “Bōshū-soko-dara”); Gilbert & Hubbs 1916:166 (additional measurements of holotype); Okada & Matsubara 1938:449 (in key; Japan); Kuroda 1951:391 (listed; Suruga Bay); Kuroda 1952:176 (1 spec. from Suruga Bay); Matsubara 1955:1309 (in key; Japan).

Coryphaenoides (Coryphaenoides) marginatus: Okamura 1970b: table 1 (listed; Japan).

Diagnosis. Pelvic-fin rays 6–9 (usually 8). Snout short, slightly protruding beyond upper jaw. Tip and lateral angles of snout armed with stout tubercles; scales along head ridges not coarsely modified. Mouth small, posterior margin of upper jaw not extending beyond vertical through midorbit; upper-jaw length 28–32% HL; lateral corner of mouth moderately restricted by skin folds. Outermost gill slit greatly restricted by skin folds, length 3–9% HL. Barbel moderately short, stout at base, length 5–14% HL. Teeth all small, conical, in wide bands in both jaws, none especially enlarged. Body scales covered with short, greatly reclined, needle-like spinules in tightly packed parallel rows; tip of last spinule in each row scarcely extending beyond posterior scale margin. Transverse scale rows below first dorsal-fin midbase 5.5–8.5. Top of snout fully scaled; underside of head with broad naked band above upper lip. Second spinous ray of first dorsal fin moderately to greatly prolonged, height of first dorsal fin equal to or greater than HL (101–219% HL); serration along its leading edge rudimentary in adults; first dorsal-fin rays II, 9–11. Pelvic-fin bases closer together, their interspace less than base length of each fin. Body generally dusky brown overall.

Material examined. 45 specimens. **Holotype of *Coryphaenoides awae*:** CAS-SU 8547 (sex undetermined, 117 mm HL, 622+ mm TL), west of southern Chiba Pref. (Awa), Japan, date unknown. **Non-types: Japan:** BSKU 27128 (1 sex undetermined, 33.2 mm HL, 186+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.3833°N, 127.5150°E, 650 m, F/V *Yuryo-maru*, No. 8, tr. 21, bottom trawl, coll. Y. Kinoshita and S. Hagino, 5 Feb. 1978; BSKU 27925 (1 female, 42.6 mm HL, 237+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 28.7500°N, 127.1167°E, 520–542 m, F/V *Ryoan-maru*, No. 28, tr. 58, bottom trawl, coll. Y. Kinoshita and S. Hagino, 16 Mar. 1978; BSKU 71614 (1 male, 38.4 mm HL, 230+ mm TL), BSKU 71615 (1 sex undetermined, 26.9 mm HL, 152+ mm TL), Tosa Bay, 33.1950°N, 133.6750°E, 528–537 m, FRV *Kotaka-maru*, cr. 97-5-500, otter trawl, 13 May 1997; BSKU 69280 (1 sex undetermined, 49.6 mm HL, 254+ mm TL), Tosa Bay, 33.1417°N, 133.6433°E, 528–537 m, FRV *Kotaka-maru*, cr. 97-5-500, otter trawl, 13 May 1997; BSKU 23284 (1 male, 73.4 mm HL, 445 mm TL), Mimase fish market, bottom trawl, date unknown; BSKU 29492 (1 female, 61.7 mm HL, 348+ mm TL), Tosa Bay, 32.9667°N, 133.5333°E, 605 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, tr. T-19, bottom trawl, coll. O. Okamura *et al.*, 21 Dec. 1979; BSKU 103049 (1 male, 66.7 mm HL, 378 mm TL), BSKU 103052 (1 female, 105 mm HL, 541 mm TL), BSKU 103055 (1 female, 102 mm HL, 512+ mm TL), Suruga Bay, 350 m, coll. A. Fukui and M. Takami, 13 Oct. 2009; BSKU 103067 (1 female, 81.0 mm HL, 444+ mm TL), BSKU 103070 (1 female, 75.1 mm HL, 409+ mm TL), Suruga Bay, 500 m, coll. A. Fukui and M. Takami, 2 Oct. 2009; BSKU 103060 (1 male, 60.6 mm HL, 360+

mm TL), BSKU 103061 (1 sex undetermined, 54.5 mm HL, 323+ mm TL), BSKU 103062 (1 male, 60.5 mm HL, 349+ mm TL), BSKU 103063 (1 sex undetermined, 61.9 mm HL, 355+ mm TL), BSKU 103064 (1 male, 64.5 mm HL, 381+ mm TL), BSKU 103066 (1 male, 73.4 mm HL, 450 mm TL), Suruga Bay, 500 m, coll. A. Fukui and M. Takami, 3 Aug. 2009; BSKU 103057 (1 female, 80.1 mm HL, 396+ mm TL), BSKU 103059 (1 male, 85.5 mm HL, 452+ mm TL), Suruga Bay, coll. A. Fukui and M. Takami, 11 Jun. 2009; BSKU 103033 (1 female, 92.9 mm HL, 449+ mm TL), Suruga Bay, coll. A. Fukui and M. Takami, 12 Dec. 2007; BSKU 103047 (1 male, 71.6 mm HL, 420+ mm TL), BSKU 103048 (1 female, 71.4 mm HL, 392 mm TL), Suruga Bay, coll. A. Fukui and M. Takami, 13 Jun. 2007; BSKU 103044 (1 female, 89.1 mm HL, 463 mm TL), BSKU 103045 (1 male, 44.0 mm HL, 252+ mm TL), BSKU 103046 (1 male, 68.1 mm HL, 413 mm TL), Suruga Bay, coll. A. Fukui and M. Takami, 16 2009; BSKU 103034 (1 male, 71.6 mm HL, 404+ mm TL), BSKU 103035 (1 female, 96.5 mm HL, 536 mm TL), BSKU 103036 (1 female, 81.1 mm HL, 450 mm TL), BSKU 103037 (1 sex undetermined, 73.7 mm HL, 421+ mm TL), BSKU 103038 (1 male, 52.8 mm HL, 324+ mm TL), BSKU 103039 (1 male, 60.4 mm HL, 362 mm TL), BSKU 103040 (1 male, 79.4 mm HL, 461+ mm TL), Suruga Bay, coll. A. Fukui and M. Takami, 23 Jul. 2009; BSKU 110135 (1 sex undetermined, 72.0 mm HL, 405 mm TL), Suruga Bay, 34.7619°N, 138.4811°E, 196–387 m, F/V *Hinode-maru*, sta. 5, bottom trawl, coll. N. Nakayama and R. Misawa, 23 Apr. 2013; BSKU 110113 (1 male, 69.4 mm HL, 404+ mm TL), BSKU 110114 (1 male, 74.0 mm HL, 439 mm TL), BSKU 110115 (1 female, 57.4 mm HL, 336 mm TL), BSKU 110116 (1 male, 59.3 mm HL, 369 mm TL), BSKU 110117 (1 male, 77.0 mm HL, 457 mm TL), BSKU 110118 (1 male, 70.3 mm HL, 422+ mm TL), Suruga Bay, 34.7489°N, 138.4664°E, 200–450 m, F/V *Hinode-maru*, sta. 6, bottom trawl, coll. N. Nakayama and R. Misawa, 23 Apr. 2013; BSKU 110121 (1 female, 96.4 mm HL, 525 mm TL), BSKU 110122 (1 male, 59.7 mm HL, 378+ mm TL), BSKU 110123 (1 male, 63.5 mm HL, 381+ mm TL), Suruga Bay, 34.7127°N, 138.4553°E, 262–434 m, F/V *Hinode-maru*, sta. 1, bottom trawl, coll. N. Nakayama and R. Misawa, 23 Apr. 2013; FRLM 1780 (1 sex undetermined, 74.6 mm HL), 20 km southwest of Inohana, Kumano-nada, 900–1100 m, fish trap, 29 Nov. 1978.

Counts and measurements. See Table 9.

Size. Attains about 62 cm TL (CAS-SU 8547, holotype of *Coryphaenoides awae*, 622+ mm TL, west of southern Chiba Pref., Japan).

Variation. Serrations along the second spinous ray of the first dorsal fin are better developed in smaller specimens than in larger ones. Those of adults are somewhat rudimentary, giving a moderately smooth appearance to the leading edge.

Sexual dimorphism and dichromatism. Females

FIGURE 119. Fresh specimens of *Coryphaenoides marginatus*. (A–C) BSKU 103046, male, 68.1 mm HL, 413 mm TL; (D–F) BSKU 103048, female, 71.4 mm HL, 392 mm TL; (G) BSKU 103052, female, 105 mm HL, 541 mm TL. All specimens from Suruga Bay, Shizuoka Pref., Pacific. (A, D, G) Lateral views; (B, E) dorsal and (C, F) ventral views of the head and trunk. [Photos: N. Nakayama]

attain a much larger body size than males, with the largest female measuring 541 mm in TL (vs. 461+ mm in males). Compared with females, males have a narrower cheek (orbit–preopercle distance 37–41% HL vs. 40–44%), a wider interorbital (20–24% HL vs. 16–20%), and a larger posterior nostril (7–11% HL vs. 6–8%). Sexual dichromatism is also evident, with females having a much paler body than males (Fig. 119A–C vs. D–G). In females, the first dorsal and pelvic fins are mostly pale except for dark apical bands, whereas in males, those fins are uniformly blackish.

Development. Early life stage was fully described by Fukui *et al.* (2008), with a complete ontogenetic series of eggs and larvae.

Distribution. Known only from Japan and Korea at depths of 115–1165 m (Kim *et al.* 2009b; this study; Appendix 3–6F). Shao *et al.* (2008a, 2008b) reported this species from Taiwan, but their specimens are currently referred to as *C. microps* (Smith & Radcliffe in Radcliffe, 1912) (see Iwamoto *et al.* 2015). Abundant in southern Japan, but rare in the east of the Boso Peninsula (139.89°E).

Remarks. *Coryphaenoides marginatus* was originally described from a single specimen collected from Tokyo, Japan (probably originally from off Enoshima Island, Sagami Bay; Amaoka 2007:28). Although the holotype is thought to be in the Naturhistorisches Museum, Vienna (NMW; Fricke *et al.* 2020), it is now missing among that collection (K. Amaoka pers. com.). For a full description see Okamura (1970a).

Iwamoto in Randall & Lim (2000) listed *C. marginatus* from the South China Sea, but its occurrence

needs to be verified by voucher specimens. Although Shao *et al.* (2008a, 2008b) recorded *C. marginatus* from Taiwan, their specimens were re-identified as *C. microps* by Iwamoto *et al.* (2015). *Coryphaenoides marginatus* appears to be restricted to Japan and Korea.

Nomenclatural discussion. Jordan & Gilbert (in Jordan & Starks 1904) described *Coryphaenoides awae* as a new species based on a single specimen collected from off Nanaura, Awa, southern part of Chiba Pref., Japan (Fig. 120), suggesting its close relationship with *C. marginatus*. According to the original description, *C. awae* differs from *C. marginatus* in having a larger and broader head (HL 5.3 in TL vs. ca. 5.75; head width 1.83 in HL vs. 2.25), a deeper body (depth 5.3 in TL vs. ca. 7.5), a smaller orbit (4.33 in HL vs. 3.6), a shorter snout (3.4 in HL vs. 3.83), and more spinule rows on the body scales (13–17 vs. 12). Jordan & Gilbert (in Jordan & Starks 1904) further distinguished the two species by head squamation: in *C. awae*, the underside of the head is naked along margins of the interopercle and mandibular rami, and above the side of the upper lip, whereas the naked areas of *C. marginatus* are confined to inner margins of the mandibular rami. As far as I know, no additional specimens of *C. awae* have been recorded since the original description.

Okamura (1970a) was the first who recognized the two nominal species to be synonymous, although he did not examine the holotype of *C. awae*. All subsequent authors have followed his conclusion, and *C. awae* is currently regarded as a junior synonym of *C. marginatus* (e.g., Kuroda 1971; Iwamoto 1990; Kim *et al.* 2009b). To reconfirm this synonymy, the holotype of *C. awae* was examined in this study (sex was not determined to avoid

FIGURE 120. Preserved specimen of *Coryphaenoides marginatus*. CAS-SU 8547, holotype of *C. awae*, 117 mm HL, 622+ mm TL, west of southern Chiba Pref., Pacific. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

TABLE 9. Morphometric measurements and meristic counts* for *Coryphaenoides marginatus*.

Nominal species	<i>C. marginatus</i>			<i>C. awae</i>
Type status	NT			HT
Sex	Males	Females	Unknown	Unknown
Number of specimens	<i>n</i> = 22	<i>n</i> = 14	<i>n</i> = 8	<i>n</i> = 1
Measurements (mm)				
TL	230+–461+	237+–541	152+–421+	622
HL	38.4–85.5	42.6–105	26.9–74.6	117
% of HL (% of PRL)				
Snout length	29–32 (39–45)	27–30 (37–42)	28–32 (37–45)	28 (39)
Orbit diameter	26–31 (35–43)	22–28 (31–39)	26–30 (37–43)	22 (31)
Postorbital length	43–48 (60–66)	45–51 (62–70)	42–46 (60–64)	50 (69)
Postrostral length	70–73 (–)	71–74 (–)	71–75 (–)	73 (–)
Orbit–preopercle distance	37–41 (51–57)	40–44 (55–61)	38–43 (53–60)	43 (59)
Suborbital width	11–14 (15–20)	11–14 (15–20)	11–13 (15–18)	14 (20)
Upper-jaw length	28–32 (39–43)	30–32 (41–45)	30–32 (40–45)	31 (42)
Length of rictus	20–24 (28–33)	22–25 (30–34)	22–25 (30–36)	22 (30)
Length of PMX tooth band	16–21 (22–29)	18–21 (24–30)	18–21 (24–29)	20 (27)
Preoral length	15–19 (20–26)	14–17 (19–24)	15–18 (20–25)	11 (15)
Dist. tip and lat. angle of snout	14–19 (20–26)	13–17 (19–24)	15–20 (21–28)	12 (16)
Snout width	24–28 (34–39)	21–27 (29–37)	24–29 (33–41)	20 (28)
Internasal width	16–19 (22–26)	15–17 (20–24)	16–20 (22–28)	14 (20)
Interorbital width	20–24 (27–33)	16–20 (21–29)	17–22 (23–32)	19 (27)
Body width over P. bases	44–58 (60–79)	45–52 (62–73)	40–48 (56–65)	56 (77)
Body depth at 1D. origin	75–87 (103–124)	80–91 (109–125)	69–86 (97–120)	97 (133)
Body depth at A. origin	69–82 (95–113)	70–79 (96–110)	55–74 (77–103)	81 (111)
Pre-V. length	110–122 (153–170)	110–120 (155–165)	107–115 (148–161)	113 (155)
Pre-anus length	154–171 (214–238)	152–173 (211–242)	146–164 (205–230)	161 (221)
Pre-A. length	162–182 (226–253)	159–179 (219–250)	150–170 (211–238)	171 (234)
Isthmus–V. distance	41–48 (57–67)	40–47 (55–66)	39–45 (55–63)	46 (63)
Isthmus–anus distance	85–99 (119–141)	80–101 (111–142)	78–95 (109–133)	95 (130)
Isthmus–A. distance	94–111 (131–154)	87–109 (121–152)	82–101 (115–141)	105 (145)
V.–A. distance	52–67 (73–94)	48–65 (67–92)	45–58 (63–79)	63 (86)
Anus–A. distance	5–11 (7–15)	6–11 (8–15)	4–8 (6–11)	– (–)
V. length	52–73 (73–102)	50–65 (70–88)	50–85 (70–114)	– (–)
P. length	52–71 (72–98)	54–66 (75–90)	49–64 (67–87)	59 (82)
Pre-1D. length	121–129 (167–180)	118–129 (164–178)	118–127 (163–178)	129 (178)
Height of 1D.	101–191 (139–265)	119–199 (161–276)	127–219 (172–308)	– (–)
Length of 1D. base	27–36 (37–50)	27–32 (38–44)	26–31 (37–42)	30 (41)
Interdorsal length	11–51 (14–70)	11–58 (15–81)	10–44 (13–61)	18 (25)
Length of gill slit	3–7 (5–10)	5–8 (6–11)	4–9 (6–12)	5 (8)
Length of posterior nostril	7–11 (9–15)	6–8 (8–10)	6–11 (8–15)	8 (10)
Barbel length	7–12 (10–17)	7–13 (10–19)	5–14 (8–19)	12 (16)
Counts				
1D. rays	II,9–11	II,9–11	II,9–11	II,9
P. rays	i18–i23	i19–i25	i16–i23	i21
V. rays	6–8	7–8	7–9	7–8
Outer GR on first arch	3–5	3–5	3–5	2
Inner GR on first arch	9–11	10–11	9–11	10
Outer GR on second arch	9–11	10–11	9–12	10
Inner GR on second arch	8–11	9–10	8–10	8
Longitudinal scales	32–43	36–43	34–42	39
Scales below 1D. origin	7.5–10.5	7.5–10.5	7.5–10.5	9.5
Scales below 1D. midbase	5.5–8.5	5.5–8.5	5.5–8.5	6.5
Scales below 2D. origin	6.5–10	7–9.5	7–9.5	8.5
Scales above A. origin	18.5–25.5	18.5–26.5	20.5–24.5	–

...See the next page for the table footer

*Abbreviations: 1D.—first dorsal fin; 2D.—second dorsal fin; A.—anal fin; GR—gill rakers; HT—holotype; NT—non-type; P.—pectoral fin; PMX—premaxillary; V.—pelvic fin

further damage to the specimen), and it was compared with additional materials of *C. marginatus*.

The holotype of *C. awae* (117 mm HL, 622 mm TL) is distinctly larger than the *C. marginatus* specimens examined (26.9–105 mm HL, 152–541 mm TL), and this made a comparison of morphometric characters dubious. However, despite their size differences, data of *C. awae* generally lie within range of variation of *C. marginatus* (Table 9). The holotype of *C. awae* differs from the specimens of *C. marginatus* examined in having a shorter preoral (11% HL vs. 14–19%), a narrower internasal (14% vs. 15–20%), and a deeper body (depth below first dorsal-fin origin 97% HL vs. 69–91%). However, the above differences are attributable to size-related variation, considering allometric growth of these characters in *C. marginatus* (Fig. 121). Also, no significant differences were found in meristic characters between the two species, except for outer gill rakers on the first arch (2 in *C. awae* vs. 3–5 in *C. marginatus*). However, the subtle difference in gill rakers seems to reflect ontogenetic variation, and the count range for *C. marginatus* may extend to two when larger specimens become available for study. In large individuals of *C. marginatus* (especially females), the underside of the head is broadly naked above the upper lip and along the margins of the interopercle, preopercle, and mandibular rami. This condition agrees well with the description of *C. awae* given by Jordan & Gilbert (*in* Jordan & Starks 1904), and thus the two species cannot be distinguished by head squamation. Due to the lack of adequate comparative materials, the possibility of there being two species cannot be fully discussed in this study. However, there is little reason to doubt their conspecificity, and *C. awae* is therefore regarded here as a junior synonym of *C. marginatus*.

Relationships. *Coryphaenoides marginatus* belongs to the subgenus *Coryphaenoides* (*sensu* Iwamoto 1990). It falls into a group of species that Gilbert & Hubbs (1920:413) defined by the combination of “a produced dorsal spine, a deep and sharply compressed body, and a dorsal contour horizontal behind the first dorsal fin”. According to Shcherbachev & Iwamoto (1995:300), this group is further characterized by having “a relatively small, restricted mouth opening; low, slightly protruding snout; small teeth in bands lacking a prominently enlarged series; small barbel; and highly restricted gill openings”. Other members of the group are: *C. macrolophus* (Alcock, 1889), *C. microps*, *C. semiscaber* Gilbert & Hubbs, 1920, and *C. tydemani* (Weber, 1913). Shcherbachev & Iwamoto (1995) provided comparisons of these species except *C. marginatus*. This group is called here the *C. marginatus* group.

FIGURE 121. Scatter diagrams showing relationships of the (A) preoral length, (B) internasal width, and (C) body depth at the first dorsal-fin origin (as % HL) to HL (mm) in *Coryphaenoides marginatus*. Orange squares indicate the holotype of *C. awae*.

Comparisons. *Coryphaenoides marginatus* is readily distinguished from other members of the *C. marginatus* group by spinulation on the body scales. In *C. marginatus*, spinules on the body scales are distinctly short, with the last spinule in each spinule row barely overlapping the posterior scale margin (Fig. 122A–B); the spinules of the other four species are much longer, and the tips of the last spinules extend well beyond that level (Shcherbachev & Iwamoto 1995; this study; see

also Fig. 122C–D). It further differs from all in having a broad naked band above the upper lip (vs. ventral surfaces of snout and suborbital shelf completely scaled or only narrowly naked; Shcherbachev & Iwamoto 1995; this study). *Coryphaenoides marginatus* also differs from *C. macrolophus* in that the preopercle is broadly rounded posteroventrally (vs. sharply angulated at posteroventral corner; Shcherbachev & Iwamoto 1995:303); from *C. microps* in having a larger orbit (22–31% HL vs. 19–22%) and a shorter postorbital (42–51% HL vs. 50–55%), and in that the mandibular rami are only narrowly scaled medially (vs. almost completely scaled, except for anterior portions) and the preopercle is naked along its ventral margin (vs. almost completely scaled); from *C. semiscaber* in having a shorter pelvic fin (50–85% HL vs. 95–96%; Shcherbachev & Iwamoto 1995:299, fig. 4); and from *C. tydemani* in having a longer isthmus–pelvic distance (39–48% HL vs. 31–37%) and a lower first dorsal fin (101–219% HL vs. 238–276%).

In Japanese waters, *C. marginatus* is collected with *C. nasutus* Günther, 1877, with which it might be confused in the field. However, the two species are readily distinguished from each other by the positions of the pelvic fins. In *C. marginatus*, the pelvic fins are inserted closer together and the interpelvic distance is less than the base length of each fin (Figs. 119C, F, 120C), whereas in *C. nasutus*, the fins are well separated by a broad gap that is greater than their base length (Figs. 125C, 126C). The two species are further distinguished by squamation characters. *Coryphaenoides marginatus* differs from *C. nasutus* in that spinules on the body scales are arranged in tightly packed parallel rows (Fig. 122A vs. convergent rows, Fig. 122E). The top of the snout is fully scaled in *C. marginatus*, whereas that of *C. nasutus* is narrowly naked along the leading edges. Serrations of the first dorsal fin are fewer and less developed in *C. marginatus* than in *C. nasutus* (number of serrations 0–13 vs. 20–35). Females of *C. marginatus* differ notably from *C. nasutus* in that the distal margins of the first dorsal and pelvic fins are prominently outlined in black (vs. fins uniformly dark in *C. nasutus*). The two species are difficult to distinguish by morphometric characters, although *C. marginatus* tends to have a shorter interdorsal space (10–58% HL vs. 34–114%) and a higher first dorsal fin (101–219% HL vs. 78–125%). Regarding meristic characters, *C. marginatus* differs from *C. nasutus* in modal counts of pelvic-fin rays [8 (range 6–9), vs. 9 (8–10)] and gill rakers on the outer side of the second arch [10 (9–12) vs. 9 (8–10)].

***Coryphaenoides microps* (Smith & Radcliffe in Radcliffe, 1912)**

[No Japanese name]

(Figs. 122C–D, 123–124; Appendix 3-7A)

Macrourus microps Smith & Radcliffe in Radcliffe, 1912:116, pl. 25, fig. 2 [original description; holotype: USNM 72933, from Atulayan Island, eastern coast of Luzon Island, *Albatross* sta. 5470, 13°37'30"N, 123°41'09"E, in 560 ftm (1024 m); 4 paratypes from northern Luzon].

Coryphaenoides marginatus (not Steindachner & Döderlein 1887): Shao 1993:171 (compiled; Taiwan); Shao *et al.* 2008b: table 2 (36 spec. listed from northeastern, southeastern and southwestern Taiwan, and South China Sea).

Coryphaenoides microps: Chiou *et al.* 2004a: table 1 (listed; Taiwan); Shao *et al.* 2008b: table 2 [1 spec. listed from southwestern Taiwan (as first Taiwanese record)]; Kim *et al.* 2009b:108, fig. 3 [brief description; 14 spec. from Korea, Taiwan, and Philippines (including holotype); new Korean record from Korea]; Iwamoto *et al.* 2015:70 (brief description; 53 spec. from northeastern and southwestern Taiwan, and Philippines).

Diagnosis. Pelvic-fin rays 7–8. Snout short, slightly protruding beyond upper jaw. Tip and lateral angles of snout armed with stout tubercles; scales along head ridges not coarsely modified. Mouth small, posterior margin of upper jaw not reaching vertical through midorbit; upper-jaw length 30–31% HL; lateral corner of mouth moderately restricted by skin folds. Outermost gill slit greatly restricted, length 4–7% HL. Barbel short, slender, length 5–10% HL. Teeth all small, conical, slender in wide bands in both jaws, none especially enlarged. Body scales covered with short, greatly reclined, needle-like spinules in tightly packed subparallel rows; tip of last spinule in each row extending distinctly beyond posterior scale margin. Transverse scale rows. Top of snout fully scaled; underside of snout narrowly naked above upper lip. Second spinous ray of first dorsal fin notably prolonged, height of first dorsal fin distinctly greater than HL (198–300% HL); serration along its leading edge rudimentary in adults; first dorsal-fin rays. Pelvic-fin bases closer together, their interspace less than base length of each fin. Body dusky brown overall.

Material examined. 5 specimens. **Taiwan:** ASIZP 66413 (1, 43.7 mm HL, 241+ mm TL), Dong-gang fish market, Pingtung, bottom trawl, coll. P.-F. Lee, 7 Oct. 2005; ASIZP 63792 (1, 78.5 mm HL, 367+ mm TL), off Kaohsiung, 22.3667°N, 120.1000°E, 821 m, otter trawl, coll. H.-M. Yeh, 29 Aug. 2002; ASIZP 66099 (1, 52.4 mm HL, 298 mm TL), ASIZP 66109 (1, 80.4 mm HL, 324+ mm TL), off Kaohsiung, 22.3333°N, 120.1000°E, 687 m, ORE-type beam trawl, coll. P.-F. Lee, 11 Aug. 2005; *FAKU 204442 (1, 37.0 mm HL, 201+ mm TL), Dong-gang fish market, Pingtung, bottom trawl, coll. N. Nakayama, 27 Feb. 2017.

Counts and measurements. Based on 4 specimens (43.7–80.4 mm HL, 241+–367+ mm TL). Counts: first dorsal-fin rays II,9–10; pectoral-fin rays i19–i21; pelvic-fin rays 7–8; gill rakers on first arch (outer/inner) 4–5/9–10, on second arch 9–10/8–9; longitudinal scales 33–41;

FIGURE 122. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of three species of the *Coryphaenoides marginatus* group. (A–B) *C. marginatus*, BSKU 103048, 71.4 mm HL; (C–D) *C. microps*, ASIZP 66413, 43.7 mm HL; (E–F) *C. nasutus*, BSKU 103042, 78.5 mm HL. (A, C, E) Views from above; (B, D, F) oblique views. [Photos: NSMT]

transverse scale rows below first dorsal-fin origin 10–11, below first dorsal-fin midbase 7.5–9, below second dorsal-fin origin 7–8.5, above anal-fin origin 25–28.

The following measurements are in % of HL,

followed by those in % of PRL in parentheses: snout length 28–30 (38–42); orbit diameter 19–22 (26–31); postorbital length 50–55 (70–74); postrostral length 71–74; orbit–preopercle distance 41–46 (57–63); suborbital

width 12–14 (17–20); upper-jaw length 30–31 (42–43); length of rictus 21–23 (29–31); length of premaxillary tooth band 18–20 (26–27); preoral length 11–14 (16–20); distance between tip and lateral angle of snout 12–16 (17–22); snout width 21–25 (28–35); internasal width 15–17 (21–23); interorbital width 18–21 (24–29); body width over pectoral-fin bases 44–58 (62–81); body depth at first dorsal-fin origin 78–86 (109–121); body depth at anal-fin origin 70–83 (98–116); prepelvic length 118–134 (165–182); preanus length 172–179 (241–244); preanal length 177–187 (248–254); isthmus–pelvic distance 43–46 (60–62); isthmus–anus distance 89–101 (124–141); isthmus–anal distance 94–107 (131–150); pelvic–anal distance 55–65 (77–91); pelvic-fin length 53–62 (75–84); pectoral-fin length 64–68 (90–92); predorsal length 117–124 (165–174); height of first dorsal fin 198–300 (269–421); length of first dorsal-fin base 27–29 (37–41); interdorsal length 12–24 (17–33); length of gill slit 4–7 (5–9); length of posterior nostril 6–8 (9–11); barbel length 5–10 (7–14).

Size. Attains at least 46 cm TL (ASIZP 66111, 464+ mm TL, South China Sea; see Iwamoto *et al.* 2015:70).

Distribution. Known from the Philippines, Taiwan, and Korea, at depths of 115–1165 m (Radcliffe 1912; Shcherbachev & Iwamoto 1995; Shao *et al.* 2008a, 2008b). No specimens have been collected from Japanese waters (Appendix 3-7A). Common in Taiwan.

Remarks. For a full description see Shcherbachev & Iwamoto (1995). *Coryphaenoides microps* appears to be a sister species to *C. marginatus* Steindachner & Döderlein, 1887 known from Japan and Korea. The two species are parapatric, with their distributions only slightly overlapping in the Sea of Japan off Korea (Appendix 3-6F and 3-7A).

Relationships and comparisons. *Coryphaenoides microps* belongs to the *C. marginatus* group (see the Relationships of *C. marginatus*), and is most similar to *C. marginatus*. However, *C. microps* readily differs from *C. marginatus* in that the spinules on the body scales are arranged in tightly packed, narrowly convergent rows (vs. tightly packed parallel rows), with the tip of the last spinule in each row extending distinctly beyond the posterior scale margin (vs. scarcely reaching this level) (Fig. 122C–D vs. A–B). *Coryphaenoides microps* further differs from that species in having a smaller orbit (19–22% HL vs. 22–31%) and a longer postorbital (50–55% HL vs. 42–51%). In the four *C. microps* specimens examined, the underside of the head was almost fully scaled along the infraorbital, preopercular, and mandibular canals, and naked areas were apparent only on anterior portions of the mandibular rami and a narrow band above the upper lip. In contrast, the *C. marginatus* specimens examined had broad naked areas above the upper lip and along margins of the interopercle, preopercle, and mandibular rami. *Coryphaenoides microps* is unlikely to be confused with females of *C. marginatus* by having a uniformly dark

body (vs. paler ventrally) and blackish fins (vs. first dorsal and pelvic fins pale except for dark distal margins).

Coryphaenoides nasutus Günther, 1877

[Japanese name: Hana-sokodara]

(Figs. 122E–F, 125–126; Appendix 3-7B)

Coryphaenoides nasutus Günther, 1877:440 [original description; lectotype (designated herein): BMNH 1887.12.7.78, from “south of Yeddo” (off Cape Omaezaki), *Challenger* sta. 235, in 565 fth (1034 m); 1 paralectotype from type locality]; Gilbert & Hubbs 1916:168 (brief description; 45 spec. from Hyuga-nada, off Cape Shionomisaki, Enshu-nada, southwest of Hokkaido, south of Izu Peninsula, and Sagami Bay); Jordan & Hubbs 1925:327 (spec. from Misaki); Okada & Matsubara 1938:449 (in key; Japan); Kamohara 1950:279 (listed; Kochi Pref.); Kamohara 1952:98 (selected counts and measurements; 1 spec. from Kochi Pref.); Matsubara 1955:1310 (in key; Japan; Japanese name: “Hana-sokodara”); Kamohara 1958:73 (listed; Kochi Pref.); Kamohara 1964:95 (listed; Kochi Pref.); Ueno 1965:19 (listed; Pacific off Hokkaido); Wheeler & O’Riordan 1969:96 (comments on type specimens); Okamura 1970a:140, pl. XXXI, text-fig. 56 (description; biological notes; 38 spec. from Pacific from off Hokkaido to Mimase and East China Sea); Ueno 1971:101 (listed; Hokkaido); Kuroda 1971:56 [listed from Suruga Bay after Okamura (1970a)]; Tominaga & Uyeno 1981:489 (listed; Japan); Okamura 1982:165, 351, fig. 99 [brief description; 8 spec. from Tosa Bay; photo based on BSKU spec. (catalog no. unknown)]; Shiogaki 1982:33 (listed; Aomori Pref.); Sawada 1983:113, 195, fig. 64 (brief description; 5 spec. from Pacific off Tohoku; photo based on HUMZ spec. (catalog no. unknown)); Ohta 1983: photo 28, tables A–B (listed; Suruga Bay; in situ observation); Yatou 1984:219, 364, fig. 155 [brief description; 10 spec. from Okinawa Trough; photo based on BSKU spec. (catalog no. unknown)]; Okamura 1984b:95, pl. 82, fig. F (compiled); Okamura 1988:95, pl. 82, fig. F (compiled); Iwamoto 1990:216, fig. 496 (synopsis); Maeda & Maruyama 1991:366 (listed; Hokkaido); Nakabo 1993:365 (in key; Japan); Amaoka *et al.* 1995:103, fig. 153 (compiled; northern Japan); Shinohara *et al.* 1996:169 (2 spec. listed from Pacific off Tohoku); Okamura 1997:126, fig. 3 (compiled); Shinohara & Matsuura 1997:291 (listed; Suruga Bay); Nakabo 2000:429 (in key; Japan); Shinohara *et al.* 2001:305 (5 spec. listed from Tosa Bay); Nakabo 2002:429 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Maeda & Tsutsui 2003:488 (listed; Hokkaido); Shiogaki *et al.* 2004:55 (listed; Pacific off Aomori Pref.); Shinohara *et al.* 2005:417 (listed; Ryukyu Islands); Suetsugu & Ohta 2005: table 3 (listed; Enshu-nada); Senou *et al.* 2006:420 (listed; Sagami Sea); Kitagawa *et al.* 2008:41, unnumbered fig. (brief description; spec. from Pacific off Tohoku); Shao *et al.* 2008b: table 2 (5 spec. listed from northeastern and southeastern Taiwan; first Taiwanese record); Matsuura *et al.* 2009:24 (1 spec. from Miyagi Pref.); Shinohara *et al.* 2009:708 (2 spec.

FIGURE 123. Fresh specimen of *Coryphaenoides microps*. FAKU 204442, 37.0 mm HL, 201+ mm TL, Dong-gang fish market, southwestern Taiwan, South China Sea. Lateral view. [Photo: N. Nakayama]

FIGURE 124. Preserved specimen of *Coryphaenoides microps*. ASIZP 63792, 78.5 mm HL, 367+ mm TL, off Kaohsiung, northern Taiwan, South China Sea, 821 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

listed from Pacific off Tohoku); Amaoka *et al.* 2011:130, fig. 165 (compiled; Hokkaido); Nakabo & Kai 2013:505 (in key; Japan); Ikeda & Nakabo 2015:320, pl. 65, figs. 9–13 (brief description; 3 spec. from Kii Channel off Wakayama Pref.); Iwamoto *et al.* 2015:71, fig. 10 (brief description; 9 spec. from northeastern and southeastern Taiwan, South China Sea, and Japan); Amaoka *et al.* 2020:163, fig. 221 (listed; Hokkaido); Motomura 2020:39 (listed; Japan).

Macrurus nasutus: Günther 1887:132, pl. XXX, fig. B (description; 11 spec. from Enshu-nada and Sagami Bay).

[?] *Macrurus altipinnis*: Günther 1887:138 (description; in part, 1 spec., BMNH 1887.12.7.91; Enshu-nada).

Macrourus nasutus: Jordan & Snyder 1901:120 (listed; Japan);

Jordan & Gilbert in Jordan & Starks 1904:616 (brief description; 16 spec. from Sagami Bay and “off the shore of Izu”); Franz 1910:26 (5 spec. from Aburatsubo); Jordan *et al.* 1913:417 (listed; Japan; new Japanese name: “Hana-naga-sokodara”); Jordan & Thompson 1914:306 [selected counts; 3 spec. from Misaki (FMNH 57384, 2 spec.) and Sagami Bay]; Kamohara 1938:72 (selected counts and measurements; 1 spec. from Kochi Pref.).

Coryphaenoides (Coryphaenoides) nasutus: Okamura 1970b: table 1 (listed; Japan).

Diagnosis. Pelvic-fin rays 8–10 (usually 9). Snout conical, moderately protruding beyond upper jaw. Tip

and lateral angles of snout armed with prominent enlarged tubercles; scales along head ridges not coarsely modified. Mouth small, posterior margin of upper jaw not extending beyond vertical through midorbit; upper-jaw length 28–32% HL; lateral corner of mouth moderately restricted by skin folds. Outermost gill slit greatly restricted, length 4–8% HL. Barbel short, slender, length 5–8% HL. Teeth all small, conical, in wide bands in both jaws, none especially enlarged. Body scales covered with short, greatly reclined, needle-like spinules in tightly packed convergent rows; tip of last spinule in each row scarcely extending beyond posterior scale margin. Transverse scale rows below first dorsal-fin midbase 4.5–7.5. Dorsal and ventral surfaces of snout broadly naked along leading edges. Height of first dorsal fin 78–125% HL; second spinous ray moderately prolonged, finely serrated along its leading edge; first dorsal-fin rays II,9–12. Pelvic fin well separated, interspace greater than base length of each fin. Body dusky brown overall.

Material examined. 32 specimens. **Lectotype of *Coryphaenoides nasutus*** (designated herein): BMNH 1887.12.7.78 (81.2 mm HL, 337+ mm TL), off Cape Omaezaki, Enshu-nada, Japan, 34.1167°N, 138.0000°E, 565 ftm (1034 m), *Challenger* sta. 235, trawl, 4 Jun. 1875. **Paralectotype of *C. nasutus*** (designated herein): BMNH 1887.12.7.79 (1, 72.2 mm HL, 365 mm TL), collected with lectotype. **Non-types: Japan:** BSKU 26668 (1, 97.8 mm HL, 429+ mm TL), west of Yaku-shima Island, Okinawa Trough, 30.0300°N, 128.3650°E, 900 m, F/V *Yuryo-maru*, No. 8, tr. 16-1, bottom trawl, coll. Y. Kinoshita and S. Hagino, 3 Feb. 1978; BSKU 28493 (1, 95.6 mm HL, 455+ mm TL), BSKU 28495 (1, 88.7 mm HL, 437+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.7333°N, 128.0500°E, 815 m, F/V *Ryoan-maru*, No. 28, tr. 74, bottom trawl, coll. Y. Kinoshita and S. Hagino, 20 Mar. 1978; BSKU 28076 (1, 105 mm HL, 477+ mm TL), BSKU 28079 (1, 98.0 mm HL, 478+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 28.8833°N, 127.3000°E, 820–830 m, F/V *Ryoan-maru*, No. 28, tr. 60, bottom trawl, coll. Y. Kinoshita and S. Hagino, 16 Mar. 1978; BSKU 47158 (1, 73.9 mm HL, 361+ mm TL), BSKU 47159 (1, 71.8 mm HL, 358+ mm TL), BSKU 47163 (1, 64.7 mm HL, 335+ mm TL), BSKU 47164 (1, 75.3 mm HL, 376+ mm TL), off Tosa Bay, 32.8953°N, 133.6777°E, 996–1010 m, R/V *Tansei-maru*, cr. KT-89-06, sta. T1, beam trawl, 8 Nov. 1989; BSKU 95207 (1, 63.0 mm HL, 237+ mm TL), BSKU 95213 (1, 61.5 mm HL, 315+ mm TL), BSKU 95215 (1, 58.1 mm HL, 279+ mm TL), BSKU 95219 (1, 60.1 mm HL, 325+ mm TL), BSKU 95221 (1, 53.6 mm HL, 240+ mm TL), BSKU 95226 (1, 34.6 mm HL, 185+ mm TL), Tosa Bay, 750 m, FRV *Kotaka-maru*, otter trawl, 13 Sept. 1988; BSKU 43456 (1, 67.2 mm HL, 351+ mm TL), south-southwest of Okino-shima Island, 32.5450°N, 132.4750°E, 1075–1092 m, R/V *Tansei-maru*, cr. KT-86-16, sta. C, 3-m beam trawl, 2 Nov. 1986; BSKU 103041 (1, 64.0 mm HL, 337+

mm TL), BSKU 103042 (1, 78.5 mm HL, 383+ mm TL), Suruga Bay, coll. A. Fukui and M. Takami, 10 Sept. 2007; BSKU 103030 (1, 69.3 mm HL, 342+ mm TL), Suruga Bay, coll. A. Fukui and M. Takami, 14 Nov. 2007; BSKU 103031 (1, 69.9 mm HL, 329+ mm TL), Suruga Bay, coll. A. Fukui and M. Takami, 3 Dec. 2008; BSKU 97868 (1, 56.9 mm HL, 286+ mm TL), BSKU 97872 (1, 82.3 mm HL, 422+ mm TL), BSKU 97873 (1, 55.2 mm HL, 288+ mm TL), BSKU 97887 (1, 74.9 mm HL, 381 mm TL), BSKU 97888 (1, 88.2 mm HL, 419+ mm TL), BSKU 97890 (1, 70.4 mm HL, 361+ mm TL), BSKU 97893 (1, 77.2 mm HL, 373+ mm TL), BSKU 97900 (1, 46.8 mm HL, 242+ mm TL), BSKU 97901 (1, 48.2 mm HL, 209+ mm TL), BSKU 97903 (1, 41.5 mm HL, 212+ mm TL), off Kamaishi, 39.1444°N, 142.3748°E, 847–935 m, R/V *Tansei-maru*, cr. KT-89-3, sta. SR89, 3-m beam trawl, 10 Mar. 1989.

Counts and measurements. Based on 32 specimens (34.6–105 mm HL, 185+–478+ mm TL). Counts: first dorsal-fin rays II,9–12; pectoral-fin rays i19–i24; pelvic-fin rays 8–10; gill rakers on first arch (outer/inner) 3–5/8–11, on second arch 8–10/8–9; longitudinal scales 37–45; transverse scale rows below first dorsal-fin origin 6.5–9.5, below first dorsal-fin midbase 4.5–7.5, below second dorsal-fin origin 6.5–9, above anal-fin origin 19.5–27.5.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 28–33 (38–47); orbit diameter 25–31 (34–42); postorbital length 41–49 (57–67); postrostral length 69–75; orbit–preopercle distance 37–44 (53–60); suborbital width 11–15 (16–21); upper-jaw length 28–32 (39–43); length of rictus 21–29 (29–38); length of premaxillary tooth band 17–22 (24–30); preoral length 14–22 (20–31); distance between tip and lateral angle of snout 16–21 (21–30); snout width 26–32 (35–46); internasal width 18–22 (24–30); interorbital width 17–25 (24–34); body width over pectoral-fin bases 40–63 (57–90); body depth at first dorsal-fin origin 65–86 (91–122); body depth at anal-fin origin 53–76 (74–109); prepelvic length 103–125 (141–178); preanus length 142–168 (196–240); preanal length 147–179 (206–249); isthmus–pelvic distance 34–45 (47–61); isthmus–anus distance 70–94 (97–134); isthmus–anal distance 63–104 (88–145); pelvic–anal distance 41–68 (57–97); anus–anal distance 4–13 (6–18); pelvic-fin length 40–83 (57–117); pectoral-fin length 45–56 (62–80); predorsal length 110–122 (154–174); height of first dorsal fin 78–125 (104–175); length of first dorsal-fin base 23–32 (33–44); interdorsal length 34–114 (48–159); length of gill slit 4–8 (6–12); length of posterior nostril 5–10 (7–14); barbel length 5–8 (6–11).

Size. Attains about 52 cm TL (FAKU 102252, 524 mm TL, off Fukushima Pref., Japan).

Distribution. Restricted to Japan and Taiwan. Known from the Pacific off Japan northward to Muroran (42.32°, Hokkaido), Okinawa Trough, and northern South China Sea, at depths of 450–2056 m (Shao *et al.* 2008a, 2008b;

FIGURE 125. Fresh specimen of *Coryphaenoides nasutus*. BSKU 103042, 78.5 mm HL, 383+ mm TL, Suruga Bay, Shizuoka Pref., Pacific. (A) Lateral view; (B) dorsal and (C) ventral view of head and trunk. [Photos: N. Nakayama]

FIGURE 126. Lectotype of *Coryphaenoides nasutus*, BMNH 1887.12.7.78, 81.2 mm HL, 337+ mm TL, off Cape Omaezaki, Enshu-nada, Pacific, 1034 m depth. (A) Lateral view; (B) dorsal and (C) ventral view of head and trunk. Preserved condition. [Photos: N. Nakayama]

this study; Appendix 3-7B). Abundant on the upper slope of the Pacific off southern Japan.

Comments on type specimens. *Coryphaenoides nasutus* was originally described by Günther (1877) from the “south of Yeddo” (= Tokyo), Japan. Although

Günther (1877) did not indicate how many specimens were involved in the original description, he (1887:132) subsequently noted that five specimens were collected from the above locality (*Challenger* sta. 235). Two syntypes (BMNH 1887.12.7.78–79, 337+–365 mm TL; Fig. 126)

were examined in this study, whereas the remaining three are presumed to be lost. Günther (1887) also reported six additional specimens from “off Inoshima” (= Eno-shima Island), Sagami Bay, Japan (*Challenger* sta. 232), but these specimens do not have any type status. While Bailly *et al.* (1999) considered MNHN 1890-123 (from *Challenger* sta. 232) as a syntype of this species, this specimen is also obviously a non-type. One of the syntypes examined (BMNH 1887.12.7.78; Fig. 126) is herein designated as the lectotype of *C. nasutus*, and thus the other specimens used in the original description automatically become paralectotypes of the species.

Remarks. For a full description see Okamura (1970a). Although Okamura (1982:165, 351) described the number of branchiostegal rays as seven, this is likely a mistype of six. Among the FAKU collection is a single specimen of *C. nasutus* from off Minamidaito-jima Island (FAKU 13367, 1 spec., 29.2 mm HL, 137+ mm TL). However, the register of FAKU indicates its locality with a question mark. This record was excluded from the distribution of the species in this study. *Coryphaenoides nasutus* appears to be restricted to the upper continental slope along Japan and Taiwan, and its occurrence on the Philippine Plate is doubtful.

Relationships and comparisons. *Coryphaenoides nasutus* belongs to the subgenus *Coryphaenoides* (sensu Iwamoto 1990), and closely resembles *C. hoskynii* (Alcock, 1890) known from the Indian Ocean. However, the two species are readily distinguished by spinulation of the body scales: in *C. nasutus*, the tip of the last spinule in each row barely overlaps the posterior scale margin (Fig. 122E–F), whereas those of *C. hoskynii* extend well beyond this level. *Coryphaenoides nasutus* further differs from *C. hoskynii* in having a narrower snout (internasal width 18–22% HL vs. 24–33%) and much smaller scales (longitudinal scales 37–45 vs. 31–34, scale rows below the first dorsal-fin midbase 4.5–7.5 vs. 3.5–4.5, below the second dorsal-fin origin 6.5–9 vs. 4–5). [Data for *C. hoskynii* are from Shcherbachev & Iwamoto (1995).]

Among Japanese congeners, *C. nasutus* is most similar to *C. marginatus* Steindachner & Döderlein, 1887 in general appearance, but the two species are clearly separable by the positions of the pelvic fins, body-scale spinulation, head squamation, and fin color (see the Comparisons of *C. marginatus*).

***Coryphaenoides pectoralis* (Gilbert, 1891)**

[Japanese name: Mune-dara]

(Figs. 127–128; Appendix 3-7C)

Macrurus (Malacocephalus) pectoralis Gilbert, 1891:563 [original description; 2 syntypes: USNM 47235 and 47236, from off Washington, in 685–877 ftn (1253–1604 m)].

Macrurus (Nematonurus) magnus Gill & Townsend, 1897:234 (original description; holotype: USNM 48770, from southwest

of Pribilof Islands, Bering Sea).

Ateleobranchium pterotum Gilbert & Burke, 1912:94, fig. 37 [original description; holotype: USNM 74398, from off Avatcha Bay, east coast of Kamchatka, *Albatross* sta. 4798, 52°37.5'N, 158°50.0'E, in 300 ftn (549 m)].

Nematonurus pectoralis: Gilbert & Hubbs 1916:161 (new combination; first record from Japan; 3 spec. from Okhotsk Sea and south of Hokkaido; comparison with Alaskan population); Okada & Matsubara 1938:449 (in key; Japan; Japanese name: “Mune-dara”); Matsubara 1955:1308 (in key; Japan); Ueno 1965:19 (listed; Hokkaido); Matsubara 1965:504 (compiled; Japan); Maruyama 1971:33 (listed; Iwate Pref.); Hikita 1981:78, fig. 1 (spec. from Okhotsk Sea off Shiretoko Peninsula; erroneously noted Japanese name as “Himo-dara”; photo spec. represents *C. pectoralis*).

Coryphaenoides pectoralis: Taranetz 1937:169 (new combination; listed; Soviet Far East); Lindberg 1959 (listed; southern Okhotsk Sea); Okamura 1970a:118, pl. XXV, text-fig. 49 (description; 14 spec. from southeast of Hokkaido, Pacific off Hachinohe, and Kominato); Ueno 1971:101 (listed; Hokkaido); Tominaga & Uyeno 1981:489 (listed; Japan); Shiogaki 1982:33 (listed; Aomori Pref.); Sawada 1983:109, 193, 251, 325, figs. 60, 140 [brief description; 7 spec. from Pacific off Tohoku and Okhotsk Sea; photos based on HUMZ 78178 and HUMZ 78604 (not 78064)]; Amaoka & Nakaya 1983:44, fig. 60 (compiled; northern Japan); Okamura 1984b:95, pl. 82, fig. A (compiled); Okamura 1988:95, pl. 82, fig. A (compiled); Maeda & Maruyama 1991:366 (listed; Hokkaido); Nakabo 1993:362 (in key; Japan); Hori 1996:28 (listed; Ibaraki Pref.); Okamura 1997:125, fig. 4 (compiled); Funabashi 1998:85 (listed; Ibaraki Pref.); Nakabo 2000:426 (in key; Japan); Zama 2001:39 (listed; Miyagi Pref.); Nakabo 2002:426 (in key; Japan); Shiogaki *et al.* 2004:55 (listed; Pacific off Aomori Pref.); Senou *et al.* 2006:420 (listed; Sagami Sea); Kitagawa *et al.* 2008:40, unnumbered fig. (brief description; spec. from Pacific off Tohoku); Balanov *et al.* 2009:672 (14 spec. listed from Pacific off Tohoku; midwater captures); Shinohara *et al.* 2009:708 (4 spec. listed from Pacific off Tohoku); Endo *et al.* 2010:273, figs. 2–5 (larval description; 1 spec. from northwestern Pacific off Hokkaido); Amaoka *et al.* 2011:128, fig. 160 (compiled; Hokkaido); Nakabo & Kai 2013:502 (in key; Japan); Amaoka *et al.* 2020:160, fig. 214 (listed; Hokkaido); Motomura 2020:39 (listed; Japan).

Coryphaenoides (Nematonurus) pectoralis: Okamura 1970b: table 1 (listed; Japan).

Ateleobranchium [sic] *pterotum*: Ueno 1971:101 (listed; northern Kurile Islands; new Japanese name: “Nodoguro-hige”).

Albatrossia pectoralis: Iwamoto 1990:110, fig. 236 (synopsis); Endo *et al.* 1993:219, fig. 2 (larval description; 4 spec. from North Pacific); Amaoka *et al.* 1995:101, fig. 149 (compiled; northern Japan); Shinohara *et al.* 1996:168 (2 spec. listed from Pacific off Tohoku); Maeda & Tsutsui 2003:488 (listed; Hokkaido); Motomura 2020:39 (listed; Japan).

Diagnosis. Pelvic-fin rays usually 7 (rarely 5–6 or 8).

Snout conical, moderately protruding beyond upper jaw. Tip and lateral angles of snout lacking prominent tubercles; scales along head ridges not coarsely modified. Mouth large, posterior margin of upper jaw extending to vertical through hind rim or orbit or beyond; upper-jaw length 40–44% HL; lateral corner of mouth not restricted by skin folds. Outermost gill slit moderately wide, length 16–22% HL. Barbel short, slender, length 5–8% HL. Teeth prominent, conical, slender, in 2 irregular series in premaxillary, 1 distinct row in dentary. Body scales covered with short, reclined, needle-like spinules in widely divergent rows; tip of last spinule in each row scarcely extending beyond posterior scale margin (except in small specimens). Transverse scale rows below first dorsal-fin midbase 9.5–12.5. Top of snout fully scaled; ventral surfaces of head almost completely scaled except for underside of snout. Interdorsal space less than first dorsal-fin base length. First dorsal fin small, low, its height 43–50% HL; second spinous ray weakly serrated along its leading edge; first dorsal-fin rays II,8–10. Pelvic fin small, not reaching anal-fin origin when depressed. Body pale to dark brown.

Material examined. 18 specimens. **Japan:** BSKU 85148 (1, 83.1 mm HL, 490+ mm TL), BSKU 85149 (1, 79.5 mm HL, 469+ mm TL), BSKU 85150 (1, 71.0 mm HL, 375+ mm TL), Pacific off Tohoku, 700 m, T/V *Tanshu-maru*, sta. D-700, beam trawl, coll. H. Endo, 18 Apr. 1998; BSKU 19016 (1, 108 mm HL, 617+ mm TL), southeast of Oshima Island, Shichito-Iojima Ridge, 34.5750°N, 139.6583°E, 1560–1570 m, FRV *Soyo-maru*, sta. 51, beam trawl, 10 Nov. 1958; BSKU 97798 (1, 87.1 mm HL, 472+ mm TL), off Iwaki, 36.9398°N, 141.7007°E, 967–1219 m, FRV *Wakataka-maru*, sta. 51, otter trawl, 5 Nov. 1995; BSKU 96460 (1, 34.6 mm HL, 192+ mm TL), off Miyako, 39.5342°N, 143.6835°E, 3150–3201 m, R/V *Tansei-maru*, cr. KT-08-27, sta. M-4, 3-m ORE beam trawl, coll. H. Endo and N. Nakayama, 22 Oct. 2008; BSKU 97808 (1, 214 mm HL, 1010+ mm TL), BSKU 97809 (1, 135 mm HL, 797+ mm TL), BSKU 97810 (1, 125 mm HL, 719+ mm TL), BSKU 97811 (1, 151 mm HL, 828+ mm TL), off Kuji, 40.0613°N, 142.6825°E, 1005–1016 m, R/V *Tansei-maru*, cr. KT-93-15, sta. M3, 3-m ORE beam trawl, coll. Y. Machida, 18 Nov. 1993; BSKU 97799 (1, 104 mm HL, 617+ mm TL), BSKU 97800 (1, 65.8 mm HL, 440+ mm TL), off Kitakami River, 38.5318°N, 142.3370°E, 772–783 m, FRV *Wakataka-maru*, sta. 32, otter trawl, 28 Oct. 1995; BSKU 97806 (1, 92.8 mm HL, 603 mm TL), south-southeast of Akkeshi, 42.6052°N, 144.9938°E, 1113–1161 m, R/V *Tansei-maru*, cr. KT-93-15, sta. T8, 3-m ORE beam trawl, coll. Y. Machida, 16 Nov. 1993; *BSKU 114809 (1, 50.6 mm HL, 280+ mm TL), south of Kushiro, 42.6542°N, 144.3770°E, 780 m, FRV *Wakataka-maru*, sta. D-700, otter trawl, coll. R. Misawa, 7 Jun. 2014; *BSKU 114822 (1, 105 mm HL, 610+ mm TL), southeast of Tokachi, 42.3317°N, 143.9585°E, 719 m, FRV *Wakataka-maru*, sta. F-700, otter trawl, coll. R. Misawa, 10 Jun. 2014; *BSKU 114903 (1,

226 mm HL, 1100+ mm TL), *BSKU 114905 (1, 161 mm HL, 859+ mm TL), east-southeast of Cape Erimomisaki, 42.2820°N, 143.8012°E, 572 m, FRV *Wakataka-maru*, sta. G-550, otter trawl, coll. R. Misawa, 9 Jun. 2014. **Bering Sea:** BSKU 35224 (1, 147 mm HL, 821+ mm TL), 58.6633°N, 177.9167°E, 735–800 m, 26 Jun. 1979.

Counts and measurements. Based on 14 specimens (34.6–214 mm HL, 192+–1010+ mm TL). Counts: first dorsal-fin rays II,8–10; pectoral-fin rays i14–i19; pelvic-fin rays 5–8; gill rakers on first arch (outer/inner) 5–8/11–16, on second arch 11–13/10–13; transverse scale rows below first dorsal-fin origin 11.5–15, below first dorsal-fin midbase 9.5–12.5, below second dorsal-fin origin 7.5–12, above anal-fin origin 29–43.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 26–27 (32–36); orbit diameter 19–30 (26–38); postorbital length 49–57 (62–75); postrostral length 74–79; orbit–preopercle distance 40–46 (51–60); suborbital width 8–12 (11–16); upper-jaw length 40–44 (52–56); length of rictus 34–38 (44–50); length of premaxillary tooth band 29–33 (38–43); preoral length 9–12 (11–16); distance between tip and lateral angle of snout 10–12 (13–16); snout width 16–20 (20–26); internasal width 15–18 (20–25); interorbital width 23–27 (30–35); occipital width 10–14 (13–18); body width over pectoral-fin bases 34–72 (46–94); body depth at first dorsal-fin origin 55–82 (74–105); body depth at anal-fin origin 41–75 (54–99); prepelvic length 96–105 (124–138); preanus length 142–164 (189–217); preanal length 146–177 (195–226); isthmus–pelvic distance 36–44 (47–58); isthmus–anus distance 83–109 (111–139); isthmus–anal distance 88–115 (118–147); pelvic–anal distance 51–79 (68–101); anus–anal distance 3–7 (5–10); pelvic-fin length 40–53 (52–68); pectoral-fin length 51–66 (66–84); predorsal length 99–113 (132–148); height of first dorsal fin 43–50 (55–66); length of first dorsal-fin base 21–30 (28–40); interdorsal length 11–20 (14–27); length of gill slit 16–22 (22–29); length of posterior nostril 5–8 (7–10); barbel length 5–8 (6–11).

Size. *Coryphaenoides pectoralis* is by far the largest species of grenadiers. According to biological information summarized by Orlov & Tokranov (2008), it attains about 220 cm TL (Toponogov & Kurennoi 1986). The largest specimen examined is HUMZ 69383 collected from the Pacific off Hokkaido, measuring 158 cm TL.

Variation. In the smaller specimens examined, the body scales are covered with short, reclined, needle-like spinules giving a more or less spiny appearance to body surfaces (Fig. 128A–B). The spinules become weaker with increasing body size, and those of the larger specimens examined were always difficult to distinguish (Fig. 128C–D).

Sexual dimorphism. Orlov & Tokranov (2008) reported that *C. pectoralis* collected from the northwestern Pacific off the northern Kuril Islands exhibits sexually dimorphic differences in body size: females attain a larger

FIGURE 127. Fresh specimens of *Coryphaenoides pectoralis*. (A) BSKU 96460, 34.6 mm HL, 192+ mm TL, off Miyako, Iwate Pref., Pacific, 3150–3201 m depth; (B) BSKU 114809, 50.6 mm HL, 280+ mm TL, south of Kushiro, Hokkaido, Pacific; (C) BSKU 114822, 105 mm HL, 601+ mm TL, southeast of Tokachi, Hokkaido, Pacific, 719 m depth; (D) BSKU 114905, 161 mm HL, 859+ mm TL, east-southeast of Cape Erimomisaki, Hokkaido, Pacific, 572 m depth; (E–G) BSKU 114903, 226 mm HL, 1100+ mm TL, collected with BSKU 114905 (E, reversed). (A–E) Lateral views; (F) dorsal and (G) ventral views of the head and trunk. [Photos: (A) H. Endo and N. Nakayama; (B–G) N. Nakayama]

size than males (135 cm maximum TL vs. 97 cm) with the body weight reaching up to 11.8 kg (vs. 3.6 kg).

Development. Larval morphology was described by Endo *et al.* (1993, 2010), and Endo (1996, 2014) [see also Ambrose (1996)]. Based on a pelagic larva with a nearly complete tail, Endo *et al.* (2010) reported the reliable

numbers of second dorsal- and anal-fin rays, and total vertebrae of this species as 156, 137, and $14 + 117 = 131$ respectively.

Distribution. Widely distributed in the North Pacific from Japan to the United States, including the Okhotsk and Bering Seas as well as the Emperor Seamounts

FIGURE 128. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of *Coryphaenoides pectoralis*. (A–B) BSKU 85150, 71.0 mm HL; (C–D) BSKU 97811, 151 mm HL. (A, C) Views from above; (B, D) oblique views. [Photos: NSMT]

(Iwamoto & Stein 1974; Borets 1986; Iwamoto 1990; Mecklenburg *et al.* 2002; this study). Depth range varied from 140 to 3201 m, but usually shallower than 1500 m (Orlov & Tokranov 2008; this study). Arkhipkin *et al.* (2010) recently reported an 815 mm TL specimen from off the Falkland Islands, in the southwestern Atlantic (as *Albatrossia pectoralis*), suggesting its antipodal migration through the Drake Passage [this specimen was previously reported as *Albatrossia* cf. *pectoralis* by Brickle & Laptikhovsky (2002)]. In Japan, known from off the Pacific coasts southward to the Izu Peninsula (138.96°E) and the southern Okhotsk Sea (Appendix 3-7C). Depth range off Japan 152–3201 m. Common, especially in the north of Choshi (35.74°N; Chiba Pref.).

Remarks. *Coryphaenoides pectoralis* has been well described by previous authors, and thus there is no need for repetition here. For a full description see Okamura (1970a) and Iwamoto & Stein (1974). This species is among one of the most abundant continental slope fishes found in the North Pacific (Orlov & Tokranov 2008). The biology of

this species off the Kuril and Kamchatka Islands was well investigated by Orlov & Tokranov (2008).

Relationships. There has been no general consensus about the generic affiliation of this common grenadier. Gilbert (1891) originally described the species under the name *Macrurus* (*Malacocephalus*) *pectoralis*. Subsequently, it was transferred to *Nematonurus* Günther, 1887 (originally described as a subgenus of *Macrourus* Bloch, 1786) by Goode & Bean (1896), whereas Jordan & Gilbert (*in* Jordan & Evermann 1898) established its own genus *Albatrossia*. As far as I know, Fowler (1936) was the first who synonymized *Albatrossia* with *Coryphaenoides*, and Taranetz (1937) subsequently listed this species as a species of *Coryphaenoides*. Okamura (1970a) provided the species' description under *Coryphaenoides*, although *Nematonurus* retained subgeneric status; he regarded *Albatrossia* as a junior synonym of *Coryphaenoides*. Iwamoto & Stein (1974) followed Okamura's (1970a) conclusion, but Iwamoto & Sazonov (1988) latter resurrected *Albatrossia* as a separate

genus distinct from *Coryphaenoides*. During the past three decades, some authors considered the species to be the only representative of *Albatrossia* (e.g., Iwamoto 1990; Endo *et al.* 1993; Shinohara *et al.* 1996; Mecklenburg *et al.* 2002), while others regarded the species as a member of *Coryphaenoides* (e.g., Okamura 1984b, 1988; Sawada 1983; Nakabo 1993, 2000, 2002; Senou *et al.* 2006; Endo *et al.* 2010; Nakabo & Kai 2013). The difference of opinion is largely due to unique morphological features of this species. Despite their close similarity in general appearance, it readily differs from all other species of *Coryphaenoides* in having several autapomorphic features including: a horizontally elongate otolith with a comb-like ventral margin; 2 rete-gas gland complexes in a reduced swim bladder; teeth in 2 irregular series in the premaxillary, and a distinct 1 row in the dentary.

Recent molecular analyses, by contrast, showed that the species is deeply nested within a clade of *Coryphaenoides*, suggesting *Albatrossia* to be a junior synonym of that taxon (Wilson *et al.* 1991; Wilson 1994; Morita 1999; Wilson & Attia 2003; Roa-Varón & Ortí 2009). However, their results also imply that *Albatrossia* might be recognized as a subgenus of *Coryphaenoides*. According to the most comprehensive tree based on the RAG1 and mitochondrial ribosomal genes (Roa-Varón & Ortí 2009: fig. 3), the subgenus *Coryphaenoides* (*sensu* Iwamoto 1990) is divided into two clades, with “*pectoralis*” belonging to one that excludes *C. rupestris* Gunnerus, 1765 (type species of *Coryphaenoides*). Their tree also shows that “*pectoralis*” forms a small clade with *C. acrolepis* (Bean, 1884) and *C. cinereus* (Gilbert, 1896), although their close relationship does not seem to be supported by morphological characters. It is obvious that further investigations are necessary to determine whether *Albatrossia* deserves subgeneric status within *Coryphaenoides*. If so, there is an additional question around how many species are included in this group. In the present study, *Albatrossia* is tentatively regarded as a subgenus of *Coryphaenoides*, and *C. pectoralis* is considered to be its only representative.

Comparisons. *Coryphaenoides pectoralis* is highly distinctive among the genus in its unique dentition: the premaxillary teeth are arranged in two irregular series, whereas those in the dentary are in one distinct row. The small swim bladder of *C. pectoralis*, with two rete-gas gland complexes, also distinguishes the species from the other members of *Coryphaenoides* [vs. swim bladder well developed with 4–7 rete-gas gland complex (fide Iwamoto & Sazonov 1988:38)]. It further differs from all other congeners in having a horizontally elongate otolith with a comb-like ventral margin (vs. otolith moderately high, not comb-like ventrally). For more comparisons with Japanese congeners, see the key to species provided in this study.

***Coryphaenoides rudis* Günther, 1878**

[Japanese name: Daikoku-hige]

(Figs. 129–130; Appendix 3-7D)

Coryphaenoides rudis Günther, 1878:24 [original description; lectotype: BMNH 1889.12.7.74, from west of Kermadec Islands, *Challenger* sta. 171, in 600 fth (1097 m); in part, see the Comments on type specimens]; Gilbert & Hubbs 1916:144 (lectotype designation; listed); Shao *et al.* 2008b: table 2 (1 spec. listed from South China Sea; first record from Taiwan); Iwamoto *et al.* 2015:73, fig. 11 (brief description; 2 spec. from South China Sea and southeastern Taiwan); Nakayama & Endo 2016a:7, figs. 2j–l, 3 (first record from Japan; 6 spec. from Hyuga-nada, Suruga Bay, and Minami-torishima Island; re-examination of lectotype; comments on type specimens; new Japanese name: “Daikoku-hige”); Motomura 2020:39 (listed; Japan).

Macrourus paradoxus Smith & Radcliffe in Radcliffe, 1912:115, pl. 25, fig. 1 [original description; holotype: USNM 72932, from off eastern Palawan Island, 9°13'00"N, 118°51'15"E, in 1105 fth (2021 m)].

Nematonurus macrocephalus Maul, 1951:17, figs. 3, 4c [original description; holotype: MMF 898 (mounted), from off Madeira; 2 paratypes in BMNH and MMF].

Coryphaenoides macrocephalus: Iwamoto 1990:212, fig. 490 (synopsis).

Diagnosis. Pelvic-fin rays 9–10. Snout bluntly pointed, slightly protruding beyond upper jaw. Tip and lateral angles of snout armed with small scute-like scales; scales along head ridges slightly enlarged and thickened. Mouth large, posterior margin of upper jaw extending slightly beyond vertical through midorbit; upper-jaw length 35–42% HL; lateral corner of mouth not restricted by skin folds. Outermost gill slit greatly restricted, length 5–7% HL. Barbel long, slender, length 14–23% HL. Premaxillary teeth in broad tapered bands with outer series notably enlarged; dentary teeth in narrow tapered band with inner series enlarged. Body scales covered with long, reclined, needle-like spinules in tightly packed convergent rows or quincunx order; tip of last spinule in each row extending beyond posterior scale margin (except for larger adults). Transverse scale rows below first dorsal-fin midbase 6.5–9. Head fully scaled, lacking naked areas on snout. Height of first dorsal fin less than HL (76–84% HL); second spinous ray serrated along its leading edge; first dorsal-fin rays II, 9–10. Outer pelvic-fin ray slightly prolonged, its tip usually extending beyond anal-fin origin. Body uniformly dark brown.

Material examined. 7 specimens. **Lectotype of *Coryphaenoides rudis*:** BMNH 1887.12.7.74 (168 mm HL, 814+ mm TL), west of Kermadec Islands, 28.5500°S, 177.8333°E, 600 fth (1098 m), *Challenger* sta. 171, trawl, 15 Jul. 1874. **Non-types: Japan:** MSM 93-18 (1, 62.6 mm HL, 336 mm TL), Suruga Bay, F/V *Nishimiya-maru*, crab trap, 10 Mar. 1993; BSKU 49468 (1, 127 mm HL,

599+ mm TL), BSKU 49570 (1, 64.8 mm HL, 322+ mm TL), BSKU 49577 (1, 75.7 mm HL, 384+ mm TL), off Miyazaki, Hyuga-nada, 32.3083°N, 132.1817°E, 1453–1481 m, R/V *Shinkai-maru*, 3 Apr. 1991; HUMZ 72029 (1, 248 mm HL, 1079+ mm TL), HUMZ 72030 (1, 269 mm HL, 1200+ mm TL), between Shichito-Iojima Ridge and Marcus Island, 23.9333°N, 150.6000°E, 1100–1160 m, 10 Oct. 1977.

Counts and measurements. Based on 7 specimens (62.6–269 mm HL, 322+–1200+ mm TL). Counts: first dorsal-fin rays II,9–10; pectoral-fin rays i19–i23; pelvic-fin rays 9–10; gill rakers on first arch (outer/inner) 1–5/9–11, on second arch 8–10/9–11; longitudinal scales 31–40; transverse scale rows below first dorsal-fin origin 8.5–11, below first dorsal-fin midbase 6.5–9, below second dorsal-fin origin 5.5–8.5, above anal-fin origin 29–36; pyloric caeca 13–14.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 24–30 (31–40); orbit diameter 18–23 (23–31); postorbital length 53–61 (71–78); postrostral length 74–

79; orbit–preopercle distance 48–59 (64–75); suborbital width 13–15 (18–20); upper-jaw length 35–42 (47–55); length of rictus 29–36 (39–46); length of premaxillary tooth band 25–29 (33–37); preoral length 9–15 (12–21); distance between tip and lateral angle of snout 10–15 (13–20); snout width 19–25 (24–33); internasal width 18–23 (23–31); interorbital width 28–31 (37–40); body width over pectoral-fin bases 37–67 (51–87); body depth at first dorsal-fin origin 86–96 (111–124); body depth at anal-fin origin 64–85 (84–108); prepelvic length 103–115 (136–149); preanus length 155–156 (200–204); preanal length 160–177 (206–231); isthmus–pelvic distance 42–52 (55–68); pelvic–anal distance 60–75 (77–98); pelvic-fin length 44–76 (60–98); pectoral-fin length 49–62 (63–81); predorsal length 113–119 (146–155); height of first dorsal fin 76–84 (97–111); length of first dorsal-fin base 24–30 (31–40); interdorsal length 20–38 (26–48); length of gill slit 5–7 (6–10); length of posterior nostril 6–8 (8–10); barbel length 14–23 (18–31).

Size. Attains at least 120 cm TL (Iwamoto & Graham 2001; this study). The specimen HUMZ 72030 collected

FIGURE 129. *Coryphaenoides rudis*. (A) BSKU 49577, 75.7 mm HL, 384+ mm TL, Hyuga-nada, Pacific, 1453–1481 m depth; (B) BSKU 49468, 127 mm HL, 599+ mm TL, collected with BSKU 49577; (C–E) HUMZ 72029, 248 mm HL, 1079+ mm TL, between Shichito-Iojima Ridge and Marcus Island, Pacific. (A–C) Lateral views in fresh condition; (D) dorsal and (E) ventral views of the head and trunk in preserved condition. [Photos: reproduced from Nakayama & Endo (2016a: fig. 4a–e) with permission from the Ichthyological Society of Japan]

FIGURE 130. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of *Coryphaenoides rudis*. (A) BSKU 49570, 64.8 mm HL; (B) BSKU 49468, 127 mm HL; (C) BMNH 1887.12.7.74, lectotype, 168 mm HL; (D) HUMZ 72029, 248 mm HL. Views from above. [Photos: (A) N. Nakayama; (B–D) reproduced from Nakayama & Endo (2016a: fig. 2j–l) with permission from the Ichthyological Society of Japan]

from off the Shichito-Iojima Ridge is one of the largest specimens of the species (1200+ mm TL).

Distribution. Widespread in tropical and temperate waters of the world's oceans at depths of 980–2400 m (Sazonov & Iwamoto 1992; Shcherbachev & Iwamoto 1995; Iwamoto & Williams 1999; Merrett & Iwamoto 2000; Iwamoto & Graham 2001; Shao *et al.* 2008a, 2008b; Melo *et al.* 2010; Nakayama & Endo 2016a). In Japan, so far known from Hyuga-nada, Suruga Bay, and the Shichito-Iojima Ridge off Minami-torishima Island, at depths of 1100–1481 m (Nakayama & Endo 2016a; Appendix 3-7D). Rare.

Remarks, relationships, and comparisons. *Coryphaenoides rudis* is a widespread species, being a senior synonym of both *C. paradoxus* (Smith & Radcliffe in Radcliffe, 1912) and *C. macrocephalus* (Maul, 1951). For further morphological details see Iwamoto and Sazonov (1988; as *C. paradoxus*). The species was recently recorded from Japan by Nakayama & Endo (2016a) based

on the specimens examined here. They considered that Franz's (1910) record of the species from Abratsubo, Kanagawa Pref., Japan, represented a misidentification of a species of a different genus.

Coryphaenoides rudis is a member of the subgenus *Coryphaenoides* (sensu Iwamoto 1990). Among Japanese congeners, it is most similar to *C. soyoae* Nakayama & Endo, 2016, but differs from the latter in having body scales covered with spinules arranged in tightly packed convergent rows or quincunx order (Fig. 130 vs. narrowly divergent rows; Fig. 133), fewer pelvic-fin rays (9–10 vs. 11), smaller body scales (transverse scale rows 8.5–11 vs. 7–7.5 below first dorsal-fin origin; 6.5–9 vs. 4–5 below first dorsal-fin midbase), a broader interorbital space (28–31% HL vs. 24–27%), and a lower first dorsal fin (76–84% HL vs. 87%), and in lacking a naked area on the underside of the snout (vs. present).

***Coryphaenoides soyoae* Nakayama & Endo, 2016**
(Figs. 131–133; Appendix 3-7E)
[Japanese name: Kuro-hige]

Japanese name: “Kuro-hige”); Motomura 2020:39 (listed; Japan).

Coryphaenoides soyoae Nakayama & Endo, 2016a:2, figs. 1, 2a–c (original description; holotype BSKU 20297, from Shichito-Iojima Ridge, in 2740 m; 1 paratype from off Onahama; new

Diagnosis. Pelvic-fin rays 11. Snout bluntly pointed, only slightly protruding beyond upper jaw. Terminal snout scute absent; scales on lateral angles of snout and infraorbital ridge slightly enlarged and thickened.

FIGURE 131. Fresh specimen of *Coryphaenoides soyoae*. NSMT-P 78018, paratype, 73.7 mm HL, 365+ mm TL, off Onahama, Ibaraki, 2948–2991 m depth. Lateral view. [Photo: reproduced from Nakayama & Endo (2016a: fig. 1d) with permission from the Ichthyological Society of Japan]

FIGURE 132. Holotype of *Coryphaenoides soyoae*. BSKU 20297, 120 mm HL, 578+ mm TL, between Izu and Ogasawara Islands, Shichito-Iojima Ridge, Pacific, 2740 m depth. (A) Lateral, (B) dorsal, and (C) ventral views. Preserved condition [Photos: reproduced from Nakayama & Endo (2016a: fig. 1a–c) with permission from the Ichthyological Society of Japan]

FIGURE 133. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of *Coryphaenoides soyoae*. (A–B) BSKU 20297, holotype, 120 mm HL; (C) NSMT-P 78018, paratype, 73.7 mm HL. (A, C) Views from above; (B) oblique view. [Photos: reproduced from Nakayama & Endo (2016a: fig. 2a–c) with permission from the Ichthyological Society of Japan]

Dorsal contour of head prominently humped over nape, with distinct depression above orbits. Orbit diameter 20–21% HL. Mouth large, posterior margin of upper jaw extending to below hind 1/3 of orbit or beyond; upper-jaw length 35–40% HL; lateral corner of mouth not restricted by skin folds. Outermost gill slit greatly restricted, length 7–8% HL. Barbel well developed, slender, length 11–15% HL. Teeth in tapered bands in both jaws, with outer premaxillary series distinctly enlarged. Body scales covered with long, reclined, needle-like spinules in slightly divergent rows; tip of last spinule in each row extending well beyond posterior scale margin; middle row not especially higher than adjacent rows. Transverse scale rows below first dorsal-fin midbase 4–5. Snout fully scaled dorsally, but naked ventrally. Interdorsal space slightly less than first dorsal-fin base length. Height of first dorsal fin less than HL (87% HL); second spinous ray serrated along its leading edge; first dorsal-fin rays II,9–10. Pectoral-fin rays i19–i22. Outer pelvic-fin ray

greatly prolonged, extending well beyond anal-fin origin. Body uniformly blackish.

Material examined. 2 specimens. **Holotype** of *Coryphaenoides soyoae*: BSKU 20297 (120 mm HL, 578+ mm TL), between Izu and Ogasawara Islands, Shichito-Iojima Ridge, Japan, 30.0000°N, 140.5417°E, 2740 m, FRV *Soyo-maru*, sta. 4, bottom trawl, 18 Nov. 1972. **Paratype** of *C. soyoae*: NSMT-P 78018 (1, 73.7 mm HL, 365+ mm TL), off Onahama, 36.8050°N, 142.3717°E, 2948–2991 m, FRV *Soyo-maru*, cr. SO-07-02, benthos net, coll. H. Saito and M. Okanishi, 2 Aug. 2007.

Counts and measurements. Based on 2 specimens (73.7–120 mm HL, 365+–578+ mm TL). Counts: first dorsal-fin rays II,9–10; pectoral-fin rays i19–i22; pelvic-fin rays 11; gill rakers on first arch (outer/inner) 5–6/11–12, on second arch 8–9/10; longitudinal scales 34–37; transverse scale rows below first dorsal-fin origin 7–7.5, below first dorsal-fin midbase 4–5, below second dorsal-fin origin 7, above anal-fin origin 26.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 27–29 (36–39); orbit diameter 20–21 (26–28); postorbital length 55–56 (74–75); postrostral length 74; orbit–preopercle distance 49 (66); suborbital width 13–14 (18); upper-jaw length 35–40 (47–53); length of rictus 29–32 (38–43); preoral length 12–14 (16–19); distance between tip and lateral angle of snout 14–15 (19–20); snout width 23–24 (31–33); internasal width 21–23 (28–31); interorbital width 24–27 (33–36); body width over pectoral-fin bases 48–56 (65–75); body depth at first dorsal-fin origin 85–90 (114–121); body depth at anal-fin origin 70–76 (94–101); prepelvic length 110–116 (148–156); preanus length 152–159 (204–214); preanal length 158–165 (212–223); isthmus–pelvic distance 45–46 (60–62); pelvic–anal distance 43–52 (58–69); pelvic-fin length 66–84 (89–113); pectoral-fin length 49 (66); predorsal length 109–116 (147–156); height of first dorsal fin 87 (117); length of first dorsal-fin base 27–28 (36–38); interdorsal length 23–24 (30–32); length of gill slit 7–8 (9–11); length of posterior nostril 6 (7–8); barbel length 11–15 (15–20).

Size. Attains about 58 cm TL (BSKU 20297, holotype, 578+ mm TL, Shichito-Iojima Ridge, Japan).

Distribution. So far known only the Pacific off Onahama, Fukushima, and the Shichito-Iojima Ridge, at depth of 2740–2991 m (Appendix 3-7E). Very rare.

Remarks, relationships, and comparisons. *Coryphaenoides soyoe* was recently described by Nakayama & Endo (2016a), to which readers can refer for a full description of the species. In Japanese waters, *C. soyoe* is most closely similar to *C. rudis* Günther, 1878, but they can be distinguished from each other by the combination of the body-scale spinulation, head squamation, numbers of transverse scale rows, interdorsal length, and height of the first dorsal fin (see the Remarks, relationships, and comparisons for the latter species).

Coryphaenoides yaquinae Iwamoto & Stein, 1974

[Japanese name: Shinkai-yoroidara]

(Figs. 99B, 100C–D, 134; Appendix 3-7F)

Coryphaenoides yaquinae Iwamoto & Stein, 1974:34, figs. 15A, 20 (original description; holotype: USNM 211795, from Tufts Abyssal Plain, R/V *Yaquina* sta. TP-3, 44°39.9'N, 133°37.2'W, in 3724 m; 2 paratypes from off California); Endo & Okamura 1992:433, fig. 1 (first record from Japan; brief description; 28 spec. from northwestern Pacific; new Japanese name: “Shinaki-yoroidara”); Nakabo 1993:363 (in key; Japan); Okamura 1997:124, fig. (compiled); Nakabo 2000:427 (in key; Japan); Nakabo 2002:427 (in key; Japan); Senou 2008:367, fig. 31.35 (in situ observation; Japan Trench); Shinohara *et al.* 2009:708 (listed; Pacific off Tohoku); Nakabo & Kai 2013:503 (in key; Japan); Amaoka *et al.* 2020:162, fig. 218 (listed; Hokkaido); Motomura 2020:39 (listed; Japan).

[?] *Coryphaenoides* sp.: Senou 2008:367, fig. 31.36 (in situ observation; Chishima Trench).

Diagnosis. Pelvic-fin rays usually 10, rarely 9 or 11. Snout rounded in adults (pointed in young specimens), slightly protruding beyond upper jaw. Tip and lateral angles of snout armed with small scute-like scales; scales along head ridges slightly thickened but not especially enlarged. Mouth large, posterior margin of upper jaw almost reaching vertical through hind rim of orbit; upper-jaw length 37–41% HL; lateral corner of mouth not restricted by skin folds. Outermost gill slit moderately wide, length 18–22% HL. Barbel long, stout at base, length 13–21% HL. Teeth conical, sharply pointed; premaxillary teeth in 2–3 irregular rows or narrow band, with outer series enlarged; mandibular teeth in 2 irregular rows near symphysis, uniserial row posteriorly. Body scales covered with short, reclined, needle-like to knife-like spinules in subparallel to narrowly divergent rows; tip of last spinule in each row extending to posterior scale margin or slightly beyond. Transverse scale rows below first dorsal-fin midbase 6–7.5. Top of snout narrowly naked posterior to leading edges; underside of head almost completely naked. Interdorsal space greater than first dorsal-fin base length. Height of first dorsal fin less than HL (62–82% HL); second spinous ray not prolonged, with serrated leading edge; first dorsal-fin rays II, 8–10. Body uniformly dark brown.

Material examined. 23 specimens. **Japan:** BSKU 49883 (1, 59.6 mm HL, 325 mm TL), BSKU 49884 (1, 77.0 mm HL, 412 mm TL), Japan Trench, 5500 m, R/V *Hakuho-maru*, cr. KH-91-6, sta. 18, 6 Nov. 1991; NSMT-P 78028 (1, 133 mm HL, 685+ mm TL), off Kinakzan, 4000 m, FRV *Soyo-maru*, benthos net, coll. H. Saito, 7 Aug. 2007; NSMT-P 78022 (1, 82.7 mm HL, 441+ mm TL), NSMT-P 78023 (1, 118 mm HL, 610+ mm TL), NSMT-P 78024 (1, 115 mm HL, 565+ mm TL), NSMT-P 78025 (1, 90.2 mm HL, 537+ mm TL), off Kushiro, 5700 m, FRV *Soyo-maru*, sta. Kago-6, coll. H. Saito, 1–2 Aug. 2007; NSMT-P 19033 (1, 69.5 mm HL, 371+ mm TL), off Aogashima Island, Izu Trench, 31.1750°N, 141.7017°E, 5100 m, 30 Jul. 1976; NSMT-P 30198 (1, 86.3 mm HL, 467+ mm TL), NSMT-P 30201 (1, 86.4 mm HL, 468+ mm TL), east of Shichito-Iojima Ridge, western Pacific, 30.0967°N, 146.9267°E, 6160 m, R/V *Hakuho-maru*, fish trap, 11 May 1982; BSKU 35735 (1, 99.3 mm HL, 534+ mm TL), BSKU 35736 (1, 91.5 mm HL, 521+ mm TL), BSKU 35737 (1, 88.7 mm HL, 488+ mm TL), east of Japan Trench, 38.5650°N, 145.2617°E, 5350–5370 m, R/V *Hakuho-maru*, cr. KH-81-4, sta. 11, 1 Aug. 1981; BSKU 37301 (1, 92.8 mm HL, 495+ mm TL), east of Cape Inozaki, east of Japan Trench, 35.8333°N, 145.2000°E, 4100 m, 1981; BSKU 35726 (1, 69.1 mm HL, 377+ mm TL), BSKU 35727 (1, 73.7 mm HL, 396 mm TL), BSKU 35729 (1, 78.0 mm HL, 448+ mm TL), BSKU 35730 (1, 86.8 mm HL, 478+ mm TL), Kesennuma, Japan Trench, 38.5550°N, 144.3233°E, 6380–6450 m, R/V *Hakuho-*

FIGURE 134. Fresh specimens of *Coryphaenoides yaquinae*. (A) BSKU 49884, 77.0 mm HL, 412 mm TL, KH-91-6 sta. 18 (reversed); (B) BSKU 49883, 59.6 mm HL, 325 mm TL, KH-91-6 sta. 18. Lateral views. [Photos: BSKU]

maru, cr. KH-81-4, sta. 12, 31 Jul. 1981; BSKU 49930 (1, 67.3 mm HL, 311+ mm TL), Japan Trench, 38.7342°N, 144.1233°E, 5500 m, R/V *Hakuho-maru*, cr. KH-91-6, sta. 16-2, 18 Nov. 1991; *BSKU 35734 (1, 107 mm HL, 553+ mm TL), east of Japan Trench, 38.5650°N, 145.2617°E, 5350–5370 m, R/V *Hakuho-maru*, cr. KH-81-4, sta. 11, 1 Aug. 1981; BSKU 35720 (1, 69.0 mm HL, 356+ mm TL), off Kamaishi, Japan Trench, 39.1883°N, 143.9067°E, 5010–5109 m, R/V *Hakuho-maru*, cr. KH-81-4, sta. 10, 25 Jul. 1981; BSKU 19997 (1, 70.1 mm HL, 362+ mm TL), BSKU 19999 (1, 88.4 mm HL, 498+ mm TL), east of Japan Trench, 33.9750°N, 142.8550°E, 5200 m, FRV *Soyo-maru*, beam trawl, 4 Jun. 1970.

Counts and measurements. Based on 22 specimens (59.6–133 mm HL, 311+–685+ mm TL). Counts: first dorsal-fin rays II,8–10; pectoral-fin rays i18–i20; pelvic-fin rays 9–11; gill rakers on first arch (outer/inner) 6–10/11–13, on second arch 10–13/11–13; longitudinal scales 29–38; transverse scale rows below first dorsal-fin origin 6.5–10.5, below first dorsal-fin midbase 6–7.5, below second dorsal-fin origin 7–10, above anal-fin origin 26–34; pyloric caeca 9.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 25–30 (32–41); orbit diameter 19–22 (25–29); postorbital length 54–59 (72–77); postrostral length 74–78; orbit–preopercle distance 44–49 (58–66); suborbital width 10–13 (13–17); upper-jaw length 37–41 (49–56); length of rictus 33–37 (44–50); length of premaxillary tooth band 27–33 (35–43); preoral length 10–16 (12–22); distance between tip and lateral angle of snout 11–15 (14–20); snout width 20–25 (25–33); internasal width 16–20 (21–26); interorbital width 24–29 (32–38); occipital width 1–6 (2–8); body width over pectoral-fin bases 44–74 (57–96); body depth at first dorsal-fin origin 78–99 (104–128); body depth at anal-fin origin 60–80 (80–103); prepelvic length 99–122 (127–163); preanus length 168–201 (219–261); preanal length 175–205 (230–266); isthmus–pelvic distance 36–50 (47–64); isthmus–anus distance 101–135 (134–178); isthmus–anal distance 109–140 (144–185); pelvic–anal distance 71–101 (96–134); anus–anal distance 5–11 (6–14); pelvic-fin length 66–86 (85–112); pectoral-fin length 60–77 (78–101); predorsal length 111–125 (148–162); height of first dorsal fin 62–82 (81–111); length of first

dorsal-fin base 24–35 (32–45); interdorsal length 39–58 (52–75); length of gill slit 18–22 (23–29); length of posterior nostril 4–7 (6–9); barbel length 13–21 (18–28); length of pyloric caecum 35 (45).

Size. Attains about 77 cm TL (Wilson & Waples 1983; Mecklenburg *et al.* 2002).

Distribution. Known from the North and Central Pacific below a depth of about 3400 m (Iwamoto & Stein 1974; Wilson & Waples 1983; Iwamoto 1990; Endo & Okamura 1992; Mecklenburg *et al.* 2002; this study). In Japan, known only from the Japan, Izu-Ogasawara, and Kuril-Kamchatka Trenches at depths of 4000–7433 m (Appendix 3-7F). Moderately common in its depth range.

Remarks. For further morphological information see the original description given by Iwamoto & Stein (1974); see also Wilson & Waples (1983) who provided a detailed comparison between *C. yaquinae* and *C. armatus* (Hector, 1875). *Coryphaenoides yaquinae* is restricted in the Pacific Ocean, and was first reported from Japan by Endo & Okamura (1992). The previous deepest record for the species was 6380–6450 m, based on by five specimens collected from the Japan Trench (BSKU 35726–35730, 69.0–86.8 mm HL, 377+–478+ mm TL; Endo & Okamura 1992). Using a baited camera lander, Jamieson *et al.* (2009) also observed the species at 6945 m in the same area. One of the specimens examined (NSMT-P 98009, 115 mm HL, 585+ mm TL) was collected from the Japan Trench at a depth of 7340–7433 m; this represents a new deepest record for *C. yaquinae*.

Relationships and comparisons. *Coryphaenoides yaquinae* belongs to the subgenus *Nematonurus* (sensu Iwamoto 1990), and is most similar to *C. armatus*. For a comparison between the two species see the Relationships and comparisons for the latter species.

Genus *Hymenocephalus* Giglioli, 1884

[Japanese name: Sujidara-zoku]

Hymenocephalus Giglioli, 1884:228 (type species: *Hymenocephalus italicus* Giglioli, 1884, by original designation).

Mystacourus Günther, 1887:139 [as subgenus of *Macrurus*; type species: *Hymenocephalus italicus* Giglioli, 1884, by subsequent designation by Jordan & Evermann (1898)].

Papyrocephalus Gilbert & Hubbs, 1920:539 (as subgenus of *Hymenocephalus*; type species: *Hymenocephalus aterrimus* Gilbert, 1905, by original designation).

Diagnosis. Anus immediately anterior to anal-fin origin, surrounded by narrow periproct. Ventral striae, consisting of fine, parallel black lines over silvery layer on each side of isthmus, shoulder girdle, most of chest, and posterior to pelvic fins. Gular region covered with fine striations of black epithelium. Light organ developed as long tube, extending from anus to slightly anterior to pelvic-fin bases;

small lens at each end of light organ. Second spinous ray of first dorsal fin smooth along its leading edge. Pelvic-fin rays 7–15, outer ray slender, not expanded or ribbon-like distally. Teeth all small, in narrow villiform bands in both jaws. Trunk moderately deep, laterally compressed. Head bones paper-thin, fragile in most species; head skin membranous, translucent. Nasal bones weakly ossified, not forming stout, dorsoventrally flattened processes. Chin barbel present or absent. Body scales large, highly deciduous; radial grooves developed anteriorly; spinules rudimentary, very short, erect, needle-like or narrowly triangular, arranged in quincunx order; reticulate structure absent. Branchiostegal rays 7. Most species small, usually less than 20 cm TL. Body with silvery reflection when fresh (in most species). [From Iwamoto & Graham (2001), Iwamoto *et al.* (2011), McMillan & Iwamoto (2014), and Schwarzhans (2014).]

Remarks. *Hymenocephalus* comprises 27 species, six of which are known from Japanese and adjacent waters (Nakayama *et al.* 2015a). Most species are small and less than 20 cm TL (Iwamoto *et al.* 2015). Due to the fragile nature of the head and body, specimens of *Hymenocephalus* are easily damaged when trawled, which often makes species identification difficult. Species of the genus typically occur at upper slope depths of the world's oceans, but are strangely absent from the northeastern Pacific (McMillan & Iwamoto 2014; Schwarzhans 2014; Nakayama *et al.* 2015a). *Hymenogadus* Gilbert & Hubbs, 1920 and *Spicomacrus* Okamura, 1970 were previously recognized as subgenera of *Hymenocephalus* (e.g., Okamura 1970a; Iwamoto 1990), but these taxa are now considered to deserve full generic status (Iwamoto *et al.* 2011, 2015; Schwarzhans 2014). Accordingly, *Hymenocephalus* currently includes two subgenera, viz., *Hymenocephalus* and *Papyrocephalus* Gilbert & Hubbs, 1920.

Key to species of *Hymenocephalus* from Japan and adjacent waters

- 1a** Pelvic-fin rays 7–9; ventral half of tail immaculate or only slightly peppered with small black melanophores, dorsal half usually pale, darker in some specimens; chin barbel variously developed, length longer than 1/3 orbit diameter (14–73% PRL)..... **2**
- 1b** Pelvic-fin rays 11–12; dorsal and ventral halves of tail uniformly peppered with small black melanophores; chin barbel absent or rudimentary, if present, length distinctly shorter than 1/3 orbit diameter (less than 5% PRL)..... **5**
- 2a** Barbel length greater than 1/2 PRL (53–73% PRL), its tip extending to about vertical through posterior margin of orbit or beyond; orbit diameter 35–43% PRL **3**
- 2b** Barbel length less than 1/2 PRL (14–52% PRL), its tip not reaching vertical through posterior margin of orbit; orbit diameter 43–59% PRL **4**
- 3a** Pelvic-fin rays 8; snout depressed, its tip barely protruding

- beyond upper jaw; dorsal half of tail pale to dusky; interorbital width 18–27% PRL.....*H. longibarbis*
- 3b** Pelvic-fin rays 9; snout not depressed, its tip protruding distinctly beyond upper jaw; dorsal half of tail abruptly darker posteriorly; interorbital width 21% PRL.....*H. yamasakiorum*
- 4a** Barbel length 36–52% PRL; pelvic-fin length 95–103% PRL; ostial and caudal colliculi of otolith separated.....*H. hachijoensis*
- 4b** Barbel length 14–23% PRL; pelvic-fin length 70–90% PRL; ostial and caudal colliculi of otolith fused.....*H. striatissimus*
- 5a** Gill rakers on second arch (outer/inner) 24–26/22–25; suborbital width 16–17% PRL.....*H. lethoneus*
- 5b** Gill rakers on second arch (outer/inner) 18–22/15–21; suborbital width 22–26% PRL.....*H. papyraceus*

Hymenocephalus hachijoensis Okamura, 1970

[Japanese name: Hachijo-sokodara]

(Figs. 135–136; Appendix 3-8A)

Hymenocephalus striatissimus hachijoensis Okamura, 1970a:50, pl. XVI (original description; as subspecies of *H. striatissimus* Jordan & Gilbert in Jordan & Starks, 1904; holotype: BSKU 14171, from off Hachijo-jima Island, 40°03'6"N, 133°16'5"E, in 570 m; 1 paratype collected with holotype; new Japanese name: "Hachijo-sokodara"); Okamura 1970b: table 1 (listed; Japan); Tominaga & Uyeno 1981:489 (listed; Japan).

Hymenocephalus longiceps (not Smith & Radcliffe in Radcliffe 1912): Yatou 1982:141, 246, fig. 85 (brief description; in part, ZMUT KP416; Kyushu-Palau Ridge).

Hymenocephalus hachijoensis: Okamura 1984b:92, pl. 80, fig. F (compiled); Okamura 1988:92, pl. 80, fig. F (compiled); Sazonov & Iwamoto 1992:62 (description; 3 additional spec. from Emperor Seamounts and Kyushu-Palau Ridge; comparisons with similar congeners); Nakabo 1993:355 (in key; Japan); Sazonov 1994:101 (brief description); Nakabo 2000:419 (in key; Japan); Nakabo 2002:419 (in key; Japan); Nakabo & Kai 2013:495 (in key; Japan); Schwarzhans 2014:49, fig. 23 (description; comparison; in key); Nakayama *et al.* 2015:505, figs. 1–5 (comparative material; 2 additional spec. from Hachijo-jima Island and Kyushu-Palau Ridge); Motomura 2020:39 (listed; Japan).

Diagnosis. A species of *Hymenocephalus* with 7 or 8 pelvic-fin rays; barbel moderately long, length 36–52% PRL, its tip not reaching vertical through hind margin of orbit when depressed; snout high, not depressed, length 25–30% PRL; orbit diameter 43–52% PRL; interorbital width 31–34% PRL; first dorsal-fin rays II,9–11; pectoral-fin rays i13–i15; gill rakers on first arch (outer/inner) 12–15/18–22, on second arch 18–20/16–20; pyloric caeca 23; dorsal half of trunk prominently blackish, but the dark area abruptly ending posterior to end of first dorsal-fin base; ventral half of tail immaculate or only slightly peppered

with small black melanophores; caudal vertebrae barely visible from outside; ostial and caudal colliculi of otolith not fused.

Material examined. 8 specimens. **Holotype of *Hymenocephalus striatissimus hachijoensis*:** BSKU 14171 (32.4 mm HL, 196+ mm TL), northeast of Hachijo-jima Island, Shichito-Iojima Ridge, Japan, 33.2681°N, 140.0517°E, 570 m, coll. O. Okamura, 21 Apr. 1968.

Paratype of *H. s. hachijoensis*: BSKU 14172 (22.6 mm HL, 128+ mm TL), collected with holotype. **Non-types:** **Japan:** ZMUT KP416 (1, 25.0 mm HL, 106+ mm TL), Kyushu-Palau Ridge, date unknown; BSKU 19308 (1, 24.7 mm HL, 133+ mm TL), off Hachijo-jima Island, Shichito-Iojima Ridge, 33.1367°N, 139.9900°E, 440 m, FRV *Soyo-maru*, sta. B3, beam trawl, 8 Dec. 1962; *FAKU 202929 (1, 22 mm HL, 140+ mm TL), FAKU 202930 (1, 23.1 mm HL, 141+ mm TL), *FAKU 202944 (1, 19 mm HL, 133+ mm TL), Nishiura fish market, 200–300 m, bottom trawl, coll. N. Nakayama, 21 May 2016. **Emperor Seamounts:** ZMMGU P.18243 (1, 32.2 mm HL, 200 mm TL), ca. 32°N, 173°E, *Mys Yunony*, tr. 86, trawl, 6 Sept. 1979.

Counts and measurements. Based on 6 specimens (22.6–32.4 mm HL, 106+–200+ mm TL). Counts: first dorsal-fin rays II,9–11; pectoral-fin rays i13–i15; pelvic-fin rays 7–8; gill rakers on first arch (outer/inner) 12–15/18–22, on second arch 18–20/16–20; longitudinal scales 17; transverse scale rows below first dorsal-fin origin 3.5, below first dorsal-fin midbase 3, below second dorsal-fin origin 2; pyloric caeca 23.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 21–24 (25–30); orbit diameter 34–42 (43–52); postorbital length 44–49 (54–60); postrostral length 80–85; orbit–preopercle distance 40–49 (49–58); suborbital width 8–10 (10–12); upper-jaw length 51–59 (61–70); length of rictus 47–53 (57–62); length of premaxillary tooth band 43–49 (53–58); preoral length 11–17 (14–20); distance between tip and lateral angle of snout 14–17 (17–21); snout width 24–31 (29–38); internasal width 19–24 (23–29); interorbital width 26–27 (31–34); body width over pectoral-fin bases 35–46 (43–57); body depth at first dorsal-fin origin 64–80 (80–94); body depth at anal-fin origin 49–54 (61–67); prepelvic length 102–113 (123–140); preanus length 169 (199); preanal length 166–172 (203–213); isthmus–pelvic distance 49–62 (58–76); isthmus–anal distance 123–124 (151–156); pelvic–anal distance 69–74 (85–88); pelvic-fin length 76–87 (95–103); pectoral-fin length 48–61 (56–75); predorsal length 100–110 (123–135); height of first dorsal fin 78–85 (92–107); length of first dorsal-fin base 33–42 (41–49); interdorsal length 63–88 (79–109); length of gill slit 31–34 (36–42); length of posterior nostril 4 (6); barbel length 30–42 (36–52).

Size. To about 20 cm TL (ZMMGU P.18243, 200 mm TL, Emperor Seamounts).

Distribution. So far known only from Japan

(Appendix 3-8A) and the Emperor Seamounts at depths of 200–710 m (Sazonov 1994; Schwarzhans 2014; this study). Rare.

Remarks. For further morphological information see the original description (Okamura 1970a) and a supplementary description given by Schwarzhans (2014).

Hymenocephalus hachijoensis was initially described as a subspecies of *H. striatissimus* Jordan & Gilbert in Jordan & Starks, 1904 based on two specimens collected from off Hachijo-jima Island, Shichito-Iojima Ridge, Japan, at a depth of 570 m (Fig. 136). Subsequently, it was recognized as a full species by Okamura (1984b, 1988), who considered *H. hachijoensis* to have a much larger head, a longer barbel (about equal to the orbit diameter), and a smaller orbit as compared with *H. striatissimus*. Based on examinations of three additional specimens from the Emperor Seamounts and Kyushu-Palau Ridge, Sazonov & Iwamoto (1992:62) and Sazonov (1994) confirmed a clear separation between the two species. Another specimen from the Kyushu-Palau Ridge (Fig. 135A) originally recorded as *H. longiceps* by Yatou (1982) was re-identified as *H. hachijoensis* by Nakayama *et al.*

(2015a); they also reported one additional specimen from the type locality (BSKU 19308, 24.7 mm HL, 133+ mm TL). Recently, three additional specimens were obtained at the Nishiura fish market in Aichi Pref., southern Japan; these specimens were originally collected from off Izu-oshima Island, at a depth of 200–300 m (FAKU 202929–202930, 202944, 19–23.1 mm HL, 133+–141+ mm TL; Fig. 135B). *Hymenocephalus hachijoensis* appears to be restricted to oceanic elevations (such as seamounts and ridges) in the northwestern Pacific.

Relationships and comparisons. *Hymenocephalus hachijoensis* belongs to the *H. antraeus* group as defined by Schwarzhans (2014). This group comprises the following four species: *H. antraeus* Gilbert & Cramer, 1897 endemic to Hawaii; *H. hachijoensis*; *H. heterolepis* (Alcock, 1889) restricted to the Andaman Sea and the Bay of Bengal; and *H. punt* Schwarzhans, 2014 recently described from the Gulf of Aden. According to Schwarzhans (2014), *H. hachijoensis* is readily distinguished from other species of the *H. antraeus* group by having lower counts of pelvic-fin rays (7–8 vs. 10–12) and a longer barbel [30–42% HL (based on this study) vs. ≤15%].

In Japan, *H. hachijoensis* is most similar to *H.*

FIGURE 135. Fresh specimens of *Hymenocephalus hachijoensis*. (A) ZMUT KP416, 25.0 mm HL, 106+ mm TL, Kyushu-Palau Ridge, Pacific, 685–710 m depth; (B) FAKU 202929, 147+ mm TL, Nishiura fish market, Aichi Pref., Pacific. Lateral views. [Photos: (A) ZMUT; (B) N. Nakayama]

FIGURE 136. Holotype of *Hymenocephalus striatissimus hachijoensis* (= *H. hachijoensis*). BSKU 14171, 32.4 mm HL, 196+ mm TL, northeast of Hachijo-jima Island, Shichito-Iojima Ridge, Pacific, 570 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

striatissimus in general appearance, but differs notably in having an otolith with two colliculi (vs. single in *H. striatissimus*; see also Schwarzhans 2014: figs. 23C–D and 32D–F). It further differs from *H. striatissimus* in its longer barbel (36–52% PRL vs. 14–23%) and pelvic fin (95–103% PRL vs. 70–90%). *Hymenocephalus hachijoensis* somewhat resembles *H. yamasakiorum* Nakayama, Endo & Schwarzhans, 2015 in physiognomy, but differs in having lower counts of pectoral- (i13–i15 vs. i16–i17) and pelvic-fin rays (7–8 vs. 9), and higher counts of pyloric caeca (23 vs. 15). These two species are also distinguished from each other by the snout length (25–30% PRL in *H. hachijoensis* vs. 37% in *H. yamasakiorum*) and barbel length (36–52% PRL vs. 73%).

***Hymenocephalus lethonemus* Jordan & Gilbert in Jordan & Starks, 1904**

[Japanese name: Itodara]

(Figs. 137–138; Appendix 3-8B)

Hymenocephalus lethonemus Jordan & Gilbert in Jordan & Starks, 1904:615, unnumbered fig. [original description; holotype: USNM 50936, from Sagami Bay, *Albatross* sta. 3697, in 120–265 ftn (220–485 m); paratypes from Tokyo market and Sagami Bay]; Jordan *et al.* 1913:413, fig. 383 (listed; Japan; new Japanese name: “Ito-dara”); Gilbert & Hubbs 1916:188 (brief description; 22 spec. from East China Sea, off Cape Shionomisaki, Suruga Bay, and Sagami-nada); Okada & Matsubara 1938:452 (in key; Japan); Matsubara 1955:1315 (in key; Japan); Kamohara 1964:96 (listed; Kochi Pref.); Matsubara 1965:507 (compiled; Japan); Okamura 1970a:54, pl. II, text-fig. 23 (description; biological notes; 83 spec. from Pacific off southern Japan from Sagami Bay to Kochi and East

China Sea); Okamura 1970b: table 1 (listed; Japan); Kuroda 1971:56 [listed from Suruga Bay after Okamura (1970a)]; Tominaga & Uyeno 1981:489 (listed; Japan); Yatou 1982:143, 347, fig. 87 [brief description; 7 spec. from Kyushu-Palau Ridge and Tosa Bay; photo based on BSKU spec. (catalog no. unknown)]; Ohta 1983: table A (listed; Suruga Bay); Okamura 1984a:201, 358, fig. 142 (brief description; 13 spec. from Okinawa Trough; photo based on BSKU 29790); Okamura 1984b:92, pl. 80, fig. G (compiled); Okamura 1988:92, pl. 80, fig. G (compiled); Nakabo 1993:355 (in key; Japan); Shinohara *et al.* 1996:169 (25 spec. listed from Pacific off Tohoku); Okamura 1997:128, fig. 2 (compiled); Shinohara & Matsuura 1997:291 (listed; Suruga Bay); Nakabo 2000:419 (in key; Japan); Shinohara *et al.* 2001:305 (19 spec. listed from Tosa Bay); Nakabo 2002:419 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Chiou *et al.* 2004a:44, fig. 12, table 1 (brief description; 13 spec. from Da-xi; first record from Taiwan); Shinohara *et al.* 2005:417 (1 spec. listed from Ryukyu Islands); Suetsugu & Ohta 2005: tables 2–3 (listed; southeast of Boso Peninsula and Enshu-nada); Senou *et al.* 2006:421 (listed; Sagami Sea); Shinohara & Williams 2006:551, fig. 2K (listed; Sagami Sea); Kitagawa *et al.* 2008:39, unnumbered fig. (brief description; spec. from Pacific off Tohoku); Shao *et al.* 2008b: table 2 (21 spec. listed from northeastern and southwestern Taiwan, and South China Sea); Shinohara *et al.* 2009:708 (listed; Pacific off Tohoku); Motomura *et al.* 2010:77, fig. 34 (5 spec. listed from Yakushima Island, Kagoshima Pref.); Furuhashi *et al.* 2010: table 2 (5 spec. listed from northern Okinawa Trough); Nakabo & Kai 2013:495 (in key; Japan); Schwarzhans 2014:42, fig. 18 (description; comparison; in key); Iwamoto *et al.* 2015:75 (brief description; 36 spec. from northeastern and southwestern Taiwan, and Philippines); Motomura 2020:39 (listed; Japan).

Hymenocephalus striatissimus (not Jordan & Gilbert in Jordan & Starks 1904): Okamura 1997:128, fig. 3 [in part, photo spec. (BSKU 41914)].

Diagnosis. A species of *Hymenocephalus* with 11 (rarely 12) pelvic-fin rays; chin barbel absent or rudimentary, length less than 1% PRL; snout high, not depressed, length 37–44% PRL; orbit diameter 43–46% PRL; interorbital width 38–41% PRL; orbit–preopercle distance 55–60% PRL; suborbital width 16–17% PRL; first dorsal-fin rays II,11; pectoral-fin rays i14–i17; gill rakers on first arch (outer/inner) 17–20/24–26, on second arch 24–26/22–25; numerous tiny black melanophores densely and uniformly scattered over trunk and tail, giving dusky appearance to body; caudal vertebrae barely visible externally; ostial and caudal colliculi of otolith not fused.

Material examined. 8 specimens. **Holotype** of *Hymenocephalus lethonemus*: USNM 50936 (25.9 mm HL, 140 mm TL), off Manazuru, Sagami Bay, Japan, 35.1333°N, 139.1667°E, 120–265 ftm (220–485 m), *Albatross* sta. 3697, 8-ft Tanner beam trawl, 5 Mar. 1900. **Paratypes** of *H. lethonemus*: USNM 51455 (3, 15.7–23.9 mm HL, 86+–126+ mm TL, counts and measurements taken on 1 of 3), off Numazu, Suruga Bay, 35.0333°N, 138.7833°E, 63–75 ftm (116–138 m), *Albatross* sta. 3707, 8-ft Tanner beam trawl, 8 Mar. 1900. **Non-type: Japan:** BSKU 109012 (1, 27.2 mm HL, 112+ mm TL), off Amami, East China Sea, 28.5513°N, 127.0375°E, 621 m, T/V *Nagasaki-maru*, cr. N365, sta. J6, tr. 3, 3-m ORE beam trawl, coll. N. Nakayama, 16 Nov. 2012; *BSKU 44762 (1, 30.7 mm HL, 155+ mm TL), Tosa Bay, 600 m, FRV *Kotaka-maru*, otter trawl, 16 May 1988; BSKU 13348 (1, 30.4 mm HL, 169+ mm TL), west of Muroto, Tosa Bay, 350–450 m, 14 May 1968; BSKU 44766 (1, 26.5 mm HL, 153+ mm TL), Tosa Bay, 600 m, FRV *Kotaka-maru*, otter trawl, 16 May 1988.

Counts and measurements. Based on 5 specimens (23.9–30.4 mm HL, 112+–169+ mm TL). Counts: first dorsal-fin rays II,11; pectoral-fin rays i14–i17; pelvic-fin rays 11; gill rakers on first arch (outer/inner) 17–20/24–26, on second arch 24–26/22–25; longitudinal scales 17–19; transverse scale rows below first dorsal-fin origin 4.5–5, below first dorsal-fin midbase 2–2.5, below second dorsal-fin origin 3–3.5.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 28–32 (37–44); orbit diameter 32–35 (43–46); postorbital length 45–46 (60–63); postrostral length 74–76; orbit–preopercle distance 41–45 (55–60); suborbital width 12–13 (16–17); upper-jaw length 51–54 (68–71); length of rictus 44–45 (59–60); length of premaxillary tooth band 39 (52–53); preoral length 14–17 (19–23); distance between tip and lateral angle of snout 18–22 (23–30); snout width 25–30 (33–41); internasal width 19–24 (26–32); interorbital width 28–31 (38–41); occipital width 13–14 (17–19); body width over pectoral-fin bases 41–55 (54–73); body

depth at first dorsal-fin origin 66–68 (89–92); body depth at anal-fin origin 41–49 (54–66); prepelvic length 100–121 (132–162); preanus length 151–158 (199–212); preanal length 152–161 (200–215); isthmus–pelvic distance 43–57 (58–77); isthmus–anus distance 87 (117); isthmus–anal distance 89–109 (120–143); pelvic–anal distance 46–58 (60–76); pelvic-fin length 82–92 (110–120); pectoral-fin length 55–63 (73–84); predorsal length 96–105 (129–143); height of first dorsal fin 74–81 (97–109); length of first dorsal-fin base 30–36 (40–47); interdorsal length 59–66 (78–90); length of gill slit 30–33 (39–45); length of posterior nostril 4–7 (5–9); barbel length 0–1 (0–1).

Size. To about 19 cm TL (BSKU 33501, 185+ mm TL, Okinawa Trough, Japan).

Development. Early life stage was briefly described by Endo (2014), based on three specimens (7.5–9.5 mm HL) collected from Sagami Bay.

Distribution. Restricted to Japan and Taiwan. Known from the Pacific off Japan northward to the Oshika Peninsula (38.32°N), Okinawa Trough, northern Taiwan, Izu Islands, and Kyushu–Palau Ridge, at depths of 116–1000 m (Chiou *et al.* 2004a; Shao *et al.* 2008a, 2008b; Schwarzhans 2014; Iwamoto *et al.* 2015; Appendix 3-8B). This species is well represented on the upper continental slope along the Pacific coasts of Japan, and is one of the numerically dominant grenadiers in Tosa Bay, Kumano-nada, and off the Boso Peninsula (Suetsugu & Ohta 2005; Horikawa 2009: appendix 3).

Remarks. For further morphological information see the original description (Jordan & Gilbert in Jordan & Starks 1904) and Okamura (1970a); otolith morphology was detailed by Schwarzhans (2014).

Hymenocephalus lethonemus was described on the basis of specimens collected from Sagami Bay (type locality) and “Tokyo market”, Japan (Fig. 138). Although Weber (1913) reported *H. lethonemus* from Indonesian waters, Gilbert & Hubbs (1920:535) considered Weber’s specimens to be representatives of their new species, *H. nascens*, described from the Philippines and East Indies. Gilbert & Hubbs (1920) distinguished these taxa solely by modal counts of pelvic-fin rays [usually 12 (very rarely 11 or 13) in *H. nascens* vs. usually 11 (12) in *H. lethonemus*]. Subsequently, Weber & de Beaufort (1929:61) regarded *nascens* as a subspecies of *H. lethonemus*, noting that “we are no more able to discover any other difference and feel therefore not authorized to separate the specimens from the Archipelago [viz., *nascens*; note by the present author] specifically from those from Japan, known as *H. lethonemus*”. While all subsequent authors recognized each taxon as a full species (e.g., Okamura 1970a; Marshall & Iwamoto 1973; Iwamoto 1990; Iwamoto & Merrett 1997; Iwamoto & Williams 1999; Schwarzhans 2014; Iwamoto *et al.* 2015), further morphological studies are required to elucidate their taxonomic status, particularly based on well-preserved materials.

Relationships and comparisons. *Hymenocephalus*

FIGURE 137. Fresh specimens of *Hymenocephalus lethonemus*. (A) BSKU 44762, 30.7 mm HL, 155+ mm TL, Tosa Bay, Kochi Pref., Pacific, 600 m depth; (B) BSKU 109012, 27.2 mm HL, 112+ mm TL, off Amami-oshima Island, Kagoshima Pref., Okinawa Trough, East China Sea, 621 m depth. Lateral views. [Photos: (A) BSKU; (B) N. Nakayama]

FIGURE 138. Holotype of *Hymenocephalus lethonemus*. USNM 50936, 25.9 mm HL, 140 mm TL, Sagami Bay, Pacific, 220–485 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

lethonemus belongs to the *H. striatulus* group as defined by Schwarzhans (2014). In addition to *H. lethonemus*, this group includes: *H. billsam* Marshall & Iwamoto, 1973 disjunctly distributed in the northwestern Atlantic and off Brazil; *H. nascens* widespread in the East Indies and Australasian waters; and *H. striatulus* Gilbert, 1905 known only from Hawaii and the Sala y Gomez Ridge. According to Schwarzhans (2014), *H. lethonemus* substantially differs from the above three species in that its pelvic-fin rays usually number 11 [rarely 12 vs. 12–15 (rarely 11 in *H. nascens*)]. It further differs from all except *H. nascens* in its absence of the chin barbel (vs. a small but distinct barbel present in *H. billsam* and *H. striatulus*). The reader refers to Schwarzhans (2014) for further comparisons of the *H. striatulus* group.

Due to their close resemblance, *H. lethonemus* may be confused with *H. papyraceus* Jordan & Gilbert in Jordan & Starks, 1904 in Japanese waters. They share the absence of the chin barbel and similar pigmentation pattern consisting of small black melanophores densely scattered over the trunk and tail. However, *H. lethonemus* readily differs from *H. papyraceus* in having more gill rakers (24–26 vs. 18–22 on the outer side of the second arch; 22–25 vs. 15–21 on the inner side of the second arch). It further differs from *H. papyraceus* in its narrower suborbital space (16–17% PRL vs. 22–26%), shorter pectoral fin (73–84% PRL vs. 95%), longer pelvic fin (110–120% vs. 79%), and shorter, lower, and narrower snout (preoral length 19–23% PRL vs. 28–39%; snout width 33–41% PRL vs. 44–53%; internasal width 26–32% PRL vs. 34–39%).

***Hymenocephalus longibarbis* Günther, 1887**

[Japanese name: Wanidara]

(Figs. 139–140; Appendix 3-8C)

Hymenocephalus longibarbis Günther, 1887:139, pl. XVIII, fig. C [original description; holotype: BMNH 1887.12.7.94, from Fiji Islands, *Challenger* sta. 173, 19°09'35"S, 179°41'50"E, in 315 ftm (576 m)]; Schwarzhans 2014:27, fig. 8 (description; comparison; *H. longiceps* a junior synonym).

Hymenocephalus longiceps Smith & Radcliffe in Radcliffe, 1912:111 pl. 23, fig. 3 [original description; holotype: USNM 72927, from off southeastern Luzon Island, *Albatross* sta. 5459, 13°10'21"N, 123°59'54"E, in 201 ftm (368 m)]; Kamohara 1937:188 (first record from Japan; brief description; spec. from Kochi; new Japanese name: "Wanidara"); Okada & Matsubara 1938:451 (in key; Japan); Kamohara 1938:70 (brief description; spec. from Kochi Pref.); Kamohara 1950:278 (listed; Kochi and Wakayama Pref.); Chen 1951:210 (listed; Taiwan); Kamohara 1952:97 (brief description; spec. from Kochi Pref.); Matsubara 1955:1314 (in key; Japan); Kamohara 1958:73 (listed; Kochi Pref.); Kamohara 1964:96 (listed; Kochi Pref.); Matsubara 1965:507 (compiled; Japan); Okamura 1970a:45, pl. XIV (description;

9 spec. from Pacific off southern Japan off Owase and Kochi); Okamura 1970b: table 1 (listed; Japan); Tominaga & Uyeno 1981:489 (listed; Japan); Yatou 1982:140 [brief description; in part, 4 spec. from Tosa Bay; 1 spec. from Kyushu-Palau Ridge is *H. hachijoensis* (ZMUT KP416)]; Ozawa 1983:13 (listed; off Makurazaki, Kagoshima Pref., East China Sea); Okamura 1984a:199, 357, fig. 140 [brief description; 5 spec. from Okinawa Trough; photo based on BSKU spec. (catalog no. unknown)]; Okamura 1984b:92, pl. 80, fig. D (compiled); Okamura 1988:92, pl. 80, fig. D (compiled); Nakabo 1993:355 (in key; Japan); Shao 1993:171 (compiled; Taiwan); Okamura 1997:128, fig. 1 (compiled); Nakabo 2000:419 (in key; Japan); Shinohara *et al.* 2001:305 (listed; Tosa Bay); Nakabo 2002:419 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Chiou *et al.* 2004a: table 1 (listed; Taiwan); Shinohara *et al.* 2005:417 (listed; Ryukyu Islands); Shao *et al.* 2008b: table 2 (41 spec. listed from northeastern and southwestern Taiwan, and South China Sea); Furuhashi *et al.* 2010: table 2 (2 spec. listed from northern Okinawa Trough); Nakabo & Kai 2013:495 (in key; Japan); Iwamoto *et al.* 2015:75 (brief description; 47 spec. from northeastern and southwestern Taiwan, South China Sea, and Philippines); Iwatsuki *et al.* 2017:32 (listed; Hyuga-nada); Motomura 2020:39 (listed; Japan).

Diagnosis. A species of *Hymenocephalus* with 8 pelvic-fin rays; barbel very long, length 53–72% PRL, its tip extending to vertical through hind margin of orbit or beyond when depressed; snout low, depressed, length 26–33% PRL; tip of snout scarcely, if at all, protruding beyond upper jaw; orbit diameter 35–42% PRL; interorbital width 18–27% PRL; first dorsal-fin rays II,9–10; pectoral-fin rays i12–i16; gill rakers on first arch (outer/inner) 13–18/19–23, on second arch 18–22/18–21; pyloric caeca 18–25; body uniformly dark gray dorsally; lower half of tail generally immaculate; caudal vertebrae not visible externally; ostial and caudal colliculi of otolith not fused.

Material examined. 30 specimens. **Japan:** BSKU 106798 (1, 40.9 mm HL, 219+ mm TL), BSKU 106799 (1, 27.6 mm HL, 156+ mm TL), BSKU 106800 (1, 37.0 mm HL, 202+ mm TL), southeast of Shimokoshiki-jima Island, East China Sea, 31.5560°N, 129.8847°E, 415 m, F/V *Maruko-maru*, tr. 4, bottom trawl, coll. N. Nakayama *et al.*, 24 Apr. 2012; KAUM-I. 32209 (1, 37.4 mm HL, 225 mm TL), off Nomaie, Kagoshima, 31.5000°N, 129.8833°E, 370–400 m, trawl, coll. M. Yamashita and Y. Ohashi, 10 Sept. 2010; FFNU P-00623 (2, 28.6–33.3 mm HL), east of Okihirazi Bank, East China Sea, 32.2635°N, 129.5428°E, 263 m, T/V *Nagasaki-maru*, cr. N207, sta. 1, 3-m ORE beam trawl, coll. J. Hashimoto, 11 Nov. 2005; FFNU P-00707 (2, 32.8–44.8 mm HL), east of Okihirazi Bank, East China Sea, 32.2260°N, 129.5293°E, 402 m, T/V *Nagasaki-maru*, cr. N254, sta. B4-1, 3-m ORE beam trawl, coll. J. Hashimoto, 12 Mar. 2008; FFNU P-00777 (1, 35.4 mm HL), east of Hirazi Bank, East China Sea, 32.2485°N, 129.4720°E, 403 m, T/V *Nagasaki-maru*, cr.

FIGURE 139. Fresh specimens of *Hymenocephalus longibarbis*. (A) BSKU 92382, 44.7 mm HL, 239+ mm TL, Mimase fish market, Kochi Pref., Pacific; (B–C) BSKU 112441, 51.1 mm HL, 267+ mm TL, Tosa Bay off Susaki, Kochi Pref., Pacific. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

FIGURE 140. Scanning electron micrographs showing a body scale (from the chest) of *Hymenocephalus longibarbis*. BSKU 113072, 42.6 mm HL. (A) View from above; (B) oblique view. [Photos: NSMT]

N276, sta. B4, 3-m ORE beam trawl, coll. J. Hashimoto, 7 Mar. 2009; BSKU 113721 (1, 28.0 mm HL, 165+ mm TL), BSKU 113723 (1, 35.8 mm HL, 208+ mm TL), BSKU 113725 (1, 23.6 mm HL, 141+ mm TL), off Aki, Tosa Bay, 320–400 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 8 Apr. 2014; BSKU 92382 (1, 44.7 mm HL, 239+ mm TL), Mimase fish market, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama, 20 Nov. 2007; BSKU 112440 (1, 43.0 mm HL, 235+ mm TL),

BSKU 112441 (1, 51.1 mm HL, 267+ mm TL), BSKU 112443 (1, 40.4 mm HL, 223 mm TL), BSKU 112446 (1, 45.6 mm HL, 233+ mm TL), BSKU 112448 (1, 44.7 mm HL, 236+ mm TL), BSKU 112449 (1, 50.6 mm HL, 264+ mm TL), BSKU 112901 (1, 42.3 mm HL, 239+ mm TL), BSKU 112914 (1, 42.2 mm HL, 213+ mm TL), BSKU 112917 (1, 32.9 mm HL, 190+ mm TL), BSKU 113071 (1, 33.5 mm HL, 206 mm TL), BSKU 113072 (1, 42.6 mm HL, 233+ mm TL), BSKU 113075 (1, 31.5

mm HL, 190+ mm TL), off Susaki, Tosa Bay, 320–380 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 9 Mar. 2014; BSKU 113077 (1, 49.5 mm HL, 265+ mm TL), BSKU 113078 (1, 46.8 mm HL, 237+ mm TL), BSKU 113082 (1, 35.1 mm HL, 199+ mm TL), BSKU 113083 (1, 39.7 mm HL, 225+ mm TL), BSKU 113652 (1, 36.8 mm HL, 213 mm TL), off Okitsu, Tosa Bay, 380 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 11 Mar. 2014.

Counts and measurements. Based on 30 specimens (23.6–51.1 mm HL, 141+–267+ mm TL). Counts: first dorsal-fin rays II,9–10; pectoral-fin rays i12–i16; pelvic-fin rays 8; gill rakers on first arch (outer/inner) 13–18/19–23, on second arch 18–22/18–21; longitudinal scales 15–18; transverse scale rows below first dorsal-fin origin 2.5–4, below first dorsal-fin midbase 2–3, below second dorsal-fin origin 2–3.5, above anal-fin origin 6–10; pyloric caeca 18–25; rete-gas gland complexes 2.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 21–25 (26–33); orbit diameter 28–33 (35–42); postorbital length 46–52 (59–65); postrostral length 77–81; orbit–preopercle distance 41–45 (52–57); suborbital width 8–11 (10–14); upper-jaw length 52–58 (66–73); length of rictus 45–50 (57–64); length of premaxillary tooth band 42–48 (53–61); preoral length 5–12 (7–15); distance between tip and lateral angle of snout 12–19 (15–24); snout width 19–25 (25–32); internasal width 14–19 (18–24); interorbital width 14–21 (18–27); occipital width 8–11 (10–14); body width over pectoral-fin bases 45–67 (57–84); body depth at first dorsal-fin origin 65–86 (81–107); body depth at anal-fin origin 51–61 (65–78); prepelvic length 100–117 (126–151); preanus length 176 (225); preanal length 164–184 (209–233); isthmus–pelvic distance 54–68 (68–87); isthmus–anus distance 120 (154); isthmus–anal distance 122–145 (154–184); pelvic–anal distance 68–90 (87–115); pelvic-fin length 85–122 (109–153); pectoral-fin length 47–56 (58–72); predorsal length 98–109 (122–139); height of first dorsal fin 59–85 (74–107); length of first dorsal-fin base 34–47 (44–58); interdorsal length 74–100 (93–127); length of gill slit 30–35 (38–44); length of posterior nostril 1–7 (2–9); barbel length 42–56 (53–72).

Size. To about 27 cm TL (BSKU 112441, 267+ mm TL, Mimase, Japan).

Distribution. Widely distributed from Japan to Australia, including the Philippines, Indonesia, Fiji, and the New Caledonian region, at depths of 200–820 m (Günther 1887; Radcliffe 1912; Gilbert & Hubbs 1920; Weber & de Beaufort 1929; Iwamoto & Merrett 1997; Iwamoto & Williams 1999; Merrett & Iwamoto 2000; Iwamoto & Graham 2001; Shao *et al.* 2008a, 2008b; Last *et al.* 2014; Schwarzhans 2014; Iwamoto *et al.* 2015; this study). In Japan, known from off the Pacific coasts northward to the Izu Peninsula (138.96°E) and the Okinawa Trough at depths of 200–555 m (Appendix 3-8C). Usually rare, but numerous specimens were occasionally collected in Tosa Bay.

Remarks. *Hymenocephalus longibarbis* has been well described by previous authors, and thus does not need repeating here. For a full description see Gilbert & Hubbs (1920; as *H. longiceps*), Okamura (1970a; as *H. longiceps*), and Iwamoto & Merrett (1997); otolith morphology was detailed by Schwarzhans (2014).

Hymenocephalus longibarbis was described by Günther (1887) based on a single specimen collected during the “*Challenger*” Expedition from the Fiji Islands. Subsequently, Smith & Radcliffe (*in* Radcliffe 1912) described *H. longiceps* from the Albay Gulf, the southeast of Luzon, the Philippines. Although these species have been considered valid by most previous authors (e.g., Okamura 1970a; Iwamoto 1990; Iwamoto & Merrett 1997; Iwamoto & Williams 1999), Schwarzhans (2014) showed that *H. longiceps* is a junior synonym of *H. longibarbis*; this study follows his treatment.

Relationships and comparisons. *Hymenocephalus longibarbis* is one of two species of the *H. longibarbis* group as defined by Schwarzhans (2014). The only other member of this group is *H. longipes* Smith & Radcliffe *in* Radcliffe, 1912 restricted to marginal seas of the Philippines. *Hymenocephalus longibarbis* differs most notably from *H. longipes* in the presence of a well-developed barbel at the symphysis of the lower jaw (53–72% PRL vs. absent). The two species are further distinguished from each other by the color of the pelvic fin (uniformly dusky in *H. longibarbis* vs. intensely black distally in *H. longipes*).

Hymenocephalus papyraceus Jordan & Gilbert *in* Jordan & Starks, 1904

[Japanese name: Kamidara]

(Figs. 141–142; Appendix 3-8D)

Hymenocephalus papyraceus Jordan & Gilbert *in* Jordan & Starks, 1904:614, unnumbered fig. [original description; holotype: USNM 50935, from Sagami Bay, *Albatross* sta. 3697, in 120–265 fms (220–485 m)]; Jordan *et al.* 1913:412, fig. 382 (listed; Japan; new Japanese name: “Kamidara”); Okada & Matsubara 1938:452 (in key; Japan); Matsubara 1955:1315 (in key; Japan); Matsubara 1965:508 (compiled; Japan); Okamura 1970a:56, pl. XVII, text-fig. 24 [description; 1 spec. from off the east coast of southern Japan, probably in Sagami or Suruga Bays (BSKU 14205)]; Okamura 1970b: table 1 (listed; Japan); Tominaga & Uyeno 1981:489 (listed; Japan); Okamura 1984b:92, pl. 344, fig. I (compiled); Okamura 1988:92, pl. 344, fig. I (compiled); Sazonov 1993:117 (East China Sea); Senou *et al.* 2006:421 (listed; Sagami Sea); Shinohara & Williams 2006:551, fig. 2-L (listed; Sagami Sea; misspelled “*papiraceus*” in figure legend); Nakabo & Kai 2013:495 (in key; Japan); Schwarzhans 2014:37, fig. 14F–J (description; comparison; in key); Iwamoto *et al.* 2015:76, fig. 12 (brief description; 1 spec. from Da-xi; first record from Taiwan); Iwatsuki *et al.* 2017:32 (listed; Hyuga-nada);

Motomura 2020:39 (listed; Japan).

Hymenocephalus papyraceus [sic]: Nakabo 1993:355 (in key; Japan); Nakabo 2000:419 (in key; Japan); Nakabo 2002:419 (in key; Japan).

Diagnosis. A species of *Hymenocephalus* with 11 or 12 pelvic-fin rays; barbel rudimentary, length 2–4% PRL; snout high, not depressed, length 41–52% PRL; orbit diameter 33–46% PRL; interorbital width 37–52% PRL; orbit–preopercle distance 59–71% PRL; suborbital width 22–26% PRL; first dorsal-fin rays II,9–11; pectoral-fin rays i12–i15; gill rakers on first arch (outer/inner) 13–18/19–24, on second arch 18–22/15–21; small but prominent black melanophores densely scattered over trunk and tail, giving dark appearance to body; caudal vertebrae barely visible from outside; ostial and caudal colliculi of otolith not fused.

Material examined. 9 specimens. **Holotype** of *Hymenocephalus papyraceus*: USNM 50935 (25.1 mm HL, 113+ mm TL), off Manazuru, Sagami Bay, Japan, 35.1333°N, 139.1667°E, 120–265 fm (220–485 m), *Albatross* sta. 3697, 8-ft Tanner beam trawl, 5 Mar. 1900. **Non-types: Japan:** BSKU 101959 (1, 32.3 mm HL, 139+ mm TL), east of Yoron-jima Island, 27.1133°N, 128.6983°E, 870–945 m, R/V *Hakuho-maru*, cr. KH-73-02, sta. 4, 3-m beam trawl, coll. S. Ohta, 24 Feb. 1973; ZMMGU P.18258 (1, 35.0 mm HL, 125+ mm TL), west of Yaku-shima Island, Okinawa Trough, 30.5167°N, 129.3167°E, 800–826 m, R/V *Vityaz'*, cr. 25, sta. 3768, *Galatea* trawl, date unknown; BSKU 44761 (1, 29.7 mm HL, 176 mm TL), Tosa Bay, 600 m, FRV *Kotaka-maru*, otter trawl, 16 May 1988; BSKU 44292 (1, 32.2 mm HL, 137+ mm TL), BSKU 44296 (1, 34.6 mm HL, 163+ mm TL), BSKU 44297 (1, 32.9 mm HL, 129+ mm TL), Tosa Bay, 600 m, FRV *Kotaka-maru*, otter trawl, 17 Nov. 1987; BSKU 20312 (1, 28.5 mm HL, 129+ mm TL), off Atami, Sagami Bay, 35.0783°N, 139.1467°E, 450 m, FRV *Soyo-maru*, sta. T10, beam trawl, 24 Nov. 1972; BSKU 99483 (1, 35.7 mm HL, 177+ mm TL), off Pacific coast of southern Japan, date unknown.

Counts and measurements. Based on 9 specimens (25.1–35.7 mm HL, 113+–177+ mm TL). Counts: first dorsal-fin rays II,9–11; pectoral-fin rays i12–i15; pelvic-fin rays 11–12; gill rakers on first arch (outer/inner) 13–18/19–24, on second arch 18–22/15–21; transverse scale rows below first dorsal-fin origin 3.5, below first dorsal-fin midbase 4.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 30–36 (41–52); orbit diameter 24–31 (33–46); postorbital length 45–50 (63–67); postrostral length 69–75; orbit–preopercle distance 43–49 (59–71); suborbital width 16–19 (22–26); upper-jaw length 45–49 (61–69); length of rictus 38 (53); length of premaxillary tooth band 36–37 (49–50); preoral length 20–29 (28–39); distance between tip and lateral angle of snout 21–26 (28–35); snout width 32–38 (44–53);

internasal width 24–29 (34–39); interorbital width 27–36 (37–52); body width over pectoral-fin bases 29–49 (40–66); body depth at first dorsal-fin origin 66–74 (91–103); body depth at anal-fin origin 40–46 (56–64); prepelvic length 105–122 (144–168); preanus length 152–168 (207–232); preanal length 154–171 (211–237); isthmus–pelvic distance 44–54 (60–74); isthmus–anus distance 105–120 (140–165); isthmus–anal distance 92–123 (126–169); pelvic–anal distance 53–74 (72–102); pelvic-fin length 59 (79); pectoral-fin length 65 (95); predorsal length 98–107 (135–152); length of first dorsal-fin base 27–34 (36–47); interdorsal length 43–59 (58–); length of gill slit 25–28 (34–39); barbel length 1–3 (2–4).

Size. To at least 18 cm TL (BSKU 99483, 177+ mm TL, off southern Japan).

Development. Endo (2014) provided a brief description of a juvenile (9.6 mm HL) collected from Sagami Bay.

Distribution. Restricted to Japan and Taiwan (Sazonov 1993; Schwarzhans 2014; Iwamoto *et al.* 2015; this study; Appendix 3-8D). Capture depths 220–945 m. Uncommon.

Remarks. For further morphological details see the original description (Jordan & Gilbert in Jordan & Starks 1904) and a redescription given by Sazonov (1993); otolith morphology was fully described by Schwarzhans (2014).

Hymenocephalus papyraceus was originally described from a single specimen collected from Sagami Bay (Fig. 142). Subsequently, Okamura (1970a) reported one additional specimen from “off the east coast of southern Japan” (probably from Sagami or Suruga Bays). Because his specimen (BSKU 14205) is badly deteriorated, it was impossible to confirm the identification with certainty. Sazonov (1993) also provided a redescription of the species based on a specimen collected from the East China Sea off Yaku-shima Island. During the course of this study, a total of 14 specimens was newly found among the collection of Kochi University (BSKU) and the Natural History Museum and Institute, Chiba (CBM), 11 of which were used in Schwarzhans' (2014) review of the genus. Iwamoto *et al.* (2015) also recorded the species from northern Taiwan, based on a single specimen collected from the Da-xi fish market.

Relationships and comparisons. *Hymenocephalus papyraceus* belongs to the *H. aterrimus* group as defined by Schwarzhans (2014). This group is substantially consistent with the subgenus *Papyrocephalus* Gilbert & Hubbs, 1920, and comprises the following six species: *H. aterrimus* Gilbert, 1905 (type species of *Papyrocephalus*) sporadically recorded from tropical and temperate waters of the world's oceans, including the Hawaiian Islands (type locality), east and west coasts of Australia, New Caledonian region, and northwestern Atlantic; *H. barbatulus* Gilbert & Hubbs, 1920 known only from the Philippines; *H. fuscus* McMillan & Iwamoto, 2014 recently described from New Zealand; *H. nesaeae*

FIGURE 141. Fresh specimen of *Hymenocephalus papyraceus*. BSKU 44761, 29.7 mm HL, 176 mm TL, Tosa Bay, Kochi Pref., Pacific, 700 m depth. Lateral view. [Photo: BSKU]

FIGURE 142. Holotype of *Hymenocephalus papyraceus*. USNM 50935, 25.1, 113+ mm TL, Sagami Bay, Pacific, 220–485 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

Merrett & Iwamoto, 2000 so far known from Vanuatu; *H. papyraceus*; and *H. sazónovi* Schwarzhans, 2014 probably endemic to the Nazca and Sala y Gomez Ridges.

Hymenocephalus papyraceus is unlikely to be confused with *H. barbatulus* and *H. nesaeae* by having 11–12 pelvic-fin rays [vs. 7 in *H. barbatulus*, 13–14 in *H. nesaeae*, fide Schwarzhans (2014: table 1)]. This species is somewhat similar to *H. aterrimus*, *H. fuscus*,

and *H. sazónovi* in having a uniformly dark tail, but it differs notably from the latter three by its unique pigmentation. Although the ground color of the tail is immaculate in *H. papyraceus*, small black melanophores are densely scattered over its surface; this intense pigmentation gives a distinctly dark appearance to the tail. Such melanophores are also found in *H. aterrimus*, *H. fuscus*, and *H. sazónovi*, but the ground color (=

epidermis except where the melanophores are present) is also uniformly dark, and thus their tail appears to be more blackish compared with that of *H. papyraceus*. For further comparisons see McMillan & Iwamoto (2014) and Schwarzhans (2014).

In Japan, *H. papyraceus* may be confused with *H. lethonemus* Jordan & Gilbert in Jordan & Starks, 1904, but the two species are separable by several meristic and morphometric characters (see the Relationships and comparisons of the latter species).

***Hymenocephalus striatissimus* Jordan & Gilbert in Jordan & Starks, 1904**

[Japanese name: Sujidara]

(Figs. 143–145; Appendix 3-8E)

Hymenocephalus striatissimus Jordan & Gilbert in Jordan & Starks, 1904:612, unnumbered fig. [original description; holotype: USNM 50934, from Suruga Bay, *Albatross* sta. 3738, in 167 ftn (306 m); 3 paratypes from Suruga Bay]; Jordan *et al.* 1913:411, fig. 381 (listed; Japan; new Japanese name: “Suji-dara”); Gilbert & Hubbs 1916:187 (brief description; 15 spec. from East China Sea, off Cape Shionomisaki, and Suruga Bay); Izuka & Matsuura 1920:103 (spec. from “Tōkyō Market”); Okada & Matsubara 1938:451 (in key; Japan); Kamohara 1950:277 (listed; Kochi and Wakayama Pref.); Kuroda 1951:392 (listed; Suruga Bay); Uchinomi 1951:76 (listed; southern part of Kii Peninsula); Kamohara 1952:96 (brief description; Kochi Pref.); Matsubara 1955:1315 (in key; Japan); Kobayashi 1956:72 (listed; Enshu-nada); Kamohara 1958:74 (listed; Kochi Pref.); Kamohara & Okamura 1958:32, pl. 31, fig. 151 (compiled; Japan); Okada *et al.* 1959:83 (listed; Kumano-nada); Kamohara 1964:96 (listed; Kochi Pref.); Tatara *et al.* 1965:121 (listed; off Shikoku); Matsubara 1965:507 (compiled; Japan); Yatou 1982:143, 346, fig. 86 [brief description; 7 spec. from Kyushu-Palau Ridge and Tosa Bay; photo based on BSKU spec. (catalog no. unknown)]; Ohta 1983: table A (listed; Suruga Bay); Ozawa 1983:13 (listed; off Makurazaki, Kagoshima Pref., East China Sea); Okamura 1984a:199, 358 [brief description; 25 spec. from Okinawa Trough; photo based on BSKU 36798 (from Mimase fish market)]; Okamura 1984b:92, pl. 80, fig. E (compiled); Okamura 1988:92, pl. 80, fig. E (compiled); Nakabo 1993:355 (in key; Japan); Shao 1993:171 (compiled; Taiwan); Shinohara *et al.* 1996:169 (5 spec. listed from Pacific off Tohoku); Okamura 1997:128, fig. 3 [compiled; photo spec. is *H. lethonemus* (BSKU 41914)]; Shinohara & Matsuura 1997:291 (listed; Suruga Bay); Nakabo 2000:419 (in key; Japan); Shinohara *et al.* 2001:305 (667 spec. listed from Tosa Bay); Nakabo 2002:419 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Nakajima 2003:53 (brief description; 1 spec. from Enshu-nada); Chiou *et al.* 2004a: table 1 (listed; Taiwan); Shinohara *et al.* 2005:417 (listed; Ryukyu Islands); Kitagawa *et al.* 2008:38, unnumbered fig. (brief description; spec. from Pacific off Tohoku); Shinohara *et al.* 2009:708 (listed; Pacific off Tohoku); Furuhashi *et al.* 2010:

table 2 (49 spec. listed from northern Okinawa Trough); Nakabo & Kai 2013:495 (in key; Japan); Schwarzhans 2014:60, fig. 32 (description; comparison; in key); Iwatsuki *et al.* 2017:32 (listed; Hyuga-nada); Motomura 2020:39 (listed; Japan).

Hymenocephalus striatissimus striatissimus: Chen 1951:210 (listed; Taiwan); Okamura 1970a:48, pl. XV (description; biological notes; 90 spec. from Pacific off southern Japan from Suruga Bay to Totoro and East China Sea); Okamura 1970b: table 1 (listed; Japan); Kataoka & Tomida 1981:78 (listed; Mie Pref.); Tominaga & Uyeno 1981:489 (listed; Japan); Ohta 1983: table B (listed; Suruga Bay); Akazaki 1984:265 (listed; Aoshima, Miyazaki Pref.); Suetsugu & Ohta 2005: tables 2–3 (listed; southeast of Boso Peninsula and Enshu-nada); Shao *et al.* 2008b: table 2 (34 spec. listed from northeastern and eastern Taiwan, and South China Sea); Iwamoto *et al.* 2015:77 (brief description; 42 spec. from northeastern and eastern Taiwan, and South China Sea).

Diagnosis. A species of *Hymenocephalus* with 8 or 9 pelvic-fin rays; barbel moderately short, length 14–23% PRL, its tip not reaching vertical through hind margin of orbit when depressed; snout high, not depressed, length 26–37% PRL; orbit diameter 48–59% PRL; interorbital width 32–40% PRL; first dorsal-fin rays II, 8–10; pectoral-fin rays i11–i16; gill rakers on first arch (outer/inner) 12–18/20–24, on second arch 19–23/16–23; dorsal surface of trunk prominently blackish below first dorsal-fin base; the dark area on dorsum trunk followed by faint dusky stripe forming clusters of small melanophores; lower half of tail lacking black pigmentation; body flesh translucent posteriorly on tail (only when fresh) where caudal vertebrae visible from outside; ostial and caudal colliculi of otolith fused.

Material examined. 32 specimens. **Holotype of *Hymenocephalus striatissimus*:** USNM 50934 (currently comprising 2 spec., 16.7–17.7 mm HL, 99+–112+ mm TL, 1 of which is holotype), off Heta, Shizuoka Pref., Suruga Bay, Japan, 34.9667°N, 138.7667°E, 167 ftn (306 m), *Albatross* sta. 3738, 8-ft Tanner beam trawl, 17 Mar. 1900. **Paratype of *H. striatissimus*:** CAS-SU 8549 (1, 23+ mm HL, 113+ mm TL), Suruga Bay, *Albatross* sta. 3721 or 3738, 8-ft Tanner beam trawl, Mar. 1900. **Non-types: Japan:** BSKU 26290 (1, 32.8 mm HL, 148+ mm TL), off Senkaku Islands, Okinawa Trough, 24.6833°N, 122.5933°E, 510–615 m, F/V *Yuryo-maru*, No. 8, tr. 2, bottom trawl, coll. T. Kitajima *et al.*, 13 Jan. 1978; BSKU 77294 (1, 29.7 mm HL, 170+ mm TL), south of Muroto, 33.0790°N, 134.1645°E, 414–513 m, R/V *Tansei-maru*, cr. KT-05-29, sta. K-500, 3-m ORE beam trawl, coll. H. Endo *et al.*, 17 Nov. 2005; BSKU 98924 (1, 32.6 mm HL, 201+ mm TL), BSKU 98931 (1, 35.8 mm HL, 213+ mm TL), Tosa Bay, 33.1428°N, 133.6248°E, 450–490 m, FRV *Kotaka-maru*, otter trawl, 1 Aug. 1988; BSKU 12934 (1, 31.1 mm HL, 187+ mm TL), Tosa Bay, 350–450 m, bottom trawl, 14 May 1968; BSKU 41903 (1, 32.7 mm HL, 199+ mm TL), BSKU 41904 (1, 35.1 mm HL, 217+ mm TL),

FIGURE 143. Fresh specimens of *Hymenocephalus striatissimus*. (A) BSKU 92222, 157 mm TL, Mimase fish market, Kochi Pref., Pacific; (B–D) BSKU 113966, 20.0 mm HL, 159+ mm TL, Tosa Bay off Susaki, Kochi Pref., Pacific. (A–B) Lateral views; (C) dorsal and (D) ventral views of the head and trunk. [Photos: N. Nakayama]

mm TL), BSKU 41906 (1, 33.5 mm HL, 204+ mm TL), BSKU 41909 (1, 32.7 mm HL, 189+ mm TL), Tosa Bay, 420–450 m, FRV *Kotaka-maru*, otter trawl, 6 Jun. 1985; BSKU 58543 (1, 29.8 mm HL, 191 mm TL), Mimase fish market, bottom trawl, 1 Feb. 2002; BSKU 70088 (1, 16.2 mm HL, 67+ mm TL), Mimase fish market, bottom trawl, 15 Jan. 2004; BSKU 53083 (1, 26.8 mm HL, 174+ mm TL), Mimase fish market, bottom trawl, 18 Dec. 2000; BSKU 58427 (1, 27.1 mm HL, 160+ mm TL), Mimase fish market, bottom trawl, 3 Dec. 2001; BSKU 78828 (1, 28.4 mm HL, 170+ mm TL), Mimase fish market, bottom trawl, 6 Apr. 2006; *BSKU 92222 (1, ca. 157 mm TL), Mimase fish market, F/V *Seiryō-maru*, bottom trawl, coll. N. Nakayama, 7 Nov. 2007; BSKU 94814 (1, 23.7 mm HL, 161 mm TL), BSKU 94815 (1, 19.0 mm HL, 92+ mm TL), Mimase fish market, F/V *Kosei-maru*, bottom trawl, 28 Nov. 2007; BSKU 112461 (1, 22.6 mm HL, 150 mm TL), BSKU 112462 (1, 21.9 mm HL, 118+ mm TL), off Susaki, Tosa Bay, 320–380 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 9 Mar. 2014; BSKU 113966 (1, 29.0 mm HL, 159+ mm TL), off Susaki, Tosa

Bay, 350–365 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 23 Apr. 2014; BSKU 112514 (1, 21.6 mm HL), BSKU 112517 (1, 24.6 mm HL, 160+ mm TL), BSKU 113656 (1, 23.1 mm HL, 145+ mm TL), BSKU 113658 (1, 21.3 mm HL, 138+ mm TL), BSKU 113659 (1, 26.3 mm HL, 153+ mm TL), BSKU 113660 (1, 20.3 mm HL, 132+ mm TL), off Okitsu, Tosa Bay, 380 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 11 Mar. 2014; *BSKU 43410 (1, 31.3 mm HL, 206+ mm TL), BSKU 43412 (1, 29.4 mm HL, 133+ mm TL), south of Okino-shima Island, 32.6400°N, 132.5367°E, 440–451 m, R/V *Tansei-maru*, cr. KT-86-16, sta. A-3, 2-m beam trawl, 1 Nov. 1986; BSKU 113665 (1, 27.3 mm HL, 174+ mm TL), Numazu fish market, coll. S. Ohashi and M. Takami, 16 Feb. 2010.

Counts and measurements. Based on 30 specimens (16.2–35.8 mm HL, 67+–217+ mm TL). Counts: first dorsal-fin rays II,8–10; pectoral-fin rays i11–i16; pelvic-fin rays 8–9; gill rakers on first arch (outer/inner) 12–18/20–24, on second arch 19–23/16–23; longitudinal scales 16–19; transverse scale rows below first dorsal-fin

FIGURE 144. Possible holotype of *Hymenocephalus striatissimus*. USNM 50934 (1 or 2 spec.), 16.7 mm HL, 99+ mm TL, Suruga Bay, Shizuoka Pref., Pacific., 306 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

FIGURE 145. Scanning electron micrographs showing a body scale (from the dorsum below the interdorsal space) of *Hymenocephalus striatissimus*. BSKU 43410, 31.3 mm HL. (A) View from above; (B) oblique view. [Photos: NSMT]

origin 2.5–4.5, below first dorsal-fin midbase 2–3, below second dorsal-fin origin 2–3, above anal-fin origin 5–8.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 21–28 (26–37); orbit diameter 39–46 (48–59); postorbital length 43–47 (52–57); postrostral length 75–83; orbit–preopercle distance 38–46 (47–59); suborbital width 10–12 (12–15); upper-jaw length 49–55 (62–67); length of rictus 43–47 (52–59); length of premaxillary tooth band 38–45 (47–57); preoral length 11–18 (14–22); distance between tip and lateral angle of snout 14–20 (17–25); snout width 27–31 (33–40); internasal width 17–25 (21–33); interorbital width 26–34 (32–40); occipital width 8–13 (10–16); body width over pectoral-fin bases 38–60 (47–77); body depth

at first dorsal-fin origin 57–87 (69–107); body depth at anal-fin origin 43–56 (53–72); prepelvic length 99–118 (123–153); preanus length 167 (201); preanal length 151–175 (185–221); isthmus–pelvic distance 46–63 (55–79); isthmus–anal distance 107–131 (134–163); pelvic–anal distance 53–76 (65–96); pelvic-fin length 57–70 (70–90); pectoral-fin length 51–74 (61–92); predorsal length 99–110 (121–145); height of first dorsal fin 64–95 (79–114); length of first dorsal-fin base 29–36 (36–46); interdorsal length 66–95 (81–119); length of gill slit 29–36 (35–43); length of posterior nostril 3–6 (3–8); barbel length 11–18 (14–23).

Size. To about 22 cm TL (BSKU 41904, 217+ mm TL, Tosa Bay, Japan).

Distribution. Known from the Pacific off Japan

northward to Iwaki (37.07°N, Fukushima Pref.), Kyushu-Palau Ridge, East China Sea northward to Tsushima Island, and northern South China Sea (Shao *et al.* 2008a, 2008b; Schwarzhans 2014; Iwamoto *et al.* 2015; this study; Appendix 3-8E). Schwarzhans (2014:60) also recorded the species from the Philippines based on a single specimen collected from off Jolo Island (ZMUC P375860). Depth range varies from 150 to 1188 m, but usually shallower than 500 m. Very abundant on the upper continental slope off southern Japan.

Remarks. For further morphological details see the original description (Jordan & Gilbert in Jordan & Starks 1904) and a redescription given by Okamura (1970a); otolith morphology was fully described by Schwarzhans (2014).

Gilbert & Hubbs (1920) recognized three subspecies under *H. striatissimus*, viz., *H. s. aeger* Gilbert & Hubbs, 1920, *H. s. striatissimus*, and *H. s. torvus* Smith & Radcliffe in Radcliffe, 1912. Subsequently, Okamura (1970a) described a fourth subspecies, *H. s. hachijoensis*. All these taxa currently deserve full species status (Schwarzhans 2014).

Comments on type specimens. *Hymenocephalus striatissimus* was described from three specimens collected from Suruga Bay. In the original description, Jordan & Gilbert (in Jordan & Starks 1904) designated a 108 mm long specimen as the holotype, giving the catalog number as USNM 50934. However, this lot currently contains two specimens of about equal size (16.7–17.7 mm HL, 99+–112+ mm TL; Fig. 144), and it is uncertain which specimen represents the holotype.

Relationships and comparisons. *Hymenocephalus striatissimus* belongs to the *H. striatissimus* group as defined by Schwarzhans (2014). The other members of this group are: *H. aeger* known from Indonesia, Papua New Guinea, and off Western Australia; *H. megalops* Iwamoto & Merrett, 1997 distributed in the New Caledonian region and off Queensland; and *H. torvus* probably endemic to the Philippines. According to Schwarzhans (2014), *H. striatissimus* differs primarily from *H. aeger* in having a shorter barbel [15–20% HL (11–18% in the specimens examined in this study) vs. 20–25%]; from *H. megalops* in having a smaller orbit [37–42% HL (39–46%) vs. 41–45%]; and from *H. torvus* in its higher counts of pelvic-fin rays [8 (8–9) vs. 7]. See Schwarzhans (2014) for further comparisons of these four species.

McMillan & Iwamoto (2014) recently described *H. maculicaudus* from New Zealand; this species appears to fall into the *H. striatissimus* group. *Hymenocephalus maculicaudus* is unique within the genus in having a distinctive mid-lateral line of black melanophores on the anterior portion of the tail (vs. absent in all other congeners). It further differs from *H. striatissimus* in its lower counts of gill rakers [16–20 (modally 18) vs. 20–24 (22) on the inner side of the first arch; 15–20 (17) vs. 19–23 (21) on the outer side of the second arch].

In Japanese waters, *H. striatissimus* is most similar

to *H. hachijoensis* Okamura, 1970 and *H. yamasakiorum* Nakayama, Endo & Schwarzhans, 2015 in general appearance, but differs notably from the latter two species in having a shorter barbel (14–23% PRL vs. 36–52% in *H. hachijoensis*, 73% in *H. yamasakiorum*) and an otolith with a single colliculum (vs. ostial and caudal colliculi not fused).

***Hymenocephalus yamasakiorum* Nakayama, Endo & Schwarzhans, 2015**

[Japanese name: Oguro-sujidara]

(Fig. 146; Appendix 3-8F)

Hymenocephalus yamasakiorum Nakayama, Endo & Schwarzhans, 2015a:505, figs. 1–4 (original description; holotype: BSKU 113692, from Tosa Bay off Aki, in 320–400 m; new Japanese name: “Oguro-sujidara”); Motomura 2020:39 (listed; Japan).

Diagnosis. A species of *Hymenocephalus* with 9 pelvic-fin rays; barbel very long, length 73% PRL, its tip extending beyond vertical through hind margin of orbit when depressed; snout high, not depressed, length 37% PRL; orbit diameter 43% PRL; interorbital space 31% PRL; first dorsal-fin rays II,10; pectoral-fin rays i16–i17; gill rakers on first arch (outer/inner) 14/20, on second arch 20/19; pyloric caeca 15; trunk blackish dorsally, but the dark area abruptly ending posterior to first dorsal-fin base; dorsal half of tail prominently darker posteriorly, but ventral half completely lacking black melanophores; caudal vertebrae barely visible externally when fresh; ostial and caudal colliculi of otolith not fused.

Material examined. 1 specimen. **Holotype of *Hymenocephalus yamasakiorum*:** BSKU 113692 (1, 26.5 mm HL, 151+ mm TL), off Aki, Kochi Pref., Tosa Bay, Japan, 320–400 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 8 Apr. 2014.

Counts and measurements. Counts: first dorsal-fin rays II,10; pectoral-fin rays i16–i17; pelvic-fin rays 9; gill rakers on first arch (outer/inner) 14/20, on second arch 20/19; pyloric caeca 15.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 28 (37); orbit diameter 33 (43); postorbital length 45 (59); postrostral length 76; orbit–preopercle distance 40 (52); suborbital width 9 (12); upper-jaw length 51 (67); length of rictus 46 (60); length of premaxillary tooth band 42 (55); preoral length 12 (16); distance between tip and lateral angle of snout 16 (21); snout width 26 (35); internasal width 18 (24); interorbital width 24 (31); body width over pectoral-fin bases 46 (61); body depth at first dorsal-fin origin 63 (82); body depth at anal-fin origin 49 (64); prepelvic length 105 (138); preanal length 162 (212); isthmus–pelvic distance 49 (64); isthmus–anal distance 105 (138); pelvic–anal distance 60 (79); pelvic-fin length 78 (102); pectoral-fin length 53 (70); predorsal

FIGURE 146. Holotype of *Hymenocephalus yamasakiorum*. BSKU 113692, 26.5 mm HL, 151+ mm TL, Tosa Bay off Aki, Kochi Pref., Pacific, 320–400 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Fresh condition. [Photos: reproduced from Nakayama *et al.* (2015a: fig. 1) with permission from the Ichthyological Society of Japan]

length 102 (133); height of first dorsal fin 82 (107); length of first dorsal-fin base 35 (46); interdorsal length 77 (101); length of gill slit 30 (39); length of posterior nostril 6 (8); barbel length 55 (73); length of pyloric caecum 14 (19).

Size. At least 15 cm TL.

Distribution. So far known only from Tosa Bay off Aki at a depth of about 320–400 m (Appendix 3-8F). Very rare.

Remarks, relationships, and comparisons.

Hymenocephalus yamasakiorum was recently described from a single specimen collected from Tosa Bay, Japan (Fig. 146), and it belongs to the *H. iwamotoi* group as defined by Schwarzhans (2014) (Nakayama *et al.* 2015a). In Japanese waters, the species might be confused with *H. longibarbis* by sharing a long chin barbel (73% and 53–72% PRL in *H. yamasakiorum* and *H. longibarbis* respectively), but they are readily separable by a difference in physiognomy: in *H. yamasakiorum*, the tip of the snout is high and moderately protrudes beyond the upper jaw giving a slightly pointed appearance to the snout (Fig. 146A), whereas the snout of *H. longibarbis* is low and greatly depressed, barely extending beyond the upper jaw (Fig. 139A). Body pigmentation is also different between the two species; in *H. yamasakiorum*, the dorsal half of the tail is abruptly darker posteriorly, but is uniformly dusky in *H. longibarbis*. Regarding meristic and morphometric characters, *H. yamasakiorum* differs from *H. longibarbis*, in having more pelvic- (9 vs. 8) and pectoral-fin rays (i16–i17 vs. i12–i16), fewer pyloric caeca (15 vs. 18–25), a longer snout (37% PRL vs. 26–33%), a broader interorbital space (31% PRL vs. 18–27%), a shorter pelvic fin (102% PRL vs. 109–153%), and a narrower pelvic–anal distance

(79% vs. 87–115%). See the original description given by Nakayama *et al.* (2015a) for further morphological details and comparisons with other congeners.

Genus *Hymenogadus* Gilbert & Hubbs, 1920

[Japanese name: Oni-sujidara-zoku]

Hymenogadus Gilbert & Hubbs, 1920:521 (as subgenus of *Hymenocephalus*; type species: *Hymenocephalus gracilis* Gilbert & Hubbs, 1920, by original designation).

Diagnosis. Anus immediately anterior to anal-fin origin, surrounded by narrow periproct. Ventral striae, consisting of fine, parallel black lines over silvery layer on each side of isthmus, shoulder girdle, most of chest, and posterior to pelvic fins. Gular region covered with fine striations of black epithelium. Light organ developed as long tube, extending from anus to slightly anterior to pelvic-fin bases; small lens at each end of light organ. Second spinous ray of first dorsal fin serrated along its leading edge. Pelvic-fin rays 8, outer ray slender, not expanded or ribbon-like distally. Teeth all small, in narrow villiform bands in both jaws. Trunk shallow, cylindrical in cross section. Head bones thin, fragile; head skin membranous, translucent. Nasal bones weakly ossified, not forming stout, dorsoventrally flattened processes. Chin barbel present. Branchiostegal rays 7. Species small, less than 13 cm TL. Body with silvery reflection when fresh. [From Iwamoto *et al.* (2011), McMillan & Iwamoto (2014), and Schwarzhans (2014).]

Remarks. *Hymenogadus* was originally established as a subgenus of *Hymenocephalus* Giglioli, 1884, with the type species *H. gracilis* (Gilbert & Hubbs, 1920). Subsequently, Okamura (1970a) elevated this taxon to full generic rank, with description of another subgenus *Spicomacrurus*. In addition to *H. gracilis*, *Hymenogadus* includes *H. tenuis* (Gilbert & Hubbs, 1917) known only from the holotype collected from Hawaii.

***Hymenogadus gracilis* (Gilbert & Hubbs, 1920)**

[Japanese name: Oni-sujidara]

(Fig. 147; Appendix 3-9A)

Hymenocephalus gracilis Gilbert & Hubbs, 1920:522, fig. 31 [original description; holotype: USNM 78227, from off southern Luzon Island, *Albatross* sta. 5292, 13°28'45"N, 121°01'12"E, in 162 ftm (296 m)]; Kamohara 1936:934 (first record from Japan; 3 spec. from Kochi); Okada & Matsubara 1938:451 (in key; Japan; new Japanese name: “Oni-sujidara”); Kamohara 1938:69 (brief description; 3 spec. from Kochi Pref.; new Japanese name: “Hoso-suji-dara”); Kamohara 1950:278 (listed; Kochi Pref.); Kamohara 1952:96 (brief description; 3 spec. from Kochi Pref.); Matsubara 1955:1314 (in key; Japan); Kamohara 1958:73 (listed; Kochi Pref.); Kamohara 1964:96 (listed; Kochi Pref.); Tatara *et al.* 1965:121 (listed; off Shikoku); Shinohara *et al.* 2001:305 (142 spec. listed from Tosa Bay); Shao *et al.* 2008b: table 2 (3 spec. listed from northeastern Taiwan and South China Sea); Iwatsuki *et al.* 2017:32 (listed; Hyuga-nada).

Hymenogadus gracilis: Okamura 1970a:61, pl. XVIII, text-fig. 27 (description; 60 spec. from Pacific off southern Japan from Owase to Totoro); Okamura 1970b: table 1 (listed; Japan);

Tominaga & Uyeno 1981:489 (listed; Japan); Okamura 1984a:201, 359, fig. 143 [brief description; 5 spec. from Okinawa Trough; photo based on BSKU spec. (catalog no. unknown)]; Okamura 1984b:93, pl. 80, fig. H (compiled); Okamura 1988:93, pl. 80, fig. H (compiled); Nakabo 1993:354 (in key; Japan); Okamura 1997:128, fig. 4 (compiled); Nakabo 2000:418 (in key; Japan); Nakabo 2002:418 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Chiou *et al.* 2004a:44, fig. 13, table 1 (brief description; 2 spec. from Nan-fang-ao; first record from Taiwan); Shinohara *et al.* 2005:418 (listed; Ryukyu Islands); Nakabo & Kai 2013:494 (in key; Japan); Iwamoto *et al.* 2015:78 (brief description; 6 spec. from northeastern Taiwan and Philippines); Motomura 2020:39 (listed; Japan).

Hymenocephalus striatissimus (not Jordan & Gilbert in Jordan & Starks 1904): Suzuki & Kataoka 1997:81, pl. 32, fig. 173 (brief description; 1 spec. from Kumano-nada) [re-identified here from figure].

Diagnosis. *Hymenogadus gracilis* readily differs from its only congener, *H. tenuis* (Gilbert & Hubbs, 1917), in having a longer chin barbel (19–33% HL vs. 12%) and higher counts of pectoral-fin rays (i14–i19 vs. i12) and outer gill rakers on the second arch (13–19 vs. 10). [Data for the two species are compiled from Gilbert & Hubbs (1917), Marshall & Iwamoto (1973), McMillan & Iwamoto (2014), Schwarzhans (2014), and this study.]

Material examined. 6 specimens. **Japan:** BSKU 112920 (1, 24.6 mm HL, 131+ mm TL), off Okitsu, Tosa Bay, 280–320 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 24 Jan. 2014; BSKU 118435 (1, 22.2 mm HL, 122+ mm TL), off mouth of Niyodo River, Tosa Bay, 300 m, FRV *Tosa-Kaiyo-maru*, bottom otter trawl,

FIGURE 147. Fresh specimen of *Hymenogadus gracilis*. BSKU 91754, 20.9 mm HL, 117+ mm TL, Mimase fish market, Kochi Pref., Pacific. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

coll. by T. Ohkawa, 14 Oct. 2015; BSKU 121258 (1, 22.7 mm HL, 113+ mm TL), BSKU 121264 (1, 22.7 mm HL, 103+ mm TL), BSKU 121268 (1, 22.8 mm HL, 119+ mm TL), off mouth of Niyodo River, Tosa Bay, 200 m, FRV *Tosa-Kaiyo-maru*, bottom otter trawl, coll. by T. Ohkawa, 24 Dec. 2015; *BSKU 91754 (1, 20.9 mm HL, 117+ mm TL), Mimase fish market, bottom trawl, coll. N. Nakayama, 12 Oct. 2007.

Counts and measurements. Based on 5 specimens (22.2–24.6 mm HL, 103+–131+ mm TL). Counts: first dorsal-fin rays II, 10–11; pectoral-fin rays i15–i17; pelvic-fin rays 8; gill rakers on first arch (outer/inner) 10–13/15–17, on second arch 14–18/15–17; longitudinal scales 18–21; transverse scale rows below first dorsal-fin origin 3–3.5, below first dorsal-fin midbase 2–2.5, below second dorsal-fin origin 3–3.5, above anal-fin origin 7–9.5.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 24–28 (31–36); orbit diameter 34–36 (44–47); postorbital length 42–46 (53–58); postrostral length 77–80; orbit–preopercle distance 35–37 (45–47); suborbital width 8–9 (11–12); upper-jaw length 41–44 (54–57); length of rictus 37–39 (48–50); length of premaxillary tooth band 32–36 (41–45); preoral length 13–14 (16–18); distance between tip and lateral angle of snout 13–15 (17–20); snout width 21–25 (27–32); internasal width 18–19 (23–25); interorbital width 16–18 (20–23); occipital width 7–8 (9–11); body width over pectoral-fin bases 53–61 (68–78); body depth at first dorsal-fin origin 54–59 (70–76); body depth at anal-fin origin 44–50 (57–62); prepelvic length 95–102 (123–133); preanal length 158–173 (204–217); isthmus–pelvic distance 46–49 (60–63); isthmus–anal distance 109–127 (140–160); pelvic–anal distance 63–79 (81–99); pelvic-fin length 40–48 (52–60); pectoral-fin length 42–51 (54–64); predorsal length 108–113 (140–143); height of first dorsal fin 62–68 (77–89); length of first dorsal-fin base 32–40 (41–52); interdorsal length 57–68 (73–89); length of gill slit 22–24 (29–31); length of posterior nostril 4–7 (6–9); barbel length 21–27 (28–34).

Size. To about 13 cm TL (McMillan & Iwamoto 2014).

Distribution. Widely known from the North Atlantic, Mediterranean, southwestern Indian Ocean off Mozambique, South China Sea off the Philippines, and Pacific including Japan, Taiwan, New Zealand, and the Sala y Gomez Ridge, at depths of about 182–657 m (Marshall & Iwamoto 1973; Sazonov & Iwamoto 1992; Anderson 1996; Iwamoto & Merrett 1997; Shao *et al.* 2008a, 2008b; Last *et al.* 2014; McMillan & Iwamoto 2014; Schwarzhans 2014; this study). In Japan, known from off the Pacific coasts northward to the Izu Peninsula (138.96°E) and the Okinawa Trough at depths of 182–385 m (Appendix 3-9A). Common.

Remarks. For a full description see the original description given by Gilbert & Hubbs (1920). Additional descriptions were also provided by Okamura (1970a),

Marshall & Iwamoto (1973), and Iwamoto & Merrett (1997). Otolith morphology was described by Schwarzhans (2014). *Hymenocephalus gracilis* was widely recorded from the world's oceans, with the type locality in the South China Sea off the Philippines (see the Distribution). McMillan & Iwamoto (2014) compared the New Zealand specimens with those collected from various parts of its range (including the holotype), and showed no significant differences in major meristic and morphometric characters. This species is currently considered to have a nearly cosmopolitan distribution (McMillan & Iwamoto 2014; Schwarzhans 2014), whereas its only congener, *H. tenuis*, is represented by only the holotype collected from Hawaii.

Genus *Kumba* Marshall, 1973

[Japanese name: Kagamihige-zoku]

Kumba Marshall, 1973:616 (type species: *Kumba dentoni* Marshall, 1973, by original designation).

Parakumba Trunov, 1981:28 (type species: *Parakumba maculisquama* Trunov, 1981, by original designation).

Diagnosis. Anus separated from anal-fin origin in most species (immediately anterior to anal-fin origin in *K. musorstom* Merrett & Iwamoto, 2000); periproct narrowly developed. Small dermal window of light organ between pelvic-fin bases (no external evidence of light organ in *K. musorstom*). Pelvic-fin rays 8–13, its origin below or anterior to pectoral-fin base. Second spinous ray of first dorsal fin serrated along its leading edge. No modified scales on head ridges and tip and lateral angles of snout. Chin barbel present. Teeth small, in narrow bands in both jaws. Underside of head almost completely naked, including gular and branchiostegal membranes; dorsal surface of snout broadly naked anteriorly. Body scales covered with short, erect, needle-like spinules in widely spaced, subparallel rows; scale spinules lacking buttresses; no reticulate structure. Grooved lateral line present. Cephalic sensory pores present or absent. Branchiostegal rays 7. [Modified from Iwamoto & Sazonov (1994).]

Remarks. The genus *Kumba* was established by Marshall (1973) for his new species *K. dentoni*. This genus had been represented by only the type species, until Iwamoto & Sazonov (1994) provided a systematic revision of the group. In their revised classification, *Kumba* was considered not only to be a senior synonym of *Parakumba* Trunov, 1981, but also to include the following nine species: *K. calvifrons* Iwamoto & Sazonov, 1994; *K. dentoni*; *K. gymnorhynchus* Iwamoto & Sazonov, 1994; *K. hebetata* (Gilbert, 1905); *K. japonica* (Matsubara, 1943); *K. maculisquama* (Trunov, 1981) (type species of *Parakumba*); *K. punctulata* Iwamoto & Sazonov, 1994; and two undescribed species (these species were not named in their study, due to the immature state or

questionable status of their materials). Of these, *K. hebetata* and *K. japonica* were previously classified under *Nezumia* Jordan in Jordan & Starks, 1904 and *Ventrifossa* Gilbert & Hubbs, 1920 respectively (Iwamoto 1990).

Merrett & Iwamoto (2000) described *K. musorstom*, the latest species of the genus to be described, based on a single specimen collected from the Norfolk Ridge. This species is unique within the genus in that the anus is situated immediately anterior to the anal-fin origin, without any external evidence of the light organ anterior to the periproct. In the other species of *Kumba*, the anus is separated from the anal-fin origin, usually closer to the pelvic-fin bases, and a small dermal window of the light organ is present anterior to the periproct. The unique characters of *K. musorstom* casts doubt on the generic affiliation of this species. The position of the anus and the presence/absence of the light organ are considered to be important diagnostic characters in the generic classification of grenadiers (Iwamoto 1990). Based on the original description given by Merrett & Iwamoto (2000), *K. musorstom* seems to be closely similar to *Asthenomacrus* Sazonov & Shcherbachev, 1982, with which it shares many morphological features, but they differ in the number of pelvic-fin rays (12–13 in *K. musorstom* vs. 7–9 in *Asthenomacrus*) and the presence/absence of the serration along the leading edge of the first dorsal fin (absent vs. present). A further in-depth study of *K. musorstom* is obviously required to clarify the generic affiliation of this enigmatic species.

Kumba japonica (Matsubara, 1943)

[Japanese name: Kagami-hige]

(Figs. 148–149; Appendix 3-9B)

Lionurus japonicus Matsubara, 1943:149, fig. 9 [original description; holotype: FAKU 1951 (ex. Sigenkagaku Fish Collection no. 8), from Kumano-nada; 1 paratype probably collected with holotype; new Japanese name: “Kagami-hige”]; Matsubara 1955:1316 (in key; Japan); Okada *et al.* 1959:83 (listed; Kumano-nada); Kamohara 1964:96 (listed; Kochi Pref.).

Nezumia japonicus: Okamura 1970a:88, pl. XIX, text-fig. 39 (new combination; description; 5 spec. from Kumano-nada and Tosa Bay).

Nezumia japonica: Okamura 1970b: table 1 (listed; Japan); Kataoka & Tomida 1981:78 (listed; Mie Pref.); Tominaga & Uyeno 1981:489 (listed; Japan); Ozawa 1983:13 (listed; off Makurazaki, Kagoshima Pref., East China Sea).

Ventrifossa japonica: Okamura 1982:147, 349, fig. 91 (brief description; 10 spec. from Kyushu-Palau Ridge; photo based on BSKU 29332); Okamura 1984b:93, pl. 81, fig. F (compiled); Okamura 1988:93, pl. 81, fig. F (compiled); Nakabo 1993:357 (in key; Japan); Okamura 1997:128, fig. 8 (compiled); Nakabo 2000:421 (in key; Japan); Shinohara *et al.* 2001:306 (listed; Tosa Bay); Nakabo 2002:421 (in key;

Japan); Kitagawa *et al.* 2008:39, unnumbered fig. (compiled; Pacific off Tohoku); Shinohara *et al.* 2009:709 (listed; Pacific off Tohoku).

Kumba japonica: Iwamoto & Sazonov 1994:231, fig. 7 (Kyushu-Palau Ridge and Tosa Bay); Chiou *et al.* 2004a:45, fig. 14, table 1 (brief description; 2 spec. from Dong-gang; first record from Taiwan); Shao *et al.* 2008b: table 2 (3 spec. listed from northeastern and southwestern Taiwan); Nakabo & Kai 2013:497 (in key; Japan); Iwamoto *et al.* 2015:80 (brief description; 10 spec. from northeastern and southwestern Taiwan, and Japan); Motomura 2020:39 (listed; Japan).

Diagnosis. A species of *Kumba* with 11–13 pelvic-fin rays; naked area dorsally on head restricted anterior to line connecting lateral angles of snout; anus separated from anal-fin origin, situated posterior 1/3 of pelvic–anal distance; head firm, not inflated; cephalic sensory pores present along all canals, those on infraorbital and mandibular canals larger; chin barbel long, length 21–30% HL; snout length 25–32% HL; orbit diameter 33–40% HL; upper-jaw length 35–39% HL; interorbital width 23–28% HL; 3 (rarely 4) small, black pigmented areas near mid-length of anal-fin base (only found in smaller specimens).

Material examined. 12 specimens. **Japan:** BSKU 101176 (1, 25.1 mm HL, 159+ mm TL), Tosa Bay, 300 m, FRV *Kotaka-maru*, 4 Dec. 1997; *BSKU 117558 (1, 16.3 mm HL, 91+ mm TL), off mouth of Niyodo River, Tosa Bay, 300 m, FRV *Tosa-Kaiyo-maru*, coll. T. Ohkawa, 24 Mar. 2015; BSKU 99318 (1, 27.9 mm HL, 170+ mm TL), Tosa Bay, 400 m, 20 Mar. 1989; BSKU 43817 (1, 24.9 mm HL, 168 mm TL), BSKU 43818 (1, 25.6 mm HL, 169+ mm TL), Tosa Bay, 400 m, FRV *Kotaka-maru*, 11 Jun. 1987; FAKU 204496 (1, 24.1 mm HL, 150+ mm TL), Tosa Bay, Mimase fish market, F/V *Tsukasa-maru*, bottom trawl, coll. N. Nakayama, 4 Mar. 2017; BSKU 44189 (1, 20.9 mm HL, 140 mm TL), Mimase fish market, 7 Nov. 1987; NSMT-P 114948 (1, 29.9 mm HL, 187 mm TL), southwest of Cape Omaezaki, Enshu-nada, 34.3816°N, 138.0010°E, 475–488 m, R/V *Tansei-maru*, cr. KT-95-17, sta. EN1-4, 3-m ORE beam trawl, 10 Dec. 1995; NSMT-P 104114 (1, 28.3 mm HL, 158+ mm TL), Kumano-nada, 420–425 m, R/V *Tansei-maru*, cr. KT-95-17, sta. KN-01(2), 3-m ORE beam trawl, date unknown; BSKU 32216 (1, 22.1 mm HL, 136+ mm TL), BSKU 32217 (1, 20.3 mm HL, 128+ mm TL), BSKU 32220 (1, 20.5 mm HL, 133+ mm TL), Kyushu-Palau Ridge, 28.1117°N, 134.6656°E, 550–600 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, tr. 9, bottom trawl, coll. O. Okamura *et al.*, 19 Dec. 1979.

Counts and measurements. Based on 11 specimens (20.3–29.9 mm HL, 128+–187 mm TL). Counts: first dorsal-fin rays II, 10–12; pectoral-fin rays i18–i23; pelvic-fin rays 11–13; gill rakers on first arch (outer/inner) 8–11/11–13, on second arch 11–14/11–13; longitudinal scales 34–40; transverse scale rows below first dorsal-fin

FIGURE 148. Fresh specimens of *Kumba japonica*. (A–C) BSKU 117558, 16.3 mm HL, 91+ mm TL, Tosa Bay off Niyodo River, Kochi Pref., Pacific, 300 m depth; (D) unknown specimen, Kyushu-Palau Ridge, Pacific; (E) BSKU 43818, 25.6 mm HL, 169+ mm TL, Tosa Bay, Kochi Pref., Pacific, 400 m depth. (A, D–E) Lateral views; (B) dorsal and (C) ventral views of the head and trunk. [Photos: (A–C) N. Nakayama; (D–E) BSKU]

origin 10–11, below first dorsal-fin midbase 6.5–8, below second dorsal-fin origin 6.5–8.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout

length 25–32 (32–43); orbit diameter 33–40 (44–52); postorbital length 37–43 (50–58); postrostral length 73–79; orbit–preopercle distance 29–37 (39–48); suborbital width 13–15 (17–20); upper-jaw length 35–39 (46–51);

FIGURE 149. Scanning electron micrographs showing a body scale (from the dorsum below the interdorsal space) of *Kumba japonica*. NSMT-P 104114, 28.3 mm HL. (A) View from above; (B) oblique view. [Photos: NSMT]

length of rictus 28–34 (36–46); length of premaxillary tooth band 18–22 (24–28); preoral length 20–23 (27–32); distance between tip and lateral angle of snout 14–18 (18–25); snout width 24–31 (31–42); internasal width 19–22 (25–30); interorbital width 23–28 (30–38); body width over pectoral-fin bases 37–48 (47–65); body depth at first dorsal-fin origin 71–85 (97–111); body depth at anal-fin origin 62–74 (79–98); prepelvic length 95–105 (124–142); preanus length 126–142 (160–187); preanal length 143–170 (190–220); isthmus–pelvic distance 34–47 (46–60); isthmus–anus distance 67–91 (86–115); isthmus–anal distance 85–120 (116–155); pelvic–anal distance 46–76 (62–99); anus–anal distance 17–29 (23–38); pelvic-fin length 49–59 (62–79); pectoral-fin length 54–68 (72–86); predorsal length 110–121 (148–161); height of first dorsal fin 88–103 (116–131); length of first dorsal-fin base 28–37 (37–49); interdorsal length 50–84 (67–108); length of gill slit 14–19 (20–26); length of posterior nostril 6–11 (8–15); barbel length 21–30 (28–40).

Size. To about 19 cm TL (NSMT-P 114948, 187 mm TL, Enshu-nada, Japan).

Variation. The unique pigmentation of *K. japonica*, consisting of three (or rarely four) small, black spots along the anal-fin base, has been considered to be one of the diagnostic features of this species (Iwamoto 1990; Iwamoto & Sazonov 1994). However, these spots are often obsolete and barely discernible in large specimens (Fig. 148E).

Distribution. Known only from the northwestern Pacific from the east of Tohoku southward to the northern South China Sea off southern Taiwan, including the Kyushu-Palau Ridge, at depths of 213–710 m (Iwamoto *et al.* 2015; this study; Appendix 3-9B).

Remarks. This species was well described in the original description (Matsubara 1943) and by Okamura (1970a; as *Nezumia japonicus*), and these should be referred to for further details. It was previously allocated

to *Lionurus* Günther, 1887, *Nezumia* Jordan in Jordan & Starks, 1904, or *Ventrifossa* Gilbert & Hubbs, 1920 (see the synonymy of this species). However, based on the work of Iwamoto & Sazonov (1994:232), the extensive naked areas on the head and the distinctive spinulation of the head scales support its placement in *Kumba*.

Comments on type specimens. *Kumba japonica* was originally described by Matsubara (1943) as a new species of *Lionurus* based on two specimens collected from the Kumano-nada, Japan. Unfortunately, the holotype was destroyed during WWII (see the Comments on holotype of *C. hige*). The paratype was not found in the FAKU collection during the author's visits in 2009 and 2010, and has been presumed to be lost (see also Iwamoto *et al.* 2015:80).

Comparisons. *Kumba japonica* is distinctive within the genus, and readily differs from any other congeners in having 3–4 black spots along the anal-fin base (but often obscure in large specimens vs. completely absent) and a longer chin barbel (21–30% HL vs. $\leq 21\%$). It further differs from all but *K. gymnorhynchus* Iwamoto & Sazonov, 1994 and *K. musorstom* Merrett & Iwamoto, 2000 in its higher counts of pelvic-fin rays [11–13 (usually 12) vs. 8–9]. [Meristic and morphometric data for the other species are based on Iwamoto & Sazonov (1994: table 1) and Merrett & Iwamoto (2000:762).]

Genus *Kuronezumia* Iwamoto, 1974

[Japanese name: Suruga-nezumidara-zoku]

Kuronezumia Iwamoto, 1974:509 [type species: *Nezumia* (*Kuronezumia*) *bubonis* Iwamoto, 1974, by original designation].

Diagnosis. Anus separated from anal-fin origin, closer to pelvic-fin bases; periproct teardrop-shaped, moderately

developed. Small dermal window of light organ between pelvic-fin bases. Infraorbital ridge not connected with preopercular ridge, separated by distinct gap. Second spinous ray of first dorsal fin finely serrated along its leading edge. Pelvic fin inserted below or anterior to pectoral-fin base. Mouth moderately large, usually subterminal. Chin barbel well developed. Teeth small, in broad bands in both jaws; premaxillary band ending anterior to posterior edge of maxillary process. Tip and lateral angles of snout lacking prominent scutes in most species. Suborbital region almost vertical, without coarsely modified scales. Head fully scaled, except for narrow naked area above upper lip. Body scales not deciduous, closely adhered to body surfaces; scale spinules needle-like, densely scattered over exposed portion or tightly packed subparallel to convergent rows; buttresses of spinules rudimentary; reticulate structure developed on posterior parts of unexposed portion. Grooved lateral line present. Cephalic sensory pores present or absent. Branchiostegal rays 7. Body dark to swarthy, lacking silvery reflection when fresh. [Modified from Sazonov & Iwamoto (1992) and Iwamoto & Graham (2001).]

Remarks. *Kuronezumia* was originally established as a subgenus of *Nezumia* Jordan in Jordan & Starks, 1904, but Sazonov & Iwamoto (1992) elevated this taxon to full generic rank. Shcherbachev *et al.* (1992) provided a synopsis for all the then-known species (seven species), viz., *K. bubonis* Iwamoto, 1974 (type species); *K. dara* (Gilbert & Hubbs, 1916); *K. leonis* (Barnard, 1925); *K. macronema* (Smith & Radcliffe in Radcliffe, 1912); *K. paepkei* Shcherbachev, Sazonov & Iwamoto, 1992; *K. pallida* Sazonov & Iwamoto, 1992; and *Kuronezumia* sp. [this undescribed species was initially recognized by Iwamoto (1974), but still remains to be described]. Two species are distributed in Japanese waters, one of which is new to science [different from the one reported by Iwamoto (1974)] and is described below.

Key to species of *Kuronezumia* from Japan and adjacent waters

- 1a Pelvic-fin rays 11–12; posterior half of mandibular rami scaled; branchiostegal membranes naked; no open pores along cephalic sensory canals; oral cavity blackish.....*K. dara*
- 1b Pelvic-fin rays 15; mandibular rami almost entirely naked except for posterior margins; scaly patches present on lower portions of branchiostegal rays; cephalic sensory pores present; oral cavity pale*K. endoi* sp. nov.

Kuronezumia dara (Gilbert & Hubbs, 1916)

[Japanese name: Suruga-nezumidara]

(Figs. 150–151, 152A–B; Table 10; Appendix 3–9C)

Lionurus darus Gilbert & Hubbs, 1916:197, pl. 10, fig. 1 [original description; holotype: USNM 76867, from “Suruga Gulf”, *Albatross* sta. 5060, in 197 fth (361 m); misspelled “*durus*” in caption of pl. 10]; Okada & Matsubara 1938:453, pl. 111, fig. 2 (in key; Japan); Kuroda 1951:392 (listed; Suruga Bay; Japanese name: “Suruga-nezumidara”); Matsubara 1955:1316 (in key; Japan); Matsubara 1965:509 (compiled; Japan).

Nezumia darus: Okamura 1970a:101 [new combination; brief description after Gilbert & Hubbs (1916)]; Tominaga & Uyeno 1981:489 (listed; Japan); Okamura 1982:161, 349, fig. 95 (description; 2 spec. from Tosa Bay; photo based on BSKU 29499); Okamura 1984a:217, 363, fig. 153 (brief description; 4 spec. from Okinawa Trough; photo based on BSKU 27666); Okamura 1984b:94, pl. 81, fig. J (compiled); Okamura 1988:94, pl. 81, fig. J (compiled); Nakabo 1993:360 (in key; Japan); Okamura 1997:128, fig. 14 (compiled); Shinohara & Matsuura 1997:292 (listed; Suruga Bay); Nakabo 2000:424 (in key; Japan); Shinohara *et al.* 2001:306 (listed; Tosa Bay); Nakabo 2002:424 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Shinohara *et al.* 2005:418 (listed; Ryukyu Islands); Suetsugu & Ohta 2005: table 2 (questionable; listed; southeast of Boso Peninsula).

Nezumia dara: Okamura 1970b: table 1 (listed; Japan).

Kuronezumia dara: Shcherbachev *et al.* 1992:100 (new combination); Nakabo & Kai 2013:500 (in key; Japan); Iwamoto *et al.* 2015:82, fig. 14 (brief description; 5 spec. from southwestern Taiwan and Japan); Motomura 2020:39 (listed; Japan).

Kuronezumia darus: Shao *et al.* 2008b: table 2 (1 spec. from southwestern Taiwan; new Taiwanese record).

Diagnosis. A species of *Kuronezumia* without large, scaly, tubercular swelling anterior to anus; pelvic-fin rays 11–12; no enlarged scutes at snout tip; no open pores along cephalic sensory canals; mandibular rami scaled posteriorly; branchiostegal membranes naked; first dorsal fin uniformly black; oral cavity blackish; body scales covered with long, moderately erect, needle-like spinules densely scattered over exposed portion; first dorsal-fin rays II,9–10; pectoral-fin rays i21–i26; inner gill rakers on first arch 9–11; upper-jaw length 30–35% HL.

Material examined. 9 specimens. **Holotype** of *Lionurus darus*: USNM 76867 (19.6 mm HL, 132+ mm TL), off Fuji River, Shizuoka Pref., Suruga Bay, Japan, 35.1000°N, 138.6694°E, 197 fth (361 m), *Albatross* sta. 5060, 9-ft *Albatross-Blake* beam trawl, 13 Oct. 1906. **Non-types: Japan:** BSKU 26326 (1, 34.5 mm HL, 210+ mm TL), off Senkaku Islands, Okinawa Trough, 25.6283°N, 122.8933°E, 560–692 m, F/V *Yuryo-maru*, No. 8, tr. 3, bottom trawl, coll. T. Kitajima *et al.*, 14 Jan. 1978; BSKU 27666 (1, 49.8 mm HL, 256+ mm TL), west of Okinawa-jima Island, Okinawa Trough, 28.7333°N, 127.0167°E, 610–640 m, F/V *Ryoan-maru*, No. 28, tr. 49, bottom trawl, coll. Y. Kinoshita and S. Hagino, 11 Mar. 1978; BSKU 44828 (1, 42.0 mm HL, 220+ mm TL), Tosa Bay, 33.1755°N, 133.6498°E, 500 m, FRV *Kotaka-*

maru, 25 Apr. 1988; BSKU 45036 (1, 60.0 mm HL, 374+ mm TL), Tosa Bay, 500 m, FRV *Kotaka-maru*, 23 Aug. 1988; BSKU 30497 (1, 20.1 mm HL, 111+ mm TL), Tosa Bay, 32.9667°N, 133.5333°E, 605 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, bottom trawl, coll. O. Okamura *et al.*, 21 Dec. 1979; BSKU 29499 (1, 21.1 mm HL, 137+ mm TL), Tosa Bay, 32.9667°N, 133.5333°E, 605 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, cr. 4-Ky, tr. 19, bottom trawl, coll. O. Okamura *et al.*, 21 Dec. 1979; NSMT-P 48555 (1, 54.9 mm HL, 340 mm TL), southwest of Heta, Suruga Bay, 34.9517°N, 138.7267°E, 380–1000 m, F/V *Seishin-maru*, sta. CT-1-5, bottom trawl, 26 Oct. 1993; NSMT-P 101346 (1, 69.8 mm HL, 416+ mm TL), west of Matsuzaki, Suruga Bay, 34.7567°N, 138.6650°E, 652–676 m, R/V *Tansei-maru*, cr. KT-89-06, sta. SB26, 3-m ORE beam trawl, 17 May 1989.

Redescription. General features are shown in Figs. 150–151. Counts and measurements are given in Table 10. Trunk deep, compressed, width over pectoral-fin bases 1.7–2.7 in depth at first dorsal-fin origin. Body deepest at first dorsal-fin origin, gradually tapering to long, well compressed tail. Head moderately large, HL about 5.1–6.7 in TL. Supraoccipital crest moderately high, giving humpback appearance to dorsal contour of predorsal region (only in larger specimens). Snout short, protruding scarcely beyond upper jaw, length 0.7–1.1 times as long as orbit diameter; ventral contour almost vertical when viewed laterally; snout high, its tip situated on horizontal through midorbit or above. Orbit circular, moderate in size, greatest diameter 1.1–1.9 in postorbital length. Interorbital space flat or slightly concave in preserved specimens, width 1.1–1.6 in orbit diameter. Mouth small, subinferior, upper-jaw length 0.8–1.2 in orbit diameter; posterior margin of maxilla not reaching vertical through midorbit; lateral corner of mouth moderately restricted by lip folds; lips thick, highly papillose near tooth bands. Suborbital region deep, almost flat, lacking stout bony ridge. Preopercle large, posterior margin almost vertical, broadly rounded ventrally; preopercular ridge low, forming right angle posteroventrally. Interopercle narrowly exposed as narrow tab beyond preopercle. Gill membranes narrowly connected across, and attached mesially to isthmus, with posterior free fold. Anteroventral end of gill opening not reaching below posterior margin of lower jaw. Outermost gill slit moderately restricted by skin folds, length 1.5–2.2 in orbit diameter. Gill rakers small, tubercular, armed with short, slender spines distally; those on outer side of first arch and inner side of fourth arch much smaller than others. Gill filaments moderately long. Barbel long, slender, length 1.1–1.6 in orbit diameter.

Anus closer to pelvic-fin bases than to anal-fin origin. Periproct broad, more or less oval in shape. Light organ small, circular, separated from periproct by narrow scaled area; its anterior margin lying on line connecting outer pelvic-fin bases.

Teeth short, slender, sharp, in broad tapered bands in both jaws. Premaxillary teeth arranged in about 4–5 tooth rows anteriorly with outermost series significantly enlarged. Mandibular band with about 5 tooth rows at widest point near symphysis, none especially enlarged. Posterior ends of tooth bands falling far short of lateral corner of mouth in both jaws. All teeth deeply embedded in thick layer of gum papillae.

Body scales small, thin, adherent, covered with long, moderately erect, needle-like spinules densely scattered over exposed portion (Fig. 152A–B); tip of posteriormost spinules extending well beyond scale margin; spinules forming angle of about 45° to scale surface, giving villiform texture to body surfaces; buttresses of spinules poorly developed; reticulate structure narrowly developed on posterior parts of unexposed portion. Body fully scaled except for periproct, light organ, and fins, but pelvic fin heavily scaled proximally.

Head ridges inconspicuous, without coarsely modified scales. Terminal snout scutes absent (scales on snout tip very slightly enlarged and thickened in larger specimens); lateral angles of snout lacking coarsely modified scales. Head scales covered with short, erect, needle-like spinules in widely divergent rows. Head almost completely and uniformly scaled; naked areas confined to membrane around anterior nostril, narrow band above upper lip, and anterior half of mandibular rami; gular and branchiostegal membranes entirely naked.

Cephalic sensory canals without open pores. Grooved lateral line almost complete, often narrowly interrupted above pectoral-fin base.

First dorsal fin originating slightly posterior to vertical through pectoral-fin base; second spinous ray not especially elongate, its tip extending to base of 16th–26th second dorsal-fin ray when laid back; height of first dorsal fin 3.4–5.3 times as long as its base length; leading edge of second spinous ray beset with short, erect, conical denticles. Second dorsal fin moderately separated from first dorsal, originating above bases of 8th–12th anal-fin rays; interdorsal space 0.8–1.5 times as long as first dorsal-fin base length. Pelvic fin inserted slightly anterior to vertical through pectoral-fin base; outermost ray greatly prolonged, its tip reaching base of 9th–15th anal-fin ray when laid back.

Color when fresh (Fig. 150). Uniformly dark to dusky gray; abdomen much darker; gular and branchiostegal membranes dull black; first dorsal, pectoral, and pelvic fins uniformly black; anal fin dusky, but darker anteriorly.

Color in alcohol (Fig. 151). Almost same as in fresh condition; oral cavity blackish; gums pale; gill cavity paler anteriorly, blackish posteriorly; gular membrane dusky, branchiostegal membranes black; light organ and periproct dull black.

Size. To about 42 cm TL (NSMT-P 101346, 416+ mm TL, Suruga Bay, Japan).

Distribution. Restricted to Japan and Taiwan. Known from Suruga Bay, Tosa Bay, the Okinawa Trough,

FIGURE 150. Fresh specimens of *Kuronezumia dara*. (A) BSKU 29499, 21.1 mm HL, 137+ mm TL, Tosa Bay, Kochi Pref., Pacific, 605 m depth; (B) BSKU 44828, 42.0 mm HL, 220+ mm TL, Tosa Bay, Kochi Pref., Pacific, 500 m depth; (C) BSKU 45036, 60.0 mm HL, 374+ mm TL, Tosa Bay, Kochi Pref., Pacific, 500 m depth. Lateral views. [Photos: BSKU]

and the northern South China Sea, at depths of 280–1000 m (Shcherbachev *et al.* 1992; Shao *et al.* 2008a, 2008b; Iwamoto *et al.* 2015; this study; Appendix 3-9C). It appears to be rare wherever found.

Remarks. *Kuronezumia dara* is redescribed herein based on the holotype from Suruga Bay and eight additional specimens from Japan. Although Gilbert & Hubbs (1916) originally referred this species to *Lionurus* Günther, 1887, it was subsequently transferred to *Nezumia* Jordan *in* Jordan & Starks, 1904 by Okamura (1970a), and then shifted to *Kuronezumia* by Sazonov & Iwamoto (1992).

Relationships and comparisons. Data used in the

following comparisons are based on Shcherbachev *et al.* (1992) and this study. *Kuronezumia dara* is one of six species of the genus that is characterized by the absence of a large, scaly, bulbous swelling anterior to the anus. In this respect, *K. dara* is similar to the following five species: *K. endoi* **sp. nov.** from Japan; *K. leonis* (Barnard, 1925) widespread in temperate to subarctic waters of the South Atlantic, Indian Ocean, and southwestern Pacific; *K. macronema* (Smith & Radcliffe *in* Radcliffe, 1912) probably restricted to the Philippines; *K. paepkei* Shcherbachev, Sazonov & Iwamoto, 1992 known only from the western Indian Ocean off Kenya; and *K. pallida* Sazonov & Iwamoto, 1992 disjunctly distributed in the

FIGURE 151. Holotype of *Lionurus darus* (= *Kuronezumia dara*). USNM 76867, 19.6 mm HL, 132+ mm TL, Suruga Bay, Shizuoka Pref., Pacific, 361 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

Sala y Gomez Ridge in the southeastern Pacific (type locality) and the southeastern Indian Ocean off Western Australia.

Kuronezumia dara differs notably from all but *K. pallida* in lacking open pores along the cephalic sensory canals [vs. small but present (unknown in *K. macronema*)]. In addition, this species is unlikely to be confused with *K. leonis* as it lacks a large spiny scute at the tip of the snout (vs. present). It also differs from *K. endoi*, *K. leonis*, and *K. pallida* in that the gular and branchiostegal membranes are completely naked (vs. scaly patches are present on the lower portions of the branchiostegal membranes). *Kuronezumia dara* is further distinguished from *K. paepkei* by its higher counts of transverse scale rows (8.5–10.5 vs. 7 scales below the first dorsal-fin midbase; 10–11.5 vs. 8 scales below the second dorsal-fin origin) and lower counts of gill rakers (9–11 vs. 12 on the inner side of the first arch; 8–10 vs. 11 on the outer side of the second arch).

As discussed by Shcherbachev *et al.* (1992:102), *K. dara* is most similar to *K. macronema*, and they are difficult to distinguish by meristic and morphometric characters (Shcherbachev *et al.* 1992: table 1). According to the key to species given by Shcherbachev *et al.* (1992), *K. dara* is distinguished from *K. macronema* in having a uniformly black first dorsal fin (vs. darker basally and paler distally). Unfortunately, the present author had no opportunity to examine specimens of the latter species, and their differences should be verified in the future.

***Kuronezumia endoi* sp. nov.**

[New Japanese name: Kuro-nezumi-dara]

(Figs. 152C–D, 153–156; Table 10; Appendix 3-9D)

Diagnosis. A species of *Kuronezumia* without large, scaly, tubercular swelling anterior to anus; pelvic-fin rays 15; no

enlarged scutes at snout tip; prominent open pores present along cephalic sensory canals; mandibular rami almost entirely naked except for posterior portions; scaly patches present on lower portions of branchiostegal rays; first dorsal fin blackish; oral cavity pale; body scales covered with long, moderately reclined, needle-like spinules in convergent rows; first dorsal-fin rays II, 10–11; pectoral-fin rays i22–i27; inner gill rakers on first arch 9; upper-jaw length 31–32% HL.

Material examined. 2 specimens. **Holotype:** MSM 71-945 (32.7 mm HL, 215+ mm TL), 2 km off Fuji River, Shizuoka Pref., Suruga Bay, Japan, F/V *Asahi-maru*, bottom trawl, coll. S. Kishimoto, 23 Dec. 1971. **Paratype:** BSKU 44829 (1, 26.1 mm HL, 159+ mm TL), Tosa Bay, 33.1755°N, 133.6498°E, 500 m, FRV *Kotaka-maru*, 25 Apr. 1988.

Description. General features are shown in Figs. 153–155. Counts and measurements are given in Table 10. The following description based on the holotype, unless otherwise stated; variations found in the paratype are indicated in parentheses. Trunk deep, compressed, width over pectoral-fin bases 2.1 (2.2) in depth at first dorsal-fin origin. Body deepest at first dorsal-fin origin, gradually tapering to long, slender tail. Head moderately large, HL about 6.6 (6.1) in TL. Supraoccipital crest moderately high, giving humpback appearance to dorsal contour of predorsal region (only slightly humped). Snout short, protruding slightly beyond upper jaw, length 0.9 (0.8) times as long as orbit diameter; ventral contour only slightly reclined from vertical when viewed laterally; snout high, its tip situated on horizontal through midorbit. Orbit moderate in size, circular, greatest diameter 1.4 (1.1) in postorbital length. Interorbital space slightly concave in preserved specimens, width 1.3 in orbit diameter. Mouth small, subinferior, upper-jaw length 1.0 (1.2) in orbit diameter; posterior margin of maxilla almost reaching vertical through midorbit (anterior 1/3 of orbit);

lateral corner of mouth slightly restricted by lip folds; lips moderately thick, highly papillose near tooth bands. Suborbital region deep, only weakly ridged, lacking stout bony shelf. Preopercle large, posterior margin almost vertical, smoothly curved posteroventrally; preopercular ridge low, poorly marked, forming slight backward extension at angle. Interopercle narrowly exposed beyond preopercle. Gill membranes narrowly connected across, and attached mesially to isthmus, with slight posterior free fold. Gill opening moderately wide, extending forward to below posterior margin of lower jaw. Outermost gill slit moderately restricted by skin folds, length 2.2 (2.3) in orbit diameter. Gill rakers small, tubercular, armed with short, slender spines distally; those on outer side of first arch and inner side of fourth arch much smaller than others. Gill filaments moderately long. Barbel slender, well developed, length 1.4 (1.7) in orbit diameter.

Anus closer to pelvic-fin bases than to anal-fin origin. Periproct broad, teardrop-shaped, connected anteriorly to dermal window of light organ. Light organ small, circular, its anterior margin lying on line connecting outer pelvic-fin bases.

Teeth short, slender, sharp, arranged in tapered bands in both jaws. Premaxillary band moderately narrow, with about 4 tooth rows anteriorly; outermost series distinctly enlarged; posterior margin of the tooth band falling far short of corner of mouth. Mandibular band broad but abruptly narrowing posteriorly, with about 5–6 tooth rows at widest point near symphysis; no teeth significantly enlarged. All teeth deeply embedded in thick layer of gum papillae.

Body scales very small, thin, not highly deciduous, covered with long, moderately reclined, needle-like spinules in about 9 convergent rows (Fig. 152C–D) (5 subparallel rows); tip of last spinule in each row extending well beyond posterior scale margin; buttresses of spinules absent; reticulate structure broadly developed on posterior parts of unexposed portion. Body fully scaled except for periproct, light organ, and fins, but pelvic fin heavily scaled proximally.

Head ridges not prominently marked, without coarsely modified scales. Tip and lateral angles of snout lacking prominent scutes. Scales on head covered with cluster of short, erect, needle-like spinules. Dorsal and lateral surfaces of head completely and uniformly scaled except for narrow area around anterior nostril; underside of snout with broad, triangular naked area above upper lip (Fig. 156A); mandibular rami almost completely naked except for small scaly patches on posterior portions (Fig. 156C); gular membranes naked, but branchiostegal membranes with elongate scaly patches on bases of 4th and 5th branchiostegal rays respectively (Fig. 156C).

Cephalic sensory pores present, serially arranged along sensory canals (Fig. 156); those on underside of head prominent and easily discernible. Grooved lateral line complete throughout its length.

First dorsal fin originating slightly posterior to vertical through pectoral-fin base; second spinous ray not especially elongate, its tip extending to base of 17th (22th) second dorsal-fin ray when laid back; height of first dorsal fin 3.4 (3.5) times as long as its base length; leading edge of second spinous ray armed with short, reclined, needle-like denticles (denticles much longer and more erect). Second dorsal fin moderately separated from first dorsal, originating above 9th (12th) anal-fin ray; interdorsal space 0.8 times as long as first dorsal-fin base length. Pelvic fin inserted slightly anterior to vertical through pectoral-fin base; outermost ray prolonged as hair-line filament, its tip reaching base of 6th (13th) anal-fin ray when laid back.

Color when fresh (Fig. 153). Uniformly dark; fins intensely blackish.

Color in alcohol (Figs. 154–155). Head and body dark brown overall, with much darker abdomen; gular and branchiostegal membranes, dermal window of light organ, and periproct prominently blackish; lips dark; oral cavity and gum papillae pale; gill cavity paler anteriorly, blackish posteriorly; gill rakers, arches, and filaments yellowish brown; chin barbel generally immaculate, but slightly peppered near base; first dorsal fin dusky basally, but prominently black distally; second dorsal, pectoral, and anal fin uniformly dusky; pelvic fin darker. In paratype, ground color of head and body much darker than that in holotype; chest, abdomen, and gill apparatus distinctly blackish; gill rakers and arches with small melanophores densely scattered; first dorsal and pelvic fins uniformly black; pelvic fin blackish basally; anal fin darker anteriorly.

Size. To about 22 cm TL (MSM 71-945, holotype, 215+ mm TL, Suruga Bay, Japan).

Distribution. So far known only from Suruga Bay and Tosa Bay at depths of about 500 m (Appendix 3-9D). Very rare.

Etymology. The new species is named in honor of Dr. Hiromitsu Endo of the Laboratory of Marine Biology, Faculty of Science and Technology, Kochi University (BSKU), for his contributions to our knowledge of gadiform systematics. The Japanese name, Kuro-nezumidara, is derived from a combination of the generic name (= *Kuronezumia*) and a Japanese name for gadiform fishes (= dara).

Remarks, relationships, and comparisons. Data used in the following comparisons are based on Shcherbachev *et al.* (1992) and this study. *Kuronezumia endoi* **sp. nov.** is poorly represented in museum collections, with only two specimens being available. The new species is readily diagnosed within the genus by its higher counts of pelvic-fin rays (15 vs. ≤ 13 in other species, rarely 14 in *K. bubonis* Iwamoto, 1974). It further differs from all in that the mandibular rami are completely naked except for their very posterior portions (Fig. 156C vs. mostly or fully scaled). In addition, *K. endoi* differs notably from *K. leonis* (Barnard, 1925) in

TABLE 10. Morphometric measurements and meristic counts* for *Kuronezumia bubonis*, *K. dara*, and *K. endoi* sp. nov.

Species	<i>K. bubonis</i>		<i>K. dara</i>		<i>K. endoi</i>	
Type status	NT		HT	NT	HL	PT
Number of specimens	<i>n</i> = 3	<i>n</i> = 12**	<i>n</i> = 1	<i>n</i> = 8	<i>n</i> = 1	<i>n</i> = 1
Measurements (mm)						
TL	347+–639+	360–730	132+	111+–416+	215+	159+
HL	53.5–116	54.0–124	19.6	20.1–69.8	32.7	26.1
% of HL (% of PRL)						
Snout length	27 (35–36)	26–32	27 (34)	25–29 (33–40)	28 (37)	30 (40)
Orbit diameter	25–31 (32–41)	23–31	37 (48)	26–36 (36–48)	31 (42)	36 (48)
Postorbital length	46–55 (60–73)	49–51	40 (52)	42–50 (56–67)	43 (57)	39 (54)
Postrostral length	76–77 (–)	–	77 (–)	73–77 (–)	76 (–)	74 (–)
Orbit–preopercle distance	35–44 (46–58)	40–41	34 (44)	33–43 (44–57)	35 (46)	30 (40)
Suborbital width	18–20 (23–26)	18–20	15 (20)	15–19 (20–26)	17 (23)	18 (24)
Upper-jaw length	34–43 (45–55)	35–44	32 (41)	30–35 (39–47)	32 (43)	31 (42)
Length of rictus	28–35 (38–46)	–	25 (32)	23–30 (31–40)	24 (32)	23 (31)
Length of PMX tooth band	20–27 (26–35)	–	17 (22)	15–21 (20–28)	17 (22)	15 (21)
Preoral length	14–16 (19–21)	14–18	18 (24)	14–20 (19–26)	20 (27)	19 (26)
Dist. tip and lat. angle of snout	11 (14–15)	–	12 (15)	11–14 (15–19)	13 (17)	14 (19)
Snout width	19–20 (24–26)	–	19 (25)	18–23 (24–31)	24 (31)	26 (35)
Internasal width	18–19 (23–24)	17–20	16 (21)	17–20 (23–26)	18 (23)	22 (30)
Interorbital width	23–30 (30–39)	23–26	22 (29)	24–25 (32–34)	24 (32)	27 (37)
Body width over P. bases	47–60 (62–79)	–	42 (54)	28–58 (37–77)	52 (69)	40 (54)
Body depth at 1D. origin	97–114 (129–150)	–	82 (107)	76–102 (100–141)	110 (146)	87 (119)
Body depth at A. origin	91–108 (120–143)	–	72 (94)	59–95 (78–127)	100 (132)	79 (107)
Pre-V. length	109–111 (144–145)	–	105 (136)	93–117 (123–161)	106 (141)	103 (140)
Pre-anus length	124–125 (163–165)	–	114 (148)	105–129 (140–178)	113 (149)	109 (148)
Pre-A. length	146–147 (192–194)	–	138 (179)	128–154 (171–212)	145 (192)	126 (171)
Isthmus–V. distance	37–45 (48–60)	–	36 (47)	33–42 (44–59)	41 (54)	46 (63)
Isthmus–anus distance	50–60 (66–79)	–	– (–)	42–54 (58–74)	52 (68)	53 (72)
Isthmus–A. distance	75–81 (97–106)	–	– (–)	63–82 (82–113)	87 (116)	69 (94)
V.–A. distance	39–40 (51–53)	–	42 (54)	29–49 (38–65)	50 (66)	29 (39)
Anus–A. distance	23–26 (31–35)	–	26 (34)	17–32 (22–42)	35 (47)	18 (24)
V. length	37–55 (49–73)	–	– (–)	59–89 (76–118)	66 (88)	67 (91)
P. length	59–70 (77–93)	–	– (–)	59–68 (80–92)	63 (83)	59 (81)
Pre-1D. length	117–121 (155–157)	–	114 (148)	108–120 (144–161)	123 (162)	117 (159)
Height of 1D.	– (–)	–	– (–)	102–117 (137–156)	110 (146)	108 (147)
Length of 1D. base	33–41 (44–54)	–	27 (34)	19–31 (26–41)	32 (43)	31 (42)
Interdorsal length	27–31 (36–41)	–	39 (50)	20–38 (26–52)	27 (35)	25 (34)
Length of gill slit	16–18 (22–23)	18–20	17 (22)	14–20 (19–26)	14 (19)	16 (21)
Length of posterior nostril	6–8 (7–10)	–	8 (11)	6–9 (8–11)	6 (8)	6 (8)
Barbel length	22–25 (29–33)	20–29	23 (30)	19–25 (25–34)	22 (29)	21 (28)
Counts						
1D. rays	II,11–13	II,11–12	II,10	II,9–10	II,10	II,11
P. rays	i22–i24	i21–i24	i21–i22	i21–i26	i22	i25–i27
V. rays	12–13	11–13 (rarely 9 or 14)	11	11–12	15	15
Outer GR on first arch	5–7	6–8	8	7–9	7	9

...Continued on the next page

TABLE 10. (Continued)

Species	<i>K. bubonis</i>		<i>K. dara</i>		<i>K. endoi</i>	
Type status	NT		HT	NT	HL	PT
Number of specimens	<i>n</i> = 3	<i>n</i> = 12**	<i>n</i> = 1	<i>n</i> = 8	<i>n</i> = 1	<i>n</i> = 1
Inner GR on first arch	10	8–10	9	9–11	9	9
Outer GR on second arch	9	9–10	9	8–10	8	9
Inner GR on second arch	9	9–10	9	9–10	9	9
Longitudinal scales	44–48	44–61	53	35–48	56	58
Scales below 1D. origin	19–20	19–21	—	13–16	19	19
Scales below 1D. midbase	12–13	12–16	10.5	8.5–10.5	15.5	13
Scales below 2D. origin	14.5–15.5	12–15	11	10–11.5	16	15
Scales above A. origin	—	—	—	36	—	—

*Abbreviations: 1D.—first dorsal fin; 2D.—second dorsal fin; A.—anal fin; GR—gill rakers; HT—holotype; NT—non-type; P.—pectoral fin; PMX—premaxillary; PT—paratype; V.—pelvic fin

**Shcherbachev *et al.* (1992)

FIGURE 152. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of two species of *Kuronezumia*. (A–B) *K. dara*, BSKU 45036, 60.0 mm HL; (C–D) *K. endoi* **sp. nov.**, MSM 71-945, holotype, 32.7 mm HL. (A, C) Views from above; (B, D) oblique views. [Photos: (A–B) N. Nakayama; (C–D) NSMT]

FIGURE 153. Fresh specimen of *Kuronezumia endoi* **sp. nov.** BSKU 44829, paratype, 26.1 mm HL, 159+ mm TL, Tosa Bay, Kochi Pref., Pacific, 500 m depth. Lateral view. [Photo: BSKU]

FIGURE 154. Holotype of *Kuronezumia endoi* **sp. nov.** MSM 71-945, 32.7 mm HL, 215+ mm TL, Suruga Bay, Shizuoka Pref., Pacific. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

the absence of a large spiny scute at the tip of the snout (vs. present); from *K. dara* (Gilbert & Hubbs, 1916) and *K. pallida* Sazonov & Iwamoto, 1992 in the presence of the cephalic sensory pores (Fig. 156 vs. absent); from *K. bubonis* and an undescribed species from the South China Sea (sensu Iwamoto 1974; Shcherbachev *et al.* 1992) in the absence of a large tubercular swelling anterior to the anus (vs. present); and from all except *K. leonis*, *K. pallida*, and the above undescribed species in the presence of scaly patches on lower portions of the branchiostegal membranes (Fig. 156C vs. absent).

It is putatively most similar to *K. paepkei* Shcherbachev, Sazonov & Iwamoto, 1992 known only from the holotype collected from the western Indian Ocean off Kenya. In addition to the above-mentioned differences, *K. endoi* further differs from *K. paepkei* in having fewer gill rakers (9 vs. 12 in the inner side of the first arch; 8–9 vs. 11 on the outer side of the second arch), and smaller body scales (longitudinal scales 56–58 vs. 34–37; transverse scale rows below first dorsal-fin origin 19 vs. 13; below first dorsal-fin midbase 13–15.5 vs. 7; below second dorsal-fin origin 15–16 vs. 8).

FIGURE 155. Preserved specimen of *Kuronezumia endoi* **sp. nov.** BSKU 44829, paratype, 26.1 mm HL, 159+ mm TL, Tosa Bay, Kochi Pref., Pacific, 500 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

FIGURE 156. Diagrammatic illustrations showing the cephalic sensory system and naked areas (gray) of *Kuronezumia endoi* **sp. nov.** MSM 71-945, 32.7 mm HL. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Illustrations: N. Nakayama]

Genus *Lucigadus* Gilbert & Hubbs, 1920

[No Japanese name]

Lucigadus Gilbert & Hubbs, 1920:553 (type species: *Macrourus lucifer* Smith & Radcliffe in Radcliffe, 1912, by original designation).

Diagnosis. Anus closer to pelvic-fin bases than to anal-fin origin; periproct moderately narrow; 2 dermal windows of light organ, one immediately anterior to anus, the other between pelvic-fin bases. Infraorbital ridge not connected with preopercular ridge, separated by distinct gap. Second spinous ray of first dorsal fin serrated along its leading edge. Ventral region of body positioned well forward so that gill membranes unite below orbits. Pelvic fin inserted below opercle. Snout smoothly rounded, lacking enlarged scutes. Head fully covered with small, unmodified scales. Chin barbel well developed. Teeth in narrow tapered bands in both jaws; premaxillary band extending beyond posterior edge of maxillary process. Body scales covered with short, erect, needle-like spinules; buttresses of spinules and reticulate structure absent. Grooved lateral line present. Branchiostegal rays 7. Body with silvery reflection when fresh. [Modified from Iwamoto & Anderson (1994), Iwamoto & Merrett (1997), and Iwamoto & Okamoto (2015).]

Remarks. *Lucigadus* was previously recognized as a subgenus of *Ventrifossa* Gilbert & Hubbs, 1920 (e.g., Iwamoto 1979, 1990), but Sazonov (1985) elevated this group to full generic rank. Most records of the genus were from tropical waters of the western Pacific or subtropical waters of the Southern Hemisphere (Iwamoto & Okamoto 2015). According to Iwamoto & Merrett (1997) and Iwamoto & Okamoto (2015), *Lucigadus* currently comprises the following seven species: *L. acrolophus* Iwamoto & Merrett, 1997 known only from New Caledonia and the Norfolk Ridge; *L. borealis* Iwamoto & Okamoto, 2015 recently described from the Emperor Seamounts; *L. lucifer* (Smith & Radcliffe in Radcliffe, 1912) probably endemic to the Philippines; *L. microlepis* (Günther, 1878), a senior synonym of *L. fasciatus* (Weber, 1913) and *L. vittatus* (Weber, 1913), widespread in the southwestern Pacific; *L. nigromaculatus* (McCulloch, 1907) recorded from southeastern Australia, New Zealand, and off Chile; *L. nigromarginatus* (Smith & Radcliffe in Radcliffe, 1912) known from the Philippines, Indonesia, China, and Taiwan; and *L. ori* (Smith, 1968) disjunctly distributed off the east coast of Africa and the west coast of Australia. A nominal species, *Macruropus potronus* Pequeño, 1971, was tentatively regarded as a junior synonym of *L. nigromaculatus* by Iwamoto & Okamoto (2015). Although the genus has not been recorded from Japanese waters, *L. nigromarginatus* is known from off the northeastern coast of Taiwan.

Sazonov (1985) considered this group to be closest

to *Malacocephalus* Günther, 1862, but *Lucigadus* differs notably from the latter in that the ventral region of the body is positioned well forward so that the gill membranes unite below the orbits (vs. posterior to this level). They are further distinguished from each other by the combination of the lower jaw dentition (tapering band in *Lucigadus* vs. generally uniserial in *Malacocephalus*) and the serration along the second spinous ray of the first dorsal fin [finely serrated along its leading edge vs. entirely smooth in most species (subgenus *Malacocephalus*)]. On the other hand, *Lucigadus* is similar in general appearance to *Kuronezumia* Iwamoto, 1974, but readily differs in the absence of a reticulate structure on the body scales (vs. well developed in posterior parts of the unexposed portion) and its paler body coloration (vs. uniformly dark).

Lucigadus nigromarginatus (Smith & Radcliffe in Radcliffe, 1912)

[No Japanese name]

(Figs. 157–159; Appendix 3-9E)

Macrourus nigromarginatus Smith & Radcliffe in Radcliffe, 1912:114, pl. 24, fig. 2 [original description; holotype: USNM 72930, from near Simaluc Island, 5°33'15"N, 120°15'30"E, in 303 ftn (554 m); numerous paratypes].

Lucigadus lucifer (not Smith & Radcliffe in Radcliffe 1912): Chiou et al. 2004a:45, fig. 15 (brief description; 1 spec. from Nan-fang-ao; first record from Taiwan).

Ventrifossa lucifer (not Smith & Radcliffe in Radcliffe 1912): Chiou et al. 2004a: table 1 (listed; Taiwan).

Ventrifossa nigromarginata: Chiou et al. 2004a: table 1 (listed; first record from Taiwan).

Lucigadus nigromarginatus: Shao et al. 2008b: table 2 (2 spec. listed from northeastern and southwestern Taiwan); Iwamoto et al. 2015:84 (brief description; 9 spec. from northeastern and southwestern Taiwan, South China Sea, Philippines, and New Caledonia).

Diagnosis. A species of *Lucigadus* with 10–11 pelvic-fin rays; first dorsal fin blackish over anterior 3–4 rays; distal margin of anal fin prominently marked with black anteriorly; snout length 24–28% HL; interorbital width 22–31% HL; height of first dorsal fin 92–107% HL; barbel length 20–26% HL; scales below second dorsal-fin origin 7–9.5; branchiostegal membranes heavily scaled ventrally; body pale overall, without broad dark bands. [Modified from the key to *Lucigadus* species provided by Iwamoto & Okamoto (2015:372).]

Material examined. 2 specimens. **Taiwan:** BSKU 98982 (1, 35.2 mm HL, 175+ mm TL), Dong-gang fish market, Pingtung, bottom trawl, coll. H.-C. Ho, 16 Nov. 2007; FAKU 204468 (1, 31.0 mm HL, 156+ mm TL), Dong-gang fish market, Pingtung, bottom trawl, coll. N. Nakayama, 27 Feb. 2017.

Counts and measurements. For counts and

FIGURE 157. Fresh specimen of *Lucigadus nigromarginatus*. FAKU 204468, 31.0 mm HL, 156+ mm TL, Dong-gang fish market, southwestern Taiwan, South China Sea. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

FIGURE 158. Preserved specimen of *Lucigadus nigromarginatus*. BSKU 98982, 35.2 mm HL, 175+ mm TL, Dong-gang fish market, southern Taiwan, South China Sea, ca. 200–300 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

measurements see Iwamoto *et al.* (2015:84) and Iwamoto & Okamoto (2015: table 1).

Size. To about 20 cm TL (Iwamoto & Okamoto 2015).

Distribution. Known only from the western central Pacific including the Philippines (type locality), Indonesia, China, and Taiwan, at depths of 247–717 m (Radcliffe 1912; Gilbert & Hubbs 1920; Weber & de Beaufort 1929; Shao *et al.* 2008a, 2008b). No specimens have been collected within Japan's EEZ (Appendix 3-9E). Common in Taiwan.

Remarks and comparisons. For further morphological information see the original description given by Smith & Radcliffe (*in* Radcliffe 1912). In the key to *Lucigadus* species provided by Iwamoto &

Okamoto (2015), *L. nigromarginatus* is most similar to *L. acrolophus* Iwamoto & Merrett, 1997. According to their study, *L. nigromarginatus* differs from *L. acrolophus* in having a shorter snout (24–28% HL vs. 28–32%), a lower first dorsal fin (92–107% HL vs. 108–129%), an apical black spot on the first dorsal fin (vs. anterior third of the fin membrane blackish), and a blackish anterodistal margin of the anal fin (vs. absent).

Genus *Malacocephalus* Günther, 1862

[Japanese name: Manjudara-zoku]

Malacocephalus Günther, 1862:396 (type species: *Macrourus*

FIGURE 159. Scanning electron micrographs showing a body scale (from the dorsum below the interdorsal space) of *Lucigadus nigromarginatus*. BSKU 98982, 35.2 mm HL. (A) View from above; (B) oblique view. [Photos: N. Nakayama]

laevis Lowe, 1843, by original designation).

Pawnurus Parr, 1946:27 (as subgenus of *Malacocephalus*; type species: *Malacocephalus occidentalis* Goode & Bean, 1885, by original designation).

Diagnosis. Anus closer to pelvic-fin bases than to anal-fin origin; periproct moderately broad; 2 dermal windows of light organ, one small, immediately anterior to anus, the other large, between pelvic-fin bases. Infraorbital ridge not connected with preopercular ridge, separated by distinct gap. Leading edge of second spinous ray of first dorsal fin smooth or serrated. Chin barbel well developed. Teeth in 2 rows or narrow broad in upper jaw, with outer series greatly enlarged; lower jaw teeth arranged in uniserial row. Body scales covered with long, erect, needle-like spinules in quincunx order; buttresses of spinules scarcely developed; reticulate structure absent. Grooved lateral line present. Cephalic sensory canals lacking open pores. Pyloric caeca numerous, about 50–100. Branchiostegal rays 7. Body gray to swarthy, with silvery reflection when fresh. [Modified from Sazonov & Iwamoto (1992), Iwamoto & Merrett (1997), and Iwamoto & Graham (2001).]

Remarks. According to Iwamoto (1990), *Malacocephalus* comprises two subgenera: *Malacocephalus* and *Pawnurus* Parr, 1946. The subgenus *Malacocephalus* includes four nominal species, viz., *M. hawaiiensis* Gilbert, 1905 confined to Hawaii; *M. luzonensis* Gilbert & Hubbs, 1920 so far known only from the Philippines; *M. nipponensis* Gilbert & Hubbs, 1916 recorded from Japan and Taiwan; and *M. laevis* (Lowe, 1843) widespread in tropical and temperate waters of the world's oceans (see also the Remarks of *M. nipponensis*). On the other hand, the subgenus *Pawnurus* includes three valid species: *M. boretzii* Sazonov, 1985 endemic to the Emperor Seamounts; *M. occidentalis* Goode & Bean,

1885 distributed in both sides of the Atlantic; and *M. okamurai* Iwamoto & Arai, 1987 known only from the Atlantic coast of South America.

***Malacocephalus nipponensis* Gilbert & Hubbs, 1916**

[Japanese name: Nihon-manjudara]
(Figs. 160–162; Appendix 3-9F)

Malacocephalus nipponensis Gilbert & Hubbs, 1916:189, pl. 9, fig. 2 [original description; holotype: USNM 76866, from “off the east coast of central Hondo” (off Cape Shionomisaki), *Albatross* sta. 4967, in 244–253 fms (447–463 m)]; Okada & Matsubara 1938:452, pl. 111, fig. 1 (in key; Japan; new Japanese name: “Nihon-manju-dara”); Kamohara 1938:72 (brief description; 9 spec. from Kochi Pref.; new Japanese name: “Bôzu-dara”); Kamohara 1950:279 (listed; Kochi and Wakayama Pref.); Kamohara 1952:98 (selected counts and measurements; spec. from Kochi Pref.); Matsubara 1955:1315 (in key; Japan); Kamohara 1958:74 (listed; Kochi Pref.); Kamohara 1964:96 (listed; Kochi Pref.); Okamura 1982:145, 347, fig. 88 (brief description; 1 spec. from Tosa Bay; BSKU 29516); Okamura 1984a:203, 360, fig. 145 (brief description; 18 spec. from Okinawa Trough; photo based on BSKU 29587); Okamura 1984b:93, pl. 80, fig. J (compiled); Okamura 1988:93, pl. 80, fig. J (compiled); Nakabo 1993:356 (in key; Japan); Okamura 1997:128, fig. 6 (compiled); Nakabo 2000:420 (in key; Japan); Shinohara *et al.* 2001:306 (listed; Tosa Bay); Nakabo 2002:420 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Shinohara *et al.* 2005:418 (listed; Ryukyu Islands); Shao *et al.* 2008b: table 2 (2 spec. listed from South China Sea; first record from Taiwan); Nakabo & Kai 2013:496 (in key; Japan); Ikeda & Nakabo 2015:319, pl. 65, figs. 1–5 (brief description; spec. from Wakayama Pref.); Iwamoto *et al.* 2015:86 (brief

description; 20 spec. from northeastern and southwestern Taiwan, and South China Sea); Iwatsuki *et al.* 2017:32 (listed; Hyuga-nada).

Malacocephalus laevis (not Lowe 1843): Okada & Matsubara 1938:452 (in key; Japan; new Japanese name: “Manju-dara”); Matsubara *et al.* 1951:42 (listed; Mie Pref.); Kuroda 1951:392 (listed; Suruga Bay); Kuroda 1952:177 (1 spec. from Suruga Bay); Matsubara 1955:1315 (in key; Japan); Kamohara 1958:74 (listed; Kochi Pref.); Okada *et al.* 1959:83, pl. 2, fig. 13 (listed; Kumano-nada); Kamohara 1964:96 (listed; Kochi Pref.); Matsubara 1965:508 (compiled; Japan); Okamura 1970a:69, pl. IV, text-fig. 31 (description; biological notes; 56 spec. from Pacific off southern Japan from Choshi to Totoro); Okamura 1970b: table 1 (listed; Japan); Kataoka & Tomida 1981:78 (listed; Mie Pref.); Tominaga & Uyeno 1981:489 (listed; Japan); Shao 1993:172, fig. 38-1 (compiled; Taiwan; photo spec. is *Lucigadus nigromarginatus*); Chiou *et al.* 2004a: table 1 (listed; Taiwan); Shao *et al.* 2008b: table 2 (9 spec. listed from northeastern, eastern, and southwestern Taiwan); Motomura 2020:39 (listed; Japan).

Diagnosis. Second spinous ray of first dorsal fin smooth along its leading edge; pelvic-fin rays 8–10 (usually 9);

mandibular teeth canine-like, widely spaced, arranged in 1 distinct row; anterior dermal window of light organ large, bean-shaped, laterally expanded; body scale very small, covered with short, erect, needle-like spinules; scales below second dorsal fin without enlarged median spinules; gill membranes heavily scaled along branchiostegal rays; snout bluntly pointed in lateral view, length 26–31% HL; tip of snout about on horizontal through midorbit; mouth large, upper-jaw length 46–52% HL; interorbital space broad, width 31–37% HL; height of first dorsal fin 56–72% HL; pectoral fin short, length 51–63% HL; outer gill slight moderately wide, length 24–29% HL; leading edges of snout, supranarial ridges, and upper rim of orbits not marked with black.

Material examined. 30 specimens. **Holotype of *Malacocephalus nipponensis*:** USNM 76866 (69.1 mm HL, 400+ mm TL), off Cape Shionomisaki, Wakayama Pref., Kii Peninsula, Japan, 33.4194°N, 135.6222°E, 244–253 ftn (447–463 m), *Albatross* sta. 4967, 9-ft *Albatross-Blake* beam trawl, 29 Aug. 1906. **Non-types. Japan:** BSKU 29742 (1, 77.0 mm HL, 420+ mm TL), north-northeast of Miyako-jima Island, Okinawa Trough, 25.9833°N, 125.8500°E, 430 m, F/V *Yuryo-maru*, No. 8,

FIGURE 160. Fresh specimens of *Malacocephalus nipponensis*. (A) BSKU 112433, 46.3 mm HL, 274+ mm TL, Tosa Bay off Susaki, Kochi Pref., Pacific, 320–380 m depth; (B–D) BSKU 112450, 70.9 mm HL, 415+ mm TL, collected with BSKU 112433. (A–B) Lateral views; (C) dorsal and (D) ventral views of the head and trunk. [Photos: N. Nakayama]

FIGURE 161. Holotype of *Malacocephalus nipponensis*. USNM 76866, 69.1 mm HL, 400+ mm TL, off Cape Shionomisaki, Wakayama Pref., Pacific, 447–463 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

FIGURE 162. Scanning electron micrographs showing a body scale (from the dorsum below the interdorsal space) of *Malacocephalus nipponensis*. BSKU 13317, 65.3 mm HL. (A) View from above; (B) oblique view. [Photos: N. Nakayama]

sta. 2-T3, bottom trawl, coll. H. Maeda, 10 Oct. 1979; BSKU 13314 (1, 43.1 mm HL, 263+ mm TL), BSKU 13316 (1, 56.9 mm HL, 264+ mm TL), BSKU 13317 (1, 65.3 mm HL, 338+ mm TL), BSKU 13323 (1, 40.0 mm HL, 244+ mm TL), ca. 10–15 km west of Cape Murotomisaki, Tosa Bay, 360–435 m, 8–9 Jun. 1968; BSKU 113055 (1, 76.8 mm HL, 411+ mm TL), BSKU 113056 (1, 60.6 mm HL, 344+ mm TL), off Aki, Tosa Bay, 310–390 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 7 Apr. 2014; BSKU 45321 (1, 55.4 mm HL, 315+ mm TL), off Usa, Tosa Bay, 300–400 m, bottom trawl, 5 Nov. 1988; BSKU 10029 (1, 72.6 mm HL, 423+ mm TL), Mimase

fish market, 10 Feb. 1961; BSKU 102911 (1, 70.3 mm HL, 448+ mm TL), BSKU 102912 (1, 71.2 mm HL, 413+ mm TL), BSKU 102913 (1, 65.6 mm HL, 378+ mm TL), BSKU 102928 (1, 67.1 mm HL, 366+ mm TL), Mimase fish market, 400 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama, 18 Apr. 2010; BSKU 45406 (1, 58.2 mm HL, 345+ mm TL), Mimase fish market, bottom trawl, 22 Nov. 1988; KSHS 19471 (1, 73.7 mm HL, 441+ mm TL), KSHS 19494 (1, 78.2 mm HL, 449+ mm TL), KSHS 19495 (1, 81.5 mm HL, 483+ mm TL), Mimase fish market, coll. T. Yamakawa, 15 Apr. 1981; KSHS 5836 (1, 81.0 mm HL, 325+ mm TL), Mimase fish market, coll. T.

Yamakawa, 28 Feb. 1967; BSKU 13635 (1, 52.4 mm HL, 291+ mm TL), Mimase fish market, coll. Y. Mishima (= Y. Machida), Jan.–Mar. 1968; BSKU 93850 (1, 45.7 mm HL, 277+ mm TL), Mimase fish market, F/V *Kosei-maru*, coll. N. Nakayama, 28 Feb. 2008; BSKU 112433 (1, 46.3 mm HL, 274+ mm TL), BSKU 112450 (1, 70.9 mm HL, 415+ mm TL), off Susaki, Tosa Bay, 320–380 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 9 Mar. 2014; BSKU 113961 (1, 61.6 mm HL, 357+ mm TL), BSKU 113962 (1, 43.8 mm HL, 276+ mm TL), off Susaki, Tosa Bay, 350–365 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 23 Apr. 2014; BSKU 29516 (1, 75.4 mm HL, 431+ mm TL), Tosa Bay, 32.9833°N, 133.4833°E, 453 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, cr. 4-Ky, tr. 18, bottom trawl, coll. O. Okamura *et al.*, 21 Dec. 1979; BSKU 39954 (1, 70.0 mm HL, 389+ mm TL), off Okitsu, Tosa Bay, 270 m, bottom trawl, 30 Mar. 1984; BSKU 112512 (1, 41.6 mm HL, 257+ mm TL), off Okitsu, Tosa Bay, 380 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 11 Mar. 2014; BSKU 45376 (1, 50.7 mm HL, 281+ mm TL), off Okitsu, Tosa Bay, bottom trawl, 1 Dec. 1988; BSKU 110031 (1, 45.9 mm HL, 266+ mm TL), Suruga Bay, 34.7127°N, 138.4553°E, 262–434 m, F/V *Hinode-maru*, sta. 1, bottom trawl, coll. N. Nakayama and R. Misawa, 23 Apr. 2013.

Counts and measurements. Based on 30 specimens (40–81.5 mm HL, 244+–483+ mm TL). Counts: first dorsal-fin rays II, 10–12; pectoral-fin rays i16–i20; pelvic-fin rays 8–10; gill rakers on first arch (outer/inner) 8–12/9–12, on second arch 9–12/8–11; longitudinal scales 61–90; transverse scale rows below first dorsal-fin origin 15–22, below first dorsal-fin midbase 11.5–15.5, below second dorsal-fin origin 10.5–14.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 26–31 (33–43); orbit diameter 29–37 (39–50); postorbital length 41–47 (54–64); postrostral length 72–78; orbit–preopercle distance 41–49 (53–65); suborbital width 11–14 (14–19); upper-jaw length 46–52 (60–69); length of rictus 38–43 (50–58); length of premaxillary tooth band 28–34 (38–45); preoral length 18–21 (23–28); distance between tip and lateral angle of snout 14–17 (18–23); snout width 22–27 (29–36); internasal width 21–26 (28–33); interorbital width 31–37 (41–48); body width over pectoral-fin bases 43–65 (57–87); body depth at first dorsal-fin origin 75–92 (99–123); body depth at anal-fin origin 73–94 (96–124); prepelvic length 100–123 (128–169); preanus length 113–132 (148–179); preanal length 121–144 (158–195); isthmus–pelvic distance 32–48 (44–64); isthmus–anus distance 40–62 (55–81); isthmus–anal distance 52–74 (71–98); pelvic–anal distance 21–33 (28–44); anus–anal distance 7–17 (9–23); pelvic-fin length 30–41 (40–54); pectoral-fin length 51–63 (70–85); predorsal length 106–115 (137–156); height of first dorsal fin 56–72 (77–94); length of first dorsal-fin base 22–33 (30–44); interdorsal length 43–76 (57–101);

length of gill slit 24–29 (31–38); length of posterior nostril 4–8 (5–10); barbel length 11–27 (15–35).

Size. To 52 cm TL (Okamura 1970a).

Distribution. This species appears to be restricted to Japan and Taiwan. Known from off the Pacific coasts of Japan northward to Choshi (35.74°N; Chiba Pref.), Okinawa Trough, and northern South China Sea (Shao *et al.* 2008; this study; Appendix 3–9F). Depth range 250–550 m, plus a single specimen captured at a depth of 445–1185 m (ASZIP 65517, west of Taiwan). Common.

Remarks. The reader can refer to Gilbert & Hubbs (1916) and Okamura (1970a; as *M. laevis* Lowe, 1843) for a full description of this species. *Malacocephalus nipponensis* was originally described from a single specimen collected from off Cape Shionomisaki, Wakayama Pref., Japan (Fig. 161). Unfortunately, the holotype is in poor condition, with a damaged snout and abdomen.

The synonymies of the species of the subgenus *Malacocephalus* have been problematic over the last half century. In his review of Japanese grenadiers, Okamura (1970a) synonymized *M. nipponensis* with *M. laevis*, suggesting no significant differences between the two species. However, he (1982) subsequently considered *M. nipponensis* to be distinct from *M. laevis*, and distinguished these taxa by the presence/absence of a scaly patch on the gular membranes (absent in *M. nipponensis* vs. present in *M. laevis*) and the shape and size of the anterior dermal window (ADW) of the light organ (small and circular vs. large and triangular). According to Okamura (1984a), who examined the holotype of *M. laevis*, *M. nipponensis* further differs from that species in having a wider interorbital space (equal to or greater than the orbit diameter vs. less than the orbit diameter) and ADW situated between the pelvic-fin bases (vs. anterior to a line connecting outer pelvic-fin bases).

On the other hand, Marshall (1973:650–651) considered all nominal species valid, noting that “examination of a good series of individuals may show that *nipponensis* is identical with *hawaiiensis*” (= *M. hawaiiensis* Gilbert, 1905). Iwamoto (1979, 1990) further indicated that not only *M. nipponensis* but also *M. hawaiiensis* are closely related to *M. laevis*, suggesting that these three taxa may eventually prove to represent one widely distributed species. The separation between *M. laevis* and *M. luzonensis* Gilbert & Hubbs, 1920 was also doubted by Iwamoto & Anderson (1994) and Iwamoto & Williams (1999). In a recent study, Iwamoto *et al.* (2015) tentatively treated *M. nipponensis* as a valid taxon and applied this name to the population of Japan and Taiwan. As noted in their study, it is unknown how many species are involved in this group, and a comprehensive study is required to elucidate the taxonomic relationships of all nominal species. The present study tentatively follows Iwamoto *et al.* (2015) in regarding *M. nipponensis* as a valid species.

Genus *Nezumia* Jordan in Jordan & Starks, 1904

[Japanese name: Nezumidara-zoku]

[?] *Macruroplus* Bleeker, 1874:369 (type species: *Macrourus serratus* Lowe, 1843, by original designation). [See also Iwamoto (1990:248).]

Nezumia Jordan in Jordan & Starks, 1904:620 (type species: *Nezumia condylura* Jordan & Gilbert in Jordan & Starks, 1904, by original designation).

Diagnosis. Anus closer to pelvic-fin bases than to anal-fin origin; periproct teardrop-shaped, narrow to moderately broad. Small dermal window of light organ between pelvic-fin bases. Infraorbital ridge not connected with preopercular ridge, separated by distinct gap. Second spinous ray of first dorsal fin serrated along its leading edge. Ventral aspects of body not positioned forward, and anal-fin origin below or posterior to hind margin of first dorsal-fin base. Chin barbel present. Teeth small, in narrow to broad bands in both jaws; premaxillary band ending anterior to posterior edge of maxillary process. Tip and lateral angles of snout armed with prominent scutes. Suborbital shelf formed of stout, coarsely modified scales. Underside of snout variously naked. Spinules on body scales needle-like, triangular, or lanceolate; buttresses of spinules rudimentary; reticulate structure developed on posterior parts of unexposed portion. Grooved lateral line present (rarely interrupted anteriorly). Cephalic sensory pores usually present. Branchiostegal rays 7 (rarely 6). Body dark to dusky, lacking silvery reflection when fresh. [Modified from Sazonov & Iwamoto (1992) and Iwamoto & Williams (1999).]

Remarks. *Nezumia* comprises about 50 species (Iwamoto *et al.* 2015; this study), seven of which, including one new species, are herein recorded from Japan and adjacent waters. Some species may well be transferred to other genera if more closely studied (Iwamoto & Williams 1999; Iwamoto *et al.* 2015), and a thorough phylogenetic analysis of the seven-branchiostegal-rayed grenadiers is needed to address their generic placements. The genus shows great abundance in temperate to subtemperate areas of the world's oceans (Iwamoto & Merrett 1997; Nakayama & Endo 2015), with the highest diversity in the eastern Pacific (10 species; Merrett & Iwamoto 2000: table 4).

Key to species of *Nezumia* from Japan and adjacent waters

- 1a Pelvic-fin rays 13–17 2
- 1b Pelvic-fin rays 8–11 3
- 2a Snout length less than orbit diameter; transverse scale rows below second dorsal-fin origin 10–12.5; body scales covered with needle-like spinules in discrete parallel to slightly convergent rows *N. condylura*

- 2b Snout length longer than orbit diameter; transverse scale rows below second dorsal-fin origin 7.5; body scales covered with narrowly lanceolate spinules in tightly packed convergent rows *N. kamoharai*
- 3a Snout barely protruding beyond upper jaw, its ventral contour almost vertical 4
- 3b Snout protruding well beyond upper jaw, its ventral contour oblique 5
- 4a Transverse scale rows below second dorsal-fin origin 9–10; orbit–preopercle distance 39–41% HL; interorbital width 20–22% HL; pelvic-fin length 46–54% HL; length of gill slit 13–16% HL *N. rara* sp. nov.
- 4b Transverse scale rows below second dorsal-fin origin 8.5; orbit–preopercle distance 42–43% HL; interorbital width 23–25% HL; pelvic-fin length 69% HL; length of gill slit 17–18% HL *N. tomiyamai*
- 5a Underside of head mostly scaled, naked areas confined to underside of snout and anterior portions of mandibular rami; spinules on body scales lanceolate *N. proxima*
- 5b Underside of head completely naked including mandibular rami; spinules on body scales needle-like 6
- 6a Pelvic-fin rays 10; spinules on body scales arranged in discrete parallel rows; second spinous ray of first dorsal fin not especially prolonged, height of first dorsal fin about equal to HL *N. shinoharai*
- 6b Pelvic-fin rays 8; spinules on body scales densely scattered over exposed portion; second spinous ray of first dorsal fin greatly prolonged, height of first dorsal fin distinctly longer than HL *N. spinosa*

Nezumia condylura Jordan & Gilbert in Jordan & Starks, 1904

[Japanese name: Nezumidara]

(Figs. 163–165; Table 11; Appendix 3–10A)

Nezumia condylura Jordan & Gilbert in Jordan & Starks, 1904:620 [original description; holotype: USNM 50937, from Suruga Bay, *Albatross* sta. 3721, in 207–250 ftn (379–458 m); paratypes from Sagami Bay and Suruga Bay]; Jordan *et al.* 1913:413, fig. 384 (listed; Japan; new Japanese name: “Nezumi-dara”); Kamohara 1934a:54 (listed; Kochi); Kamohara 1934c:462 (1 spec. from Mimase, Kochi Pref.); Kamohara 1938:71 (spec. from Kochi Pref.); Okamura 1970a:85, pl. VII, text-fig. 38 (description; biological notes; 71 spec. from Pacific off southern Japan from Choshi to Totoro); Okamura 1970b: table 1 (listed; Japan); Tominaga & Uyeno 1981:489 (listed; Japan); Okamura 1982:161, 350, fig. 96 (brief description; 2 spec. from Tosa Bay; photo based on BSKU 29498); Ozawa 1983:13 (listed; off Makurazaki, Kagoshima Pref., East China Sea); Okamura 1984a:215, 362, fig. 151 (brief description; 10 spec. from Okinawa Trough; photo based on BSKU 29888); Okamura 1984b:94, pl. 81, fig. G (compiled); Okamura 1988:94, pl. 81, fig. G (compiled); Iwamoto 1990:269, fig. 625 (synopsis); Nakabo 1993:361 (in key; Japan); Shinohara *et al.* 1996:170 (2 spec. listed from

Pacific off Tohoku); Okamura 1997:128, fig. 15 (compiled); Shinohara & Matsuura 1997:292 (listed; Suruga Bay); Nakabo 2000:425 (in key; Japan); Shinohara *et al.* 2001:306 (15 spec. listed from Tosa Bay); Nakabo 2002:425 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Chiou *et al.* 2004a:45, fig. 16, table 1 (brief description; 9 spec. from Nan-fang-ao; first record from Taiwan); Shinohara *et al.* 2005:418 (listed; Ryukyu Islands); Suetsugu & Ohta 2005: tables 2–3 (listed; southeast of Boso Peninsula and Enshu-nada); Senou *et al.* 2006:421 (listed; Sagami Sea); Shinohara & Williams 2006:551 (listed; Sagami Sea); Kitagawa *et al.* 2008:39, unnumbered fig. (brief description; spec. from Pacific off Tohoku); Shao *et al.* 2008b: table 2 (25 spec. listed from northeastern and eastern Taiwan, and South China Sea); Shinohara *et al.* 2009:709 (4 spec. listed from Pacific off Tohoku); Motomura *et al.* 2010:77, fig. 35 (2 spec. listed from Yaku-shima Island, Kagoshima Pref.); Furuhashi *et al.* 2010: table 2 (51 spec. listed from northern Okinawa Trough); Nakabo & Kai 2013:501 (in key; Japan); Iwamoto *et al.* 2015:91 (brief description; 33 spec. from northeastern Taiwan, South China Sea, and Philippines); Iwatsuki *et al.* 2017:32 (listed; Hyuga-nada); Nakae *et al.* 2018:219 (3 spec. listed from Amami-oshima Island, Kagoshima Pref.); Motomura 2020:39 (listed; Japan).

Lionurus condylura: Gilbert & Hubbs 1916:195 (new combination; supplementary description; 29 spec. from Hyuga-nada, off Cape Shionomisaki, Suruga Bay, and Sagami Bay); Okada & Matsubara 1938:453 (in key; Japan); Kuroda 1951:392 (listed; Suruga Bay); Matsubara 1955:1316 (in key; Japan); Kamohara 1958:74 (listed; Kochi Pref.); Kamohara 1964:96 (listed; Kochi Pref.).

Lionurus condylurus: Kamohara 1950:278 (listed; Kochi and Wakayama Pref.); Kamohara 1952:98 (spec. from Kochi Pref.).

Nezumia propinquus (not Gilbert & Cramer 1897): Okamura 1982:163, 350, fig. 97 (first record from Japan; brief description; 6 spec. from Kyushu-Palau Ridge; photo based on BSKU 26027; new Japanese name: “Kiheri-nezumidara”); Okamura 1984b:94, pl. 81, fig. H (compiled); Okamura 1988:94, pl. 81, fig. H (compiled).

Nezumia propinqua (not Gilbert & Cramer 1897): Nakabo 1993:361 (in key; Japan); Shinohara *et al.* 1996:170 (2 spec. listed from Pacific off Tohoku); Okamura 1997:128, fig. 16 (compiled); Nakabo 2000:425 (in key; Japan); Nakabo 2002:425 (in key; Japan); Shinohara *et al.* 2009:709 (listed; Pacific off Tohoku); Nakabo & Kai 2013:501 (in key; Japan); Motomura 2020:39 (listed; Japan).

[?] *Nezumia evides* (not Gilbert & Hubbs 1920): Shao *et al.* 2008b: table 2 (5 spec. listed from eastern Taiwan and northern South China Sea); Iwamoto *et al.* 2015:92, fig. 18 (brief description; in part, 6 spec. from northeastern Taiwan and northern South China Sea).

Diagnosis. A species of *Nezumia* with 13–17 pelvic-fin rays. Snout short (length 27–30% HL), slightly protruding beyond upper jaw, ventral contour moderately steep in

lateral view; snout blunt in dorsal view; underside of head almost completely naked. Second spinous ray of first dorsal fin not especially prolonged (height of fin 91–106% HL). Body scales covered with short, moderately reclined, needle-like spinules in discrete parallel to slightly convergent rows; tip of last spinule in each row extending slightly beyond posterior scale margin; middle row of spinules not enlarged; transverse scale rows below second dorsal-fin origin 10–12.5. Cephalic sensory pores present; those on mandibular and infraorbital canals small but conspicuous. No prominent dark band encircling trunk; first dorsal fin blackish apically.

Material examined. 33 specimens. **Holotype of *Nezumia condylura*:** USNM 50937 (30.6 mm HL, 178+ mm TL), off Oigawa River, Shizuoka Pref., Suruga Bay, Japan, 34.7667°N, 138.3000°E, 207–250 ftm (379–458 m), *Albatross* sta. 3721, 8-ft Tanner beam trawl, 12 May 1900. **Paratypes of *N. condylura*:** *USNM 51472 (4, not measured), collected with holotype. **Non-types: Japan:** BSKU 109013 (1, 26.0 mm HL, 146+ mm TL), BSKU 109014 (1, 24.2 mm HL, 150+ mm TL), BSKU 109015 (1, 23.0 mm HL, 119+ mm TL), off Amami, East China Sea, 28.5686°N, 127.0419°E, 604 m, T/V *Nagasaki-maru*, cr. N365, sta. J6, tr. 4, 3-m ORE beam trawl, coll. N. Nakayama, 16 Nov. 2012; *BSKU 43418 (1, 26.1 mm HL, 176+ mm TL), south of Okino-shima Island, 32.6400°N, 132.5367°E, 440–451 m, R/V *Tansei-maru*, cr. KT-86-16, sta. A-3, 2-m beam trawl, 1 Nov. 1986; BSKU 13332 (1, 30.6 mm HL, 179+ mm TL), BSKU 13338 (1, 24.2 mm HL, 170+ mm TL), ca. 10–15 km west of Cape Murotomisaki, Tosa Bay, 360–435 m, coll. O. Okamura, 8–9 Jun. 1968; BSKU 99460 (1, 26.6 mm HL, 157+ mm TL), BSKU 99461 (1, 17.8 mm HL, 114+ mm TL), BSKU 99462 (1, 17.2 mm HL, 103+ mm TL), BSKU 99463 (1, 18.3 mm HL, 129+ mm TL), Tosa Bay, 33.1900°N, 133.6883°E, 542–585 m, FRV *Kotaka-maru*, otter trawl, coll. H. Endo, 22 Jan. 1998; BSKU 98941 (1, 31.2 mm HL, 209+ mm TL), BSKU 98942 (1, 30.8 mm HL, 188+ mm TL), Tosa Bay, 33.1428°N, 133.6248°E, 450–490 m, FRV *Kotaka-maru*, otter trawl, 1 Aug. 1988; BSKU 99126 (1, 31.9 mm HL, 205+ mm TL), BSKU 99129 (1, 25.7 mm HL, 147+ mm TL), Tosa Bay, 600 m, FRV *Kotaka-maru*, otter trawl, 17 Nov. 1987; *BSKU 44338 (1, 25.2 mm HL, 156+ mm TL), Tosa Bay, 780 m, FRV *Kotaka-maru*, otter trawl, 24 Nov. 1987; BSKU 101210 (1, 32.0 mm HL, 181+ mm TL), BSKU 44785 (1, 32.1 mm HL, 209+ mm TL), BSKU 44786 (1, 30.5 mm HL, 155+ mm TL), BSKU 44787 (1, 30.4 mm HL, 182+ mm TL), Tosa Bay, 700 m, FRV *Kotaka-maru*, otter trawl, 24 May 1988; BSKU 84384 (1, 32.4 mm HL, 200+ mm TL), Tosa Bay, 800 m, FRV *Kotaka-maru*, otter trawl, 25 Apr. 1990; NSMT-P 101248 (2, 22.3–34.9 mm HL, 146+–231+ mm TL, counts and measurements taken on 1 of 2), west of Matsuzaki, Suruga Bay, 34.7567°N, 138.6650°E, 652–676 m, R/V *Tansei-maru*, cr. KT-89-06, sta. SB26, 3-m ORE beam trawl, 17 May 1989; NSMT-

P 101432 (3, 18.2–33.6 mm HL, 124+–192+ mm TL, counts and measurements taken on 2 of 3), southeast of Miyake-jima Island, Shichito-Iojima Ridge, 34.1340°N, 139.6717°E, 520–644 m, R/V *Tansei-maru*, cr. KT-89-06, sta. ME11, 3-m ORE beam trawl, 19 May 1989; BSKU 26113 (1, 32.7 mm HL, 192+ mm TL), Kyushu-Palau Ridge, 27.9033°N, 134.6583°E, 700 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. 2-4, bottom trawl, coll. Y. Kobayashi *et al.*, 20 Feb. 1978; BSKU 26028 (1, 33.5 mm HL, 189+ mm TL), Kyushu-Palau Ridge, 27.9000°N, 134.6550°E, 650 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. 2-2, tr. 9, bottom trawl, coll. Y. Kobayashi *et al.*, 15 Feb. 1978; BSKU 26065

(1, 33.5 mm HL, 207+ mm TL), Kyushu-Palau Ridge, 27.9167°N, 134.6500°E, 700 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. 2-3, bottom trawl, coll. Y. Kobayashi *et al.*, 20 Feb. 1978.

Counts and measurements. See Table 11.

Size. To about 24 cm TL (NSMT-P 94610, 240+ mm TL, Kumano-nada, Japan).

Distribution. Known from Japan and Taiwan. Distributed in the Pacific off Japan northward to Kuji (40.19°N; Iwate Pref.), Okinawa Trough, Taiwan Strait, northern South China Sea, and Kyushu-Palau Ridge, at depths of 250–1185 m (Chiou *et al.* 2004a; Shao *et al.* 2008a, 2008b; this study; Appendix 3-10A). Very common

FIGURE 163. Fresh specimens of *Nezumia condylura*. (A) BSKU 44338, 25.5 mm HL, 156+ mm TL, Tosa Bay, Kochi Pref., Pacific, 780 m depth; (B) BSKU 109013, 26.0 mm HL, 146+ mm TL, off Amami-oshima Island, Kagoshima Pref., Okinawa Trough, East China Sea, 604 m depth; (C) BSKU 43418, 26.1 mm HL, 176+ mm TL, south of Okino-shima Island, Kochi Pref., Pacific, 440–451 m depth. Lateral views. [Photos: (A, C) BSKU; (B) N. Nakayama]

FIGURE 164. Holotype of *Nezumia condylura*. USNM 50937, 30.6 mm HL, 178+ mm TL, Suruga Bay, Shizuoka Pref., Pacific, 379–458 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

FIGURE 165. Scanning electron micrographs showing body scale (from the dorsum below the interdorsal space) of *Nezumia condylura*. BSKU 44785, 32.1 mm HL. (A) View from above; (B) oblique view. [Photos: NSMT]

on the upper slope of Pacific Japan, but seemingly rare in the East and South China Seas.

Remarks, relationships, and comparisons. For a full description see the original description given by Jordan & Gilbert (*in* Jordan & Starks 1904) and Okamura (1970a). *Nezumia condylura* was originally described from Suruga Bay, Japan. This species is quite similar to *N. propinqua* (Gilbert & Cramer, 1897), the latter of which was originally described from Hawaii and subsequently recorded from various parts of the Pacific and Indian Oceans, including Japan (Okamura 1982), South Africa (Iwamoto 1986; Iwamoto & Anderson 1994), the west

and east coasts of Australia (Iwamoto & Williams 1999; Iwamoto & Graham 2001), the New Caledonian Region (Iwamoto & Merrett 1997; Merrett & Iwamoto 2000), and the Sala y Gomez Ridge (Sazonov & Iwamoto 1992). Okamura (1982, 1984a) distinguished the two species primarily by the position of the anus (anus–anal distance less than pelvic–anus distance in *N. condylura* vs. about equal or greater in *N. propinqua*). However, because the suggested difference was quite subtle, Iwamoto (1990:270) stated that *N. condylura* might be a junior synonym of *N. propinqua*. Sazonov & Iwamoto (1992:74) also considered their conspecificity, although they noted that

TABLE 11. Morphometric measurements and meristic counts* for *Nezumia condylura*.

Type status	HT	NT	
Locality	Suruga Bay	Main islands of Japan	Kyushu-Palau Ridge
Number of specimens	<i>n</i> = 1	<i>n</i> = 21	<i>n</i> = 3
Measurements (mm)			
TL	178+	103+–231+	189+–207+
HL	30.6	17.2–34.9	32.7–33.5
% of HL (% of PRL)			
Snout length	28 (37)	27–30 (35–41)	26–29 (33–39)
Orbit diameter	33 (44)	31–37 (42–50)	31–36 (41–46)
Postorbital length	42 (56)	39–44 (52–60)	45–46 (57–63)
Postrostral length	75 (–)	73–76 (–)	74–78 (–)
Orbit–preopercle distance	37 (50)	33–38 (44–52)	35–39 (46–50)
Suborbital width	14 (19)	13–17 (17–22)	14–15 (19–20)
Upper-jaw length	32 (42)	29–34 (39–45)	28–35 (36–45)
Length of rictus	24 (32)	24–27 (32–35)	22–27 (29–35)
Length of PMX tooth band	16 (21)	14–19 (19–25)	17–18 (22–23)
Preoral length	23 (31)	22–26 (29–35)	18–21 (22–28)
Dist. tip and lat. angle of snout	17 (23)	15–19 (20–25)	15–16 (20–22)
Snout width	30 (40)	26–31 (34–42)	28–30 (37–41)
Internasal width	22 (29)	20–23 (27–32)	22–25 (28–33)
Interorbital width	22 (29)	21–27 (28–37)	23–28 (30–36)
Body width over P. bases	44 (59)	38–61 (50–81)	47–48 (62–64)
Body depth at 1D. origin	85 (114)	77–101 (105–135)	– (–)
Body depth at A. origin	73 (97)	62–95 (83–126)	– (–)
Pre-V. length	112 (149)	102–121 (136–163)	102–106 (135–142)
Pre-anus length	134 (179)	122–145 (167–191)	127–130 (163–176)
Pre-A. length	151 (201)	133–166 (181–228)	155–162 (200–214)
Isthmus–V. distance	47 (62)	35–44 (46–58)	40–41 (51–55)
Isthmus–anus distance	– (–)	57–71 (75–94)	62–70 (80–92)
Isthmus–A. distance	– (–)	68–101 (93–138)	91–106 (117–140)
V.–A. distance	47 (63)	33–63 (46–87)	55–66 (71–88)
Anus–A. distance	23 (30)	11–39 (14–53)	26–39 (35–52)
V. length	60 (80)	62–77 (83–104)	67–83 (86–112)
P. length	– (–)	53–60 (70–80)	54 (70)
Pre-1D. length	114 (153)	112–127 (148–169)	117–124 (154–160)
Height of 1D.	– (–)	91–106 (120–143)	92 (121)
Length of 1D. base	30 (41)	27–34 (37–45)	28–32 (37–41)
Interdorsal length	37 (50)	22–44 (29–59)	37–37 (48–50)
Length of gill slit	14 (19)	12–15 (16–20)	12–13 (16–17)
Length of posterior nostril	5 (7)	3–6 (5–8)	6–8 (8–11)
Barbel length	19 (25)	17–25 (22–33)	22–23 (29–31)
Counts			
1D. rays	II,10 or 11	II,10–13	II,10–12
P. rays	i22–i23	i17–i22	i19–i22
V. rays	13–14	14–17	13–15
Outer GR on first arch	9	5–10	11
Inner GR on first arch	9	8–11	8–10
Outer GR on second arch	8	7–11	9
Inner GR on second arch	8	8–10	9–10
Longitudinal scales	36	37–48	36–39
Scales below 1D. origin	11.5	11–16	10.5–12
Scales below 1D. midbase	10	7.5–11	7.5–9
Scales below 2D. origin	12	10–12.5	12–12.5
Scales above A. origin	–	ca. 36–46	–

...See the next page for the table footer

*Abbreviations: 1D.—first dorsal fin; 2D.—second dorsal fin; A.—anal fin; GR—gill rakers; HT—holotype; NT—non-type; P.—pectoral fin; PMX—premaxillary; V.—pelvic fin

the two species might be separable by the combination of the pelvic-fin length (68–81% HL in *N. condylura* vs. 97–100% in *N. propinqua*) and the number of spinule rows on the body scales (8–12 spinule rows on scales below the interdorsal space vs. 5–8 rows). This study tentatively treats *N. propinqua* as a valid species, but a future study based on more comprehensive materials may eventually show that the two taxa are conspecific.

Nezumia condylura is unlikely to be confused with other Japanese congeners in having higher counts of pelvic-fin rays (13–17) and transverse scale rows (10.5–16, 7.5–11, and 10–12.5 below the first dorsal-fin origin, first dorsal-fin midbase, and second dorsal-fin origin respectively). It further differs from all Japanese congeners in having a paler first dorsal fin, with a black apical spot (Fig. 163 vs. uniformly dark).

***Nezumia kamoharai* Okamura, 1970**

[Japanese name: Sokodara]

(Figs. 166–167; Appendix 3-10B)

Nezumia kamoharai Okamura, 1970a:91, pl. XX, text-fig. 40 (original description; holotype: BSKU 14202, from Sagami Bay; new Japanese name: “Soko-dara”); Okamura 1970b: table 1 (listed; Japan); Tominaga & Uyeno 1981:489 (listed; Japan); Okamura 1984b:94, pl. 344, fig. K (compiled); Okamura 1988:94, pl. 344, fig. K (compiled); Nakabo 1993:361 (in key; Japan); Nakabo 2000:425 (in key; Japan); Nakabo 2002:425 (in key; Japan); Senou *et al.* 2006:421 (listed; Sagami Sea); Nakabo & Kai 2013:501 (in key; Japan); Motomura 2020:39 (listed; Japan).

Diagnosis. A species of *Nezumia* with 13 pelvic-fin rays. Snout moderately long, conical, protruding distinctly beyond upper jaw, length 30% HL, ventral contour oblique in lateral view; underside of snout completely naked; ventral surfaces of head mostly scaled posterior to vertical through midorbit; mandibular rami naked, with narrow scaly patches on posterior portions; barbel length 12% HL. Second spinous ray of first dorsal fin not extremely prolonged (height of fin 99% HL). Body scales covered with short, greatly reclined, narrowly lanceolate spinules in tightly packed convergent rows; tip of last spinule in each row extending slightly beyond posterior scale margin; middle row of spinules distinctly larger than adjacent rows; scales below second dorsal-fin origin 7.5. Cephalic sensory pores present; those on mandibular and infraorbital canals small but conspicuous. No prominent dark band encircling trunk; first dorsal fin without black blotch apically.

Material examined. 1 specimen. **Holotype** of

Nezumia kamoharai: BSKU 14202 (1, 62.4 mm HL, 350+ mm TL), Sagami Bay, Kanagawa Pref., Japan, 1967.

Counts and measurements. Counts: first dorsal-fin rays II,9; pectoral-fin rays i19–i21; pelvic-fin rays 13; gill rakers on first arch (outer/inner) 8–9/10, on second arch 9/11; longitudinal scales 36; transverse scale rows below first dorsal-fin origin 10, below first dorsal-fin midbase 7.5, below second dorsal-fin origin 7.5, above anal-fin origin 25.5.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 30 (41); orbit diameter 28 (39); postorbital length 46 (64); postrostral length 73; orbit–preopercle distance 39 (53); suborbital width 15 (21); upper-jaw length 30 (41); length of rictus 24 (33); length of premaxillary tooth band 18 (24); preoral length 25 (34); distance between tip and lateral angle of snout 18 (25); snout width 27 (37); internasal width 21 (29); interorbital width 21 (29); body width over pectoral-fin bases 57 (78); body depth at first dorsal-fin origin 82 (112); body depth at anal-fin origin 69 (95); prepelvic length 116 (160); preanus length 135 (185); preanal length 153 (210); isthmus–pelvic distance 38 (52); isthmus–anus distance 57 (78); isthmus–anal distance 74 (101); pelvic–anal distance 42 (58); anus–anal distance 18 (25); pelvic-fin length 44 (60); pectoral-fin length 50 (68); predorsal length 115 (158); height of first dorsal fin 99 (135); length of first dorsal-fin base 26 (36); interdorsal length 35 (49); length of gill slit 14 (19); length of posterior nostril 4 (5); barbel length 12 (16).

Size. To at least 35 cm TL.

Distribution. So far known only from Sagami Bay (Appendix 3-10B). Depth unknown. Very rare.

Remarks. For a full description see the original description given by Okamura (1970a). *Nezumia kamoharai* was described from a single specimen collected from Sagami Bay (Fig. 166). The holotype is the only representative of this rare species and, despite intensive sampling efforts in southern Japan, no additional specimens have been collected since the original description.

Relationships and comparisons. *Nezumia kamoharai* is unlikely to be confused with any other Japanese congeners in having 13 pelvic-fin rays, a moderately long, pointed snout (30% HL), and the middle row of body-scale spinules distinctly enlarged compared with adjacent rows (Fig. 167). Elsewhere, this species appears to be most closely related to *N. longebarbata* (Roule & Angel, 1933) so far known from Madeira and the Gulf of Mexico. These two species are similar to each other in having 13 pelvic-fin rays and the underside of the head broadly naked from the snout tip to a vertical at the

FIGURE 166. Holotype of *Nezumia kamoharai*. BSKU 14202, 62.4 mm HL, 350+ mm TL, Sagami Bay, Pacific. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

FIGURE 167. Scanning electron micrographs showing body scale (from the dorsum below the interdorsal space) of *Nezumia kamoharai*. BSKU 14202, holotype, 62.4 mm HL. (A) View from above; (B) oblique view. [Photos: NSMT]

hind rim of the orbit. However, *N. kamoharai* is separable from *N. longebarbata* by having a smaller orbit (28% HL vs. 31–32%), shorter barbel (12% HL vs. 17–23%) and pectoral fin (50% HL vs. 53–57%), and slightly larger body scales (transverse scale rows below the second dorsal-fin origin 7.5 vs. 8.8). [Data for *N. longebarbata* are from Marshall & Iwamoto (1973).]

In his visit to BSKU in 2009, T. Iwamoto (CAS) informed the present author that *N. kamoharai* also shares many features with *N. obliquata* (Gilbert, 1905), a rare species confined to the Hawaiian-Emperor Seamount Chain. The latter species was originally described from

a small juvenile collected from Hawaii [USNM 51514, holotype (ca. 24 mm HL, 127+ mm TL, fide Sazonov 1994:103)], and is poorly represented in museum collections. Sazonov (1994) recorded an additional specimen of this species from Ojin Guyot in the Emperor Seamounts (ZMMGU P.18245, 60.6 mm HL, 398+ mm TL), although he considered his identification tentative due to a number of differences related to the difference in body size between the holotype and the second specimen. An examination of Sazonov's (1994) specimen revealed its close similarity with *N. kamoharai*, but the holotype of *N. kamoharai* differs from that specimen in having

shorter barbel (12% HL vs. 17%), pelvic fin (44% HL vs. 65%), and pectoral fin (50% HL vs. 59%). Spinulation of the body scales is also different between the two specimens. In the holotype of *N. kamoharai*, the middle row of spinules is distinctly larger than adjacent rows (Fig. 167), whereas in the “*N. obliquata*” specimen from the Emperor Seamounts, the middle row is not especially enlarged (Sazonov 1994: fig. 3). These two specimens are almost equal in size, and their differences appear to warrant specific separation. The status of Sazonov’s (1994) specimen is still uncertain, but it does not seem to represent either *N. obliquata* nor *N. kamoharai*. Based on the original description given by Gilbert (1905), the holotype of *N. obliquata* differs from *N. kamoharai* in having fewer pelvic-fin rays (12 vs. 13) and smaller body scales (9 below the second dorsal-fin origin vs. 7.5). To resolve the taxonomic status of these species, a further study should be done when more specimens become available.

***Nezumia proxima* (Smith & Radcliffe in Radcliffe, 1912)**

[Japanese name: Higo-sokodara]

(Figs. 168–170; Table 12; Appendix 3-10C)

Macrurus altipinnis (not Günther 1877): Günther 1887:138 (brief description; in part, BMNH 1887.12.7.92; Enshu-nada).

Macrourus nasutus (not Günther 1877): Jordan & Gilbert in Jordan & Starks 1904:614 (in part, 1 spec. from off Izu; see Gilbert & Hubbs 1916:201).

[?] *Macrourus rudis* (not Günther 1887): Franz 1910:16 (1 spec. from Aburatsubo); Jordan *et al.* 1913:417 (listed; Japan; new Japanese name: “Aka-sokodara”).

Macrourus proximus Smith & Radcliffe in Radcliffe, 1912:119, pl. 26, fig. 2 [original description; holotype: USNM 72936, from Sogod Bay, Philippines, *Albatross* sta. 5202, 10°12'00"N, 125°04'10"E, in 502 ftn (918 m)].

Lionurus proximus: Gilbert & Hubbs 1916:201 [new combination; brief description; in part, 10 spec. from East China Sea, Hyuga-nada, off Cape Shionomisaki, and Enshu-nada; 1 spec. from *Albatross* sta. 4918 (CAS-SU 22941) is *Nezumia spinosa*]; Okada & Matsubara 1938:453 (in key; Japan); Matsubara 1955:1316 (in key; Japan); Matsubara 1965:509 (compiled; Japan; new Japanese name: “Higo-sokodara”).

Lionurus abei Matsubara, 1943:146, figs. 7–8 (original description; holotype: FAKU 4909, from Kumano-nada; new Japanese name: “Kasumi-hige”); Matsubara 1955:1317, pl. 134, fig. 458 (in key; Japan); Okada *et al.* 1959:83 (listed; Kumano-nada); Matsubara 1965:509 (compiled; Japan); Kuroda 1966:29 (brief description; spec. from Suruga Bay).

Nezumia proximus: Okamura 1970a:94, pl. XXI, text-fig. 41 (description; biological notes; 17 spec. from Pacific off southern Japan from Choshi to Kumano-nada and East China Sea); Kuroda 1971:56 [listed from Suruga Bay after Okamura (1970a)].

Nezumia proxima: Okamura 1970b: table 1 (listed; Japan); Kataoka & Tomida 1981:78 (listed; Mie Pref.); Tominaga & Uyeno 1981:489 (listed; Japan); Okamura 1982:163, 351, fig. 162 (brief description; 6 spec. from Tosa Bay; photo based on BSKU 29500); Sawada 1983:107, 192, fig. 58 [brief description; 4 spec. from Pacific off Tohoku; photo based on HUMZ spec. (catalog no. unknown)]; Ohta 1983: table A (listed; Suruga Bay); Okamura 1984a:217, 362, fig. 152 (brief description; 8 spec. from Okinawa Trough; photo based on BSKU 29792); Okamura 1984b:94, pl. 81, fig. I (compiled); Okamura 1988:94, pl. 81, fig. I (compiled); Iwamoto 1990:281, fig. 645 (synopsis); Nakabo 1993:360 (in key; Japan); Hori 1996:27 (listed; Ibaraki Pref.); Shinohara *et al.* 1996:170 (2 spec. listed from Pacific off Tohoku); Okamura 1997:128, fig. 17 (compiled); Shinohara & Matsuura 1997:292 (listed; Suruga Bay); Funabashi 1998:85 (listed; Ibaraki Pref.); Nakabo 2000:424 (in key; Japan); Shinohara *et al.* 2001:306 (3 spec. listed from Tosa Bay); Nakabo 2002:424 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Shinohara *et al.* 2005:418 (listed; Ryukyu Islands); Suetsugu & Ohta 2005: table 3 (listed; Enshu-nada); Senou *et al.* 2006:421 (listed; Sagami Sea); Shao *et al.* 2008b: table 2 (2 spec. listed from eastern Taiwan and South China Sea; first Taiwanese record); Kitagawa *et al.* 2008:40, unnumbered fig. (brief description; spec. from Pacific off Tohoku); Shinohara *et al.* 2009:709 (1 spec. listed from Pacific off Tohoku); Nakabo & Kai 2013:500 (in key; Japan); Iwamoto *et al.* 2015:93, fig. 19 (brief description; 3 spec. from northeastern and southwestern Taiwan, and South China Sea); Amaoka *et al.* 2020:159, fig. 213 (erroneously listed from Hokkaido); Motomura 2020:39 (listed; Japan).

[?] *Lionurus spinosus* (not Gilbert & Hubbs 1916): Maruyama 1971:33 (listed with a question mark; Iwate Pref.).

Diagnosis. A species of *Nezumia* with 8–10 pelvic-fin rays. Snout moderately long, conical, protruding well beyond upper jaw, length 29–33% HL, ventral contour oblique in lateral view; underside of snout broadly naked; ventral surfaces of head scaled posterior to vertical through midorbit; mandibular rami mostly scaled. Second spinous ray of first dorsal fin not extremely prolonged (height of fin 92–118% HL). Body scales covered with short, greatly reclined, lanceolate spinules in tightly packed convergent rows or densely scattered over exposed portion; tip of last spinule in each row extending well beyond posterior scale margin; middle row of spinules not enlarged; scales below second dorsal-fin origin 6–8.5. Anterior dermal window of light organ (ADW) situated between inner pelvic-fin bases. Cephalic sensory pores present, but barely discernible. Broad dark band encircling trunk (often faint or absent in preserved specimens); first dorsal fin uniformly dark.

Material examined. 34 specimens. **Japan:** BSKU 99112 (1, 50.4 mm HL, 314+ mm TL), BSKU 99117 (1, 59.5 mm HL, 357+ mm TL), off Pacific coast of Tohoku, FRV *Wakataka-maru*, sta. 32 (F-800), coll. H. Endo,

FIGURE 168. Fresh specimens of *Nezumia proxima*. (A) BSKU 45547, 43.8 mm HL, 264+ mm TL, east of Katsuura, Chiba Pref., Pacific, 507–578 m depth; (B) BSKU 44837, 54.2 mm HL, 319+ mm TL, Tosa Bay, Kochi Pref., Pacific, 700–720 m depth; (C) KPM-NI 28856, 44.8 mm HL, 236+ mm TL, west-southwest of Shirahama, Wakayama Pref., Pacific, 532 m depth. Lateral views. [Photos: (A–B) BSKU; (C) H. Senou]

date unknown; BSKU 26667 (1, 61.7 mm HL, 336+ mm TL), west of Yaku-shima Island, Okinawa Trough, 30.0300°N, 128.3650°E, 900 m, F/V *Yuryo-maru*, No. 8, tr. 16-1, bottom trawl, coll. Y. Kinoshita and S. Hagino, 3 Feb. 1978; BSKU 86874 (1, 31.4 mm HL, 189+ mm TL), off Nobeoka, Hyuga-nada, 32.5163°N, 132.2175°E, 997–1030 m, R/V *Tansei-maru*, cr. KT-99-18, sta. BT-2, beam trawl, coll. H. Endo and S. Nagatomo, 16 Dec. 1999; BSKU 44837 (1, 54.2 mm HL, 319+ mm TL), Tosa Bay, 32.9886°N, 133.5978°E, 700–720 m, FRV *Kotaka-maru*, otter trawl, 21 Jun. 1988; BSKU 98937 (1, 33.3 mm HL, 197+ mm TL), BSKU 98938 (1, 47.6 mm HL, 218+ mm TL), BSKU 98939 (1, 28.4 mm HL, 147+ mm TL), Tosa Bay, 33.0452°N, 133.6448°E, 733–740 m, FRV *Kotaka-maru*, otter trawl, 25 Jan. 1989; BSKU 101591 (1, 42.1 mm HL, 254+ mm TL), BSKU 101592 (1, 49.2 mm HL, 303+ mm TL), BSKU 101593 (1, 48.0 mm HL, 280+

mm TL), BSKU 101594 (1, 27.9 mm HL, 176+ mm TL), BSKU 101595 (1, 33.3 mm HL, 203+ mm TL), BSKU 101596 (1, 39.9 mm HL, 229+ mm TL), BSKU 101597 (1, 40.9 mm HL, 253+ mm TL), BSKU 101598 (1, 35.2 mm HL, 196+ mm TL), BSKU 101599 (1, 49.9 mm HL, 302+ mm TL), Tosa Bay, 700 m, FRV *Kotaka-maru*, otter trawl, 22 May 1989; BSKU 47166 (1, 37.6 mm HL, 233+ mm TL), off Tosa Bay, 32.8953°N, 133.6777°E, 996–1010 m, R/V *Tansei-maru*, cr. KT-89-06, sta. T1, beam trawl, 8 Nov. 1989; BSKU 23147 (1, 45.1 mm HL, 274+ mm TL), Tosa Bay, 33.0350°N, 133.7500°E, 945 m, FRV *Kaiyo-maru*, cr. SK-75-2, sta. 18, tr. 3, bottom trawl, 26 Jan. 1975; BSKU 23076 (1, 46.7 mm HL, 225+ mm TL), BSKU 23181 (1, 48.1 mm HL, 291 mm TL), off Kochi, 32.9800°N, 133.8550°E, 1043 m, FRV *Kaiyo-maru*, cr. SK-75-2, sta. 19, tr. 2, bottom trawl, 26 Jan. 1975; *KPM-NI 28856 (1, 44.8 mm HL, 236+ mm TL), ca. 24 km

FIGURE 169. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of *Nezumia proxima*. (A–B) BSKU 19764, typical specimen, 57.9 mm HL; (C–D) BSKU 19891, uncertain variant, 50.8 mm HL. (A, C) Views from above; (B, D) oblique views. [Photos: NSMT]

west-southwest of Shirahama, Wakayama, 33.5795°N, 135.0995°E, 532 m, R/V *Tansei-maru*, cr. KT-10-16, sta. YK-2, box corer, coll. T. Sato, 15 Aug. 2010; BMNH 1887.12.7.92 (1, 24.0 mm HL, 149+ mm TL), off Cape Omaezaki, Enshu-nada, 34.1167°N, 138.0000°E, 565 ftn (1034 m), *Challenger* sta. 235, trawl, 4 Jun. 1875; BSKU 19762 (1, 41.6 mm HL, 211+ mm TL), BSKU 19764 (1, 57.9 mm HL), Sagami Bay, 35.1167°N, 139.4500°E, 730–790 m, FRV *Soyo-maru*, sta. T7, 23 Nov. 1966; BSKU 20250 (1, 39.2 mm HL, 254+ mm TL), west of Misaki, Sagami Bay, 35.1283°N, 139.4983°E, 450 m, FRV *Soyo-maru*, sta. 77, 30 Jun. 1972; BSKU 14184 (1, 35.7 mm HL, 221+ mm TL), BSKU 14186 (1, 29.2 mm HL, 190+ mm TL), off Hiratsuka, Sagami Bay, 35.2522°N, 139.5178°E, 510–600 m, R/V *Tansei-maru*, coll. O. Okamura, 12 Apr. 1969; BSKU 45547 (1, 43.8 mm HL, 264+ mm TL), east of Katsuura, 35.1400°N, 140.8125°E, 507–578 m, R/V *Tansei-maru*, cr. KT-89-03, sta. TB5, 13 Mar. 1989; BSKU 101193 (1, 60.3 mm HL, 374+ mm TL), off Hitachi, 900 m, FRV *Wakataka-maru*, sta. H, coll. H. Endo, 10 Oct.

1997; BSKU 97917 (1, 56.7 mm HL, 345+ mm TL), off Oshika Peninsula, 38.5318°N, 142.3370°E, 772–783 m, FRV *Wakataka-maru*, sta. 32, 28 Oct. 1995; BSKU 97906 (1, 58.7 mm HL, 375+ mm TL), BSKU 97910 (1, 51.4 mm HL, 341+ mm TL), off Otsuchi, 39.1444°N, 142.3748°E, 847–935 m, R/V *Tansei-maru*, cr. KT-89-03, sta. SR89, 10 Mar. 1989. **Uncertain variant:** BSKU 19891 (50.8 mm HL, 274+ mm TL), east-southeast of Miyake-jima Island, Shichito-Iojima Ridge, Japan, 33.9183°N, 140.0083°E, 860–870 m, FRV *Soyo-maru*, beam trawl, 1 Dec. 1968.

Description of Japanese specimens. General features are shown in Fig. 168. Counts and measurements are given in Table 12. Body long, slender, gradually tapering from first dorsal-fin origin to end of tail. Trunk short, laterally compressed, width over pectoral-fin bases 1.3–1.7 in depth below first dorsal-fin origin. Head short, HL about 5.2–6.6 in TL, its dorsal profile almost straight from tip of snout to nape. Snout moderately long, protruding well beyond upper jaw, conical in lateral view, bluntly pointed in dorsal view; snout length 0.9–1.1

FIGURE 170. Uncertain variant of *Nezumia proxima*. BSKU 19891, 50.8 mm HL, 274+ mm TL, east-southeast of Miyake-jima Island, Shichito-Iojima Ridge, Pacific, 860–870 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of head. Preserved condition. [Photos: N. Nakayama]

times as long as orbit diameter; ventral contour oblique, forming angle of about 45° to horizontal axis of head and body; tip of snout situated on horizontal through midorbit. Orbit moderately large, circular, greatest diameter 1.1–1.9 in postorbital length. Interorbital space almost flat, width 1.3–1.7 in orbit diameter. Mouth small, inferior, upper-jaw length 0.9–1.2 in orbit diameter; posterior margin of maxilla extending beyond vertical through midorbit; lateral corner of mouth slightly restricted by lip folds; lips thick, fleshy, papillose near tooth bands. Suborbital region narrow, divided into upper and lower halves by infraorbital ridge. Preopercle large, broadly rounded ventrally, posterior margin slightly inclined from vertical; preopercular ridge low, poorly marked, forming slight backward extension at angle. Interopercle narrowly exposed beyond preopercle. Gill membranes broadly connected across, and attached mesially to isthmus, with broad posterior free fold. Anteroventral end of gill opening not reaching below posterior margin of lower jaw. Outermost gill slit moderately restricted by skin folds, length 2.0–2.8 in orbit diameter. Gill rakers small, tubercular, armed with short, fine spines distally; those on outer side of first arch and inner side of fourth arch much smaller than others. Gill filaments moderately long. Barbel well developed, length 1.4–2.1 in orbit diameter.

Anus closer to pelvic-fin bases than to anal-fin origin. Periproct broad, oval in shape. Anterior dermal window of light organ small, circular, situated between inner bases of pelvic fins.

Teeth short, slender, sharp, in broad tapered bands in both jaws. Premaxillary teeth arranged in about 5 tooth

rows near symphysis, with outermost series distinctly enlarged. Mandibular band similar to that of premaxillary, with about 6 tooth rows at widest point near symphysis, but none significantly enlarged. Posterior ends of tooth bands falling short of lateral corner of mouth in both jaws. All teeth deeply embedded in thick layer of gum papillae.

Body scales small, thin, moderately deciduous, covered with short, greatly reclined, lanceolate spinules in tightly packed convergent rows or densely scattered over exposed portion (Fig. 169A–B); tip of last spinule in each row extending well beyond posterior scale margin; middle row of spinules not enlarged; buttresses of spinules absent; reticulate structure narrowly developed on posterior parts of unexposed portion. Body fully scaled except for fins, periproct, and dermal window of light organ; pelvic fins mostly naked, but basal portions covered with tiny scales.

Suborbital shelf formed of 2 rows of stout, coarsely modified scales. Snout tipped with 2 enlarged, thickened, closely adjoined tubercles; scales on lateral angles of snout modified into scute-like scales. Scales on head generally similar to those on body, except that spinules more erect and arranged in subparallel rows. Dorsal and lateral surfaces of head almost fully scaled; membrane around anterior nostril and paired Y-shaped areas dorsally posterior to leading edges of snout narrowly naked. Underside of snout almost completely naked; lower half of suborbital region mostly scaled, but narrowly naked above upper lip; anterior 1/3–1/2 of mandibular rami naked; gular and branchiostegal membranes lacking scales.

TABLE 12. Morphometric measurements and meristic counts* for *Nezumia proxima*.

Type status	NT	
Identification	Typical specimens	Uncertain variant
Number of specimens	<i>n</i> = 32	<i>n</i> = 1
Measurements (mm)		
TL	147+–375+	274+
HL	24.0–61.7	50.8
% of HL (% of PRL)		
Snout length	29–33 (39–46)	31 (44)
Orbit diameter	28–35 (40–48)	29 (40)
Postorbital length	37–45 (52–61)	44 (62)
Postrostral length	70–75 (–)	71 (–)
Orbit–preopercle distance	32–40 (44–54)	35 (50)
Suborbital width	13–16 (18–22)	14 (20)
Upper-jaw length	28–34 (39–47)	31 (44)
Length of rictus	23–28 (32–38)	27 (38)
Length of PMX tooth band	16–20 (22–28)	20 (28)
Preoral length	22–30 (30–42)	25 (35)
Dist. tip and lat. angle of snout	16–20 (21–28)	16 (23)
Snout width	25–31 (34–42)	26 (37)
Internasal width	19–23 (27–32)	24 (34)
Interorbital width	19–24 (27–32)	25 (35)
Body width over P. bases	43–70 (61–94)	59 (83)
Body depth at 1D. origin	70–92 (97–124)	79 (111)
Body depth at A. origin	58–92 (80–124)	70 (98)
Pre-V. length	103–124 (144–174)	114 (161)
Pre-anus length	121–145 (169–200)	134 (188)
Pre-A. length	146–176 (200–241)	157 (220)
Isthmus–V. distance	32–45 (44–62)	33 (46)
Isthmus–anus distance	46–69 (64–95)	51 (72)
Isthmus–A. distance	72–107 (99–147)	76 (107)
V.–A. distance	39–64 (55–89)	45 (63)
Anus–A. distance	21–38 (30–53)	26 (36)
V. length	44–71 (59–99)	69 (97)
P. length	49–61 (69–84)	56 (79)
Pre-1D. length	108–125 (147–173)	114 (160)
Height of 1D.	92–118 (126–162)	97 (136)
Length of 1D. base	23–33 (32–46)	24 (33)
Interdorsal length	31–66 (44–92)	44 (62)
Length of gill slit	11–16 (15–21)	15 (21)
Length of posterior nostril	2–6 (3–8)	9 (12)
Barbel length	15–21 (20–29)	15 (21)
Counts		
1D. rays	II,8–10	II,9
P. rays	i17–i21	i19
V. rays	8–10	9
Outer GR on first arch	5–9	7
Inner GR on first arch	8–10	10
Outer GR on second arch	8–9	9
Inner GR on second arch	8–10	9
Longitudinal scales	32–40	39
Scales below 1D. origin	7.5–10	9
Scales below 1D. midbase	5–8	5.5
Scales below 2D. origin	6–8	8.5
Scales above A. origin	19–26	21.5

... See the next page for the table footer

*Abbreviations: 1D.—first dorsal fin; 2D.—second dorsal fin; A.—anal fin; GR—gill rakers; NT—non-type; P.—pectoral fin; PMX—premaxillary; V.—pelvic fin

Cephalic sensory pores present, arranged along sensory canals; those on underside of head very slightly enlarged, more or less tubular. Free neuromasts small, tubercular, densely scattered on underside of snout and mandibular rami; similar but much smaller neuromasts serially arrayed along leading edges of snout, supranarial ridges, suborbital shelf, and supraorbital canals. Grooved lateral line incomplete anteriorly, occurring as short interrupted segments, but complete posterior to vertical through midbase of first dorsal fin.

Origin of first dorsal fin slightly posterior to vertical through pectoral- and pelvic-fin bases, the latter two on about same vertical; first dorsal fin high, its height 3.2–4.7 times as long as its base length; second spinous ray long, but not especially elongate, armed along its leading edge with short, reclined, conical denticles; its tip extending to bases of 4th–20th second dorsal-fin rays when laid back. Second dorsal fin well separated from first dorsal, originating above 5th–13th anal-fin rays; interdorsal space 1.1–2.4 times as long as first dorsal-fin base length. Outermost pelvic-fin ray slightly prolonged, hair-line distally, its tip usually extending beyond anal-fin origin when laid back.

Color when fresh (Fig. 168). Uniformly dark brown; broad dark band encircling trunk (especially prominent when body scales mostly missing; Fig. 168C).

Color in alcohol. Similar to that in fresh specimens; head and body dark brown overall, with broad dark band encircling trunk (band often indistinct in old specimens); abdomen and gill cover prominently darker in smaller specimens; lips dark; oral cavity blackish except for pale gums; gill cavity mostly black, but paler on ceratohyal, epihyal, and anterior to lower end of cleithrum; gill rakers and arches dark dusky, filaments pale; chin barbel varying from pale to dark; gular membrane blackish, but abruptly pallid laterally; branchiostegal membranes black, especially along distal margins; fins uniformly dark.

Size. To about 40 cm TL (NSMT-P 94347, 398+ mm TL, off Hitachi, Ibaraki Pref., Japan).

Distribution. Distributed in the western Pacific from the Philippines to Japan (Radcliffe 1912; Gilbert & Hubbs 1920; Shao *et al.* 2008b; this study). In Japanese waters, known from off the Pacific coasts northward to Kuji (40.19°N; Iwate Pref.) and the Okinawa Trough at depths of 361–1348 m (Appendix 3-10C). Common, especially in the Pacific off Honshu north of Choshi (35.74°N; Chiba Pref.).

Remarks. *Nezumia proxima* was originally described from the Philippines, and first recorded from Japan by Gilbert & Hubbs (1916). Matsubara (1943) described *N. abei* (as a species of *Lionurus* Günther, 1887) from a single specimen collected from the Kumano-nada, Japan,

but this species has been regarded as a junior synonym of *N. proxima* (e.g., Okamura 1970a; Iwamoto 1990; Iwamoto *et al.* 2015). Unfortunately, the holotype of *N. abei* is presumed to be lost (see the Comments on holotype of *Coelorinchus hige*).

Among the FAKU collection was a single specimen of *N. proxima* collected from off Minamidaito-jima Island (FAKU 13366, 1 spec., 59.4 mm HL, 366+ mm TL), but the register indicates its locality with a question mark. This record was not included in the distribution of this species (Appendix 3-10C). Amaoka *et al.* (2020) recently listed the species from Hokkaido, but it is obviously based on Sawada's (1983) record from the Pacific off Tohoku.

Relationships and comparisons. *Nezumia proxima* is putatively most similar to *N. namatahi* McCann & McKnight, 1980, a species known only from New Zealand and Australia, and they share the combination of 8–10 pelvic-fin rays (9–10 in *N. namatahi*), a broad naked area on the underside of the snout, a long barbel (15–21% HL in *N. proxima*, and 13–30% in *N. namatahi*), body scales covered with lanceolate to shield-shaped spinules in parallel to convergent rows, and a broad, prominent, dark band completely encircling the trunk. However, *N. proxima* readily differs from that species in that the anterior dermal window of the light organ (ADW) is situated between the inner pelvic-fin bases (vs. ADW usually distinctly posterior to a line connecting inner pelvic-fin bases). It further differs from *N. namatahi* in its tiny pores along the infraorbital and mandibular canals (vs. large and prominent). [Data for *N. namatahi* are based on Iwamoto & Williams (1999:194–195), Iwamoto & Graham (2001:489), and this study.]

Uncertain variant. A specimen collected from the Shichito-Iojima Ridge at a depth of 860–870 m (Fig. 170; Table 12) was initially regarded as a different species from *N. proxima* by having a wider interorbital space (25% HL vs. 19–24%). It further differs from the typical *N. proxima* specimens examined in having a wider internasal space (24% HL vs. 19–23%), a larger posterior nostril (9% HL vs. 2–6%), a relatively shorter barbel (15% HL vs. 15–21%), and slightly more scales below the second dorsal-fin origin (8.5 vs. 6–8). However, no significant differences were evident in squamation, spinulation of body scales (Fig. 169), and the number and arrangement of cephalic sensory pores. Although the above differences may warrant their specific separation, the paucity of materials from the Shichito-Iojima Ridge did not allow a definitive conclusion. Therefore, the specimen was tentatively identified as an uncertain variant of *N. proxima*.

***Nezumia rara* sp. nov.**

[New Japanese name: Kakure-hige]

(Figs. 171, 172A–B; Table 13; Appendix 3-10D)

[?] *Macrourus burragei* (not Gilbert 1905): Kamohara 1937:189, fig. 2 (first record from Japan; brief description; 1 spec. from Kochi; new Japanese name: “Nan’yo-dara”); Kamohara 1938:72, fig. 41 [after Kamohara (1937)].

[?] *Lionurus burragei* (not Gilbert 1905): Okada & Matsubara 1938:453 (in key; Japan); Kamohara 1950:278 (listed; Kochi Pref.); Kamohara 1952:98, fig. 96 [brief description after Kamohara (1937); Kochi Pref.]; Matsubara 1955:1316 (in key; Japan); Kamohara 1958:74 (questionable; listed; Kochi Pref.); Kamohara 1964:96 (questionable; listed; Kochi Pref.).

[?] *Nezumia burragei* (not Gilbert 1905): Okamura 1970a:100 [description after Gilbert (1905) and Kamohara (1937, 1938)]; Okamura 1970b: table 1 (listed; Japan); Tominaga & Uyeno 1981:489 (listed; Japan); Okamura 1984b:94, pl. 345, fig. C (compiled); Okamura 1988:94, pl. 345, fig. C (compiled); Nakabo 1993:360 (in key; Japan); Nakabo 2000:424 (in key; Japan); Shinohara *et al.* 2001:306 (listed; Tosa Bay); Nakabo 2002:424 (in key; Japan); Nakabo & Kai 2013:500 (in key; Japan); Motomura 2020:39 (listed; Japan).

Nezumia sp.: Sawada 1983:107, 193, fig. 59 (brief description; 1 spec. from Pacific off Tohoku; HUMZ 78437).

Diagnosis. A species of *Nezumia* with 10–11 pelvic-fin rays. Snout short, protruding scarcely beyond upper jaw, length 26–28% HL, ventral contour almost vertical in lateral view; underside of snout broadly naked; ventral surfaces of head scaled posterior to lateral angles of snout; mandibular rami mostly scaled; orbit–preopercle distance 39–41% HL; interorbital width 20–22% HL; pelvic-fin length 46–54% HL; length of outer gill slit 13–16% HL. Second spinous ray of first dorsal fin not extremely prolonged (height of fin 80–98% HL). Body scales covered with short, greatly reclined, narrowly triangular spinules in tightly packed, convergent rows or densely scattered over exposed portion; tip of last spinule in each row extending slightly beyond posterior scale margin; middle row of spinules not enlarged; scales below second dorsal-fin origin 9–10. Cephalic sensory pores present along all canals, but barely discernible. No prominent dark band encircling trunk; first dorsal fin uniformly dark.

Material examined. 3 specimens. **Holotype:** NSMT-P 100179 (73.0 mm HL, 419 mm TL), Suruga Bay, Shizuoka Pref., Japan, R/V *Tansei-maru*, cr. KT-78-19, sta. TW5, 2-m beam trawl, Dec. 1978. **Paratypes:** NSMT-P 97666 (1, 71.6 mm HL, 415 mm TL), southeast of Oshima Island, Shichito-Iojima Ridge, 621–710 m, R/V *Hakuho-maru*, cr. KH-89-test-III, 4-m beam trawl, 5 Sept. 1989; HUMZ 78437 (1, 78.9 mm HL, 447+ mm TL), off Ibaraki, 36.0583°N, 141.3333°E, 1100 m, bottom trawl, coll. K. Sakamoto, 9 Jan. 1978.

Description. General features are shown in Fig. 171. Counts and measurements are given in Table 13.

The following description based on the holotype, unless otherwise stated; variations found in paratypes are indicated in parentheses. Body moderately deep, deepest at first dorsal-fin origin, gradually tapering to long, slender tail. Trunk short, compressed, width over pectoral-fin bases 1.5 in depth below first dorsal-fin origin. Head moderately large, HL about 5.7 (5.7–5.8) in TL; dorsal profile almost straight from tip of snout to first dorsal-fin origin, without pronounced hump over nape nor distinct depression above orbits. Snout short, protruding scarcely beyond upper jaw, ventral contour almost vertical, forming angle of about 70° to horizontal axis of head and body; snout length 0.9 (0.9–1.1) times as long as orbit diameter; tip of snout situated above horizontal through midorbit, bluntly pointed in lateral and dorsal views. Orbit moderately large, circular in outline, greatest diameter 1.6 (1.7–1.9) in postorbital length. Interorbital space almost flat (slightly concave), width 1.4 (1.2–1.4) in orbit diameter. Mouth moderately small, subinferior, upper-jaw length 0.8 in orbit diameter; posterior margin of maxilla extending to vertical through hind 1.3 of orbit (midorbit); lateral corner of mouth slightly restricted by lip folds; lips thick, fleshy, papillose near tooth bands. Suborbital region narrow, divided into upper and lower halves by infraorbital ridge. Preopercle large, hind margin slightly inclined forward, forming broad lobe posteroventrally; preopercular ridge low, with slight backward extension at posteroventral corner. Interopercle narrowly exposed beyond preopercle. Gill membranes broadly connected across, and attached mesially to isthmus, with broad posterior free fold. Anteroventral end of gill opening not reaching below posterior margin of lower jaw. Outermost gill slit moderately restricted by skin folds, length 1.8 (2.0) in orbit diameter. Gill rakers developed into small, ridge-like tubercles, their tips armed with short, fine spines; rakers on outer side of first arch and inner side of fourth arch much smaller than others. Gill filaments moderately long. Barbel long, slender, filamentous at tip, length 1.7 (1.3) in orbit diameter.

Anus closer to pelvic-fin bases than to anal-fin origin. Periproct broad, teardrop-shaped. Anterior dermal window of light organ small, circular, situated between inner bases of pelvic fins.

Teeth short, slender, conical, in tapered bands in both jaws. Premaxillary teeth arranged in about 5–6 tooth rows near symphysis, with outermost series distinctly enlarged. Mandibular band with about 5 tooth rows at widest point near symphysis, and no teeth significantly enlarged. Posterior ends of tooth bands falling short of lateral corner of mouth in both jaws. All teeth deeply embedded in thick layer of gum papillae.

Body scales small, thin, not especially deciduous, covered with short, greatly reclined, narrowly triangular spinules in tightly packed convergent rows or densely scattered over exposed portion (Fig. 172A–B); tip of last spinule in each row extending slightly beyond posterior scale margin; middle row of spinules not enlarged;

buttresses of spinules absent; reticulate structure narrowly developed on posterior parts of unexposed portion. Body fully scaled except for fins, periproct, and dermal window of light organ; proximal portion of pelvic fins heavily covered with tiny scales.

Suborbital shelf armed with stout, coarsely modified scales. Snout tipped with 2 enlarged, thickened, closely adjoined tubercles; scales on lateral angles of snout modified into scute-like scales. Other scales on head covered with short, erect, triangular spinules in parallel to divergent rows. Dorsal and lateral surfaces of head almost fully scaled; membrane around anterior nostril and paired Y-shaped areas dorsally posterior to leading edges of snout narrowly naked. Underside of snout almost completely naked; lower half of suborbital region mostly scaled, but narrowly naked above upper lip; anterior 1/3 of mandibular rami naked; gular and branchiostegal membranes lacking scales.

Cephalic sensory pores present along sensory canals; those on underside of head very slightly enlarged, more or less tubular. Free neuromasts densely scattered over naked areas on underside of snout and mandibular rami, and serially arranged along leading edges of snout, supranarial ridges, suborbital shelf, and supraorbital canals; those on underside of snout largest. Naked area on underside of snout densely covered with short, hair-like papillae. Grooved lateral line incomplete anteriorly,

occurring as short interrupted segments, but complete posterior to vertical through midbase of first dorsal fin.

Origin of first dorsal fin slightly posterior to vertical through pectoral-fin base; first dorsal fin high, its height 3.4 (2.7–3.0) times as long as its base length; second spinous ray long, but not especially elongate, armed along its leading edge with short, reclined, conical, narrowly spaced denticles; its tip extending to base of 11th (8th) second dorsal-fin ray when laid back. Second dorsal fin well separated from first dorsal, originating above 10th (9th–10th) anal-fin ray; interdorsal space 1.4 times as long as first dorsal-fin base length. Pelvic fin inserted below pectoral-fin base; outermost ray moderately elongate with filamentous tip, reaching base of 4th (3rd) anal-fin ray when laid back.

Color when fresh. Unknown.

Color in alcohol (Fig. 171). Head and body uniformly dark overall, with much darker abdomen; orbit barely marked with narrow dark rim; lips broadly delineated in dark brown; septum between nostrils much darker than surroundings; oral cavity blackish including gum papillae; gill cavity mostly black, but paler on ceratohyal, epihyal, and anterior to lower end of cleithrum; gill rakers and arches pale to dusky, filaments orange brown; chin barbel mostly pale with dusky base; gular membrane blackish; fins uniformly dark, but outermost pelvic-fin ray paler.

FIGURE 171. Holotype of *Nezumia rara* sp. nov. NSMT-P 100179, 73.0 mm HL, 419 mm TL, Suruga Bay, Shizuoka Pref., Pacific. (A) Lateral, (B) dorsal, and (C) ventral views. Preserved condition. [Photos: N. Nakayama]

FIGURE 172. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of two species of *Nezumia*. (A–B) *N. rara* sp. nov., NSMT-P 97666, paratype, 71.6 mm HL; (C–D) *N. tomiyamai*, HUMZ 71963, 68.7 mm HL. (A, C) Views from above; (B, D) oblique views. [Photos: N. Nakayama]

Size. To about 45 cm TL (HUMZ 78437, 447+ mm TL, off Ibaraki Pref., Japan).

Distribution. So far known only from Japan: Suruga Bay, off the Izu Islands, and Kashima-nada, at depths of 621–1100 (Appendix 3-10D). Very rare.

Etymology. The specific epithet is derived from the Latin adjective *rarus*, meaning “sparsely occurring”, in reference to the species’ rare occurrence. The Japanese name, Kakure-hige, also alludes to the rarity of the species (kakure, meaning “hiding”, and hige, meaning “grenadiers”).

Remarks. Sawada (1983) reported “*Nezumia* sp.” based on a single specimen collected from off Ibaraki, Japan (HUMZ 78437, 78.9 mm HL, 477+ mm TL). His specimen is herein designated as a paratype of the new species. See also the Comments on the Japanese record of *Nezumia burragei* (given below).

Relationships and comparisons. *Nezumia rara* sp. nov. is very similar to *N. tomiyamai* (Okamura, 1963), with which it shares most of the diagnostic features

separating the latter species from its congeners. No differences were found in meristic characters, except for transvers scale rows below the second dorsal-fin origin (9–10 in *N. rara* vs. 8.5 in *N. tomiyamai*; Table 13). However, regarding morphometric characters, the two species are different in many aspects, including the postorbital length (46–49% HL in *N. rara* vs. 49–51% in *N. tomiyamai*), postrostral length (74–75% vs. 77%), orbit–preopercle distance (39–41% vs. 42–43%), upper-jaw length (33–36% vs. 36–37%), interorbital width (20–22% vs. 23–25%), pelvic-fin length (46–54% vs. 69%), pectoral-fin length (61–63% vs. 58%), and length of gill slit (13–16% vs. 17–18%) (Table 13). These differences are subtle, but in combination, they appear to warrant their specific separation. In fact, the general appearance of the two species is quite different from each other (Fig. 171 vs. Figs. 178–179), which results from the differences in the above-mentioned characters.

TABLE 13. Morphometric measurements and meristic counts* for *Nezumia rara* **sp. nov.** and *N. tomiyamai*.

Species	<i>N. rara</i>		<i>N. tomiyamai</i>		
	HT	PT	HT	NT	
Type status					
Number of specimens	<i>n</i> = 1	<i>n</i> = 2	<i>n</i> = 1	<i>n</i> = 2	
References	This study	This study	Ori. descr.**	This study	This study
Measurements (mm)					
TL	419	415–447+	350	321+	300+–372+
HL	73.0	71.6–78.9	58.7	58.4	58.4–68.7
% of HL (% of PRL)					
Snout length	27 (36)	26–28 (35–37)	21	27 (35)	27 (35)
Orbit diameter	29 (39)	26–28 (35–38)	30	28 (37)	31 (40)
Postorbital length	46 (62)	47–49 (63–66)	49	49 (63)	51 (66)
Postrostral length	74 (–)	75 (–)	–	77 (–)	77 (–)
Orbit–preopercle distance	41 (56)	39–41 (52–56)	–	42 (55)	43 (57)
Suborbital width	16 (22)	15 (20)	15	15 (20)	16 (20)
Upper-jaw length	36 (48)	33–34 (45)	34	36 (47)	37 (48)
Length of rictus	28 (37)	28 (38)	–	28 (36)	29 (38)
Length of PMX tooth band	23 (31)	19–21 (26–28)	–	22 (29)	24 (31)
Preoral length	22 (30)	16–19 (22–25)	–	20 (26)	21 (27)
Dist. tip and lat. angle of snout	15 (20)	12–15 (16–19)	–	15 (19)	12 (16)
Snout width	23 (31)	24 (32)	–	25 (33)	23 (30)
Internasal width	21 (28)	19 (25–26)	–	21 (27)	23 (29)
Interorbital width	21 (28)	20–22 (26–29)	24	23 (30)	25 (33)
Body width over P. bases	59 (80)	57 (77)	–	42 (55)	53 (69)
Body depth at 1D. origin	90 (121)	86 (116)	–	73 (95)	85 (111)
Body depth at A. origin	79 (106)	69–81 (93–108)	–	63 (81)	70 (91)
Pre-V. length	115 (156)	132 (177)	–	104 (136)	113 (147)
Pre-anus length	137 (185)	– (–)	132	130 (170)	128 (167)
Pre-A. length	159 (214)	166 (223)	–	155 (201)	155 (202)
Isthmus–V. distance	40 (54)	42 (56)	48	44 (57)	36 (47)
Isthmus–anus distance	– (–)	– (–)	–	– (–)	52 (67)
Isthmus–A. distance	– (–)	– (–)	–	– (–)	80 (105)
V.–A. distance	47 (64)	41 (55)	–	53 (69)	48 (63)
Anus–A. distance	26 (36)	– (–)	–	28 (37)	28 (36)
V. length	54 (74)	46 (61)	–	– (–)	69 (90)
P. length	63 (86)	61 (81)	–	– (–)	58 (75)
Pre-1D. length	117 (159)	106–113 (142–152)	–	119 (154)	118 (154)
Height of 1D.	98 (133)	80–95 (107–127)	–	78 (102)	90 (117)
Length of 1D. base	29 (39)	29–32 (39–43)	–	26 (34)	37 (48)
Interdorsal length	41 (56)	40–45 (54–60)	–	46 (60)	40 (53)
Length of gill slit	16 (21)	13–14 (18–19)	–	18 (23)	17 (23)
Length of posterior nostril	3 (4)	4 (5)	–	7 (9)	6 (8)
Barbel length	17 (23)	19 (26)	21	15 (20)	20 (26)
Counts					
1D. rays	II,10	II,10–11	II,10	II,10	II,12
P. rays	i21	i20–i22	20***	i20	i21
V. rays	11	10–11	10	10	11
Outer GR on first arch	8	7–9	–	8	7–8
Inner GR on first arch	10	9	11	11	8–9
Outer GR on second arch	10	9	–	10	8–9
Inner GR on second arch	10	9–10	–	10	10–11
Lateral-line scales	39	37	–	39	36
Scales below 1D. origin	10.5	10–12	–	10.5	10.5
Scales below 1D. midbase	8.5	7.5–9.5	7–7.5	7.5	9.5

...Continued on the next page

TABLE 13. (Continued)

Species	<i>N. rara</i>		<i>N. tomiyamai</i>		
Type status	HT	PT	HT	NT	
Number of specimens	<i>n</i> = 1	<i>n</i> = 2	<i>n</i> = 1	<i>n</i> = 2	
References	This study	This study	Ori. descr.**	This study	This study
Scales below 2D. origin	9	9–10	9	8.5	8.5
Scales above A. origin	–	–	–	–	30

*Abbreviations: 1D.—first dorsal fin; 2D.—second dorsal fin; A.—anal fin; GR—gill rakers; HT—holotype; NT—non-type; P.—pectoral fin; PMX—premaxillary; PT—paratype; V.—pelvic fin

**Morphometric values given here are calculated from those given in the original description (Okamura 1963a)

***Probably i19

***Nezumia shinoharai* Nakayama & Endo, 2012**

[Japanese name: Shinohara-hige]

(Figs. 173–174, 175A; Appendix 3-10E)

Nezumia shinoharai Nakayama & Endo, 2012:62, figs. 1–3, 4A.

(original description; holotype: NSMT-P 97667, from Pacific off Boso Peninsula, in 627–673 m; 1 paratype from Tosa Bay; new Japanese name: “Shinohara-hige”); Nakabo & Kai 2013:1874 (comments on Japanese records); Motomura 2020:39 (listed; Japan).

Diagnosis. A species of *Nezumia* with 10 pelvic-fin rays. Snout moderately long, conical, protruding well beyond upper jaw, length 31–34% HL, ventral contour oblique in lateral view; underside of head almost completely naked. Second spinous ray of first dorsal fin not extremely prolonged (height of fin 99% HL). Body scales covered with short, greatly reclined, needle-like spinules in discrete parallel rows; tip of last spinule in each row extending well beyond posterior scale margin; middle row of spinules not enlarged; scales below second dorsal-fin origin 8–9. Cephalic sensory pores well developed especially on mandibular and infraorbital canals. No prominent dark

band encircling trunk; first dorsal fin uniformly dark with pale tip.

Material examined. 2 specimens. **Holotype of *Nezumia shinoharai*:** NSMT-P 97667 (38.0 mm HL, 244+ mm TL), east of Boso Peninsula, Chiba Pref., Japan, 35.1048°N, 140.8367°E, 627–673 m, R/V *Tansei-maru*, cr. KT-95-05, sta. TB16, 3-m ORE beam trawl, 25 Apr. 1995. **Paratype of *N. shinoharai*:** BSKU 44790 (1, 33.3 mm HL, 198+ mm TL), Tosa Bay, 33.0517°N, 133.6152°E, 700 m, FRV *Kotaka-maru*, 24 May 1988.

Counts and measurements. Based on 2 specimens (33.3–38 mm HL, 198+–244+ mm TL). Counts: first dorsal-fin rays II,9; pectoral-fin rays i20–i24; pelvic-fin rays 10; gill rakers on first arch (outer/inner) 9/10–11, on second arch 10/10–11; longitudinal scales 38–39; transverse scale rows below first dorsal-fin origin 10.5, below first dorsal-fin midbase 7.5, below second dorsal-fin origin 8–9.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 31–34 (45–49); orbit diameter 29–30 (43–43); postorbital length 39–42 (57–60); postrostral length 68–69; orbit–preopercle distance 34–35 (50); suborbital width 14–15

FIGURE 173. Fresh specimen of *Nezumia shinoharai*. BSKU 44790, paratype, 33.3 mm HL, 198+ mm TL, Tosa Bay, Kochi Pref., Pacific, 700 m depth. Lateral view. [Photo: reproduced from Nakayama & Endo (2012: fig. 2A)]

FIGURE 174. Holotype of *Nezumia shinoharai*. NSMT-P 97667, 38.0 mm HL, 244+ mm TL, east of Boso Peninsula, Chiba Pref., Pacific, 627–673 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of head. Preserved condition. [Photos: (A) reproduced from Nakayama & Endo (2012: fig. 1); (B–C) N. Nakayama]

(21–22); upper-jaw length 26–27 (38–40); length of rictus 19–20 (28–29); length of premaxillary tooth band 14 (20); preoral length 29–30 (42–44); distance between tip and lateral angle of snout 18–19 (27); snout width 25–26 (36–37); internasal width 22–22 (32–32); interorbital width 23–25 (33–36); body width over pectoral-fin bases 42–47 (61–68); body depth at first dorsal-fin origin 71–80 (104–116); body depth at anal-fin origin 66 (96); prepelvic length 102–105 (147–154); preanus length 113–120 (163–175); preanal length 134–142 (193–208); isthmus–pelvic distance 35–36 (51–52); pelvic–anal distance 37–41 (53–61); anus–anal distance 23–24 (33–34); pelvic-fin length 62–65 (90–96); pectoral-fin length 52–55 (77–79); predorsal length 113–116 (166–167); height of first dorsal fin 99 (143); length of first dorsal-fin base 23–28 (34–40); interdorsal length 39–41 (58–59); length of gill slit 12–14 (17–20); length of posterior nostril 8 (11); barbel length 11–12 (15–17).

Size. To about 24 cm TL (NSMT-P 97667, holotype, 244+ mm TL, east of Boso Peninsula, Japan).

Distribution. So far known only from off the Boso Peninsula and Tosa Bay at depths of 627–700 m (Nakayama & Endo 2012; Appendix 3-10E). Very rare.

Remarks, relationships, and comparisons. *Nezumia*

shinoharai was recently described by Nakayama & Endo (2012) based on two specimens collected from southern Japan (Figs. 173–174). Apart from the type specimens, no additional materials are available for this study. It belongs to the *N. spinosa* group as defined by Nakayama & Endo (2015), in which the following six species are included: *N. aspidentata* Iwamoto & Merrett, 1997 from New Caledonia; *N. holocentra* (Gilbert & Cramer, 1897) from Hawaii; *N. infranudis* Gilbert & Hubbs, 1920 from the Celebes and Timor Seas; *N. merretti* Iwamoto & Williams, 1999 from the eastern Indian Ocean; *N. shinoharai*; and *N. spinosa* (Gilbert & Hubbs, 1916) from the Indo-West Pacific.

Nezumia shinoharai readily differs from other species of that group by its unique spinulation on the body scales: in *N. shinoharai*, spinules are arranged in discrete parallel rows (Fig. 175A), whereas those of the other five species are in tightly packed subparallel or convergent rows, or densely scattered over the exposed portion of the scales. It also differs from all but *N. aspidentata* in having 10 pelvic-fin rays (vs. 8–9 or 11). For further details see Nakayama & Endo (2012, 2015).

FIGURE 175. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of two species of the *Nezumia spinosa* group. (A) *N. shinoharai*, NSMT-P 97667, holotype, 38.0 mm HL; (B) *N. spinosa*, USNM 76868, holotype, 46.4 mm HL. Views from above. [Photos: reproduced from Nakayama & Endo (2012: fig. 4A, E)]

***Nezumia spinosa* (Gilbert & Hubbs, 1916)**

[Japanese name: Goten-hige]

(Figs. 175B, 176–177; Appendix 3-10F)

Lionurus spinosus Gilbert & Hubbs, 1916:199, pl. 10, fig. 2 [original description; holotype: USNM 76868, from “Eastern Sea” (East China Sea), *Albatross* sta. 4915, in 427 ftm (781 m)]; Okada & Matsubara 1938:453 (in key; Japan); Matsubara 1955:1317 (in key; Japan); Matsubara 1965:510 (compiled; Japan; new Japanese name: “Goten-hige”).

Lionurus proximus (not Smith & Radcliffe in Radcliffe 1912): Gilbert & Hubbs 1916:201 [in part, 1 spec. (CAS-SU 22941) from East China Sea, *Albatross* sta. 4918].

Nezumia spinosa: Okamura 1984b:94, pl. 345, fig. A (compiled); Okamura 1988:94, pl. 345, fig. A (compiled); Nakabo 1993:360 (in key; Japan); Nakabo 2000:424 (in key; Japan); Nakabo 2002:424 (in key; Japan); Suetsugu & Ohta 2005: table 3 (questionable; listed; Enshu-nada); Shao *et al.* 2008b: table 2 (3 spec. listed from southwestern Taiwan and South China Sea; first Taiwanese record); Nakabo & Kai 2013:500 (in key; Japan); Iwamoto *et al.* 2015:94, fig. 20 (brief description; 6 spec. from southwestern Taiwan and South China Sea); Motomura 2020:39 (listed; Japan).

Diagnosis. A species of *Nezumia* with 8 pelvic-fin rays. Snout moderately long, conical, protruding well beyond upper jaw, length 27–33% HL, ventral contour oblique in lateral view; underside of head almost completely naked. Second spinous ray of first dorsal fin extremely prolonged (height of fin 120–138% HL). Body scales covered with short, greatly reclined, needle-like spinules densely scattered over exposed portion; posteriormost spinules extending well beyond scale margin; no spinules especially enlarged; scales below second dorsal-fin

origin 7.5–8.5. Cephalic sensory pores well developed especially on mandibular and infraorbital canals. No prominent dark band encircling trunk; first dorsal fin uniformly dark.

Material examined. 9 specimens. **Holotype of *Lionurus spinosus*:** USNM 76868 (46.4 mm HL, 262+ mm TL), southwest of Kamikoshiki-jima Island, Kagoshima Pref., Japan, East China Sea, 31.5167°N, 129.4250°E, 427 ftm (781 m), *Albatross* sta. 4915, 9-ft Agassiz beam trawl, 12 Aug. 1906. **Non-types: Japan:** CAS-SU 22941 (1, 19.6 mm HL, 117+ mm TL), west of Yaku-shima Island, Okinawa Trough, 30.3667°N, 129.1417°E, 660 m, *Albatross* sta. 4918, 13 Aug. 1906; SFU 1840 (1, 37.0 mm HL, 176+ mm TL), South China Sea, Apr. 1982. **Australia:** CSIRO H1971-01 (1, 40.8 mm HL, 199+ mm TL), off Queensland, 16.9167°S, 151.5667°E, 880 m, 6 Dec. 1985; CSIRO H1961-01 (1, 40.4 mm HL, 197+ mm TL), off Queensland, 18.9000°S, 150.4833°E, 1005–1013 m, 25 Nov. 1985; CSIRO H2549-10 (1, 48.8 mm HL, 254+ mm TL), off Western Australia, 21.8500°S, 113.7667°E, 650–685 m, 24 Jan. 1991; CSIRO H2580-04 (1, 47.5 mm HL, 249+ mm TL), off Western Australia, 27.1000°S, 112.3667°E, 713–714 m, 31 Jan. 1991. **Timor Sea:** BSKU 16687 (1, 35.5 mm HL, 183+ mm TL), BSKU 16688 (1, 19.4 mm HL, 130 mm TL), 9.4500°S, 127.9767°E, 690–850 m, R/V *Hakuho-maru*, cr. KH-72-01, coll. O. Okamura, 18–9 Jan. 1972.

Counts and measurements. Based on 9 specimens (19.4–48.8 mm HL, 117+–262+ mm TL). Counts: first dorsal-fin rays II,8–10; pectoral-fin rays i18–i22; pelvic-fin rays 8; gill rakers on first arch (outer/inner) 4–11/9–11, on second arch 8–11/9–11; longitudinal scales 37–40; transverse scale rows below first dorsal-fin origin 7.5–12, below first dorsal-fin midbase 5.5–9, below second dorsal-fin origin 7.5–8.5, above anal-fin origin 22–23.

FIGURE 176. Holotype of *Lionurus spinosus* (= *Nezumia spinosa*). USNM 76868, 46.4 mm HL, 262+ mm TL, southwest of Kamikoshiki-jima Island, Kagoshima Pref., East China Sea, 781 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of head. Preserved condition. [Photos: N. Nakayama]

FIGURE 177. Preserved juvenile specimen of *Nezumia spinosa*. CAS-SU 22941, 19.6 mm HL, 117+ mm TL, west of Yaku-shima Island, Kagoshima Pref. East China Sea, 660 m depth. Lateral view. [Photo: N. Nakayama]

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 27–33 (36–46); orbit diameter 27–32 (38–43); postorbital length 41–47 (57–63); postrostral length 71–75; orbit–preopercle distance 33–40 (46–53); suborbital width 11–19 (15–26); upper-jaw length 27–33 (35–45); length of rictus 20–26 (27–34); length of premaxillary tooth band 14–21 (20–28); preoral length 18–28 (24–39); distance between tip and lateral angle of snout 11–20 (15–26); snout width 20–27 (27–36); internasal width 16–20 (21–28); interorbital width 19–22 (25–31); body width over pectoral-fin bases 41–58 (55–82); body depth at first dorsal-fin origin 72–82 (102–115); body depth at anal-fin origin 63–77 (88–103); prepelvic length 104–124 (138–170); preanus length 118–137 (156–188); preanal length 141–156 (194–216); isthmus–pelvic distance 33–40 (45–54); isthmus–anus distance 45–49 (63–65); isthmus–anal distance 60–78 (84–106); pelvic–anal distance 32–43 (44–60); anus–anal distance 16–29 (22–40); pelvic-fin length 55–88 (78–124); pectoral-fin length 51–121 (71–162); predorsal length 110–124 (154–168); height of first dorsal fin 120–138 (165–195); length of first dorsal-fin

base 23–30 (32–42); interdorsal length 34–56 (46–75); length of gill slit 10–14 (13–19); length of posterior nostril 6–11 (8–15); barbel length 9–14 (12–20).

Size. To at least 26 cm TL (Iwamoto & Williams 1999). The holotype is the largest specimen of this species, measuring 262+ mm TL.

Distribution. Widely distributed in the Indo-West Pacific from South Africa, eastward to Australia and New Caledonia, and northward to Japan, at depths of 420–1258 m (Gilbert & Hubbs 1920; Iwamoto & Anderson 1994; Iwamoto & Merrett 1997; Iwamoto & Williams 1999; Shao *et al.* 2008a, 2008b; Last *et al.* 2014; Iwamoto *et al.* 2015; Nakayama & Endo 2015). Very rare in Japanese waters, but not in other parts of its range. The Japanese records based on only two *Albatross* specimens collected from the Okinawa Trough in 660–781 m (Appendix 3-10F).

Remarks. For a full description see the original description given by Gilbert & Hubbs (1916). *Nezumia spinosa* was originally described from a single specimen collected from the East China Sea off Japan (Fig. 176). Subsequently, it has been widely recorded from the Indo-West Pacific, including the South China Sea off Taiwan

(Shao *et al.* 2008a, 2008b; Iwamoto *et al.* 2015), the Philippines (Gilbert & Hubbs 1920), Western Australia (Iwamoto & Williams 1999), New Caledonia and Fiji (Iwamoto & Merrett 1997; Merrett & Iwamoto 2000), Timor Sea (Nakayama & Endo 2015), and eastern Indian Ocean off South Africa (Iwamoto & Anderson 1994). However, *N. spinosa* appears to include multiple species, as previous authors have mentioned (see below). Gilbert & Hubbs (1920:554) indicated that four small specimens from the Philippines differ from the holotype in having a larger head and a deeper body, suggesting the presence of a second species. Similarly, Iwamoto & Williams (1999:204, table 6) highlighted notable differences in meristic and morphometric characters between African and Australian populations of *N. spinosa*, particularly in the mode of first dorsal-fin ray counts (II,8 vs. II,9 in the African and Australian populations respectively), preoral length (28–35% HL vs. 18–29%), and postorbital length (37–42% vs. 43–46%). They implied that the African population might represent an undescribed species, noting “isolation has resulted in significant divergence between the populations on each side of the Indian Ocean”.

The Japanese records of *N. spinosa* are based on only two specimens: one is the holotype and the other is a small specimen (CAS-SU 22941, 19.6 mm HL, 117+ mm TL; Fig. 177) collected from the type locality. This second specimen was initially reported by Gilbert & Hubbs (1916) as *N. proxima* (Smith & Radcliffe in Radcliffe, 1912), but it was subsequently re-identified as *N. spinosa* by Iwamoto & Merrett (1997). Although Okamura (1970a) considered *N. spinosa* as a junior synonym *N. proxima*, he (Okamura 1984a, 1984b) resurrected this species after an examination of its holotype. In his list of fishes from Iwate Pref., Maruyama (1971) listed *N. spinosa* with a question mark (as *Lionurus spinosus*). Although the whereabouts of his specimen(s) is unknown, it probably represents *N. proxima*, considering the common occurrence of this species along the Pacific coasts of northern Japan.

Relationships and comparisons. *Nezumia spinosa* belongs to the *N. spinosa* group as defined by Nakayama & Endo (2015) (see the Remarks, relationships, and comparisons of *N. shinoharai*), and is most similar to *N. infranudis* Gilbert & Hubbs, 1916. The latter species is so far represented by the holotype from the Celebes Sea and an additional specimen recently reported from the Timor Sea (Nakayama & Endo 2015). *Nezumia spinosa* is readily distinguished from that species by having lower counts of pelvic-fin rays (8 vs. 11).

Nezumia tomiyamai (Okamura, 1963)

[Japanese name: Tomiyama-hige]

(Figs. 172C–D, 178–179; Table 13; Appendix 3-11A)

Lionurus tomiyamai Okamura, 1963a:31, figs. 8–10 (original

description, holotype ZMUT 52045, from “off eastern coast of Japan”; new Japanese name: “Tomiyamahige”).

Nezumia tomiyamai: Okamura 1970a:98, pl. XXII, text-fig. 42 [description after Okamura (1963)]; Okamura 1970b: table 1 (listed; Japan); Tominaga & Uyeno 1981:489 (listed; Japan); Okamura 1984b:94, pl. 345, fig. B (compiled); Okamura 1988:94, pl. 345, fig. B (compiled); Nakabo 1993:360 (in key; Japan); Nakabo 2000:424 (in key; Japan); Nakabo 2002:424 (in key; Japan); Nakabo & Kai 2013:500 (in key; Japan); Amaoka *et al.* 2020:159, fig. 212 (photograph of holotype; erroneously listed from Hokkaido); Motomura 2020:39 (listed; Japan).

Nezumia sp.: Okamura 1982:306 (listed; 1 spec., BSKU 26116; Kyushu-Palau Ridge).

Diagnosis. A species of *Nezumia* with 10–11 pelvic-fin rays. Snout short, protruding scarcely beyond upper jaw, length 27% HL, ventral contour almost vertical in lateral view; underside of snout broadly naked; ventral surfaces of head scaled posterior to vertical through lateral angles of snout; mandibular rami mostly scaled; orbit–preopercle distance 42–43% HL; interorbital width 23–25% HL; pelvic-fin length 69% HL; length of outer gill slit 17–18% HL. Second spinous ray of first dorsal fin not extremely prolonged (height of fin 78–90% HL). Body scales covered with short, greatly reclined, narrowly triangular spinules in tightly packed convergent rows or densely scattered over exposed portion; tip of last spinule in each row extending slightly beyond posterior scale margin; middle row of spinules not enlarged; scales below second dorsal-fin origin 8.5. Cephalic sensory pores present along all canals, but barely discernible. No dark band encircling trunk; first dorsal fin uniformly blackish.

Material examined. 3 specimens. **Holotype** of *Lionurus tomiyamai*: ZMUT 52045 (58.4 mm HL, 321+ mm TL), off east coast of Japan, no further information. **Non-types: Japan:** BSKU 26116 (1, 58.4 mm HL, 300+ mm TL), Kyushu-Palau Ridge, 27.9033°N, 134.6583°E, 700 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. 2-4, bottom trawl, coll. Y. Kobayashi *et al.*, 20 Feb. 1978; HUMZ 71963 (1, 68.7 mm HL, 372+ mm TL), Shichito-Iojima Ridge, 26.6333°N, 140.8833°E, 580–1100 m, 1 Nov. 1977.

Counts and measurements. See Table 13.

Size. To about 37 cm TL (HUMZ 71963, 372+ mm TL, Shichito-Iojima Ridge, Japan).

Distribution. So far known only from Japanese waters. Specimens associated with reliable locality information were collected from only the Kyushu-Palau and Shichito-Iojima Ridges at depths of 580–1100 m (Appendix 3-11A). Very rare.

Remarks. This species was fully described by Okamura (1963a), which readers can refer to for further detail. During the course of this study, one additional specimen of this rare species was found among the HUMZ collection (HUMZ 71963, 68.7 mm HL, 372+ mm TL,

FIGURE 178. Holotype of *Lionurus tomiyamai* (= *Nezumia tomiyamai*). ZMUT 52045, 58.4 mm HL, 321+ mm TL, off east coast of Japan, Pacific. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

FIGURE 179. Preserved specimen of *Nezumia tomiyamai*. HUMZ 71963, 68.7 mm HL, 372+ mm TL, Shichito-Iojima Ridge, Pacific, 580–1100 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

from Shichito-Iojima Ridge; Fig. 179). Okamura *et al.* (1982:306) reported “*Nezumia* sp.” from the Kyushu-Palau Ridge. Unfortunately, his specimen (BSKU 26116, 58.4 mm HL, 300+ mm TL) is in poor condition, but it appears to be a representative of *N. tomiyamai*. Amaoka *et al.* (2020) recently listed the species from Hokkaido, but it is obviously based on Sawada’s (1983) record from the Pacific off Tohoku. It should be noted that their photographed specimen represents the holotype collected from “off eastern coast of Japan” (not from Hokkaido; Okamura 1963a).

Comments on holotype. *Nezumia tomiyamai* was

originally described by Okamura (1963a) based on a single specimen collected from “off eastern coast of Japan” (no further information). Although he designated the specimen “No. 52045, Zoological Institute, Faculty of Science, Tokyo University” as the holotype of this species, the specimen associated with this catalog number was not found during the author’s visit to the Zoological Museum of the University of Tokyo (ZMUT) in September 2011. However, among the collection was an uncataloged specimen that agrees well with the original description of *N. tomiyamai* (Fig. 178). This specimen does not bear a catalog number, but it has a cloth tag with the word

“higedara” meaning grenadiers (= hige) and gadiform fishes (= dara). A close examination of this specimen along with Okamura’s (1963a) original description has convinced the author that the specimen represents the holotype of *N. tomiyamai*. No significant differences were found in meristic characters between the ZMUT specimen and the original description (Table 13). Other characters, including dentition, squamation, spinulation of the body scales, and fin position, are also identical between the two. Some differences in morphometric characters, especially in the snout length (27% HL in this study vs. 21% in the original description) and barbel length (15% HL vs. 21%) (Table 13), seem to be attributed to different methods of making measurements. According K. Sakamoto (pers. com., ZMUT), the holotype of *N. tomiyamai* was probably sent to Okamura prior to registration of the specimen.

Relationships and comparisons. *Nezumia tomiyamai* is characterized by having 10 or 11 pelvic-fin rays, and falls into “Group E” or “Group F” in the key to species of *Nezumia* given by Iwamoto (1990), who subdivided the genus into seven groups (A–G) based on the number of pelvic-fin rays. Among species of these two groups, *N. tomiyamai* is similar to the following three species: *N. africana* (Iwamoto, 1970) known from the west coast of Africa; *N. atlantica* (Parr, 1946) from the eastern Atlantic; and *N. kensmithi* Wilson, 2001 from seamounts in the northeastern Pacific. They are closely similar to each other in sharing the following characters: snout short, bluntly pointed, with ventral profile almost vertical; body scales covered with lanceolate or triangular spinules in convergent rows (or densely scattered over exposed portion); underside of snout heavily scaled posterior to lateral angles; and body uniformly dark, lacking prominent dark band encircling trunk.

Nezumia tomiyamai readily differs from these three species in that spinules on the body scales are much smaller, and their rows are more numerous as compared with the latter three species (Fig. 172C–D). It is also distinguished by a combination of morphometric characters, including the suborbital width (15–16% HL vs. 12–14%, 12–15%, and 12–14% in *N. africana*, *N. atlantica*, and *N. kensmithi* respectively), preoral length (20–21% HL vs. 15–19%, 10–13%, and 15–21% respectively), internasal width (21–23% HL vs. 18–19%, 15–18%, and 17–20% respectively) and interorbital width (23–25% HL vs. 19–22%, 17–22%, and 19–24% respectively). *Nezumia tomiyamai* further differs from *N. africana* in having more pelvic-fin rays (10–11 vs. 8–9), from *N. atlantica* in having small but prominent pores along the cephalic sensory canals (vs. probably closed, except for the preopercular and postorbital canals), and from *N. kensmithi* in having fewer scales below the second dorsal-fin origin (8.5 vs. 9–10).

The eastern Pacific *N. stelgidolepis* (Gilbert, 1890) and *N. pudens* Gilbert & Thompson in Thompson, 1916 are similar in general appearance to *N. tomiyamai*.

However, *N. stelgidolepis* is unlikely to be confused with *N. tomiyamai* in having prominent scaly patches on lower portions of the branchiostegal membranes (vs. completely naked), and lower counts of transverse scale rows [8–10 vs. 10.5 below the first dorsal-fin origin; 6–7 vs. 7.5–9.5 below the first dorsal-fin midbase; 7–8 (rarely 9) vs. 8.5 below the second dorsal-fin origin]. *Nezumia pudens* readily differs from *N. tomiyamai* in that the ventral surface of the snout is completely scaled except for a narrow naked band above the upper jaw (vs. broadly naked). It appears to be further distinguished from *N. tomiyamai* by having a narrower interorbital space (22–23% HL vs. 23–25%) and a longer chin barbel (20–25% HL vs. 15–20). [Data for *N. stelgidolepis* and *N. pudens* are based on Iwamoto (1979) and this study].

In the original description of *N. tomiyamai*, Okamura (1963a) considered this species to be most similar to *N. burragei* (Gilbert, 1905), a rare species confined to Hawaii [see also the Comments on Japanese record of *Nezumia burragei* (given below)]. However, it differs notably from *N. burragei* in that the mandibular rami are scaled except for anterior portions (vs. completely naked) and the lower surface of the suborbital area is heavily scaled (vs. the underside of the head is broadly naked from the snout tip to a vertical at the posterior margin of the upper jaw (see Gilbert 1905: fig. 260). It further differs from that species in having a wider gill slit (17–18% HL vs. 13%), higher counts of pelvic-fin rays (10–11 vs. 9), and lower counts of scales (longitudinal scales 36–39 vs. 42; scales below the first dorsal-fin origin 10.5 vs. 15.5; below the first dorsal-fin midbase 7.5–9.5 vs. 11; below the second dorsal-fin origin 8.5 vs. 13).

Comments on Japanese record of *Nezumia burragei* (Fig. 180)

Kamohara (1937) recorded *N. burragei* (Gilbert, 1905) from Japan based on a single 350 mm long specimen obtained at the Mimase fish market, Kochi. This is a very rare species, which elsewhere is known only from the holotype from Hawaii at a depth of 543–603 m (USNM 51641, 44.2 mm HL, 254+ mm TL; examined here). The occurrence of *N. burragei* in Japan is quite dubious, considering that no additional specimens have been obtained despite the relatively shallow occurrence of the species and extensive trawl surveys in southern Japan over the past several decades. Citing Kamohara’s (1937) record, all subsequent authors have believed *N. burragei* to occur in Japanese waters (e.g., Matsubara 1955; Okamura 1970a, 1984b, 1988; Nakabo 1993, 2000, 2002; Shinohara *et al.* 2001; Nakabo & Kai 2013). Unfortunately, the voucher specimen of Kamohara (1937) was apparently lost during the WWII, which makes it difficult to reconfirm the Japanese record of this species. However, Kamohara’s (1937) specimen appears to be

FIGURE 180. Drawing of a species of *Nezumia* illustrated by Toshiji Kamohara (deposited at BSKU). The illustrated specimen, which was destroyed during WWII, was the basis for the first Japanese record of *N. burragei* by Kamohara (1937). [Photo: BSKU]

attributable to a different species of *Nezumia*, due to notable differences between his figure (Fig. 180) and the holotype of *N. burragei* (see Gilbert 1905: fig. 260). Kamohara (1937) illustrated his specimen as having the ventral surface of the snout only narrowly naked above the upper lip, and the posterior margin of the naked area not extending beyond a vertical through the anterior rim of the orbit. In contrast, the ventral surface of the snout and the lower half of the suborbital area are completely naked in the holotype of *N. burragei*, with the hind margin of the naked area extending to above the posterior margin of the upper jaw. The difference between the two specimens is also evident in the dorsal contour of the head. Kamohara's (1937) figure shows that the predorsal contour of his specimen was prominently humped over the nape, whereas in the holotype, the dorsal contour of the head is straight from above the orbit to the first dorsal-fin origin.

It is most likely that the record of Kamohara (1937) represents a misidentification of *N. rara* **sp. nov.** described here. Many features of his specimen, including fin-ray counts, head squamation, and the general appearance, agree well with those of the new species. Another possibility is that his specimen was a representative of *N. proxima* (Smith & Radcliffe in Radcliffe, 1912), a moderately common species at upper and mid-slope depths in the Pacific off southern Japan. Although Kamohara extensively surveyed the deep demersal fishes of Tosa Bay and adjacent waters, he did not record *N. proxima* during his career. *Nezumia burragei* appears to be confined to the Hawaiian Archipelago, and is unlikely to occur in Japanese waters. Therefore, this species is excluded from the list of Japanese grenadiers in this study.

Comments on Japanese record of *Nezumia propinqua*

Okamura (1982) first reported *N. propinqua* (Gilbert & Cramer, 1897) from Japan based on six specimens collected from the Kyushu-Palau Ridge. Unfortunately, these specimens are in poor condition, with the tip of the snout damaged in most specimens. According to Okamura (1982, 1984a, 1984b), *N. propinqua* differs from *N. condylura* in having the anus closer to the pelvic-fin bases than to the anal-fin origin (vs. closer to anal-fin origin), a much broader periproct, and the anterior dermal window of the light organ situated between the inner bases of the pelvic fin (vs. “between pelvic bases”). However, these minor differences are likely to be attributable to intraspecific variation, considering that his specimens (“216–250 mm TL”) are much larger than the *N. condylura* specimens he examined (“152–200 mm”). Counts and measurements of Okamura’s “*N. propinqua*” lie within, or significantly overlap with those of the *N. condylura* specimens examined in this study (Table 11). Accordingly, his specimens are herein re-identified as *N. condylura*. Apart from that, no additional specimens of *N. propinqua* have been reported from Japan, and thus this species is excluded from the list of Japanese grenadiers in this study.

Genus *Odontomacrus* Norman, 1939

[Japanese name: Kurobozudara-zoku]

Odontomacrus Norman, 1939:49 (type species: *Odontomacrus murrayi* Norman, 1939, by original designation).

Phalacromacrus Maul & Koefoed, 1950:971 (type species: *Phalacromacrus pantherinus* Maul & Koefoed, 1950, by original designation).

Diagnosis. Anus midway between pelvic-fin bases and anal-fin origin, preceded by small dermal window of light organ. Second spinous ray of first dorsal fin smooth along its leading edge. Body deep, well compressed laterally. Snout broadly rounded, barely protruding beyond upper jaw. Mouth large, terminal; posterior margin of premaxillary extending beyond vertical drawn through hind rim of orbit. Chin barbel absent. Posterior margin of gill cover notably incised between opercle and subopercle. Teeth large, fang-like, arranged in 1 distinct row in both jaws. Body scales adherent, covered with long, erect, needle-like spinules in quincunx order; buttresses of spinules greatly developed, connected with those of adjacent spinules; reticulate structure absent. Lateral line short, ending at about level of second dorsal-fin origin. Cephalic sensory pores large, prominent. Swim bladder reduced. Pelvic-fin rays 8–11. Branchiostegal rays 6. Color uniformly black. [From Nakayama *et al.* (2015b).]

Remarks. *Odontomacrus* is a monotypic genus, so far known only from its type species, *O. murrayi* Norman, 1939. This species is one of few grenadiers whose vertical

distribution is confined to midwater depths of the open oceans (Marshall 1964, 1965; Marshall & Iwamoto 1973; Iwamoto 1990; Iwamoto & Graham 2001; Nakayama *et al.* 2015b). Based on only small specimens collected from the northeastern Atlantic, Maul & Koefoed (1950) described *Phalacromacrus pantherinus* as a new genus and species of macrourids. However, Marshall (1964) revealed *P. pantherinus* to represent a juvenile stage of *O. murrayi*. *Odontomacrus* is a distinctive genus that is unlikely to be confused with other genera, owing to its unique morphological characters adapted to pelagic life. Such features include the following: uniformly black color of the body; a long, string-like tail; a large, terminal mouth; sharp, fang-like teeth arranged in one distinct row in both jaws; and prominent open pores along the cephalic sensory canals.

Odontomacrus murrayi Norman, 1939

[Japanese name: Kurobozudara]

(Figs. 181–182; Appendix 3-11B)

Odontomacrus murrayi Norman, 1939:49, text-fig. 18 (original description; holotype: BMNH 1939.5.24.685, from Arabian Sea, in 0–2500 m); Nakayama *et al.* 2015b:136, figs. 1–2 (first record from Japan; description; 1 spec. from Pacific off Tohoku; new Japanese name: “Kurobozudara”); Motomura 2020:39 (listed; Japan).

Phalacromacrus pantherinus Maul & Koefoed, 1950:972, fig. 1 (original description; holotype: ZMUB 4473, from northeastern Atlantic, 2600 m wire out; paratypes deposited in ZMUB and MMF).

Diagnosis. As for genus.

Material examined. 1 specimen. **Japan:** BSKU 104866 (1, 30.0 mm HL, 325+ mm TL), off Kinkazan, 37.9665°N, 143.2269°E, 168–204 m, FRV *Kaiyo-maru*, cr. KY-12-07, sta. 6, coll. K. Uchikawa *et al.*, 16 Dec. 2009.

Counts and measurements. Based on 1 specimen. Counts: first dorsal-fin rays II,9; pectoral-fin rays i8; pelvic-fin rays 8; gill rakers on first arch (outer/inner) 8–9/11, on second arch 9–10/10–11.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 34 (47); orbit diameter 24 (33); postorbital length 49 (68); postrostral length 72; orbit–preopercle distance 52 (72); suborbital width 23 (31); upper-jaw length 70 (98); length of rictus 66 (91); length of premaxillary tooth band 55 (76); internasal width 21 (30); interorbital width 32 (44); occipital width 12 (16); body width over pectoral-fin bases 30 (42); body depth at first dorsal-fin origin 111 (154); body depth at anal-fin origin 99 (137); prepelvic length 103 (143); preanus length 118 (163); preanal length 136 (189); isthmus–pelvic distance 31 (43); isthmus–anus distance 39 (55); isthmus–anal distance 60 (83); pelvic–anal distance 37 (52); anus–anal

FIGURE 181. Fresh specimen of *Odontomacrus murrayi*. BSKU 104866, 30.0 mm HL, 325+ mm TL, off Oshika Peninsula, Miyagi Pref., Pacific, 168–204 m depth. Lateral view. [Photo: reproduced from Nakayama *et al.* (2015b: fig. 1) with permission from the Japanese Society of Systematic Zoology]

FIGURE 182. Scanning electron micrographs showing a body scale (from the dorsum below the interdorsal space) of *Odontomacrus murrayi*. BSKU 104866, 30.0 mm HL. (A) View from above; (B) oblique view. [Photos: reproduced from Nakayama *et al.* (2015b: fig. 2) with permission from the Japanese Society of Systematic Zoology]

distance 25 (35); pelvic-fin length 44 (62); pectoral-fin length 84 (116); predorsal length 105 (146); height of first dorsal fin 52 (71); length of first dorsal-fin base 42 (59); interdorsal length 63 (87); length of gill slit 23 (32); length of posterior nostril 6 (8); barbel length 0 (0).

Size. Attains at least 50 cm TL (Marshall 1973).

Distribution. Widely distributed in tropical and temperate waters of the world's oceans except for the central and eastern Pacific. In Japan, known only from the Pacific off Tohoku at a midwater depth of 168–204 m (Appendix 3-11B). Generally confined to mesopelagic and bathypelagic depths (see also Nakayama *et al.* 2015b). Very rare in the northwestern Pacific.

Development. For a full description of juveniles see Maul & Koefoed (1950; original description of

Phalacromacrus pantherinus) and Endo (1997). The ontogeny of juveniles was also detailed by Marshall (1964).

Remarks. *Odontomacrus murrayi* was recently recorded from Japan by Nakayama *et al.* (2015b), who provided a full description of the Japanese specimen, with comments on the geographic and bathymetric distribution of this species.

Genus *Pseudocetonurus* Sazonov & Shcherbachev, 1982

[No Japanese name]

Pseudocetonurus Sazonov & Shcherbachev, 1982:712 (type

species: *Pseudocetonurus septifer* Sazonov & Shcherbachev, 1982, by original designation).

Diagnosis. Anus surrounded by narrow periproct, situated midway between anal-fin origin and pelvic-fin bases. Small dermal window of light organ anterior to periproct. Pelvic-fin rays 8–10. Infraorbital ridge not connected with preopercular ridge, separated by distinct gap. Second spinous ray of first dorsal fin smooth along its leading edge. Head large, with broad interorbital space (33–44% HL). Chin barbel present but small, length less than 10% HL. Teeth small, in narrow tapered band in upper jaw, uniserial in lower jaw. Head fully covered with small, unmodified scales. Body scales covered with long, erect, needle-like spinules in quincunx order; buttresses of scale spinules scarcely developed; no reticulate structures on scale surface. Grooved lateral line absent or present. Cephalic sensory canals notably broad, but lacking open pores. Pyloric caeca short, 22–34. Branchiostegal rays 7. Body uniformly dark. [Modified from Sazonov & Shcherbachev (1982) and Sazonov & Iwamoto (1992).]

Remarks. *Pseudocetonurus* was established by Sazonov & Shcherbachev (1982) for their new species *P. septifer* described only from the holotype collected from the Sala y Gomez Ridge in the southeastern Pacific. Sazonov & Iwamoto (1992) subsequently reported additional specimens of this species from near the type locality (Nazca Ridge) as well as from the Hawaiian Islands. Chiou *et al.* (2004a) newly recorded *P. septifer* from Taiwan based on a single specimen obtained at the Da-xi fish market (originally collected from the southwestern part of the East China Sea). However, their specimen was re-identified as *P. cf. septifer* by Iwamoto *et al.* (2015).

***Pseudocetonurus* sp. cf. *septifer* Sazonov & Shcherbachev, 1982**

[No Japanese name]

(Figs. 183–184; Appendix 3-11C)

Pseudocetonurus septifer (not Sazonov & Shcherbachev 1982): Chiou *et al.* 2004a:46, fig. 17, table 1 (brief description; 1 spec. from Da-xi; first record from Taiwan); Shao *et al.* 2008b: table 2 (1 spec. listed from northeastern Taiwan).

Pseudocetonurus cf. *septifer*: Iwamoto *et al.* 2015:97 (brief description; 1 spec. from northeastern Taiwan).

Diagnosis. A species of *Pseudocetonurus* with grooved lateral line on body and pale oral cavity.

Material examined. 1 specimen. **Taiwan:** ASIZP 61237 (1, 33.2 mm HL, 144+ mm TL), Da-xi fish market, Yilan, northeastern Taiwan, bottom trawl, coll. M.-L. Qiu, 1 May 2001.

Counts and measurements. Based on 1 specimen. Counts: first dorsal-fin rays II,9; pectoral-fin rays i18;

pelvic-fin rays 8; gill rakers on first arch (outer/inner) 12/18, on second arch 15/17, below first dorsal-fin midbase 6.5.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 28 (37); orbit diameter 29 (37); postorbital length 49 (64); postrostral length 77; orbit–preopercle distance 53 (69); suborbital width 19 (25); upper-jaw length 46 (61); length of rictus 39 (51); length of premaxillary tooth band 36 (46); preoral length 19 (25); distance between tip and lateral angle of snout 21 (28); snout width 34 (44); internasal width 34 (44); interorbital width 37 (48); body width over pectoral-fin bases 33 (44); pectoral-fin length 61 (80); predorsal length 103 (135); length of first dorsal-fin base 22 (28); interdorsal length 47 (61); length of gill slit 27 (36); length of posterior nostril 5 (7); barbel length 2 (3).

Size. To about 14 cm TL.

Distribution. So far known only from the Da-xi fish market in northern Taiwan. Capture depth unknown, but probably between 100 and 650 m (Appendix 3-11C). Very rare.

Remarks. This specimen was originally reported as *P. septifer* Sazonov & Shcherbachev, 1982 by Chiou *et al.* (2004a), being the first record of the species from Taiwan. However, Iwamoto *et al.* (2015:97) re-identified this specimen as *P. cf. septifer*, based on the presence of a grooved lateral line (vs. absent in *P. septifer*). It further differs from *P. septifer* in that the oral cavity is paler dorsally (vs. blackish). The Taiwanese specimen represents an undescribed species that will be formally described later by the present author.

Genus *Pseudonezumia* Okamura, 1970

[Japanese name: Nihon-sokodara-zoku]

Pseudonezumia Okamura 1970a:38 (type species: *Pseudonezumia japonicus* Okamura, 1970, by original designation).

Paracetonurus Marshall, 1973:615 (type species: *Macrourus parvipes* Smith & Radcliffe in Radcliffe, 1912, by original designation).

Diagnosis. Anus immediately anterior to anal-fin origin, surrounded by narrow but distinct periproct. Ventral light organ absent. Infraorbital ridge not connected with preopercular ridge, separated by distinct gap. Pelvic-fin rays usually 6 (rarely 5 or 7), fin origin distinctly anterior to vertical through pectoral-fin base. Second spinous ray of first dorsal fin weakly serrated along its leading edge. Head bones weak, but not very flexible. Mouth moderately small, posterior margin of upper jaw not reaching vertical through hind rim of orbit. Nasal fossa small. Chin barbel present. Teeth in narrow tapered bands in both jaw; none especially enlarged. Head completely scaled, but underside of snout narrowly to broadly naked; gular and branchiostegal membranes

FIGURE 183. Preserved specimen of *Pseudocetonurus* sp. cf. *septifer*. ASIZP 61237, 33.2 mm HL, 144+ mm TL, Da-xi fish market, northeastern Taiwan, East China Sea. (A) Lateral view; (B) dorsal and (C) ventral views of head. [Photos: N. Nakayama]

FIGURE 184. Scanning electron micrographs showing a body scale (from the dorsum below the interdorsal space) of *Pseudocetonurus* sp. cf. *septifer*. ASIZP 61237, 33.2 mm HL. (A) View from above; (B) oblique view. [Photos: NSMT]

occasionally with scaly patches. No modified scales along head ridges; tip and lateral angles of snout lacking prominent scutes. Body scales covered with long, erect, needle-like spinules in quincunx order; buttresses of scale spinules well developed; reticulate structure developed over entire surface of unexposed portion. Scales along second dorsal and anal fins not enlarged. Grooved lateral line interrupted, occurring as short segments. Cephalic sensory canals broad, without open pores. Branchiostegal rays usually 7, rarely 8. Body lacking prominent silvery reflections when fresh.

Remarks. The present author is preparing a revision of *Pseudonezumia*, with which *Paracetonurus* Marshall, 1973 will be synonymized. The genus currently includes

the following five described species: *P. cetonuropsis* (Gilbert & Hubbs, 1916) confined to Japan; *P. flagellicauda* (Koefoed, 1927) distributed in the northeastern Atlantic as well as the Madagascar Plateau in the southwestern Indian Ocean; *P. japonica* Okamura, 1970 known only from Japan (type species of *Pseudonezumia*); *P. parvipes* (Smith & Radcliffe in Radcliffe, 1912) known from the East Indies (type species of *Paracetonurus*); and *P. pusilla* (Sazonov & Shcherbachev, 1982) sporadically recorded from the Indo-West Pacific.

Key to species of *Pseudonezumia* from Japan and adjacent waters

- 1a Preoral length 26–29% HL; caudal depth at base of 40th anal-fin ray 24–29% HL.....*P. cetonuropsis*
1b Preoral length 20–26% HL; caudal depth at base of 40th anal-fin ray 27–34% HL.....*P. japonica*

Pseudonezumia cetonuropsis (Gilbert & Hubbs, 1916)

[Japanese name: Chidori-hige]

(Figs. 185–186, 187A–B; Appendix 3-11D)

Lionurus cetonuropsis Gilbert & Hubbs, 1916:202, pl. 11, fig. 1 [original description; holotype: USNM 76869, from “off east coast of Hondo” (off Cape Omaezaki), *Albatross* sta. 5084, in 918 fth (1690 m)]; Okada & Matsubara 1938:453 (in key; Japan); Matsubara 1955:1316 (in key; Japan); Matsubara 1965:509 (compiled; Japan; new Japanese name: “Chidori-hige”).

Nezumia cetonuropsis: Okamura 1970a:103 [new combination; description after Gilbert & Hubbs (1916)]; Okamura 1970b: table 1 (listed; Japan).

Paracetonurus cetonuropsis: Tominaga & Uyeno 1981:489 (listed; Japan).

Pseudonezumia cetonuropsis: Okamura 1984b:95, pl. 345, fig. D (compiled); Okamura 1988:95, pl. 345, fig. D (compiled); Nakabo 1993:356 (in key; Japan); Okamura 1997:128, fig. 18 (compiled); Nakabo 2000:420 (in key; Japan); Shinohara *et al.* 2001:306 (listed; Tosa Bay); Nakabo 2002:420 (in key; Japan); Nakabo & Kai 2013:496 (in key; Japan); Iwatsuki *et al.* 2017:32 (listed; Hyuga-nada); Motomura 2020:39 (listed; Japan).

Diagnosis. Underside of snout broadly naked above upper lip; spinules on body scales supported by narrowly developed buttresses; transverse scale rows below first dorsal-fin origin 9–11, below first dorsal-fin midbase 7–9; orbit diameter 22–25% HL; preoral length 26–29% HL; postorbital length 42–45% HL; orbit–preopercle distance 35–41% HL; interorbital width 31–37% HL; barbel length 5–11% HL; caudal depth at base of 40th anal-fin ray 24–29% HL; body uniformly dusky to paler brown, except for darker abdomen and black prepect.

Material examined. 16 specimens. **Holotype of *Lionurus cetonuropsis*:** USNM 76869 (43.8 mm HL), off Cape Omaezaki, Shizuoka Pref., Enshu-nada, Japan, 34.0000°N, 137.8278°E, 918 fth (1679 m), *Albatross* sta. 5084, 9-ft Agassiz beam trawl, 20 Oct. 1906. **Non-types: Japan:** BSKU 43443 (1, 39.4 mm HL, 213+ mm TL), southeast of Okino-shima Island, 32.5233°N, 132.4650°E, 1642–1651 m, R/V *Tansei-maru*, cr. KT-86-19, sta. D-1, tr. 1, 3-m beam trawl, 16 Dec. 1999; BSKU 86837 (1, 30.6 mm HL, 207+ mm TL), south-southwest of Okino-shima Island, 32.3038°N, 132.3592°E, 1678–1690 m, R/V *Tansei-maru*, cr. KT-99-18, sta. BT-5, beam trawl,

coll. H. Endo and S. Nagatomo, 16 Dec. 1999; BSKU 57408 (1, 50.8 mm HL, 316 mm TL), Bungo Channel, no further information; BSKU 86863 (1, 46.0 mm HL, 321+ mm TL), Kumano-nada, 33.7500°N, 136.5955°E, 2035–2043 m, R/V *Tansei-maru*, cr. KT-99-18, sta. BT-8, beam trawl, coll. H. Endo and S. Nagatomo, 20 Dec. 1999; BSKU 18802 (1, 44.8 mm HL, 293 mm TL), BSKU 18803 (1, 38.4 mm HL, 221+ mm TL), southeast of Owase, Kumano-nada, 33.8117°N, 136.5800°E, 2050 m, FRV *Soyo-maru*, sta. 18, beam trawl, 27 Jun. 1955; *BSKU 47908 (1, 54.9 mm HL, 345+ mm TL), Kumano-nada, 33.7085°N, 136.6092°E, 2034–2036 m, R/V *Tansei-maru*, cr. KT-09-13, sta. T1, 31 Aug. 1990; BSKU 86851 (1, 42.6 mm HL, 296 mm TL), Kumano-nada, 33.7332°N, 136.5788°E, 2035–2051 m, R/V *Tansei-maru*, cr. KT-99-18, sta. BT-7, beam trawl, coll. H. Endo and S. Nagatomo, 20 Dec. 1999; BSKU 86866 (1, 42.3 mm HL, 226+ mm TL), Kumano-nada, 33.8333°N, 136.5433°E, 1947–2023 m, R/V *Tansei-maru*, cr. KT-99-18, sta. BT-9, beam trawl, coll. H. Endo and S. Nagatomo, 20 Dec. 1999; BSKU 86847 (1, 40.6 mm HL, 292 mm TL), Kumano-nada, 33.6362°N, 136.5063°E, 2072–2074 m, R/V *Tansei-maru*, cr. KT-99-18, sta. BT-6, beam trawl, coll. H. Endo and S. Nagatomo, 20 Dec. 1999; *NSMT-P 78831 (2, 47.0–48.8 mm HL, 314–346 mm TL), Kumano-nada, 33.7650°N, 136.6967°E, 2019–2054 m, R/V *Tansei-maru*, cr. KT-86-06, sta. KN6, trawl, 28 May 1986; NSMT-P 19019 (1, 45.4 mm HL, 230+ mm TL), east of Hachijo Island, Shichito-Iojima Ridge, 33.1550°N, 140.7933°E, 1700 m, 25 Jul. 1975; NSMT-P 101338 (2, 49.0–50.3 mm HL, 298+–321+ mm TL), southeast of Okinawa-jima Island, Ryukyu Trench, 26.2840°N, 128.4505°E, 1943–2013 m, R/V *Hakuho-maru*, cr. KH-05-01, sta. PS-07, 4-m ORE beam trawl, 24 May 2005.

Counts and measurements. Based on 13 specimens (30.6–50.8 mm HL, 207+–321+ mm TL). Counts: first dorsal-fin rays II,8–10; pectoral-fin rays i15–i20; pelvic-fin rays 6; gill rakers on first arch (outer/inner) 8–11/10–11, on second arch 10/11; longitudinal scales 51–63; transverse scale rows below first dorsal-fin origin 9–11, below first dorsal-fin midbase 7–9, below second dorsal-fin origin 9–11.5.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 34–39 (50–60); orbit diameter 22–25 (34–39); postorbital length 42–45 (64–67); postrostral length 65–68; orbit–preopercle distance 35–41 (52–60); suborbital width 15–18 (23–28); upper-jaw length 27–31 (41–46); length of rictus 23–24 (33–36); length of premaxillary tooth band 13–17 (20–25); preoral length 26–29 (39–45); distance between tip and lateral angle of snout 21–22 (31–34); snout width 34–38 (50–56); internasal width 29–32 (42–48); interorbital width 31–37 (46–55); body width over pectoral-fin bases 38–54 (58–80); body depth at first dorsal-fin origin 67–80 (100–119); body depth at anal-fin origin 60–72 (89–106); caudal depth at base of 40th anal-fin ray 24–29 (35–42);

FIGURE 185. Fresh specimen of *Pseudonezumia cetonuropsis*. BSKU 47908, 54.9 mm HL, 345+ mm TL, Kumano-nada, Pacific, 2034–2036 m depth. Lateral view. [Photo: BSKU]

FIGURE 186. Holotype of *Lionurus cetonuropsis* (= *Pseudonezumia cetonuropsis*). USNM 76869, 43.8 mm HL, Enshu-nada off Cape Omaezaki, Shizuoka Pref., Pacific, 1679 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of head. Preserved condition. [Photos: N. Nakayama]

prepelvic length 100–117 (150–174); preanus length 123–140 (183–208); preanal length 127–147 (190–218); isthmus–pelvic distance 23–27 (34–41); isthmus–anus distance 46–57 (69–84); isthmus–anal distance 53–62 (79–93); pelvic–anal distance 29–42 (44–62); anus–anal distance 5–11 (7–17); pelvic-fin length 28–47 (41–70); pectoral-fin length 47–56 (72–85); predorsal length 105–115 (156–170); height of first dorsal fin 65–78 (96–116); length of first dorsal-fin base 19–23 (28–34); interdorsal length 32–42 (48–63); length of gill slit 12–15 (18–23); length of posterior nostril 5–10 (8–15); barbel length 5–11 (8–16).

Size. To about 32 cm TL (BSKU 86863, 321+ mm TL, Kumano-nada, Japan).

Distribution. So far known from only Japanese waters (Appendix 3-11D). Depth range 1642–2074 m.

Remarks. For further morphological details see the

original description of the species given by Gilbert & Hubbs (1916). *Pseudonezumia cetonuropsis* was initially described as a new species of *Lionurus* Günther, 1887 based on two specimens collected from off Cape Omaezaki, Enshu-nada, Japan (Fig. 186). Gilbert & Hubbs (1916) also reported one damaged juvenile collected with the holotype; this second specimen was not found during the author's visits to the Smithsonian Institution in 2011 and 2017.

Maruyama (1971) listed *P. cetonuropsis* from Iwate Pref., northern Honshu (as *Lionurus cetonuropsis*), but with a question mark. Because his list was primarily based on specimens collected from local fish markets, this record appears to represent a different species of grenadiers. The bathymetric distribution of *P. cetonuropsis* is confined to bathyal depths below 1600 m, where commercial fisheries have not been conducted.

Shao *et al.* (2008a, 2008b) recorded *P. cetonuropsis*

FIGURE 187. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of two species of *Pseudonezumia*. (A–B) *P. cetonuropsis*, NSMT-P 78831, 48.8 mm HL; (C–D) *P. japonica*, BSKU 56976, 50.5 mm HL. (A, C) Views from above; (B, D) oblique views. [Photos: N. Nakayama]

from Taiwan based on three specimens collected from the northern South China Sea (ASIZP 66945, 20.5–32.9 mm HL, 135+–161+ mm TL). These specimens were re-identified as *P. pusilla* (Sazonov & Shcherbachev, 1982) by Iwamoto *et al.* (2015).

Relationships and comparisons. *Pseudonezumia cetonuropsis* is unlikely to be confused with *P. flagellicauda* (Koefoed, 1927) owing to its uniformly dark color (vs. lower jaw and naked area on underside of snout prominently black, and tail abruptly blackish posteriorly), shorter postorbital (42–45% HL vs. 47–53%), narrower cheek (orbit–preopercle distance 35–41% HL vs. 40–48%), and lower counts of transverse scale rows (9–11 vs. 11–15 below first dorsal-fin origin; 7–9 vs. 10.5–13.5 below first dorsal-fin midbase). In addition, this species is readily distinguished from *P. parvipes* (Smith & Radcliffe in Radcliffe, 1912) and *P. pusilla* by having a broad naked area on the underside of the snout (vs. naked area absent or represented by only a narrow ventromedial streak above the upper lip). It further differs from *P. flagellicauda* and

P. pusilla in having a shorter chin barbel (5–11% HL vs. 13–18% in *P. flagellicauda*, 11–23% in *P. pusilla*), and from *P. pusilla* in having a smaller orbit (22–25% HL vs. 28–37%).

Differences between *P. cetonuropsis* and *P. japonica* Okamura, 1970 are subtle, and the species are difficult to distinguish from each other. The possible diagnostic character recognized in this study is the preoral length: 26–29% HL in *P. cetonuropsis* vs. 20–26% in *P. japonica*. Accordingly, the snout of *P. cetonuropsis* is usually much higher than that of *P. japonica*, if specimens are in good condition (Fig. 185 vs. Fig. 188). A slight difference in the caudal depth also suggests a separation between the two species, with *P. cetonuropsis* having a slenderer tail compared with *P. japonica* (depth at base of 40th anal-fin ray 24–29% HL vs. 27–34%). However, these differences are very slight, and may be eliminated if more specimens become available for comparison; the taxonomic status of the two taxa requires further investigation.

***Pseudonezumia japonica* Okamura, 1970**

[Japanese name: Nihon-sokodara]

(Figs. 187C–D, 188–189; Appendix 3-11E)

Pseudonezumia japonicus Okamura, 1970a:39, pl. XIII, text-fig. 20 [original description; holotype: BSKU 14203, from “off Choshi”, Chiba Pref., 36°19'08"N, 141°14'00"E (off Mito, Ibaraki Pref.), in 1690 m; new Japanese name: “Nihon-sokodara”].

Pseudonezumia japonica: Okamura 1970b: table 1 (listed; Japan); Tominaga & Uyeno 1981:489 (listed; Japan); Okamura 1984b:95, pl. 81, fig. K (compiled); Okamura 1988:95, pl. 81, fig. K (compiled); Nakabo 1993:356 (in key; Japan); Nakabo 2000:420 (in key; Japan); Nakabo 2002:420 (in key; Japan);

Nakabo & Kai 2013:496 (in key; Japan); Motomura 2020:39 (listed; Japan).

Diagnosis. Underside of snout broadly naked above upper lip; spinules on body scales supported by narrowly developed buttresses; transverse scale rows below first dorsal-fin origin 8.5–12.5, below first dorsal-fin midbase 7–12; orbit diameter 22–28% HL; preoral length 20–26% HL; postorbital length 42–48% HL; orbit–preopercle distance 34–43% HL; interorbital width 29–36% HL; barbel length 8–13% HL; caudal depth at base of 40th anal-fin ray 27–34% HL; body uniformly dusky to paler brown, except for darker abdomen and black periproct.

Material examined. 31 specimens. **Holotype of**

FIGURE 188. Fresh specimen of *Pseudonezumia japonica*. BSKU 56976, 50.5 mm HL, 273+ mm TL, off Kamaishi, Iwate Pref., Pacific, 1949–2005 m depth. Lateral view. [Photo: H. Endo]

FIGURE 189. Holotype of *Pseudonezumia japonicus* (= *P. japonica*). BSKU 14203, 46.0 mm HL, 303+ mm TL, off Mito, Ibaraki Pref., Pacific, 1690 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

Pseudonezumia japonicus: BSKU 14203 (46.0 mm HL, 303 mm TL), off Mito, Ibaraki Pref., Japan, 36.3189°N, 141.2333°E, 1690 m, 3 Dec. 1967. **Non-types: Japan**: BSKU 56944 (1, 47.2 mm HL, 325 mm TL), BSKU 56945 (1, 43.6 mm HL, 258+ mm TL), BSKU 56946 (1, 46.7 mm HL, 290+ mm TL), BSKU 56947 (1, 44.9 mm HL, 280+ mm TL), BSKU 56948 (1, 48.7 mm HL, 274+ mm TL), BSKU 56949 (1, 38+ mm HL, 274+ mm TL), BSKU 56950 (1, 48.8 mm HL, 302+ mm TL), BSKU 56951 (1, 42.6 mm HL, 306 mm TL), BSKU 56952 (1, ca. 39 mm HL, 253+ mm TL), BSKU 56953 (1, 43.8 mm HL, 214+ mm TL), BSKU 56954 (1, 39.9 mm HL, 287+ mm TL), BSKU 56955 (1, 36.5 mm HL, 244+ mm TL), BSKU 56956 (1, 43.5 mm HL, 285+ mm TL), BSKU 56957 (1, 47.3 mm HL, 289+ mm TL), BSKU 56958 (1, 49.5 mm HL, 332+ mm TL), BSKU 56959 (1, 48.5 mm HL, 286+ mm TL), BSKU 56960 (1, 47.0 mm HL, 224+ mm TL), BSKU 56961 (1, 44.4 mm HL, 307+ mm TL), BSKU 56963 (1, 47.8 mm HL, 331+ mm TL), BSKU 56964 (1, 47.4 mm HL, 215+ mm TL), BSKU 56965 (1, ca. 45 mm HL, 288+ mm TL), BSKU 56966 (1, 43.8 mm HL, 273+ mm TL), BSKU 56967 (1, 43.1 mm HL, 297 mm TL), BSKU 56969 (1, 40.5 mm HL, 272 mm TL), BSKU 56971 (1, ca. 36 mm HL, 218+ mm TL), BSKU 56972 (1, 32.9 mm HL, 165+ mm TL), BSKU 56973 (1, ca. 36 mm HL, 231+ mm TL), BSKU 56974 (1, 41.7 mm HL, 270+ mm TL), BSKU 56975 (1, ca. 43 mm HL, 195+ mm TL), off Hitachi, 36.4205°N, 141.5420°E, 1845–1948 m, FRV *Wakataka-maru*, sta. 62, coll. H. Endo, 10 Nov. 1995; BSKU 56976 (1, 50.5 mm HL, 273+ mm TL), off Kamaishi, 39.3797°N, 143.1282°E, 1949–2005 m, FRV *Wakataka-maru*, sta. 59, coll. H. Endo, 7 Nov. 1995.

Counts and measurements. Based on 31 specimens (31.9–50.5 mm HL, 165+–332+ mm TL). Counts: first dorsal-fin rays II,8–10; pectoral-fin rays i14–i19; pelvic-fin rays 6; longitudinal scales 49–61; transverse scale rows below first dorsal-fin origin 8.5–12.5, below first dorsal-fin midbase 7–12, below second dorsal-fin origin 9–12; pyloric caeca 8–10.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 33–38 (48–58); orbit diameter 22–28 (31–40); postorbital length 42–48 (63–69); postrostral length 66–71; orbit–preopercle distance 34–43 (51–61); suborbital width 15–19 (24–28); upper-jaw length 25–33 (37–49); preoral length 20–26 (29–39); snout width 32–40 (47–58); internasal width 26–34 (38–49); interorbital width 29–36 (43–54); body width over pectoral-fin bases 30–56 (45–82); body depth at first dorsal-fin origin 63–89 (101–130); body depth at anal-fin origin 58–76 (86–112); caudal depth at base of 40th anal-fin ray 27–34 (40–49); prepelvic length 93–108 (134–164); preanus length 120–135 (170–198); preanal length 128–142 (182–209); isthmus–pelvic distance 23–31 (33–46); isthmus–anus distance 49–62 (69–90); isthmus–anal distance 57–71 (80–104); pelvic–anal distance 34–44 (50–66); anus–anal distance 5–12 (8–18); pelvic-fin

length 28–54 (40–81); pectoral-fin length 32–64 (47–91); predorsal length 103–119 (158–174); height of first dorsal fin 71–83 (102–120); length of first dorsal-fin base 19–27 (29–39); interdorsal length 33–48 (48–); length of gill slit 12–17 (19–24); length of posterior nostril 6–11 (9–17); barbel length 8–13 (12–19).

Size. To about 33 cm TL (BSKU 56958, 332+ mm TL, off Hitachi, Ibaraki Pref., Japan).

Distribution. Restricted to Japan (Appendix 3-11E). Depth range 1453–2005 m.

Remarks. For further morphological information see the original description given by Okamura (1970a). *Pseudonezumia japonica* was described from a single specimen collected from off Mito, Ibaraki Pref., Japan (Fig. 189), as a new genus and species of grenadiers. Okamura (1970a) considered this species to be quite different from other grenadiers in having eight branchiostegal rays (vs. six or seven). However, an examination of additional specimens from the type locality showed that the number is usually seven, and only rarely eight.

Relationships and comparisons. *Pseudonezumia japonica* is closely similar to *P. cetonuropsis* (Gilbert & Hubbs, 1916) also known from Japan. These two species are tentatively regarded as valid in this study, but a further investigation based on additional material may arrive at a different conclusion, namely, *P. japonica* is synonymized with *P. cetonuropsis* (see the Relationships and comparisons of the latter species).

Genus *Sphagemacrus* Fowler, 1925

[No Japanese name]

Sphagemacrus Fowler, 1925:3 (as subgenus of *Macruroplus*; type species: *Macrurus hirundo* Collett, 1986, by original designation).

Grenurus Parr, 1946:46 (type species: *Grenurus grenadae* Parr, 1946, by original designation).

Diagnosis. Anus closer to anal-fin origin than to pelvic-fin bases. Periproct broadly developed, extending about 2/3 of distance between anal-fin origin and pelvic-fin base. Small dermal window of light organ immediately anterior to anus, its anterior margin well posterior to pelvic-fin bases. Infraorbital ridge not connected with preopercular ridge, separated by distinct gap. Second spinous ray of first dorsal fin serrated along its leading edge. Ventral aspects of body positioned forward so that anal-fin origin distinctly anterior to hind margin of first dorsal-fin base. Snout scarcely protruding beyond upper jaw, its ventral profile almost vertical. Chin barbel present. Teeth in bands in both jaws; premaxillary band ending anterior to posterior edge of maxillary process. Tip and lateral angles of snout tipped with prominent scutes; suborbital shelf formed of stout, coarsely modified scales. Body scales covered with needle-like spinules in parallel to slightly

divergent rows; buttresses of spinules rudimentary; reticulate structure broadly developed. Underside of snout variously naked. Grooved lateral line present. Pyloric caeca usually less than 30. Branchiostegal rays 7. Body generally dark, lacking silvery reflection when fresh. [Modified from Iwamoto & Merrett (1997) and Iwamoto & Graham (2001).]

Remarks. *Sphagemacrurus* includes six nominal species, all of which are currently treated as valid (Iwamoto 1990), viz., *S. decimalis* (Gilbert & Hubbs, 1920), *S. gibber* (Gilbert & Cramer, 1897), *S. grenadae* (Parr, 1946), *S. hirundo* (Collett, 1896), *S. pumiliceps* (Alcock, 1894), and *S. richardi* (Weber, 1913). Citing a personal communication from Yuri Y. Sazonov (ZMMGU), Iwamoto & Merrett (1997:548–549) suggested that half of these species are of dubious validity. They also implied that the *S. pumiliceps* specimens they reported from the New Caledonian region may represent another species or include two different taxa. This elusive genus needs a through revision (Iwamoto & Merrett 1997; Iwamoto *et al.* 2015).

Sphagemacrurus pumiliceps (Alcock, 1894)

[No Japanese name]

(Fig. 190; Appendix 3-11F)

Macrurus pumiliceps Alcock, 1894:125 [original description; 2 syntypes ZSI F13561–F13562, from Laccadive Sea (south of India); 7°05'45"N, 75°04'E, *Investigator* sta. 150, 719 ftn (1315 m)].

Sphagemacrurus pumiliceps: Shao *et al.* 2008b: table 2 (7 spec. listed from eastern Taiwan and South China Sea); Iwamoto *et al.* 2015:100 (brief description; 6 spec. from northeastern Taiwan and South China Sea).

Diagnosis. Orbit diameter longer than snout length, about 1.0–1.2 times into orbit–preopercle distance, 33–37% HL; underside of snout mostly covered with small, irregularly shaped scales; pale, naked area anterior to margin of upper lip; scales mostly deciduous except on nape and head; teeth small, slender, with curved tips, 4–5 rows at symphysis of premaxillary, 2–3 rows at symphysis of mandible; pelvic-fin rays 11–13; total inner gill rakers on first arch 8–11; body color variously pale to dusky brown on nape and flanks, darker grayish brown on head. [From Iwamoto & Williams (1999).]

Counts and measurements. For selected counts and measurements see Iwamoto *et al.* (2015:100).

Size. To about 26 cm TL (Iwamoto & Williams 1999).

Distribution. Widely distributed in the Indo-West Pacific at depths of 732–1880 m (Iwamoto & Williams 1999; Iwamoto *et al.* 2015). Not known from Japanese waters (Appendix 3-11F), but recorded from the East China Sea off northeastern Taiwan.

Remarks. For a description and comparisons with congeners see Gilbert & Hubbs (1920; as *Lionurus pumiliceps*), Weber & de Beaufort (1929; as *L. pumiliceps*), Iwamoto & Williams (1999), and Iwamoto *et al.* (2015).

Genus *Spicomacrurus* Okamura, 1970

[Japanese name: Yaridara-zoku]

Spicomacrurus Okamura, 1970a:63 (subgenus of *Hymenogadus*; type species: *Hymenocephalus kuronumai* Kamohara, 1938, by original designation).

Diagnosis. Anus immediately anterior to anal-fin origin,

FIGURE 190. Drawing of a syntype of *Macrurus pumiliceps* (= *Sphagemacrurus pumiliceps*). [Reproduced from Alcock 1895: pl. XVI, fig. 3]

surrounded by narrow periproct. Ventral striae, consisting of fine, parallel black lines over silvery layer on each side of isthmus, shoulder girdle, most of chest, and posterior to pelvic fins. Gular region covered with netlike pattern of black epithelium. Light organ developed as long tube, extending from anus to slightly anterior to pelvic-fin bases; small lens at each end of light organ. Second spinous ray of first dorsal fin smooth along its leading edge. Pelvic-fin rays 8–10, outer ray greatly elongate and flattened with narrow ribbon-like flange on free distal portion. Teeth all small, in narrow villiform bands in both jaws. Trunk cylindrical or weakly compressed. Head bones not fragile. Nasal bones stout, forming dorsoventrally flattened, lateral and medial processes. Chin barbel present. Body scales large, highly deciduous, covered with needle-like spinules; radial grooves developed anteriorly; reticulate structure absent. Branchiostegal rays 7. All species small, less than 25 cm TL. [From Iwamoto *et al.* (2011) and Schwarzhans (2014).]

Remarks. *Spicomacrurus* was originally established by Okamura (1970a) as a monotypic subgenus of *Hymenocephalus* Giglioli, 1884, with the type species being *S. kuronumai* (Kamohara, 1938). Iwamoto *et al.* (2011) elevated this group to full generic rank after their discovery of two new species from the southwestern Pacific. In addition to *S. kuronumai*, the genus includes the following three species known only from the Southern Hemisphere: *S. adelscotti* Iwamoto & Merrett, 1997 recorded from Fiji (type locality) and Western Australia; *S. dictyogadus* Iwamoto, Shao & Ho, 2011 represented by only the holotype collected from Vanuatu; and *S. mccoskeri* Iwamoto, Shao & Ho, 2011 so far known from the Chesterfield Plateau in the Coral Sea.

Spicomacrurus kuronumai (Kamohara, 1938)

[Japanese name: Yari-dara]

(Figs. 191–192; Appendix 3-12A)

Hymenocephalus kuronumai Kamohara, 1938:70, fig. 40 (original description; holotype from Mimase fish market; new Japanese name: “Yari-dara”); Kamohara 1950:278 (listed; Kochi and Wakayama Pref.); Matsubara *et al.* 1951:42 (listed; Mie Pref.); Kamohara 1952:97, fig. 95 (description; 1 spec. from Mimase fish market); Matsubara 1955:1315 (in key; Japan); Kamohara 1958:73 (listed; Kochi Pref.); Okada *et al.* 1959:83 (listed; Kumano-nada); Kamohara 1961c:9, pl. VII, fig. 1 (invalid designation of neotype; BSKU 4333 from Mimase); Kamohara 1964:96 (listed; Kochi Pref.); Shinohara *et al.* 2001:305 (1 spec. listed from Tosa Bay); Shao *et al.* 2008b: table 2 (1 spec. listed from northeastern Taiwan; first Taiwanese record).

Hymenogadus kuronumai: Okamura 1970a:64, pl. III, text-fig. 28 (new combination; description; biological notes; 79 spec. from Pacific off southern Japan from Miya to Shibushi); Okamura 1970b: table 1 (listed; Japan); Kataoka & Tomida 1981:78

(listed; Mie Pref.); Tominaga & Uyeno 1981:489 (listed; Japan); Ozawa 1983:13 (listed; off Makurazaki, Kagoshima Pref., East China Sea); Okamura 1984a:203, 359, fig. 144 [brief description; 6 spec. from Okinawa Trough; photo based on BSKU 40016 (from off Kochi Pref.)]; Okamura 1984b:93, pl. 80, fig. I (compiled); Okamura 1988:93, pl. 80, fig. I (compiled); Nakabo 1993:354 (in key; Japan); Okamura 1997:128, fig. 5 (compiled); Shinohara & Matsuura 1997:292, pl. 2-C (listed; Suruga Bay); Suzuki & Kataoka 1997:81, pl. 32, fig. 174 (brief description; 1 spec. from off Owase); Nakabo 2000:418 (in key; Japan); Nakabo 2002:418 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Nakajima 2003:52 (brief description; 1 spec. from Enshu-nada); Shinohara *et al.* 2005:418 (listed; Ryukyu Islands); Furuhashi *et al.* 2010: table 2 (20 spec. listed from northern Okinawa Trough).

Spicomacrurus kuronumai: Iwamoto *et al.* 2011:518 (in key; elevation of *Spicomacrurus* to full generic status); Nakabo & Kai 2013:494 (in key; Japan); Ikeda & Nakabo 2015:319, pl. 64, figs. 5–10 (brief description; spec. from Wakayama Pref.); Iwamoto *et al.* 2015:103 (brief description; 2 spec. from northeastern Taiwan); Motomura 2020:39 (listed; Japan).

Diagnosis. A species of *Spicomacrurus* with 8–9 (usually 8) pelvic-fin rays; nasal bones closely adjoined mesially, forming 3 broad horizontally flattened processes; trunk shallow, cylindrical in cross section, width over pectoral-fin bases 0.9–1.3 in depth below first dorsal-fin origin; snout short, low, but distinctly protruding beyond upper jaw, length 27–31% HL; interorbital width 16–23% HL; upper-jaw length 37–41% HL; barbel short, but well developed, length 11–17% HL; pelvic-fin length 47–69% HL, distal portion of outer pelvic-fin ray distinctly expanded; body scales large, transverse rows below midbase of first dorsal fin 2.5–3.5, below second dorsal-fin origin 2–3; longitudinal scales over distance equal to predorsal length 17–22.

Material examined. 30 specimens. **Japan:** BSKU 4333 [invalid neotype of *Hymenocephalus kuronumai* designated by Kamohara (1961); 25.0 mm HL, 116+ mm TL], Mimase fish market, Kochi Pref., Japan, coll. T. Kamohara, 30 Nov. 1954; BSKU 109035 (1, 35.9 mm HL, 182+ mm TL), west of Sanpo-sone, 32.872°N, 129.0732°E, 351–389 m, T/V *Nagasaki-maru*, cr. N365, sta. B4, 3-m ORE beam trawl, coll. N. Nakayama, 19 Nov. 2012; BSKU 13356 (1, 36.6 mm HL, 166+ mm TL), BSKU 13358 (1, 39.1 mm HL, 177+ mm TL), BSKU 13360 (1, 29.7 mm HL, 132+ mm TL), BSKU 13363 (1, 35.7 mm HL, 195+ mm TL), BSKU 13366 (1, 33.2 mm HL, 167+ mm TL), BSKU 13369 (1, 36.1 mm HL, 180+ mm TL), BSKU 13373 (1, 32.7 mm HL, 161+ mm TL), BSKU 13375 (1, 35.8 mm HL, 169+ mm TL), BSKU 13376 (1, 37.0 mm HL, 183+ mm TL), BSKU 13377 (1, 37.8 mm HL, 159+ mm TL), BSKU 13378 (1, 29.6 mm HL, 139+ mm TL), BSKU 13379 (1, 33.5 mm HL, 157+ mm TL), BSKU 13380 (1, 34.4 mm HL, 175+ mm TL),

FIGURE 191. Fresh specimen of *Spicomacrurus kuronumai*. BSKU 109035, 35.9 mm HL, 182+ mm TL, west of Sanpo-sone, Okinawa Trough, East China Sea, 351–389 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

FIGURE 192. Scanning electron micrographs showing body scales of *Spicomacrurus kuronumai*. BSKU 13375, 35.8 mm HL. (A–B) Scales from the dorsum below the interdorsal space; (C–D) from the chest. (A, C) Views from above; (B, D) oblique views. [Photos: NSMT]

BSKU 13381 (1, 40.2 mm HL, 201+ mm TL), BSKU 13383 (1, 40.0 mm HL, 193+ mm TL), BSKU 13385 (1, 37.5 mm HL, 155+ mm TL), BSKU 13387 (1, 22.8 mm HL, 117+ mm TL), BSKU 13389 (1, 35.3 mm HL, 170+ mm TL), west of Muroto, Tosa Bay, 360–435 m, coll. O. Okamura, 8 Jun. 1968; BSKU 12946 (1, 22.4 mm HL, 106+ mm TL), Tosa Bay, 350–450 m, bottom trawl, 14 May 1968; BSKU 4365 (1, 24.3 mm HL, 126+ mm TL), Mimase fish market, coll. T. Kamohara, 5 Dec. 1954; KSHS 297 (1, 23.2 mm HL, 120+ mm TL), KSHS 298 (1, 23.8 mm HL, 108+ mm TL), Mimase fish market, coll. T. Yamakawa, 4 Dec. 1960; BSKU 40016 (1, 33.9 mm HL, 156+ mm TL), off Okino-shima Island, Kochi, 250 m, bottom trawl, 18 Apr. 1984; BSKU 13117 (1, 26.8 mm HL, 149+ mm TL), BSKU 13235 (1, 34.1 mm HL, 166+ mm TL), BSKU 13259 (1, 34.6 mm HL, 162+ mm TL), BSKU 13260 (1, 33.8 mm HL, 157+ mm TL), BSKU 13261 (1, 33.5 mm HL, 173+ mm TL), east of Cape Ashizurimisaki, Tosa Bay, 420–555 m, coll. Y. Mishima (= Y. Machida), 2 Jun. 1968.

Counts and measurements. Based on 30 specimens (22.4–40.2 mm HL, 106+–201+ mm TL). Counts: first dorsal-fin rays II, 10–12; pectoral-fin rays i17–i21; pelvic-fin rays 8–9; gill rakers on first arch (outer/inner) 6–10/11–15, on second arch 11–17/10–15; longitudinal scales 17–22; transverse scale rows below first dorsal-fin origin 3–4.5, below first dorsal-fin midbase 2.5–3.5, below second dorsal-fin origin 2–3, above anal-fin origin 7.5–9.5.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 27–31 (34–43); orbit diameter 27–37 (38–48); postorbital length 42–47 (54–65); postrostral length 72–78; orbit–preopercle distance 32–36 (43–50); suborbital width 8–11 (10–14); upper-jaw length 37–41 (49–56); length of rictus 33–37 (43–50); length of premaxillary tooth band 27–32 (35–43); preoral length 16–19 (20–26); distance between tip and lateral angle of snout 18–21 (23–29); snout width 29–35 (39–46); internasal width 17–20 (23–26); interorbital width 16–23 (21–31); occipital width 7–9 (9–13); body width over pectoral-fin bases 39–62 (52–84); body depth at first dorsal-fin origin 49–64 (65–88); body depth at anal-fin origin 42–55 (55–75); prepelvic length 93–106 (121–145); preanal length 152–173 (198–234); isthmus–pelvic distance 44–50 (58–69); isthmus–anal distance 100–118 (130–159); pelvic–anal distance 56–72 (74–98); pelvic-fin length 47–69 (63–95); pectoral-fin length 56–68 (76–93); predorsal length 96–107 (127–146); height of first dorsal fin 63–79 (86–107); length of first dorsal-fin base 29–38 (40–52); interdorsal length 62–86 (82–119); length of gill slit 19–25 (25–34); length of posterior nostril 4–8 (5–11); barbel length 11–17 (15–23).

Size. To about 21 cm TL (FAKU 24937, 211+ mm TL, Owase, Japan).

Distribution. Restricted to Japan and Taiwan. Known from off the Pacific coasts of southern Japan northward to

the Izu Peninsula (138.96°E), northern Okinawa Trough, and northern Taiwan, at depths of 205–680 m (Shao *et al.* 2008b; this study; Appendix 3-12A). Usually rare, but many individuals occasionally collected by bottom trawls.

Comments on type specimen. *Spicomacrurus kuronumai* was originally described as a new species of *Hymenocephalus* Giglioli, 1884 based on a single specimen collected from the Mimase fish market, Kochi, Japan. Unfortunately, the holotype was destroyed by a fire during WWII. In his list of type specimens deposited in BSKU, Kamohara (1961c) designated a neotype of this species (BSKU 4333, 25.0 mm HL, 116+ mm TL). However, his designation is considered to be invalid, according to Art. 75.2 of the ICZN (1999).

Remarks. For a full description see Okamura (1970a; as a species of *Hymenogadus* Gilbert & Hubbs, 1920). Although Iwamoto & Merrett (1997) reported *S. kuronumai* (as a species of *Hymenocephalus*) from the Coral Sea, their specimens currently comprise the type series of *S. mccoskeri* Iwamoto, Shao & Ho, 2011. Last *et al.*'s (2014) record of *S. kuronumai* from the western Coral Sea also seems to be referred to as *S. mccoskeri*.

Comparisons. Among the other three congeners (see the Remarks of the genus), *S. kuronumai* is most similar to *S. adelscotti* Iwamoto & Merrett, 1997 known from Fiji (type locality) and Western Australia. The two species readily differ from *C. dictyogadus* Iwamoto, Shao & Ho, 2011 and *S. mccoskeri* in that the nasal bones are closely adjoined mesially (vs. broadly separated by the rostral cartilage), and by having a wider interorbital space (16–23% HL vs. 8–15%) and a smaller mouth (upper-jaw length 37–42% HL vs. 46–50%) [data for *S. dictyogadus* and *S. mccoskeri* are from Iwamoto *et al.* (2011)]. According to the key to species of *Spicomacrurus* given by Iwamoto *et al.* (2011:518), *S. kuronumai* differs from *S. adelscotti* in having eight pelvic-fin rays (vs. nine), with its outer ray distinctly flattened and expanded distally (vs. only slightly flattened and expanded). However, the *S. kuronumai* specimens examined rarely had eight pelvic-fin rays, and the outer ray was not flattened nor expanded distally, when its fleshy flange was completely missing (see Iwamoto *et al.* 2011: fig. 1B vs. 1D). Nevertheless, *S. kuronumai* readily differs from that species in its longer barbel (11–17% HL vs. 9%) and higher counts of outer gill rakers (those on the lower limb of the second arch 9–14 vs. 7). According to Schwarzhans (2014:20, fig. 4), the two species are also separable by otolith morphology: the otolith of *S. kuronumai* is not strongly depressed dorsoventrally, with its height being 1.45–1.6 in its longitudinal length (vs. 1.75).

Genus *Trachonurus* Günther, 1887

[Japanese name: Watahige-zoku]

Trachonurus Günther, 1887:124 (subgenus of *Macrurus*; type species: *Coryphaenoides villosus* Günther, 1887, by original designation).

Diagnosis. Anus surrounded by broad, extensive periproct that extending most of distance between anal-fin origin and pelvic-fin bases; periproct abutting anal-fin origin. Infraorbital ridge not connected with preopercular ridge, separated by distinct gap. Pelvic fin with 6–7 soft rays, its origin distinctly posterior to vertical through pectoral-fin base (usually below midbase of first dorsal fin). Second spinous ray of first dorsal fin slender, lacking serrations along its leading edge. Chin barbel present. Teeth in narrow band in upper jaw, with outer series usually enlarged; lower jaw teeth in 2–3 rows or narrow band. Scales not deciduous, closely adhered to body surfaces. Body scales adherent, covered with long, erect, needle-like spinules in quincunx pattern; buttresses well developed; reticulate pattern developed over entire scale surface. Scales along anterior parts of second dorsal and anal fins enlarged. Grooved lateral line present or absent. Head almost fully scaled; gular and branchiostegal membranes heavily scaled in most species; no scutes on head ridges and tip and lateral angles of snout. Cephalic sensory canals lacking open pores. Pyloric caeca short and thick, usually less than 15. Branchiostegal rays 7. Body uniformly dark, lacking prominent silvery reflection. [Modified from Sazonov & Iwamoto (1992) and Iwamoto & Graham (2001).]

Remarks. Although *Trachonurus* is a distinctive genus of the family, there is much discussion on its species-level classification. It is still uncertain how many species are included in this group, and there appears to be some species still remaining to be described (Sazonov & Iwamoto 1992; Iwamoto 1997; Iwamoto & McMillan 1997; Iwamoto & Merrett 1997; Iwamoto & Williams 1999). The following six species are currently regarded as valid, one of which is known from Japanese waters: *T. gages* Iwamoto & McMillan, 1997 distributed in Australia and New Zealand; *T. robinsi* Iwamoto, 1997 probably endemic to the Philippines; *T. sentipellis* Gilbert & Cramer, 1897 recorded from the Hawaiian Islands, Australia, the New Caledonian region, and Taiwan; *T. sulcatus* (Goode & Bean, 1885), a senior synonym of *T. asperimus* (Vaillant, 1888), widely distributed in the Atlantic; *T. villosus* (Günther, 1877) (type species) probably confined to the northwestern Pacific from Japan to the Philippines (see the Remarks of this species); and *T. yiwardaus* Iwamoto & Williams, 1999 so far known from the west and south coasts of Australia.

Trachonurus villosus (Günther, 1877)

[Japanese name: Wata-hige]

(Figs. 193–195; Appendix 3–12B)

Coryphaenoides villosus Günther, 1877:441 [original description; holotype: BMNH 1887.12.7.105, from “south of Yeddo” (off Eno-shima Island), *Challenger* sta. 232, in 345 ftn (631 m)].

Macrurus villosus: Günther 1887:142–143, pl. XXXVI, fig. B [description; in part, 1 spec., BMNH 1887.12.7.105 (holotype); remaining 1 spec. probably represents different species (fide Iwamoto 1997:945)].

Trachonurus villosus: Jordan & Snyder 1901:120 (listed; Japan); Jordan & Gilbert in Jordan & Starks 1904:621 [after Günther (1877, 1887)]; Jordan *et al.* 1913:420 (listed; Japan; new Japanese name: “Wata-hige”); Gilbert & Hubbs 1916:205 (description; 4 spec. from off Cape Shionomisaki and south-southwest of Izu Island); Kamohara 1931b:544 (questionable; listed; Kochi); Kamohara 1934a:54 (questionable; listed; Kochi); Okada & Matsubara 1938:453 (in key; Japan); Kamohara 1938:73 (questionable; spec. from Kochi Pref.); Kamohara 1950:279 (questionable; listed; Kochi Pref.); Kamohara 1952:99 (questionable; spec. from Kochi Pref.); Matsubara 1955:1317 (in key; Japan); Kamohara 1958:74 (questionable; listed; Kochi Pref.); Kamohara 1964:96 (questionable; listed; Kochi Pref.); Okamura 1970a:106, pl. XXIII, text-fig. 44 (description; 1 spec. from East China Sea); Okamura 1970b: table 1 (listed; Japan); Tominaga & Uyeno 1981:489 (listed; Japan); Okamura 1984b:94, pl. 81, fig. L (compiled); Okamura 1988:94, pl. 81, fig. L (compiled); Sazonov & Iwamoto 1992:77 (unnecessary lectotype designation; synonymy; comments on *Challenger* spec.; 2 additional spec. from Sala y Gomez Ridge); Nakabo 1993:356 (in key; Japan); Okamura 1997:128, fig. 7 (compiled); Nakabo 2000:420 (in key; Japan); Shinohara *et al.* 2001:306 (listed; Tosa Bay); Nakabo 2002:420 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Shinohara *et al.* 2005:418 (listed; Ryukyu Islands); Shao *et al.* 2008b: table 2 (5 spec. listed from South China Sea); Nakabo & Kai 2013:496 (in key; Japan); Iwamoto *et al.* 2015:104, fig. 26 (brief description; 4 spec. from southwestern Taiwan and South China Sea); Motomura 2020:39 (listed; Japan).

Diagnosis. A species of *Trachonurus* with grooved lateral line; pelvic-fin rays 7; gular and lower portions of branchiostegal membranes heavily scaled; outer series of premaxillary tooth band distinctly enlarged; pectoral-fin rays i13–i17 (usually i15); inner gill rakers on first arch 10–12; body scales small, longitudinal scales 32–46; scales below first dorsal-fin midbase 4.5–8, below second dorsal-fin origin 5–9.

Material examined. 10 specimens. **Holotype** of *Coryphaenoides villosus*: *BMNH 1887.12.7.105 (26.0 mm HL, 193+ mm TL), off Eno-shima Island, Kanagawa Pref., Sagami Bay, Japan, 35.1833°N, 139.4667°E, 345 ftn (631 m), *Challenger* sta. 232, trawl and dredge, 12 Mar. 1875 (only photographed and briefly examined in

FIGURE 193. Fresh specimens of *Trachonurus villosus*. (A) BSKU 23083, 45.2 mm HL, 290+ mm TL, off Kochi, Kochi Pref., Pacific, 1043 m depth; (B) BSKU 43458, 79.7 mm HL, 463+ mm TL, south-southwest of Okino-shima Island, Kochi Pref., Pacific, 1075–1092 m depth; (C) HUMZ 71883, 110 mm HL, 675+ mm TL, off Minami-torishima Island, Pacific. Lateral views. [Photos: (A–B) BSKU; (C) HUMZ]

this study). **Non-types: Japan:** BSKU 26555 (1, 25.8 mm HL, 114+ mm TL), off Miyako-jima Island, Okinawa Trough, 26.4633°N, 124.8483°E, 680–770 m, F/V *Yuryo-maru*, No. 8, tr. 10, bottom trawl, coll. S. Kishida and Y. Kinoshita, 23 Jan. 1978; NSMT-P 93826 (1, 57.7 mm HL, 362+ mm TL), northern Okinawa Trough, 917–919 m, R/V *Tansei-maru*, cr. KT-93-09, 3-m beam trawl, 18 Jun. 1877; BSKU 23083 (1, 45.2 mm HL, 290+ mm TL), off Kochi, 32.9800°N, 133.8550°E, 1043 m, FRV *Kaiyo-maru*, cr. SK-75-2, sta. 9, tr. 2, bottom trawl, 26 Jan. 1975; BSKU 23055 (1, 60.1 mm HL, 383+ mm TL), Tosa Bay, 33.0350°N, 133.7500°E, 945 m, FRV *Kaiyo-maru*, cr. SK-

75-2, sta. 18, tr. 3, bottom trawl, 26 Jan. 1975; SNFR uncat. (1, 56.3 mm HL, 328+ mm TL), Tosa Bay, 750 m, FRV *Kotaka-maru*, otter trawl, 13 Sept. 1988; BSKU 43458 (1, 79.7 mm HL, 463+ mm TL), south-southwest of Okino-shima Island, 32.5450°N, 132.4750°E, 1075–1092 m, R/V *Tansei-maru*, cr. KT-86-16, sta. C, 3-m beam trawl, 2 Nov. 1986; BSKU 20536 (1, 37.2 mm HL, 229+ mm TL), south of Izu Peninsula, 34.3333°N, 138.8167°E, FRV *Soyo-maru*, sta. 30, 5 Oct. 1965; HUMZ 71882 (1, 124 mm HL, 495+ mm TL), HUMZ 71883 (1, 110 mm HL, 675+ mm TL), off Minami-torishima Island, 23.8667°N, 150.6667°E, 10 Oct. 1977.

FIGURE 194. Holotype of *Coryphaenoides villosus* (= *Trachonurus villosus*). BMNH 1887.12.7.105, 26.0 mm HL, 193+ mm TL, Sagami Bay, Pacific, 631 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and body. [Photos: N. Nakayama]

FIGURE 195. Scanning electron micrographs showing a body scale (from the dorsum below the interdorsal space) of *Trachonurus villosus*. BSKU 43458, 79.7 mm HL. (A) View from above; (B) oblique view. [Photos: N. Nakayama]

Counts and measurements. Based on 9 specimens (25.8–124 mm HL, 114–675 mm TL). Counts: first dorsal-fin rays II,7–9; pectoral-fin rays i13–i17; pelvic-fin rays 7; gill rakers on first arch (outer/inner) 4–9/10–12, on second arch 9–14/10–12; longitudinal scales 32–46; transverse scale rows below first dorsal-fin origin 6.5–8.5, below first dorsal-fin midbase 4.5–8, below second dorsal-fin origin 5–9, above anal-fin origin 22–33.5.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 24–29 (31–40); orbit diameter 24–33 (32–42); postorbital length 46–55 (60–71); postrostral length 73–79 (–); orbit–preopercle distance 30–39 (39–51); suborbital width 10–14 (14–18); upper-jaw length 31–38 (41–49); length of rictus 26–33 (35–42); length of premaxillary tooth band 20–26 (26–34); preoral length

12–16 (16–21); distance between tip and lateral angle of snout 10–16 (13–21); snout width 18–24 (23–31); internasal width 19–23 (25–32); interorbital width 32–38 (43–51); body width over pectoral-fin bases 38–66 (50–90); body depth at first dorsal-fin origin 66–104 (89–135); body depth at anal-fin origin 52–93 (68–119); prepelvic length 114–134 (149–181); preanus length 137–165 (180–214); preanal length 144–172 (189–230); isthmus–pelvic distance 55–65 (72–88); isthmus–anus distance 74–95 (98–123); isthmus–anal distance 82–108 (108–140); pelvic–anal distance 26–46 (34–58); anus–anal distance 8–15 (10–20); pelvic-fin length 29–45 (38–60); pectoral-fin length 42–54 (56–71); predorsal length 109–121 (143–164); height of first dorsal fin 51–65 (67–87); length of first dorsal-fin base 16–23 (21–32); interdorsal length 27–45 (36–61); length of gill slit 14–20

(19–25); length of posterior nostril 5–10 (7–14); barbel length 9–12 (11–16).

Size. Attains about 68 cm TL (HUMZ 71883, 675 mm TL, off Minami-torishima Island, Japan).

Distribution. Recorded from the northwestern Pacific off Japan and Taiwan as well as from the southeastern Pacific at the Sala y Gomez Ridge; depth range 514–1240 m (Sazonov & Iwamoto 1992; Shao *et al.* 2008a, 2008b; this study). In Japanese waters, known from the Pacific off southern Japan northward to the Miura Peninsula (139.62°E), Okinawa Trough, and off Minami-torishima Island, at depths of 631–1187 m (Appendix 3-12B). It appears to be rare wherever found.

Comments on type specimen. Günther (1877) originally described the species (as *Coryphaenoides villosus*) based on a single small specimen (BMNH 1887.12.7.105, 26 mm HL, 193+ mm TL; Fig. 194) collected from “south of Yeddo”, Japan. Ten years after the original description, he (1887) reported an additional specimen from the south of the Philippines (BMNH 1887.12.7.106, 45 mm HL, 246+ mm TL). Sazonov & Iwamoto (1992:78) erroneously considered the Philippines specimen as a syntype of *T. villosus*, designating the Japanese specimen as the lectotype of this species. This act was probably due to their recognition that the Philippines specimen “appears to be a different species with a longer upper jaw, longer barbel, longer first gill slit, wider 1D.–2D. interspace, and larger scales”. According to Iwamoto (1997:945), this specimen is unable to be allocated to any known species, partly owing to its poor condition. The above Philippines specimen is not included in the type series of *T. villosus*, being designated as a paralectotype of *Coryphaenoides asper* in this study (see the Comments on type specimens of *C. asper*).

Remarks. Sazonov & Iwamoto (1992) reported *T. villosus* from the Sala y Gomez Ridge in the southeastern Pacific, but noted that their specimens differ from the holotype from Japan in many aspects, including the orbit diameter, suborbital width, orbit–preopercle distance, upper-jaw length, and the number of pelvic-fin rays. A further study is required to confirm the conspecificity of these two populations.

Comparisons. The species of *Trachonurus* are very similar to each other in general appearance, with most species having few diagnostic features that allow ready identification (Iwamoto 1997:942). Among the other five species of the genus (see the Remarks of the genus), *T. gagates* Iwamoto & McMillan, 1997 may not be confused with any other congeners in the absence of a grooved lateral line on the body (vs. present). However, the remaining four species are quite similar to *T. villosus*, and they are most likely to be misidentified without a detailed examination of morphology. Based on published information, *T. villosus* differs from *T. sentipellis* Gilbert & Cramer, 1897 in that the outer premaxillary teeth are distinctly enlarged (vs. scarcely or not at all enlarged, fide Iwamoto & Williams

1999:213), and from *T. robinsi* Iwamoto, 1997 in having fewer spinules on the body scales (see Iwamoto 1997: fig. 2). According to Iwamoto (1997:948), *T. villosus* can be distinguished from *T. sulcatus* (Goode & Bean, 1885) in its slightly lower dimensions of the upper-jaw length, suborbital width, and snout length. However, these differences become negligible when considering variation of the *T. villosus* specimens examined in this study. The differences between *T. villosus* and *T. yiwardaus* Iwamoto & Williams, 1999 were not discussed in the original description of the latter (Iwamoto & Williams 1999), and should be confirmed by a direct comparison of the two species in the future. As noted by Iwamoto (1997:948), there remains much work to be done on *Trachonurus*, and additional characters are obviously necessary to further characterize each species. A molecular approach may be helpful to clarify the problems within this difficult group of grenadiers.

Genus *Ventrifossa* Gilbert & Hubbs, 1920

[Japanese name: Misaki-sokodara-zoku]

Ventrifossa Gilbert & Hubbs, 1920:543 (type species: *Coryphaenoides garmani* Jordan & Gilbert in Jordan & Starks, 1904, by original designation).

Atherodus Gilbert & Hubbs, 1920:544 (subgenus of *Ventrifossa*; type species: *Optonurus atherodon* Gilbert & Cramer, 1897, by original designation).

Lucigadella Gilbert & Hubbs, 1920:552 (subgenus of *Ventrifossa*; type species: *Macrourus nigromarginatus* Smith & Radcliffe in Radcliffe, 1912, by original designation).

Sokodara Iwamoto, 1979:153 (subgenus of *Ventrifossa*; type species: *Coryphaenoides misakius* Jordan & Gilbert in Jordan & Starks, 1904, by original designation).

Diagnosis. Anus closer to pelvic-fin bases than to anal-fin origin; periproct teardrop-shaped, narrowly developed. Small dermal window of light organ between pelvic-fin bases. Infraorbital ridge not connected with preopercular ridge, separated by distinct gap. Leading edge of second spinous ray of first dorsal fin smooth or serrated. Pelvic fin inserted below or anterior to pectoral-fin base. Chin barbel well developed. Premaxillary teeth in narrow tapered band, with outer series enlarged; posterior margin of premaxillary tooth band extending beyond hind edge of maxillary process; mandibular teeth small, in 2–3 rows or narrow band. Spinules on body scales needle-like or broadly triangular, arranged in quincunx order; buttresses of spinules scarcely developed; reticulate structure absent. Head fully covered with small, unmodified scales; snout tipped with small modified scute in some species (subgenus *Sokodara* Iwamoto, 1979). Grooved lateral line present. Cephalic sensory canals lacking open pores. Pyloric caeca numerous, about 30–75. Branchiostegal rays 7. Body dark to swarthy, with silvery reflection

when fresh in most species. [Modified from Sazonov & Iwamoto (1992), Iwamoto & Merrett (1997), and Iwamoto & Graham (2001).]

Remarks. *Ventrifossa* includes more than 25 species (Iwamoto & Graham 2001), eight of which are herein recorded from the study area. Iwamoto & Williams (1999:219) noted the absence of synapomorphies uniting all the species of the genus. Four subgenera were previously recognized (Iwamoto 1979): *Atherodus* Gilbert & Hubbs, 1920; *Lucigadus* Gilbert & Hubbs, 1920; *Sokodara* Iwamoto, 1979; and *Ventrifossa*. Of these, *Lucigadus* now deserves full generic status (see the Remarks of that taxon). Generally, most species of *Ventrifossa* are too similar to one another to allow ready identification in the field. In addition, almost intact specimens are often needed for reliable identification to the species level (Iwamoto *et al.* 2015). According to Iwamoto (1990:292), the genus is most similar to *Malacocephalus* Günther, 1862.

Key to species of *Ventrifossa* from Japan and adjacent waters

- 1a Second spinous ray of first dorsal fin smooth along its leading edge; outer series of premaxillary tooth band distinctly enlarged; mandibular teeth arranged in 2 irregular rows *V. macroptera*
- 1b Second spinous ray of first dorsal fin serrated along its leading edge; outer series of premaxillary tooth band only slightly enlarged 2
- 2a Snout moderately high, protruding distinctly beyond upper jaw; suborbital shelf abruptly and greatly constricted anteriorly; small but prominent tubercular scute at tip of snout; barbel short, length 3–16% HL (usually $\leq 12\%$); body scales very small, longitudinal scales over distance equal to predorsal length 58–73 3
- 2b Snout low, protruding only slightly beyond upper jaw; suborbital shelf not greatly constricted anteriorly; no scute-like scales at tip of snout; barbel well developed, length 16–38% HL; body scales moderately small, longitudinal scales over distance equal to predorsal length 35–56 4
- 3a Oral cavity black; barbel length 9–16% HL; internasal width 21–23% HL *V. johnborum*
- 3b Oral cavity pale; barbel length 3–7% HL; internasal width 24–29% HL *V. misakia*
- 4a Body scales available for identification 5
- 4b Body scales not available for identification 6
- 5a Spinules on body scales broadly triangular **V. garmani*
- 5b Spinules on body scales needle-like 6
- 6a Transverse scale rows below second dorsal-fin origin 8.5–11.5 *V. longibarata*
- 6b Transverse scale rows below second dorsal-fin origin 6–7.57
- 7a Prominent black spot on first dorsal fin 8
- 7b First dorsal fin uniformly dusky 9
- 8a Anal fin immaculate anteriorly; interorbital width 27–29%

- HL; upper-jaw length 38–40% HL; inner gill rakers on second arch 12–13 *V. nigrodorsalis*
- 8b Anal fin narrowly black along anterodistal margin; interorbital width 20–27% HL; upper-jaw length 42–47% HL; inner gill rakers on second arch 14–18 *V. rhipidodorsalis*
- 9a Snout width 25–29% HL; suborbital width 11–13% HL **V. garmani*
- 9b Snout width 30–40% HL; suborbital width 13–17% HL *V. saikaiensis*

Ventrifossa garmani (Jordan & Gilbert in Jordan & Starks, 1904)

[Japanese name: Sagami-sokodara]

(Figs. 196–198; Appendix 3-12C)

Coryphaenoides garmani Jordan & Gilbert in Jordan & Starks, 1904:610, unnumbered fig. [original description; holotype: USNM 50933, from Sagami Bay, *Albatross* sta. 3695, in 110–259 fth (202–474 m); paratypes from Sagami-nada and Suruga Bay]; Franz 1910:26 [brief description; 3 spec. from “Sagamibai” (= Sagami Bay)]; Jordan *et al.* 1913:416, fig. 386 (listed; Japan; new Japanese name: “Sagami-sokodara”); Jordan & Thompson 1914:306 [2 spec. from Boshu (= Chiba) and Misaki; FMNH 57245 and FMNH 57387]; Kamohara 1931b:544 (listed; Kochi); Kamohara 1934a:54 (listed; Kochi); Kamohara 1938:72 (selected counts and measurements; spec. from Kochi Pref.); Kamohara 1950:279 (listed; Kochi and Wakayama Pref.); Kuroda 1951:391 (listed; Suruga Bay); Uchinomi 1951:76 (listed; southern part of Kii Peninsula); Kamohara 1952:98 (selected counts and measurements; spec. from Kochi Pref.).

Lionurus garmani: Gilbert & Hubbs 1916:193 (new combination; brief description; 30 spec. from East China Sea, off Cape Shionomisaki, Pacific off Tohoku, Suruga Bay, and Sagami Bay).

Ventrifossa garmani: Okada & Matsubara 1938:452 (in key; Japan); Matsubara 1955:1315 (in key; Japan); Kobayashi 1956:72 (listed; Enshu-nada); Kamohara 1958:74 (listed; Kochi Pref.); Kamohara & Okamura 1958:32, pl. 31, fig. 152 (compiled; Japan); Kamohara 1961b:68, pl. 68, fig. 2 (compiled; Japan); Kuroda 1962:7, fig. 2 (description of fresh color; 3 spec. from Suruga Bay); Kamohara 1964:96 (listed; Kochi Pref.); Matsubara 1965:508 (compiled; Japan); Okamura 1970a:74, pl. V, text-fig. 33 (description; biological notes; 131 spec. from Pacific off southern Japan from Choshi to Totoro); Okamura 1970b: table 1 (listed; Japan); Nakajima 1975:55 (listed; Enshu-nada); Tominaga & Uyeno 1981:489 (listed; Japan); Okamura 1982:145, 348, fig. 89 [brief description; 6 spec. from Kyushu-Palau Ridge and Tosa Bay; photo based on BSKU spec. (catalog no. unknown)]; Ohta 1983: table A (listed; Suruga Bay); Ozawa 1983:13 (listed; off Makurazaki, Kagoshima Pref., East China Sea); Okamura 1984a:213, 360, fig. 148 [brief description; 6 spec. from Okinawa Trough; photo based on BSKU 36046 (from Mimase fish market)]; Okamura 1984b:93, pl. 81, fig.

E (compiled); Akazaki 1984:265 (listed; Aoshima, Miyazaki Pref.); Okamura 1988:93, pl. 81, fig. E (compiled); Iwamoto 1990:300, fig. 678 (synopsis); Nakabo 1993:359 (in key; Japan); Shao 1993:172 (compiled; Taiwan); Shinohara *et al.* 1996:170 (2 spec. listed from Pacific off Tohoku); Suzuki & Kataoka 1997:82, pl. 32, fig. 176 (brief description; 1 spec. from Kumano-nada); Okamura 1997:128, fig. 9 (compiled); Shinohara & Matsuura 1997:292 (listed; Suruga Bay); Nakabo 2000:423 (in key; Japan); Shinohara *et al.* 2001:306 (79 spec. listed from Tosa Bay); Nakabo 2002:423 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Nakajima 2003:53 (brief description; 1 spec. from Enshu-nada); Chiou *et al.* 2004a: table 1 (listed; Taiwan); Fukui & Tsuchiya 2005:311, figs. 1–2 (larval description; 3 spec. from Suruga Bay and adjacent water); Shinohara *et al.* 2005:418 (12 spec. listed from Ryukyu Islands); Senou *et al.* 2006:421 (listed; Sagami Sea); Shao *et al.* 2008b: table 2 (2 spec. listed from southwestern Taiwan); Shinohara *et al.* 2009:709 (listed; Pacific off Tohoku); Furuhashi *et al.* 2010: table 2 (217 spec. listed from northern Okinawa Trough); Jeong *et al.* 2010:317, fig. 1 (first record from Korea; brief description; 1 spec. from Ulsan); Nakabo & Kai 2013:499 (in key; Japan); Fukui *et al.* 2015:182, fig. 12 (4 spec. from off Kuro-shima Island, Kagoshima Pref., East China Sea); Ikeda & Nakabo 2015:320, pl. 65, figs. 6–8 (brief description; 2 spec. from Wakayama Pref.); Iwamoto *et al.* 2015:109, fig. 28 (brief description; 8 spec. from southwestern Taiwan and Japan); Iwatsuki *et al.* 2017:32 (listed; Hyuga-nada); Sonoyama *et al.* 2020:30, fig. 4G (1 spec. listed from Yamaguchi Pref., Sea of Japan); Motomura 2020:39 (listed; Japan).

[?] *Lionurus darus* (not Gilbert & Hubbs 1916): Kuroda 1941:80, fig. 10 (brief description; 2 spec. from Suruga Bay) [re-identified here from figure].

[?] *Malacocephalus nipponensis* (not Gilbert & Hubbs 1916): Suzuki & Kataoka 1997:82, pl. 32, fig. 175 (brief description; 1 spec. from Kumano-nada) [re-identified here from figure].

Diagnosis. Second spinous ray of first dorsal fin finely serrated along its leading edge; first dorsal and anal fins uniformly pale to dusky; anal fin uniformly dusky; prominent black spot absent on front of premaxillary ascending process; oral cavity pale; median rostral ridge lacking narrow blackish streak; leading edges of snout, supranarial ridges, and suborbital shelves only faintly marked with black; dorsal parts of trunk and tail dark, sharply demarcated from paler lateral and ventral parts of body (especially prominent in fresh specimens); snout low, slightly protruding beyond upper jaw, preoral length 14–22% HL; no scute-like scale at snout tip; body scales covered with short, moderately erect, broadly triangular spinules in quincunx order; spinules on scales along second dorsal-fin base not enlarged; body scales moderately small, transverse scale rows below midbase of first dorsal fin 5.5–7.5, below second dorsal-fin origin 5.5–7; longitudinal scales 35–44; area of spinuleless scales posterior to first dorsal-fin base almost absent or only

narrowly developed; outer series of premaxillary teeth slightly enlarged; mandibular teeth arranged in narrow tapered band; suborbital shelf not greatly constricted anteriorly; inner gill rakers on first arch 16–19, outer gill rakers on second arch 15–18; orbit large, greatest diameter 32–38% HL; suborbital width 11–13%; upper-jaw length 41–47% HL; interorbital space broad, width 27–33% HL; barbel moderately long, length 20–33% HL; pectoral-fin length 52–66% HL.

Material examined. 40 specimens. **Holotype of *Coryphaenoides garmani*:** USNM 50933 (51.7 mm HL, 285+ mm TL), off Chigasaki, Kanagawa Pref., Sagami Bay, Japan, 35.2500°N, 139.4167°E, 110–259 ftm (202–474 m), *Albatross* sta. 3695, 8-ft Tanner beam trawl, 4 Mar. 1900. **Paratypes of *C. garmani*:** CAS-SU 8548 (5, 30.6–47.1 mm HL, 191+–279+ mm TL), southeast of Cape Manazurumisaki, Sagami Bay, 35.1333°N, 139.1667°E, 120–153 ftm (220–280 m), *Albatross* sta. 3697 or 3698, 8-ft Tanner beam trawl, 5 Mar. 1905; *USNM 51415 (9, 20.4–45.7 mm HL, 83+–249+ mm TL), off Heta, Suruga Bay, 34.9667°N, 138.7667°E, 161–167 ftm (295–306 m), *Albatross* sta. 3737, tangles, 17 Mar. 1900. **Non-types:** **Japan:** BSKU 106845 (1, 42.0 mm HL, 225+ mm TL), southeast of Shimokoshiki-jima Island, East China Sea, 31.5675°N, 129.8930°E, 393 m, F/V *Maruko-maru*, tr. 2, bottom trawl, coll. N. Nakayama *et al.*, 24 Apr. 2012; BSKU 106763 (1, 39.7 mm HL, 212+ mm TL), BSKU 106764 (1, 33.9 mm HL, 187+ mm TL), southeast of Shimokoshiki-jima Island, East China Sea, 31.5658°N, 129.8915°E, 380 m, F/V *Maruko-maru*, tr. 1, bottom trawl, coll. N. Nakayama *et al.*, 24 Apr. 2012; BSKU 106820 (1, 37.6 mm HL, 199+ mm TL), southeast of Shimokoshiki-jima Island, East China Sea, 31.5652°N, 129.8878°E, 407 m, F/V *Maruko-maru*, tr. 3, bottom trawl, coll. N. Nakayama *et al.*, 24 Apr. 2012; BSKU 50878 (1, 32.5 mm HL, 160+ mm TL), off Nomaie, Kagoshima, East China Sea, F/V *Maruko-maru*, bottom trawl, coll. M. Yamashita and Y. Ohashi, 10 Sept. 2010; BSKU 109054 (1, 49.3 mm HL, 286+ mm TL), west of Sanpo-sone, 32.2872°N, 129.0732°E, 351–389 m, T/V *Nagasaki-maru*, cr. N365, sta. B4, 3-m ORE beam trawl, coll. N. Nakayama, 19 Nov. 2012; BSKU 109053 (1, 18.0 mm HL, 108+ mm TL), east of Kasayama Bank, 32.3884°N, 129.0505°E, 304–312 m, T/V *Nagasaki-maru*, cr. N365, sta. A3, 3-m ORE beam trawl, coll. N. Nakayama, 19 Nov. 2012; BSKU 109020 (1, 51.4 mm HL, 300+ mm TL), west of Okikasayama Bank, 32.1503°N, 129.0072°E, 499–504 m, T/V *Nagasaki-maru*, cr. N365, sta. C5, 3-m ORE beam trawl, coll. N. Nakayama, 19 Nov. 2012; BSKU 27287 (1, 58.8 mm HL, 325+ mm TL), BSKU 27290 (1, 53.7 mm HL, 314+ mm TL), west-northwest of Amami-oshima Island, Okinawa Trough, 28.8033°N, 127.0617°E, 425 m, F/V *Yuryo-maru*, No. 8, tr. 25, bottom trawl, coll. Y. Kinoshita and S. Hagino, 6 Feb. 1978; *BSKU 101604 (1, 41.8 mm HL), south of Muroto, 33.0790°N, 134.1645°E, 414–513 m, R/V *Tansei-maru*, cr. KT-05-29, sta. K-500,

FIGURE 196. Fresh specimens of *Ventrifossa garmani*. (A) BSKU 109020, 51.4 mm HL, 300+ mm TL, east of Okikasayama Bank, Okinawa Trough, East China Sea, 499–504 m depth; (B–D) BSKU 112453, 54.6 mm HL, 306+ mm TL, Tosa Bay off Susaki, Kochi Pref., Pacific, 320–380 m depth. (A–B) Lateral views; (C) dorsal and (D) ventral views of the head and trunk. [Photos: N. Nakayama]

beam trawl, coll. H. Endo *et al.*, 17 Nov. 2005; BSKU 71693 (1, 24.8 mm HL, 153+ mm TL), Mimase fish market, bottom trawl, 13 Feb. 2004; BSKU 72788 (1, 27.4 mm HL, 166+ mm TL), Mimase fish market, bottom trawl, 20 Feb. 2004; BSKU 112435 (1, 51.2 mm HL, 292+ mm TL), BSKU 112453 (1, 54.6 mm HL, 306+ mm TL), BSKU 112454 (1, 44.9 mm HL, 260+ mm TL), BSKU 112456 (1, 37.4 mm HL, 182+ mm TL), BSKU 112459 (1, 47.8 mm HL, 271+ mm TL), off Susaki, Tosa Bay, 320–380 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 9 Mar. 2014; BSKU 113653 (1, 40.5 mm HL, 249+ mm TL), BSKU 113654 (1, 47.0 mm HL, 283+ mm TL), off Okitsu, Tosa Bay, 380 m, F/V *Kosei-maru*, bottom trawl, coll. N. Nakayama *et al.*, 11 Mar. 2014; BSKU 113667 (1, 33.5 mm HL, 196+ mm TL), Numazu fish market, coll. S. Ohashi and M. Takami, 16

Feb. 2010; BSKU 110089 (1, 40.4 mm HL, 219+ mm TL), BSKU 110434 (1, 35.1 mm HL, 212+ mm TL), BSKU 110435 (1, 42.3 mm HL, 253+ mm TL), Suruga Bay, 34.7489°N, 138.4664°E, 200–450 m, F/V *Hinode-maru*, sta. 6, bottom trawl, coll. N. Nakayama and R. Misawa, 23 Apr. 2013; BSKU 110016 (1, 45.1 mm HL, 258+ mm TL), Suruga Bay, 34.7080°N, 138.4587°E, 200–450 m, F/V *Hinode-maru*, sta. 2, bottom trawl, coll. N. Nakayama and R. Misawa, 23 Apr. 2013.

Counts and measurements. Based on 30 specimens (18–58.8 mm HL, 108+–325+ mm TL). Counts: first dorsal-fin rays II,9–12; pectoral-fin rays i17–i23; pelvic-fin rays 8–9; gill rakers on first arch (outer/inner) 9–14/16–19, on second arch 15–18/14–18; longitudinal scales 35–44; transverse scale rows below first dorsal-fin origin 7.5–11, below first dorsal-fin midbase 5.5–7.5,

FIGURE 197. Holotype of *Coryphaenoides garmani* (= *Ventrifossa garmani*). USNM 50933, 51.7 mm HL, 285+ mm TL, Sagami Bay, Pacific, 202–474 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

FIGURE 198. Scanning electron micrographs showing a body scale (from the dorsum below the interdorsal space) of *Ventrifossa garmani*. BSKU 101604, 41.8 mm HL. (A) View from above; (B) oblique view. [Photos: N. Nakayama]

below second dorsal-fin origin 5.5–7, above anal-fin origin 21.5.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 26–29 (32–39); orbit diameter 32–38 (42–49); postorbital length 41–46 (52–61); postrostral length 74–80; orbit–preopercle distance 40–45 (52–60); suborbital width 11–13 (15–18); upper-jaw length 41–47 (55–62); length of rictus 35–39 (46–52); length of premaxillary tooth band 30–34 (38–46); preoral length 14–22 (18–27); distance between tip and lateral angle of snout 15–18 (19–23); snout width 25–29 (33–39); internasal width 20–

25 (26–33); interorbital width 27–33 (35–43); occipital width 14–19 (18–25); body width over pectoral-fin bases 38–61 (49–78); body depth at first dorsal-fin origin 79–99 (100–129); body depth at anal-fin origin 50–89 (63–116); prepelvic length 101–123 (131–159); preanus length 109–129 (144–171); preanal length 129–148 (165–196); isthmus–pelvic distance 41–58 (54–77); isthmus–anus distance 47–68 (62–90); isthmus–anal distance 67–90 (87–119); pelvic–anal distance 22–38 (28–49); anus–anal distance 12–26 (16–34); pelvic-fin length 38–44 (50–57); pectoral-fin length 52–66 (69–87); predorsal length 103–118 (136–154); height of first dorsal fin 70–87 (95–113);

length of first dorsal-fin base 26–35 (34–46); interdorsal length 50–81 (65–108); length of gill slit 24–30 (31–39); length of posterior nostril 3–6 (5–7); barbel length 20–33 (27–43).

Size. To about 33 cm TL (BSKU 27287, 325+ mm TL, Okinawa Trough, Japan).

Development. Larval morphology was described by Fukui & Tsuchiya (2005).

Distribution. Restricted to Japan, Korea, and Taiwan. Known from off the Pacific coasts of Japan northward to Rikuzen-Takata (39.03°N; Iwate Pref.), Okinawa Trough, northern South China Sea, and Sea of Japan off Hagi (Yamaguchi Pref.) as well as the Korean Peninsula, at depths of 110–980 m (Shao *et al.* 2008a, 2008b; Jeong *et al.* 2010; Sonoyama *et al.* 2020; this study; Appendix 3-12C). A dominant species of *Ventrifossa* found along the Pacific coasts of southern Japan as well as in the East China Sea (but rare in Taiwanese and Korean waters).

Remarks. For a full description see Okamura (1970a). *Ventrifossa garmani* was originally described by Jordan & Gilbert (*in* Jordan & Starks 1904; as a species of *Coryphaenoides* Gunnerus, 1765) based on specimens collected from Sagami Bay, Japan (Fig. 197). This is by far the most common species of *Ventrifossa* at upper slope depths in the Pacific off southern Japan and the Okinawa Trough in the East China Sea (Horikawa 2009: table 1; Furuhashi *et al.* 2010: table 2; author's pers. observ.).

Relationships and comparisons. *Ventrifossa garmani* belongs to the subgenus *Ventrifossa* as redefined by Iwamoto (1979), and is unlikely to be mistaken for any other congeners, if body scales are available for identification. In *V. garmani*, spinules on the body scales are broadly triangular (Fig. 198), whereas those of other species are needle-like (see Figs. 201, 205, 208). Among other Japanese congeners of the subgenus *Ventrifossa*, this species is readily distinguished from *V. longibarbata* Okamura, 1982 and *V. rhipidodorsalis* Okamura, 1984 by having a broader interorbital space (27–33% HL vs. 21–26% and 20–27% in *V. longibarbata* and *V. rhipidodorsalis* respectively). It further differs from *V. longibarbata* in having much larger scales on the body (scale rows below the first dorsal-fin midbase 5.5–7.5 vs. 8.5–11.5; below the second dorsal-fin origin 5.5–7 vs. 7–9.5) and a shorter chin barbel (20–33% HL vs. 31–38%); and from *V. rhipidodorsalis* by lacking a prominent black spot on the first dorsal fin (vs. present). *Ventrifossa garmani* is distinguished from *V. saikaiensis* Okamura, 1984 by having a paler suborbital shelf (vs. blackish), a narrower snout (snout width 25–29% HL vs. 30–40%), and a shallower suborbital space (11–13% HL vs. 13–17%).

Ventrifossa johnborum Iwamoto, 1982

[New Japanese name: Sumikui-sokodara]

(Figs. 199–200, 201A; Table 14; Appendix 3-12D)

Ventrifossa johnborum Iwamoto, 1982:55, fig. 1 (original description; holotype: AMS I. 15602-002, collected from mouth of *Diceratias bispinosus*, floating at surface near Papua New Guinea, Bismarck Sea; 1 paratype from Philippines and 2 non-types from South China Sea).

Ventrifossa misakia (not Jordan & Gilbert *in* Jordan & Starks 1904): Okamura 1984a:213, 361, fig. 149 (brief description; 1 spec. from Okinawa Trough; BSKU 28595); Okamura 1984b:93, pl. 344, fig. J (compiled); Okamura 1988:93, pl. 344, fig. J (compiled).

Diagnosis. Second spinous ray of first dorsal fin weakly serrated along its leading edge; first dorsal and anal fins uniformly dusky to dark; small black spot present on front of premaxillary ascending process; lining of oral cavity intensely black; head ridges not marked with black; body uniformly pale to dark; snout moderately high, protruding distinctly beyond upper jaw, preoral length 19–25% HL; small, stout, scute-like scale at snout tip; body scales covered with short, moderately erect, needle-like spinules in quincunx order; spinules on scales along second dorsal-fin base not enlarged; body scales very small, transverse scale rows below midbase of first dorsal fin 10–13, below second dorsal-fin origin 8.5–11; longitudinal scales 60–72; area of spinuleless scales posterior to first dorsal-fin base almost absent or only narrowly developed; outer series of premaxillary teeth slightly enlarged; mandibular teeth arranged in narrow tapered band; suborbital shelf abruptly narrowing anteriorly; inner gill rakers on first arch 12–15, outer gill rakers on second arch 11–14; orbit large, greatest diameter 30–38% HL; upper-jaw length 36–41% HL; interorbital space broad, width 26–30% HL; chin barbel short, length 9–16% HL; pectoral-fin length 52% HL.

Material examined. 9 specimens. **Holotype** of *Ventrifossa johnborum*: AMS I.15602-002 (85.3 mm HL, 365+ mm TL), Bismarck Sea, floating at surface near Tanga Island, Papua New Guinea (in mouth of ceratioid anglerfish *Diceratias bispinosus*), 30 Apr. 1967. **Non-types: Japan:** BSKU 28595 (1, 47.9 mm HL, 222+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.9500°N, 128.0000°E, 585–610 m, F/V *Ryoanmaru*, No. 28, tr. 76, bottom trawl, coll. Y. Kinoshita and S. Hagino, 20 Mar. 1978. **Southwestern Pacific:** CAS 90669 (1, 95.6 mm HL, 468+ mm TL), Norfolk Ridge, 24.6923°S, 168.6813°E, 800–1240 m, *HALIPRO* 2, sta. BT 64, coll. T. Iwamoto *et al.*, 19 Nov. 1996; CAS 90672 (3, 78.5–97.2 mm HL, 337+–461+ mm TL), Norfolk Ridge, 24.6938°S, 168.6497°E, 720–1048 m, *HALIPRO* 2, sta. BT 62, coll. T. Iwamoto *et al.*, 18 Nov. 1996; CAS 86488 (1, 46.6 mm HL, 243+ mm TL), New Caledonia, 23.1164°S, 166.8533°E, 825 m, *BIOCAL*, sta. CP 32, 29 Aug. 1985; CAS 90665 (1, 81.4 mm HL, 412+ mm TL), Lord Howe Rise, 24.2803°S, 161.7508°E, 970–1063 m, *HALIPRO* 2, sta. BT 101, coll. T. Iwamoto *et al.*, 26 Nov. 1996; CAS 86457 (1, 41.7 mm HL, 152+ mm TL), east

FIGURE 199. Holotype of *Ventrifossa johnboborum*. AMS I.15602-002, 85.3 mm HL, 365+ mm TL, floating at the surface near Tanga Island, Papua New Guinea, Bismarck Sea (in a mouth of the ceratioid anglerfish *Diceratias bispinosus*). (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

FIGURE 200. Preserved specimen of *Ventrifossa johnboborum*. BSKU 28595, 47.9 mm HL, 222+ mm TL, northwest of Amami-oshima Island, Kagoshima Pref., Okinawa Trough, East China Sea, 585–610 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

of Bellona Plateau, 21.0150°S, 160.8383°E, 745–825 m, *MUSORSTOM 5*, sta. CC 390, 22 Oct. 1986.

Counts and measurements. See Table 14.

Size. Attains about 48 cm TL (Sazonov & Iwamoto 1992; Iwamoto & Williams 1999; Iwamoto & Graham 2001).

Distribution. Widely distributed in the Indian and Pacific Oceans, including the Philippines, South China

Sea, New Caledonian region, Australia (Western Australia, Queensland, and New South Wales), and Sala y Gomez Ridge, at depths of about 400–1100 m (Iwamoto 1982; Sazonov & Iwamoto 1992; Iwamoto & Merrett 1997; Sazonov & Shcherbachev 1997; Iwamoto & Williams 1999; Merrett & Iwamoto 2000; Iwamoto & Graham 2001; Last *et al.* 2014). The Japanese record is based on a single specimen collected from the Okinawa Trough in

FIGURE 201. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of (A) *Ventrifossa johnboborum* and (B–H) *V. misakia*. (A) BSKU 28595, 47.9 mm HL; (B) BSKU 78801, 36.9 mm HL; (C) BSKU 78807, 21.6 mm HL; (D) BSKU 98219, 57.6 mm HL; (E) NSMT-P 93117, 66.3 mm HL; (F) BSKU 26068, paratype of *V. fusca*, 98.2 mm HL; (G) HUMZ 95098, 103 mm HL; (H) BSKU 26067, holotype of *V. fusca*, 112 mm HL. Views from above. [Photos: N. Nakayama]

585–610 m (Appendix 3-12D). Rare in the northwestern Pacific, but common in the Southern Hemisphere.

Remarks. This species has been adequately described by previous authors, and there is no need for repetition here. For further morphological details see the original description given by Iwamoto (1982). Sazonov & Iwamoto (1992) also provided an excellent description of the species, based on the holotype and additional specimens from the southeastern Pacific.

Ventrifossa johnboborum was originally described on the basis of the holotype collected from the Bismarck Sea (type locality; Fig. 199), and three other specimens (not designated as paratypes) from the Philippines and South China Sea, of which a Philippines specimen was previously

reported as *V. misakia* Jordan & Gilbert in Jordan & Starks, 1904 by Gilbert & Hubbs (1920). Subsequently, it has been widely recorded from the Southern Hemisphere, viz., the Freda Seamount in the western Indian Ocean (Shcherbachev 1987), west and east coasts of Australia (Iwamoto & Williams 1999; Iwamoto & Graham 2001), New Caledonian region (Iwamoto & Merrett 1997; Merrett & Iwamoto 2000), and Sala y Gomez Ridge in the southeastern Pacific (Sazonov & Iwamoto 1992). However, pronounced geographic variations in several counts and measurements, including pelvic-fin rays, gill rakers, barbel length, upper-jaw length, and pelvic–anal distance, are also known within the species (Sazonov & Iwamoto 1992; Iwamoto & Merrett 1997; Sazonov &

TABLE 14. Morphometric measurements and meristic counts* for *Ventrifossa johnborum* and *V. misakia*.

Species	<i>V. johnborum</i>			<i>V. misakia</i>			<i>V. fusca</i>		
	Nominal species			Nominal species			Nominal species		
	Type status			Type status			Type status		
Number of specimens	HT	NT	PT	HT	NT	PT	HT	NT	PT
	<i>n</i> = 1	<i>n</i> = 8	<i>n</i> = 8	<i>n</i> = 1	<i>n</i> = 8	<i>n</i> = 8	<i>n</i> = 1	<i>n</i> = 19	<i>n</i> = 3
Measurements (mm)									
TL	365+	152+–468+		334+		165+–371+	521+	147+–628+	455+–485+
HL	85.3	41.7–97.2		67.5		54.1–76.2	112	25.3–103	98.2–106
% of HL (% of PRL)									
Snout length	29 (38)	27–31 (35–42)		27 (35)		27–30 (34–42)	27 (36)	26–29 (34–39)	27–28 (35–38)
Orbit diameter	31 (41)	30–38 (41–48)		35 (44)		32–37 (43–51)	29 (38)	30–40 (39–51)	29–30 (40)
Postorbital length	44 (59)	40–45 (51–59)		45 (58)		41–47 (56–65)	48 (62)	39–48 (50–62)	46–47 (61–62)
Postrostral length	75 (–)	73–77 (–)		78 (–)		69–82 (–)	77 (–)	74–78 (–)	74–77 (–)
Orbit–preopercle distance	41 (55)	39–44 (51–60)		45 (58)		41–49 (56–65)	44 (58)	39–45 (50–60)	46 (60–62)
Suborbital width	12 (16)	12–13 (15–18)		12 (16)		12–14 (16–19)	13 (17)	12–14 (16–19)	13 (18)
Upper-jaw length	41 (55)	36–39 (48–53)		41 (52)		37–42 (49–60)	40 (52)	37–40 (48–53)	40 (53–55)
Length of rictus	35 (46)	30–33 (40–44)		32 (42)		32–34 (41–45)	35 (45)	31–34 (40–45)	33 (43–44)
Length of PMX tooth band	28 (38)	22–25 (29–34)		25 (32)		23–28 (31–40)	26 (34)	23–25 (29–34)	25 (32–33)
Preoral length	19 (25)	22–25 (29–32)		21 (27)		20–26 (27–35)	21 (27)	20–26 (26–34)	21–23 (27–31)
Dist. tip and lat. angle of snout	16 (22)	16–21 (22–29)		16 (20)		17–19 (21–26)	15 (20)	15–19 (20–24)	16–17 (22)
Snout width	28 (37)	25–28 (34–38)		26 (33)		25–28 (33–37)	24 (32)	25–30 (33–39)	27 (36)
Internasal width	23 (31)	21–23 (28–31)		25 (32)		24–27 (32–37)	25 (33)	24–29 (32–38)	25–26 (33–34)
Interorbital width	30 (39)	26–29 (34–39)		30 (39)		28–31 (38–41)	30 (39)	29–33 (38–44)	31–32 (40–43)
Body width over P. bases	48 (64)	37–53 (48–72)		45 (58)		43–46 (55–63)	51 (67)	31–57 (40–75)	47 (61–64)
Body depth at 1D. origin	– (–)	68–72 (90–99)		76 (98)		75–81 (98–112)	83 (109)	71–83 (92–111)	79–82 (104–110)
Body depth at A. origin	– (–)	61–67 (81–92)		67 (86)		63–71 (84–100)	75 (98)	56–79 (72–105)	74–76 (99–100)
Pre-V. length	93 (124)	93–115 (121–158)		110 (142)		107–116 (137–154)	111 (145)	100–117 (130–155)	103–105 (134–142)
Pre-anus length	– (–)	102–124 (134–170)		118 (152)		122–125 (152–164)	118 (154)	110–127 (143–168)	114–115 (149–155)
Pre-A. length	134 (178)	121–137 (157–188)		131 (169)		132–141 (170–183)	140 (183)	123–144 (160–191)	131–138 (171–186)
Isthmus–V. distance	34 (45)	27–37 (36–50)		35 (46)		30–35 (39–46)	36 (48)	32–37 (41–50)	37–39 (48–53)
Isthmus–anus distance	– (–)	37–45 (48–61)		44 (56)		40–43 (53–53)	46 (60)	41–50 (54–66)	47–50 (62–67)
Isthmus–A. distance	80 (106)	53–65 (68–89)		57 (74)		53–63 (70–76)	69 (90)	53–74 (69–98)	65–71 (85–96)

...Continued on the next page

TABLE 14. (Continued)

Species	<i>V. johnborum</i>			<i>V. misakia</i>			<i>V. fusca</i>		
	Nominal species			<i>V. misakia</i>			<i>V. fusca</i>		
	Type status	HT	NT	HT	NT	PT	HT	NT	PT
Number of specimens	<i>n</i> = 1	<i>n</i> = 8	<i>n</i> = 8	<i>n</i> = 1	<i>n</i> = 1	<i>n</i> = 8	<i>n</i> = 1	<i>n</i> = 19	<i>n</i> = 3
V.-A. distance	48 (64)	25–32 (32–43)	27 (34)	27 (34)	22–30 (29–38)	25–42 (32–55)	33 (44)	30–35 (39–47)	
Anus–A. distance	– (–)	14–22 (20–30)	13 (17)	13 (17)	12–18 (16–22)	12–26 (16–34)	24 (31)	17–23 (22–31)	
V. length	– (–)	35–44 (47–61)	34 (44)	34 (44)	34–38 (45–50)	27–40 (35–52)	– (–)	– (–)	
P. length	– (–)	52 (72)	58 (75)	58 (75)	48–61 (66–81)	51–64 (67–82)	– (–)	– (–)	
Pre-ID. length	106 (141)	103–109 (136–148)	105 (135)	105 (135)	97–115 (128–147)	101–110 (133–145)	108 (141)	108–111 (141–149)	
Height of 1D.	– (–)	62 (85)	69 (88)	69 (88)	61–66 (80–86)	– (–)	– (–)	– (–)	
Length of 1D. base	27 (35)	21–25 (28–34)	25 (32)	25 (32)	21–26 (29–35)	22–29 (29–38)	23 (30)	29–30 (38–40)	
Interdorsal length	42 (56)	38–52 (50–68)	43 (56)	43 (56)	34–57 (44–75)	37–62 (47–80)	49 (64)	51–58 (66–79)	
Length of gill slit	– (–)	19–21 (25–29)	21 (27)	21 (27)	20–25 (26–34)	18–24 (23–31)	22 (29)	23 (30–31)	
Length of posterior nostril	4 (5)	4–6 (6–8)	3 (4)	3 (4)	3–5 (4–6)	3–5 (4–7)	3 (4)	3–4 (4–5)	
Barbel length	16 (22)	9–12 (11–17)	3 (4)	3 (4)	5–7 (6–9)	4–7 (5–9)	6 (7)	6–7 (7–9)	
Counts									
1D. rays	II,11	II,9–11	II,11	II,11	II,10–12	II,10–13	II,10	II,10–12	
P. rays	i23	i18–i22	i20	i20	i19–i22	i19–i23	i22	i20–i23	
V. rays	9	8	8	8	8	7–9	8	8	
Outer GR on first arch	9	9–13	10	10	9–13	1–13	10	11–13	
Inner GR on first arch	12	13–15	15	15	14–15	13–16	14	15–16	
Outer GR on second arch	11	12–14	14	14	13–15	12–16	16	14–16	
Inner GR on second arch	11	13–14	14	14	14–15	13–16	14	14–15	
Longitudinal scales	60	62–72	70	70	59–73	58–70	63	61–63	
Scales below 1D. origin	21	15–17	14–15	14–15	11–16	11–16	–	15	
Scales below 1D. midbase	13	10–12	9	9	7.5–10.5	8–11	10	9–10.5	
Scales below 2D. origin	10	8.5–11	9.5	9.5	9–10	8–10	9.5	8.5–9	
Scales above A. origin	–	–	–	–	–	37	–	–	

*Abbreviations: 1D.—first dorsal fin; 2D.—second dorsal fin; A.—anal fin; GR—gill rakers; HT—holotype; NT—non-type; P.—pectoral fin; PMX—premaxillary; PT—paratype; V.—pelvic fin

Shcherbachev 1997). A further study is therefore necessary to elucidate the taxonomic status of each population of *V. johnborum* (see also Iwamoto & Graham 2001).

Based on a single specimen collected from the Okinawa Trough in the East China Sea (BSKU 28595, 47.9 mm HL, 222+ mm TL; Fig. 200), Iwamoto *et al.* (2015) newly recorded *V. johnborum* from Japan. This specimen was previously identified as *V. misakia* by Okamura (1984a). The new standard Japanese name, “Sumikui-sokodara”, is derived from the species’ characteristic black oral cavity (= sumikui) and a Japanese name for grenadiers (= sokodara).

Relationships and comparisons. According to Sazonov & Shcherbachev (1997), *V. johnborum* closely resembles *V. misakia* distributed in the northwestern Pacific from Japan to Taiwan, and *V. nasuta* (Smith, 1935) known from the southwestern Indian Ocean, and they comprise the subgenus *Sokodara* Iwamoto, 1979. *Ventrifossa johnborum* differs notably from *V. nasuta* in having higher counts of transverse scale rows (15–21 vs. 7–9 below the first dorsal-fin origin; 10–13 vs. 6–7 below the first dorsal-fin midbase; and 8.5–11 vs. 5.5–7.5 below the second dorsal-fin origin) and a shorter chin barbel (9–16% HL vs. 16–23%) [data for *V. nasuta* are from Sazonov & Shcherbachev (1997: table 2)]. On the other hand, this species is readily distinguished from *V. misakia* by its uniformly black oral cavity (vs. pale) and the presence of a small black spot on the front of the premaxillary ascending process (vs. usually absent). It further differs from that species in having a longer chin barbel (9–16% HL vs. 3–7%) and a narrower internasal space (21–23% HL vs. 24–29%).

Ventrifossa longibarbata Okamura, 1982

[Japanese name: Okina-hige]

(Figs. 202–204, 205A–B; Appendix 3-12E)

Ventrifossa longibarbata Okamura, 1982:157, fig. 94 (original description; holotype: BSKU 29494, from Tosa Bay, 32°58.0'N, 133°32.0'E, in 605 m; 4 paratypes from Suruga Bay, Tosa Bay, and Okinawa Trough, in 382–620 m; new Japanese name: “Okina-hige”); Okamura 1984a:215, 361, fig. 150 (brief description; 43 spec. from Okinawa Trough; photo based on BSKU 29891); Okamura 1984b:93, pl. 81, fig. C (compiled); Okamura 1988:93, pl. 81, fig. C (compiled); Nakabo 1993:358 (in key; Japan); Okamura 1997:128, fig. 10 [compiled; photo spec. is *V. rhipidodorsalis* (BSKU unknown)]; Shinohara & Matsuura 1997:292 (listed; Suruga Bay); Nakabo 2000:422 (in key; Japan); Shinohara *et al.* 2001:306 (26 spec. listed from Tosa Bay); Nakabo 2002:422 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Chiou *et al.* 2004a:48, fig. 20, table 1 (brief description; 4 spec. from Dong-gang; first record from Taiwan); Shinohara *et al.* 2005:418 (listed; Ryukyu Islands); Suetsugu & Ohta 2005: table 3 (listed; Enshu-nada); Shao *et al.* 2008b: table 2 (37

spec. listed from northeastern, southeastern, and southwestern Taiwan, and South China Sea); Furuhashi *et al.* 2010: table 2 (21 spec. listed from northern Okinawa Trough); Nakabo & Kai 2013:498 (in key; Japan); Iwamoto *et al.* 2015:110 (brief description; 45 spec. from northeastern and southwestern Taiwan, and South China Sea); Motomura 2020:39 (listed; Japan).

Diagnosis. Second spinous ray of first dorsal fin finely serrated along its leading edge; prominent black spot on center of first dorsal fin (often indistinct in juveniles and preserved specimens); anal fin uniformly dusky; prominent black spot absent on front of premaxillary ascending process; oral cavity pale; median rostral ridge lacking narrow blackish streak; leading edges of snout, supranarial ridges, and suborbital shelves only faintly marked with black; dorsal parts of trunk and tail dark, sharply demarcated from paler lateral and ventral parts of body (especially prominent in fresh specimens); snout low, slightly protruding beyond upper jaw, preoral length 16–20% HL; no scute-like scale at snout tip; body scales covered with short, moderately erect, needle-like spinules in quincunx order; spinules on scales along second dorsal-fin base not enlarged; body scales small, transverse scale rows below midbase of first dorsal fin 8.5–11.5, below second dorsal-fin origin 7–9.5; longitudinal scales 46–56; area of spinuleless scales posterior to first dorsal-fin base only narrowly developed; outer series of premaxillary teeth slightly enlarged; mandibular teeth arranged in narrow tapered band; suborbital shelf not greatly constricted anteriorly; inner gill rakers on first arch 14–16, outer gill rakers on second arch 13–17; orbit large, greatest diameter 34–39% HL; upper-jaw length 43–48% HL; interorbital space moderately narrow, width 21–26% HL; chin barbel long, length 31–38% HL; pectoral-fin length 55–68% HL.

Material examined. 28 specimens. **Holotype of *Ventrifossa longibarbata*:** BSKU 29494 (49.7 mm HL, 279+ mm TL), Tosa Bay, Japan, 32.9667°N, 133.5333°E, 605 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, cr. 4-Ky, tr. 19, bottom trawl, coll. O. Okamura *et al.*, 21 Dec. 1979. **Paratypes of *V. longibarbata*:** BSKU 34003 (1, 51.7 mm HL, 282+ mm TL), west of Yakushima Island, Okinawa Trough, 30.2083°N, 128.2088°E, 600–620 m, F/V *Yuryo-maru*, No. 8, sta. 3-T2, bottom trawl, coll. H. Maeda, 26 Oct. 1979; BSKU 30495 (1, 28.5 mm HL, 161+ mm TL), BSKU 30496 (1, 41.3 mm HL, 210+ mm TL), collected with holotype. **Non-types:** **Japan:** FFNU P-00770 (1, 40.3 mm HL, 244+ mm TL), east of Okihirazi Bank, East China Sea, 32.1635°N, 129.4663°E, 494 m, T/V *Nagasaki-maru*, cr. N275, sta. B5, 3-m ORE beam trawl, coll. J. Hashimoto, 3 Dec. 2008; BSKU 109021 (1, 26.9 mm HL, 141+ mm TL), west of Okikasayama Bank, 32.1503°N, 129.0072°E, 499–504 m, T/V *Nagasaki-maru*, cr. N365, sta. C5, 3-m ORE beam trawl, coll. N. Nakayama, 19 Nov. 2012;

FIGURE 202. Fresh specimens of *Ventrifossa longibarbata*. (A) BSKU 109021, 26.9 mm HL, 141+ mm TL, west of Okikasayama Bank, Okinawa Trough, East China Sea, 499–504 m depth; (B) BSKU 77922, 35.1 mm HL, 188+ mm TL, south of Muroto, Kochi Pref., Pacific, 414–513 m depth. Lateral views. [Photos: N. Nakayama]

FIGURE 203. Holotype of *Ventrifossa longibarbata*. BSKU 29494, 49.7 mm HL, 279+ mm TL, Tosa Bay, Kochi Pref., Pacific, 605 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

BSKU 28601 (1, 38.9 mm HL, 215+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.9500°N, 128.0000°E, 585–610 m, F/V *Ryoan-maru*, No. 28, tr. 76, bottom trawl, coll. Y. Kinoshita and S. Hagino, 20 Mar. 1978; BSKU 28371 (1, 41.9 mm HL, 211+ mm TL), northwest of Amami-oshima Island, Okinawa Trough,

29.7333°N, 127.8000°E, 495–510 m, F/V *Ryoan-maru*, No. 28, tr. 72, bottom trawl, coll. Y. Kinoshita and S. Hagino, 19 Mar. 1978; BSKU 27185 (1, 37.2 mm HL, 182+ mm TL), BSKU 27192 (1, 47.3 mm HL, 244+ mm TL), BSKU 27194 (1, 44.3 mm HL, 237+ mm TL), northwest of Amami-oshima Island, Okinawa Trough,

FIGURE 204. Preserved juvenile specimen of *Ventrifossa longibarbata*. NSMT-P 92270, 14.4 mm HL, 84+ mm TL, off Miyagi Pref., Pacific. Lateral view. [Photo: N. Nakayama]

29.4333°N, 127.5167°E, 570 m, F/V *Yuryo-maru*, No. 8, tr. 22, bottom trawl, coll. Y. Kinoshita and S. Hagino, 5 Feb. 1978; BSKU 27124 (1, 55.4 mm HL, 294+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.3833°N, 127.5150°E, 650 m, F/V *Yuryo-maru*, No. 8, tr. 21, bottom trawl, coll. Y. Kinoshita and S. Hagino, 5 Feb. 1978; BSKU 28003 (1, 54.7 mm HL, 283+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 28.8333°N, 127.2333°E, 700–740 m, F/V *Ryoan-maru*, No. 28, tr. 59, bottom trawl, coll. Y. Kinoshita and S. Hagino, 16 Mar. 1978; BSKU 27580 (1, 58.2 mm HL, 306+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 28.7167°N, 127.2333°E, 750–755 m, F/V *Ryoan-maru*, No. 28, tr. 48, bottom trawl, coll. Y. Kinoshita and S. Hagino, 11 Mar. 1978; BSKU 77922 (1, 35.1 mm HL, 188+ mm TL), south of Muroto, 33.0790°N, 134.1645°E, 414–513 m, R/V *Tansei-maru*, cr. KT-05-29, sta. K-500, beam trawl, coll. H. Endo *et al.*, 17 Nov. 2005; BSKU 78008 (1, 35.6 mm HL, 190+ mm TL), Tosa Bay, 33.2083°N, 133.6983°E, 526–539 m, FRV *Kotaka-maru*, otter trawl, 3 Mar. 1999; BSKU 44417 (1, 51.9 mm HL, 249+ mm TL), Tosa Bay, 600 m, FRV *Kotaka-maru*, otter trawl, 11 Jan. 1988; BSKU 101609 (1, 48.0 mm HL, 226+ mm TL), BSKU 101618 (1, 51.5 mm HL, 289+ mm TL), BSKU 101619 (1, 47.5 mm HL, 272+ mm TL), BSKU 101620 (1, 34.3 mm HL, 191+ mm TL), BSKU 101623 (1, 35.7 mm HL, 191+ mm TL), Tosa Bay, 33.1428°N, 133.6248°E, 450–480 m, FRV *Kotaka-maru*, otter trawl, 1 Aug. 1988; BSKU 12919 (1, 43.1 mm HL, 247+ mm TL), BSKU 12920 (1, 42.6 mm HL, 222+ mm TL), Tosa Bay, 350–450 m, bottom trawl, 14 May 1968; BSKU 99326 (1, 42.8 mm HL, 265 mm TL), BSKU 99327 (1, 49.5 mm HL, 288+ mm TL), BSKU 99328 (1, 34.7 mm HL, 205+ mm TL), off Kochi, 600 m, bottom trawl, 24 Mar. 1989; *NSMT-P 92270 (1, 14.4 mm HL, 84+ mm TL), off Miyagi, FRV *Wakataka-maru*, sta. E-350, coll. T.P. Satoh, 25 Oct. 2008.

Counts and measurements. Based on 27 specimens (26.9–58.2 mm HL, 141+–306+ mm TL). Counts: first dorsal-fin rays II,9–11; pectoral-fin rays i21–i26; pelvic-

fin rays 8–9; gill rakers on first arch (outer/inner) 9–14/14–16, on second arch 13–17/13–16; longitudinal scales 46–56; transverse scale rows below first dorsal-fin origin 10.5–14.5, below first dorsal-fin midbase 8.5–11.5, below second dorsal-fin origin 7–9.5.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 26–30 (34–39); orbit diameter 34–39 (45–53); postorbital length 39–44 (51–58); postrostral length 74–78; orbit–preopercle distance 37–43 (49–57); suborbital width 12–14 (15–19); upper-jaw length 43–48 (56–64); length of rictus 37–41 (48–56); length of premaxillary tooth band 31–35 (41–47); preoral length 16–20 (20–27); distance between tip and lateral angle of snout 15–18 (20–23); snout width 27–30 (35–40); internasal width 19–23 (25–31); interorbital width 21–26 (28–35); occipital width 13–16 (17–21); body width over pectoral-fin bases 37–60 (49–79); body depth at first dorsal-fin origin 77–97 (99–128); body depth at anal-fin origin 66–88 (85–118); prepelvic length 97–113 (128–153); preanus length 106–123 (141–164); preanal length 123–144 (163–191); isthmus–pelvic distance 41–52 (54–69); isthmus–anus distance 52–61 (68–82); isthmus–anal distance 70–85 (91–113); pelvic–anal distance 28–40 (37–53); anus–anal distance 17–25 (23–33); pelvic-fin length 36–43 (48–57); pectoral-fin length 55–68 (72–90); predorsal length 109–118 (141–158); height of first dorsal fin 72–87 (95–116); length of first dorsal-fin base 24–31 (31–40); interdorsal length 48–80 (62–102); length of gill slit 25–29 (32–38); length of posterior nostril 2–4 (3–6); barbel length 31–38 (40–50).

Size. To about 31 cm TL (BSKU 27580, 306+ mm TL, Okinawa Trough, Japan).

Variation. In a 14.5 mm HL specimen (NSMT-P 92270; Fig. 204), dark spots are serially arranged along the dorsal contour of the trunk and tail. These spots are prominent only when young, and are not found in larger specimens (Figs. 202–203).

Distribution. Restricted to Japan and Taiwan. Known from off the Pacific coasts of Japan northward

FIGURE 205. Scanning electron micrographs showing body scales (from the dorsum below the interdorsal space) of three species of *Ventrifossa*. (A–B) *V. longibarbata*, BSKU 27124, 55.4 mm HL; (C–D) *V. rhipidodorsalis*, BSKU 27146, paratype, 54.5 mm HL; (E–F) *V. saikaiensis*, BSKU 28006, paratype, 51.2 mm HL. (A, C, E) Views from above; (B, D, F) oblique views. [Photos: NSMT]

to Iwaki (Fukushima Pref., 37.00°N), Okinawa Trough, and northern South China Sea, at depths of 350–755 m (Okamura 1982; Chiou *et al.* 2004a; Shao *et al.* 2008a, 2008b; this study; Appendix 3-12E). Uncommon.

Remarks. For further morphological information

see Okamura (1982). *Ventrifossa longibarbata* was originally described from southern Japan, including Tosa Bay (type locality), Suruga Bay, and the Okinawa Trough. One paratype from Suruga Bay, which has been registered at the Zoological Museum of the University of Tokyo

(ZMUT), was not found during the author's visit to the museum in September 2011.

Iwamoto *et al.* (2015:111) revealed slight differences in counts and measurements between Japanese and Taiwanese populations of *V. longibarbata*, viz., the interorbital width [21–26.5% HL in the Japanese specimens they examined (21–26% in this study) vs. 25–31% HL in the Taiwanese specimens they examined], postorbital length [39–44% (39–44%) vs. 42–48%], pectoral-fin rays [i21–i26 (i21–i26) vs. i18–i21], and transverse scale rows [below the first dorsal-fin origin 12–14 (10.5–14.5) vs. 7–12; below the first dorsal-fin midbase 8.5–11.5 (8.5–11.5) vs. 4.5–7.5; and below the second dorsal-fin origin 9–10 (7–9.5) vs. 5–9.5]. These differences may indicate that the Taiwanese population represents a different species.

Relationships and comparisons. *Ventrifossa longibarbata* readily differs from other Japanese species of the subgenus *Ventrifossa* by having higher counts of transverse scale rows below the first dorsal-fin midbase (8.5–11.5 vs. ≤ 7.5). This species is unlikely to be confused with any sympatric congeners when young owing to having prominent dark spots dorsally on the trunk and tail (Fig. 204 vs. absent). It further differs from *V. rhipidodorsalis* Okamura, 1984 in having a larger orbit (34–39% HL vs. 29–35%) and a longer chin barbel (31–38% HL vs. 16–26%); and from *V. saikaiensis* Okamura, 1984 in smaller dimensions of the orbit–preopercle distance (37–43% HL vs. 42–46), snout width (27–30% HL vs. 30–40%), internasal width (19–23% HL vs. 23–30%), and interorbital width (21–26% HL vs. 26–33%). *Ventrifossa longibarbata* appears to be most similar to *V. garmani* (Jordan & Gilbert in Jordan & Starks, 1904), but is distinguished from the latter in having a narrower interorbital space (21–26% HL vs. 27–33%), a longer chin barbel (31–38% HL vs. 20–33%), and needle-like spinules on the body scales (Fig. 205A–B vs. broadly triangular, Fig. 198).

Ventrifossa macroptera Okamura, 1982

[Japanese name: Parao-sokodara]

(Figs. 206–207; Appendix 3-12F)

Ventrifossa macroptera Okamura, 1982:149, fig. 92, table 15 (original description; holotype: BSKU 32185, from Kyushu-Palau Ridge, 27°55.1'N, 134°44.8'E, in 685–710 m; 52 paratypes collected with holotype; new Japanese name: “Parao-sokodara”); Okamura 1984b:93, pl. 81, fig. A (compiled); Okamura 1988:93, pl. 81, fig. A (compiled); Iwamoto 1990:303, fig. 682 (synopsis); Nakabo 1993:358 (in key; Japan); Okamura 1997:128, fig. 12 (compiled); Nakabo 2000:422 (in key; Japan); Nakabo 2002:422 (in key; Japan); Chiou *et al.* 2004a:48, fig. 21, table 1 (brief description; 2 spec. from Dong-gang; first record from Taiwan); Shao *et al.* 2008b: table 2 (89 spec. listed from northeastern and southwestern Taiwan, and South China Sea); Nakabo & Kai 2013:498 (in

key; Japan); Iwamoto *et al.* 2015:111 (brief description; 122 spec. from northeastern and southwestern Taiwan, South China Sea, and Kyushu-Palau Ridge); Motomura 2020:39 (listed; Japan).

Ventrifossa atherodon (not Gilbert & Cramer 1897): Chiou *et al.* 2004a:46, fig. 18, table 1 (brief description; 1 spec. from Daxi; first record from Taiwan) [fide Iwamoto *et al.* (2015)].

Diagnosis. Second spinous ray of first dorsal fin smooth along its leading edge; first dorsal fin blackish; anal fin darker anteriorly; prominent black spot present on front of premaxillary ascending process; oral cavity pale; median rostral ridge lacking narrow blackish streak; leading edges of snout and suborbital shelves prominently marked with black; body dusky brown, with much darker abdomen; scale pockets narrowly outlined in dark; snout low, slightly protruding beyond upper jaw, preoral length 13–17% HL; no scute-like scale at snout tip; body scales covered with short, moderately erect, needle-like spinules in quincunx order; spinules on scales along second dorsal-fin base not enlarged; body scales moderately small, transverse scale rows below midbase of first dorsal fin 5.5–8.5, below second dorsal-fin origin 7–10; longitudinal scales 50–58; area of spinuleless scales posterior to first dorsal-fin base moderately developed; outer series of premaxillary teeth distinctly enlarged; mandibular teeth arranged in 2 irregular rows, with inner row slightly enlarged; suborbital shelf constricted anteriorly; inner gill rakers on first arch 17–19, outer gill rakers on second arch 16–18; orbit in moderate size, greatest diameter 28–33% HL; upper-jaw length 47–51% HL; interorbital space broad, width 29–33% HL; chin barbel moderately long, length 14–26% HL; pectoral fin well developed, length 69–81% HL.

Material examined. 54 specimens. **Holotype of *Ventrifossa macroptera*:** BSKU 32185 (65.8 mm HL, 338+ mm TL), Kyushu-Palau Ridge, Japan, 27.9178°N, 134.7460°E, 685–710 m, F/V *Yuryo-maru*, No. 8, tr. T8, bottom trawl, coll. O. Okamura *et al.*, 19 Dec. 1979. **Paratypes of *V. macroptera*:** *BSKU 29429 (1, 68.3 mm HL, 353+ mm TL), BSKU 32153 (1, 67.5 mm HL, 360 mm TL), *BSKU 32154 (1, 65.9 mm HL, 353+ mm TL), *BSKU 32155 (1, 67.2 mm HL, 349+ mm TL), *BSKU 32156 (1, 67.1 mm HL, 279+ mm TL), *BSKU 32157 (1, 69.9 mm HL, 307+ mm TL), *BSKU 32158 (1, 67.2 mm HL, 348+ mm TL), *BSKU 32159 (1, 68.9 mm HL, 394+ mm TL), BSKU 32160 (1, 70.2 mm HL, 328 mm TL), BSKU 32161 (1, 66.0 mm HL, 341 mm TL), BSKU 32162 (1, 61.7 mm HL, 322 mm TL), *BSKU 32163 (1, 65.2 mm HL, 257+ mm TL), BSKU 32164 (1, 59.9 mm HL, 316 mm TL), *BSKU 32165 (1, 32.2 mm HL, 348+ mm TL), *BSKU 32166 (1, 69.4 mm HL, 378+ mm TL), BSKU 32167 (1, 73.5 mm HL, 401+ mm TL), *BSKU 32168 (1, 71.0 mm HL, 391+ mm TL), BSKU 32169 (1, 68.9 mm HL, 367 mm TL), *BSKU 32170 (1, 71.8 mm HL, 331+ mm TL), *BSKU 32171 (1, 71.0 mm HL, 405+ mm TL), BSKU 32172

FIGURE 206. Fresh specimen of *Ventrifossa macroptera*. BSKU 29429, paratype, 68.3 mm HL, 353+ mm TL, Kyushu-Palau Ridge, Pacific, 685–710 m depth. Lateral view. [Photo: BSKU]

FIGURE 207. Holotype of *Ventrifossa macroptera*. BSKU 32185, 65.8 mm HL, 338+ mm TL, Kyushu-Palau Ridge, Pacific, 685–710 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

(1, 70.0 mm HL, 362 mm TL), BSKU 32173 (1, 70.7 mm HL, 403 mm TL), *BSKU 32174 (1, 67.4 mm HL, 361+ mm TL), BSKU 32175 (1, 64.5 mm HL, 326 mm TL), *BSKU 32176 (1, 60.7 mm HL, 284+ mm TL), *BSKU 32177 (1, 66.1 mm HL, 370+ mm TL), *BSKU 32178 (1, 71.0 mm HL, 367+ mm TL), BSKU 32179 (1, 66.4 mm HL, 330 mm TL), BSKU 32180 (1, 61.1 mm HL, 323 mm TL), BSKU 32181 (1, 63.6 mm HL, 327 mm TL), BSKU 32182 (1, 57.4 mm HL, 308 mm TL), *BSKU 32183 (1, 66.5 mm HL, 368+ mm TL), *BSKU 32184 (1, 67.7 mm HL, 362+ mm TL), *BSKU 32186 (1, 70.4 mm HL, 373+ mm TL), BSKU 32187

(1, 63.7 mm HL, 326 mm TL), BSKU 32188 (1, 59.7 mm HL, 270 mm TL), *BSKU 32189 (1, 62.0 mm HL, 369+ mm TL), BSKU 32192 (1, 56.9 mm HL, 290 mm TL), *BSKU 32193 (1, 58.2 mm HL, 316+ mm TL), BSKU 32194 (1, 51.4 mm HL, 262 mm TL), BSKU 32195 (1, 48.4 mm HL, 276 mm TL), BSKU 32196 (1, 50.8 mm HL, 266 mm TL), *BSKU 32197 (1, 47.9 mm HL, 256+ mm TL), BSKU 32198 (1, 47.6 mm HL, 263 mm TL), *BSKU 32199 (1, 46.1 mm HL, 249+ mm TL), BSKU 32200 (1, 54.4 mm HL, 271 mm TL), collected with holotype; BSKU 30467 (1, 61.9 mm HL, 336 mm TL), *BSKU 30468 (1, 63.5 mm HL, 345+ mm TL),

*BSKU 30469 (1, 62.4 mm HL, 335+ mm TL), Kyushu-Palau Ridge, 27.9000°N, 134.6667°E, 697–700 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. T11, bottom trawl, coll. Y. Kobayashi *et al.*, 18 Dec. 1979; *BSKU 30432 (1, 64.7 mm HL, 303+ mm TL), *BSKU 30435 (1, 69.2 mm HL, 362+ mm TL), BSKU 30436 (1, 69.8 mm HL, 372 mm TL), Kyushu-Palau Ridge, 26.7500°N, 135.3167°E, 336 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. T8, bottom trawl, coll. Y. Kobayashi *et al.*, 17 Dec. 1979. **Non-types: Japan:** BSKU 30470 (1, 73.2 mm HL, 383+ mm TL), Kyushu-Palau Ridge, 27.9000°N, 134.6667°E, 697–700 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. T11, bottom trawl, coll. Y. Kobayashi *et al.*, 18 Dec. 1979.

Counts and measurements. Based on 26 specimens (47.6–73.7 mm HL, 262+–403+ mm TL). Counts: first dorsal-fin rays II,9–11; pectoral-fin rays i20–i25; pelvic-fin rays 9–10; gill rakers on first arch (outer/inner) 11–16/17–19, on second arch 16–18/16–19; longitudinal scales 50–58; transverse scale rows below first dorsal-fin origin 8.5–10.5, below first dorsal-fin midbase 5.5–8.5, below second dorsal-fin origin 7–10, above anal-fin origin 32–39.5.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 27–31 (36–43); orbit diameter 28–33 (37–44); postorbital length 44–49 (59–65); postrostral length 72–77; orbit–preopercle distance 46–50 (59–67); suborbital width 13–16 (18–21); upper-jaw length 47–51 (62–68); length of rictus 40–44 (53–58); length of premaxillary tooth band 36–42 (48–54); preoral length 13–17 (17–22); distance between tip and lateral angle of snout 19–22 (24–29); snout width 29–33 (37–44); internasal width 23–28 (31–37); interorbital width 29–33 (38–46); occipital width 15–18 (20–23); body width over pectoral-fin bases 45–59 (61–77); body depth at first dorsal-fin origin 80–99 (105–132); body depth at

anal-fin origin 76–91 (99–121); prepelvic length 105–117 (140–156); preanus length 113–126 (147–168); preanal length 130–150 (172–200); isthmus–pelvic distance 40–51 (54–69); isthmus–anus distance 51–62 (69–82); isthmus–anal distance 67–88 (90–117); pelvic–anal distance 27–41 (36–54); anus–anal distance 15–29 (21–38); pelvic-fin length 41–54 (57–72); pectoral-fin length 69–81 (92–107); predorsal length 104–112 (138–152); height of first dorsal fin 67–78 (88–104); length of first dorsal-fin base 24–32 (32–43); interdorsal length 42–71 (57–94); length of gill slit 28–32 (38–42); length of posterior nostril 3–6 (4–8); barbel length 14–26 (18–34).

Size. To about 40 cm TL (BSKU 32173, paratype, 403+ mm TL, Kyushu-Palau Ridge, Japan).

Distribution. So far known from the Kyushu-Palau Ridge and the East and South China Seas off Taiwan at depths of 227–1320 m (Chiou *et al.* 2004a, Shao *et al.* 2008a, 2008b; Iwamoto *et al.* 2015; this study; Appendix 3-12F). Common in the Kyushu-Palau Ridge.

Remarks. For a full description see Okamura (1982). *Ventrifossa macroptera* was originally described from 53 specimens collected from the Kyushu-Palau Ridge (Figs. 206–207). In addition to the type locality, it is also known from Taiwan (Chiou *et al.* 2004a; Shao *et al.* 2008a, 2008b; Iwamoto *et al.* 2015). Chiou *et al.* (2004a) included Hawaii in the distribution of *V. macroptera*, but no specimens of this species have been collected from that area. A previous record of *V. atherodon* (Gilbert & Cramer, 1897) from Taiwan (Chiou *et al.* 2004a) represents a misidentification of *V. macroptera* (see Iwamoto *et al.* 2015:112).

Relationships and comparisons. *Ventrifossa macroptera* is the only species of the subgenus *Atherodus* Gilbert & Hubbs, 1920 known from Japan. It readily differs from other Japanese congeners in having the second spinous ray of the first dorsal fin smooth along its leading edge (vs. finely serrated), the outer series of the

FIGURE 208. Scanning electron micrographs showing a body scale (from the dorsum below the interdorsal space) of *Ventrifossa macroptera*. BSKU 32168, paratype, 71.0 mm HL. (A) View from above; (B) oblique view. [Photos: NSMT]

premaxillary tooth band is distinctly enlarged (vs. only slightly enlarged), and the mandibular teeth arranged in 2 irregular rows (vs. tapered band).

In addition to *V. macroptera*, *Atherodus* includes the following three species: *V. atherodon* distributed in the Hawaiian Archipelago and New Caledonian region; *V. macrodon* Sazonov & Iwamoto, 1992 confined to the Sala y Gomez Ridge in the southeastern Pacific; and *V. sazonomi* Iwamoto & Williams, 1999 known from Australia to Taiwan. *Ventrifossa macroptera* is readily separated from the other species of *Atherodus* by having a longer pectoral fin (69–81% HL vs. 51–68% in *V. atherodon*, 49–61% in *V. macrodon*, and 42–65% in *V. sazonomi*). This species is further distinguished from *V. macrodon* and *V. sazonomi* by having a wider interorbital space (29–33% HL vs. 20–24% and 22–28% in *V. macrodon* and *V. sazonomi* respectively) and a longer preoral (13–17% HL vs. 7–11% and 6–12% respectively). In addition, it differs notably from *V. sazonomi* in lacking enlarged spinules on the scales below the anterior portion of the second dorsal-fin base. *Ventrifossa macroptera* also differs from the two *V. atherodon* specimens examined (FMNH 2552 and FMNH 55979, 40.3–63.7 mm HL, 171+–307+ mm TL, from Hawaii) in having lower counts of outer gill rakers on the second arch (16–18 vs. 19) and many morphometric characters, including the snout length (27–31% HL vs. 25–26%), orbit diameter (28–33% HL vs. 34–35%), orbit–preopercle distance (46–50% HL vs. 42–43%), suborbital width (13–16% HL vs. 10–11%), preoral length (13–17% HL vs. 7–10%), snout width (29–33% HL vs. 24–26%), internasal width (23–28% HL vs. 19–20%), and interorbital width (29–33% HL vs. 22–23%). The body color of *V. macroptera* is uniformly dark (Figs. 206–207), whereas the *V. atherodon* specimens examined have prominent silvery reflections on the sides of the body. [Data for *V. atherodon*, *V. macrodon*, and *V. sazonomi* are based on Sazonov & Iwamoto (1992), Iwamoto & Merrett (1997), Iwamoto & Williams (1999), and this study.]

***Ventrifossa misakia* (Jordan & Gilbert in Jordan & Starks, 1904)**

[Japanese name: Misaki-sokodara]

(Figs. 201B–H, 209–212; Table 14; Appendix 3-13A)

Coryphaenoides misakius Jordan & Gilbert in Jordan & Starks, 1904:611, unnumbered fig. (original description; holotype: CAS-SU 8107, from Sagami Bay off Misaki; 10 paratypes from Sagami Bay); Jordan *et al.* 1913:415, fig. 385 (listed; Japan; new Japanese name: “Misaki-sokodara”); Jordan & Thompson 1914:306 (2 spec. from Misaki; FMNH 57091); Kamohara 1938:72 (1 spec. from Kochi Pref.); Kamohara 1950:279 (listed; Kochi and Wakayama Pref.); Matsubara *et al.* 1951:42 (listed; Mie Pref.); Kuroda 1951:391 (listed; Suruga Bay); Kamohara 1952:98 (spec. from Kochi Pref.); Kuroda 1952:176 (1 spec. from Suruga Bay).

Macrourus asper (not Günther 1877): Jordan & Thompson 1914:306, pl. XXXVIII, fig. 2 (2 spec. from Misaki).

Lionurus misakius: Gilbert & Hubbs 1916:194 (new combination; 1 spec. without precise locality data).

Ventrifossa misakia: Okada & Matsubara 1938:452 (in key; Japan); Matsubara 1955:1315 (in key; Japan); Kamohara 1958:74 (listed; Kochi Pref.); Okada *et al.* 1959:83 (listed; Kumano-nada); Kamohara 1964:96 (listed; Kochi Pref.); Matsubara 1965:508 (compiled; Japan); Okamura 1970a:78, pl. VI, text-fig. 35 (description; biological notes; 47 spec. from Pacific off southern Japan from Choshi to Mimase); Okamura 1970b: table 1 (listed; Japan); Kataoka & Tomida 1981:78 (listed; Mie Pref.); Tominaga & Uyeno 1981:489 (listed; Japan); Ohta 1983: table A (listed; Suruga Bay); Akazaki 1984:265 (listed; Aoshima, Miyazaki Pref.); Iwamoto 1990:304, fig. 684 (synopsis); Nakabo 1993:359 (in key; Japan); Shinohara *et al.* 1996:170 (5 spec. listed from Pacific off Tohoku); Sazonov & Shcherbachev 1997:529 (Kyushu-Palau Ridge; *V. fusca* a junior synonym of *V. misakia*); Shinohara & Matsuura 1997:292 (listed; Suruga Bay); Nakabo 2000:423 (in key; Japan); Shinohara *et al.* 2001:306 (1 spec. listed from Tosa Bay); Nakabo 2002:423 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Shinohara *et al.* 2005:418 (listed; Ryukyu Islands); Suetsugu & Ohta 2005: table 3 (listed; Enshu-nada); Senou *et al.* 2006:421 (listed; Sagami Sea); Shinohara & Williams 2006:551, fig. 2-M (listed; Sagami Sea); Kitagawa *et al.* 2008:38, unnumbered fig. (brief description; spec. from Pacific off Tohoku); Shinohara *et al.* 2009:709 (5 spec. listed from Pacific off Tohoku); Nakabo & Kai 2013:499 (in key; Japan); Iwamoto *et al.* 2015:112, fig. 29 (brief description; 6 spec. from South China Sea; other spec. from Japan); Iwatsuki *et al.* 2017:32 (listed; Hyuga-nada); Motomura 2020:39 (listed; Japan).

Coryphaenoides asper (not Günther 1877): Kuroda 1965 (brief description; 1 spec. from Suruga Bay) [re-identified here from description].

Ventrifossa fusca Okamura, 1982:153, fig. 93 (original description; holotype: BSKU 26067, from Kyushu-Palau Ridge, 27°55'N, 134°39'E, in 700 m; 3 paratypes collected with holotype; new Japanese name: “Karasu-hige”); Okamura 1984b:93, pl. 81, fig. B (compiled); Okamura 1988:93, pl. 81, fig. B (compiled); Nakabo 1993:358 (in key; Japan); Okamura 1997:128, fig. 13 (compiled); Nakabo 2000:422 (in key; Japan); Nakabo 2002:422 (in key; Japan); Suetsugu & Ohta 2005: table 3 (listed; Enshu-nada); Senou *et al.* 2006:421 (listed; Sagami Sea); Nakabo & Kai 2013:498 (in key; Japan).

Ventrifossa garmani (not Jordan & Gilbert 1904): Kitagawa *et al.* 2008:39, unnumbered fig. (brief description; spec. from Pacific off Tohoku) [re-identified here from figure].

Diagnosis. Second spinous ray of first dorsal fin weakly serrated along its leading edge; first dorsal and anal fins uniformly dusky to dark; front of premaxillary ascending process usually pale; lining of oral cavity pale; head ridges not marked with black; body uniformly pale to dark; snout moderately high, protruding distinctly beyond

FIGURE 209. Fresh specimens of *Ventrifossa misakia*. (A) BSKU 77921, 22.7 mm HL, 131+ mm TL, south of Muroto, Kochi Pref., Pacific, 414–513 m depth; (B) BSKU 77920, 58.8 mm HL, 274+ mm TL, collected with BSKU 77921; (C) BSKU 83747, 61.0 mm HL, 311+ mm TL, off Mito, Ibaraki Pref., Pacific, 450 m depth; (D) HUMZ 95098, 103 mm HL, 628+ mm TL, west of Amami-oshima Island, Kagoshima Pref., Okinawa Trough, East China Sea, 760–795 m depth. Lateral views. [Photos: (A–B) N. Nakayama; (C) BSKU; (D) HUMZ]

FIGURE 210. Holotype of *Coryphaenoides misakius* (= *Ventrifossa misakia*). CAS-SU 8107, 67.5 mm HL, 334+ mm TL, Misaki, Kanagawa Pref., Sagami Bay, Pacific. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

upper jaw, preoral length 20–26% HL; small, stout, scute-like scale at snout tip; body scales covered with short, moderately erect, needle-like spinules in quincunx order; spinules on scales along second dorsal-fin base not enlarged; body scales very small, transverse scale rows below midbase of first dorsal fin 7.5–11, below second dorsal-fin origin 8–10; longitudinal scales 58–73; no spinuleless scales posterior to first dorsal-fin base; outer series of premaxillary teeth slightly enlarged; mandibular teeth arranged in narrow tapered band; suborbital shelf abruptly narrowing anteriorly; inner gill rakers on first arch 13–16, outer gill rakers on second arch 12–16; orbit large, greatest diameter 29–40% HL; upper-jaw length 37–42% HL; interorbital space broad, width 28–33% HL; chin barbel short, length 3–7% HL; pectoral-fin length 48–64% HL.

Material examined. 36 specimens. **Holotype of *Coryphaenoides misakius*:** CAS-SU 8107 (67.5 mm HL, 334+ mm TL), Misaki, Kanagawa Pref., Sagami Bay, Japan, coll. D.S. Jordan and J.O. Snyder, 1900. **Paratypes of *C. misakius*:** CAS-SU 69810 (4, 59.5–72.6 mm HL, 291+–349+ mm TL, counts and measurements taken on 2 of 4), collected with holotype; USNM 51421 (4, ca. 61–76.2 mm HL, 210+–371+ mm TL), off Misaki, Sagami Bay, hook and line, coll. D.S. Jordan *et al.*, date unknown. **Holotype of *Ventrifossa fusca*:** BSKU 26067 (112 mm HL, 521+ mm TL), Kyushu-Palau Ridge, Japan, 27.9167°N, 134.6500°E, 700 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. 2-3, bottom trawl,

coll. Y. Kobayashi *et al.*, 20 Feb. 1978. **Paratypes of *V. fusca*:** BSKU 26115 (1, 477+ mm TL, heavily damaged), Kyushu-Palau Ridge, 27.9033°N, 134.6583°E, 700 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, sta. 2-4, bottom trawl, coll. Y. Kobayashi *et al.*, 20 Feb. 1978; BSKU 26068 (1, 98.2 mm HL, 485+ mm TL), BSKU 26069 (1, 106 mm HL, 455+ mm TL), collected with holotype. **Non-types: Japan:** HUMZ 95098 (1, 103 mm HL, 628+ mm TL), west of Amami-oshima, Okinawa Trough, 28.5250°N, 128.1567°E, 760–795 m, coll. K. Amaoka, 6 Nov. 1981; BSKU 99072 (1, 25.3 mm HL, 147+ mm TL), Tosa Bay, 33.1950°N, 133.6783°E, 528–537 m, FRV *Kotaka-maru*, otter trawl, 13 May 1997; BSKU 98219 (1, 57.6 mm HL, 279+ mm TL), BSKU 98220 (1, 57.1 mm HL, 289+ mm TL), BSKU 98221 (1, 56.0 mm HL, 268+ mm TL), BSKU 98222 (1, 54.9 mm HL, 262+ mm TL), Tosa Bay, 33.1428°N, 133.6248°E, 450–490 m, FRV *Kotaka-maru*, otter trawl, 1 Aug. 1988; BSKU 44127 (1, 46.4 mm HL, 224+ mm TL), Tosa Bay, 450–500 m, FRV *Kotaka-maru*, otter trawl, 5 Oct. 1987; *BSKU 77920 (1, 58.8 mm HL, 274+ mm TL), *BSKU 77921 (1, 22.7 mm HL, 131+ mm TL), *BSKU 78801 (1, 36.9 mm HL, 171+ mm TL), *BSKU 78807 (1, 21.6 mm HL, 125+ mm TL), south of Muroto, 33.0790°N, 134.1645°E, 414–513 m, R/V *Tansei-maru*, cr. KT-05-29, sta. K-500, beam trawl, coll. H. Endo *et al.*, 17 Nov. 2005; NSMT-P 93115 (1, 61.8 mm HL, 311+ mm TL), NSMT-P 93116 (1, 63.2 mm HL, 303+ mm TL), NSMT-P 93117 (1, 66.3 mm HL, 332+ mm TL), NSMT-P 93118 (1, 55.5 mm HL, 235+ mm TL), off Oh-

FIGURE 211. Fresh specimen of *Ventrifossa misakia*. BSKU 26067, holotype of *V. fusca*, 112 mm HL, 521+ mm TL, Kyushu-Palau Ridge, Pacific, 700 m depth. Lateral view. [Photo: BSKU]

FIGURE 212. Preserved specimen of *Ventrifossa misakia*. BSKU 26067, holotype of *V. fusca*, 112 mm HL, 521+ mm TL, Kyushu-Palau Ridge, Pacific, 700 m depth. (A) Lateral, (B) dorsal, and (C) ventral views. [Photos: N. Nakayama]

sima Island, Izu Islands, R/V *Tansei-maru*, cr. KT-07-31, sta. L-3-500, coll. T. Kuramochi, date unknown; BSKU 83754 (1, 54.6 mm HL, 286+ mm TL), off Fukushima, 450 m, FRV *Wakataka-maru*, sta. F2, coll. H. Endo, 6 Oct. 1997; BSKU 83755 (1, 44.8 mm HL, 226+ mm TL), off Fukushima, 350 m, FRV *Wakataka-maru*, sta. F12, coll. H. Endo, 6 Oct. 1997; BSKU 116162 (1, 37.7 mm HL,

192+ mm TL), off Minami-soma, 37.4878°N, 141.8532°E, 411–419 m, FRV *Wakataka-maru*, sta. 42 (H-400), coll. H. Endo, 3 Nov. 1995; BSKU 83757 (1, 34.1 mm HL, 166+ mm TL), off Onahama, 350 m, FRV *Wakataka-maru*, sta. G, coll. H. Endo, 16 Oct. 1997; BSKU 83734 (1, 56.4 mm HL, 258+ mm TL), BSKU 83741 (1, 58.7 mm HL, 311+ mm TL), BSKU 83744 (1, 55.9 mm HL, 286+ mm TL),

BSKU 83747 (1, 61.0 mm HL, 311+ mm TL), off Mito, 450 m, FRV *Wakataka-maru*, sta. H2, coll. H. Endo, 9 Oct. 1997.

Counts and measurements. See Table 14.

Size. Attains 63 cm TL (HUMZ 95098, 628 mm TL, Okinawa Trough, Japan).

Distribution. Restricted to Japan and Taiwan. Known from off the Pacific coasts of Japan northward to the Oshika Peninsula (38.32°N), Okinawa Trough, northern South China Sea, and Kyushu-Palau Ridge, at depths of 202–795 m (Iwamoto *et al.* 2015; this study; Appendix 3-13A). Common in the south of Choshi (35.74°N; Chiba Pref.), but seemingly rare in other parts of its range.

Remarks. For a full description see Okamura (1970a). *Ventrifossa misakia* was originally described from Sagami Bay, Japan (Jordan & Gilbert in Jordan & Starks 1904; as a species of *Coryphaenoides* Gunnerus, 1765). The original description was based on 11 specimens (Fig. 210), two of which (from *Albatross* sta. 3695) were not found among the CAS and USNM collections.

Records of “*Macrourus asper*” (Günther, 1877) by Jordan & Thompson (1914:306, pl. XXXVIII, fig. 2) and *V. garmani* (Jordan & Gilbert in Jordan & Starks, 1904) by Kitagawa *et al.* (2008:39, unnumbered fig.) can be referred to as *V. misakia*, according to their figures that clearly depict characteristic physiognomy of the latter species. Okamura (1984a) reported *V. misakia* from the Okinawa Trough, but his specimen was re-identified as *V. johnboborum* Iwamoto, 1982 (see the Remarks of the latter species).

Nomenclatural discussion. Okamura (1982) described *V. fusca* based on four large specimens (98.2–112 mm HL, 455+–521+ mm TL; Figs. 211–212) collected from the Kyushu-Palau Ridge, considering this species most similar to “*V. macronema*” (Smith & Radcliffe in Radcliffe, 1912). However, the latter species is currently classified in the genus *Kuronezumia* Iwamoto, 1974 (see Shcherbachev *et al.* 1992), and is quite different from *V. fusca* in many aspects. A close similarity between *V. fusca* and *V. misakia* was first recognized by Iwamoto (1990:305), who wrote that “no sharp differences are apparent” to distinguish the two species. In his key to species of the genus (Iwamoto 1990:295), *V. fusca* is distinguished from *V. misakia* by subtle differences in coloration of the body (dark vs. medium brown) and fins (blackish vs. dusky to pale). He (1990) also noted that *V. fusca* has a scaly patch on the gular membrane (vs. absent in *V. misakia*) and more spinules on the body scales as compared with *V. misakia*. Subsequently, Sazonov & Shcherbachev (1997) considered that these differences are attributable to size-related variations, and do not warrant their specific separation; *V. fusca* was therefore synonymized with *V. misakia* in their study. However, subsequent authors tentatively treated *V. fusca* as a valid species (Iwamoto & Merrett 1997; Nakabo 1993, 2000, 2002; Iwamoto & Williams 1999; Nakabo & Kai 2013).

Most recently, Iwamoto *et al.* (2015) followed Sazonov & Shcherbachev (1997) in regarding this species as a junior synonym of *V. misakia*.

An examination of the type series of *V. misakia* and *V. fusca*, along with many additional specimens from Japan, reconfirmed the conspecificity of these two nominal species. Firstly, all counts and measurements of the type specimens of *V. fusca* generally lie within (or greatly overlap with) the range of variation for *V. misakia* (Table 14); thus, they cannot be distinguished by meristic and morphometric characters. Secondly, as noted by Iwamoto (1990), the body scales of *V. fusca* (Fig. 201F, H) are covered with more spinules as compared with the type specimens of *V. misakia*, but the number of spinules tends to increase with growth in the latter species (Fig. 201 B–E, G). Considering the large body size of the type specimens of *V. fusca* (98.2–112 mm HL, 455+–521+ mm TL), the differences in the scale spinulation can be attributed to ontogenetic variation. A large additional specimen of *V. misakia* collected from the Okinawa Trough (HUMZ 95098, 103 mm HL), which is almost equal in size to the paratypes of *V. fusca*, has similarly numerous spinules on the body scales (Fig. 201G). Thirdly, the holotype of *V. fusca* is readily distinguished from *V. misakia* by having a small scaly patch on the gular membrane, but this feature is not found in the paratypes of this species. The presence/absence of the scales on the gular membrane is likely intraspecific variation. Fourthly, Iwamoto (1990) distinguished *V. fusca* from *V. misakia* in having darker coloration (Fig. 211), but Fig. 209 shows that the body and fins of *V. misakia* become darker with increasing size. Sazonov & Shcherbachev (1997) also reported that an additional specimen from the Kyushu-Palau Ridge (type locality of *V. fusca*) had significantly paler color than the type specimens of this species. Therefore, the differences in coloration can be attributed to ontogenetic or geographical variation. Finally, the holotype of *V. fusca* differs from *V. misakia* in that the mandibular teeth are arranged in two distinct series laterally (vs. 2–3 irregular series or a narrow band), with the inner series distinctly enlarged (Okamura 1982: fig. 93B vs. slightly enlarged). In their keys to species of Japanese grenadiers, Nakabo (1993, 2000, 2002) and Nakabo & Kai (2013) separated the two species based on this character. However, in the paratypes of *V. fusca*, the mandibular teeth are arranged in 2–3 irregular series, and the size of the teeth becomes progressively larger inwardly. The unique dentition of the holotype of *V. fusca* is herein referred to individual variation. This difference might be attributable to the larger body size of this specimen, which is by far the largest among the available materials for the two nominal species. Accordingly, there are no reasons to doubt their conspecificity, and *V. fusca* is re-confirmed here as a junior synonym of *V. misakia*.

Relationships and comparisons. According to Sazonov & Iwamoto (1992:80), *V. misakia* is most closely

related to *V. johnboborum* Iwamoto, 1982. These species are distinguished from each other by the combination of the barbel length, internasal width, and coloration of the oral cavity (see the Relationships and comparisons of the latter species). They differ from other Japanese species of *Ventrifossa* in having a small but prominent scute on the tip of the snout (vs. absent), smaller body scales (longitudinal scales ≥ 58 vs. ≤ 56), a shorter barbel ($\leq 16\%$ HL (usually $\leq 12\%$) vs. $\geq 16\%$), and a longer preoral ($\geq 19\%$ HL (usually $\geq 20\%$) vs. $\leq 22\%$ (usually $\leq 19\%$)). The two species also differ from the other Japanese congeners in that the suborbital shelf is abruptly and greatly constricted anteriorly (vs. moderately deep).

***Ventrifossa nigrodorsalis* Gilbert & Hubbs, 1920**

[No Japanese name]

(Fig. 213; Appendix 3-13B)

Ventrifossa nigrodorsalis Gilbert & Hubbs, 1920:546, fig. 36 [original description; holotype: USNM 83627, from off northern coast of Mindanao, *Albatross* sta. 5502, 8°37'37"N, 124°35'E, in 214 ftn (393 m); numerous paratypes from near Philippines and Indonesia]; Chen 1951:210 (listed; Taiwan); Iwamoto 1990:307, fig. 690 (synopsis); Shao 1993:172 (compiled; Taiwan); Chiou *et al.* 2004a: table 1 (listed; Taiwan); Shao *et al.* 2008b: table 2 (2 spec. listed from northeastern Taiwan and South China Sea); Ho & Shao 2011:36 (type catalog); Iwamoto *et al.* 2015:114 [brief description; 14 spec. from northeastern Taiwan and Philippines (including holotype)].

Diagnosis. Second spinous ray of first dorsal fin finely serrated along its leading edge; first dorsal fin with elongate black spot; anal fin pale, without narrow blackish margin anteriorly; front of premaxillary ascending process

pale; oral cavity pale; median rostral ridge lacking narrow blackish streak; leading edges of snout and suborbital shelves slightly darker in preserved specimens; dorsum not strongly demarcated in color from rest of body; snout low, slightly protruding beyond upper jaw, preoral length 19–22% HL; no scute-like scale at snout tip; body scales covered with short, moderately erect, needle-like spinules in quincunx order; spinules on scales along second dorsal-fin base not enlarged; body scales moderately small, transverse scale rows below midbase of first dorsal fin 6.5, below second dorsal-fin origin 6–7; longitudinal scales 37–41; area of spinuleless scales posterior to first dorsal-fin base narrowly developed; outer series of premaxillary teeth slightly enlarged; mandibular teeth arranged in narrow tapered band; suborbital shelf not greatly constricted anteriorly; inner gill rakers on first arch 13–15, outer gill rakers on second arch 13–14; orbit large, greatest diameter 32–35% HL; upper-jaw length 38–40% HL; interorbital space broad, width 27–29% HL; chin barbel moderately long, length 22–25% HL; pectoral-fin length 60–61% HL.

Material examined. 5 specimens. **Taiwan:** *FAKU 204466 (1, 28.7 mm HL, 182+ mm TL), Dong-gang fish market, Pingtung, bottom trawl, coll. N. Nakayama, 27 Feb. 2017. **South China Sea:** BSKU 108942 (1, 26.9 mm HL, 109+ mm TL), BSKU 108944 (1, 25.3 mm HL, 143+ mm TL), BSKU 108950 (1, 30.5 mm HL, 162+ mm TL), off Swatow, 21.7000°N, 117.5233°E, 412–430 m, R/V *Hakuho-maru*, cr. KH-73-02, sta. 44-2, 3-m beam trawl, 19 Mar. 1973; ASIZP 66263 (1, 37.5 mm HL, 202+ mm TL), 21.5833°N, 117.2833°E, 364 m, otter trawl, coll. P.-F. Lee, 17 Aug. 2005.

Counts and measurements. Based on 4 specimens (25.3–37.5 mm HL, 109+–202+ mm TL). Counts: first dorsal-fin rays II,9–10; pectoral-fin rays i19–i22; pelvic-fin rays 8–9; gill rakers on first arch (outer/inner) 10–

FIGURE 213. Fresh specimen of *Ventrifossa nigrodorsalis*. FAKU 204466, 28.7 mm HL, 182+ mm TL, Dong-gang fish market, southwestern Taiwan, South China Sea. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. [Photos: N. Nakayama]

13/13–15, on second arch 13–14/12–13; longitudinal scales 37–41; transverse scale rows below first dorsal-fin origin 7.5–8.5, below first dorsal-fin midbase 6.5, below second dorsal-fin origin 6–7.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 28–30 (36–40); orbit diameter 32–35 (43–46); postorbital length 41–44 (55–58); postrostral length 74–77; orbit–preopercle distance 39–42 (51–55); suborbital width 12–15 (16–19); upper-jaw length 38–40 (50–53); length of rictus 33–37 (43–49); length of premaxillary tooth band 26–29 (35–38); preoral length 19–22 (24–28); distance between tip and lateral angle of snout 15–18 (20–23); snout width 28–31 (36–41); internasal width 22–24 (28–31); interorbital width 27–29 (36–37); occipital width 15 (20); body width over pectoral-fin bases 42–46 (56–61); body depth at first dorsal-fin origin 78–93 (105–122); body depth at anal-fin origin 71–87 (95–113); prepelvic length 107–114 (142–149); preanus length 115–128 (151–168); preanal length 126–142 (165–185); isthmus–pelvic distance 43–50 (57–65); isthmus–anus distance 50–61 (66–80); isthmus–anal distance 64–77 (84–104); pelvic–anal distance 23–33 (29–45); anus–anal distance 12–24 (16–32); pelvic-fin length 36–40 (47–54); pectoral-fin length 60–61 (79–82); predorsal length 113–117 (150–153); height of first dorsal fin 26–94 (34–127); length of first dorsal-fin base 25–29 (34–38); interdorsal length 55–63 (73–83); length of gill slit 22–23 (28–30); length of posterior nostril 5–7 (7–9); barbel length 22–25 (29–32).

Size. To about 34 cm TL (Merrett & Iwamoto 2000; MNHN 1995-903, 251+–344+ mm TL, Vanuatu).

Distribution. Widely known from Western Australia, eastward to the New Caledonian region, and northward to the East China Sea off Taiwan, at depths of about 300–800 m (Gilbert & Hubbs 1920; Iwamoto & Merrett 1997; Iwamoto & Williams 1999; Merrett & Iwamoto 2000; Iwamoto & Graham 2001; Shao *et al.* 2008a, 2008b; Last *et al.* 2014). No specimens have been collected within Japan's EEZ (Appendix 3-13B).

Remarks. *Ventrifossa nigrodorsalis* is a widespread species, with high abundance in tropical waters of the western Pacific (Merrett & Iwamoto 2000; Iwamoto *et al.* 2015). For further morphological information see the original description (Gilbert & Hubbs 1920), Iwamoto & Merrett (1997), and Iwamoto & Williams (1999). It has not been reported from Japanese waters, but is known from the East China Sea off Taiwan (Shao *et al.* 2008a, 2008b; Iwamoto *et al.* 2015). Although Okamura (1982) reported this species from Japan based on two specimens collected from Tosa Bay, these specimens were subsequently allocated as paratypes of *V. rhipidodorsalis* Okamura, 1984. Iwamoto & Merrett (1997:561) highlighted marked color variation between three populations within the New Caledonian region, viz., the first group from the north of Fiji, the second from the Chesterfield and Bellona Plateaus, and the last from off New Caledonia (see also Iwamoto

& Merrett 1997:553). Iwamoto *et al.* (2015:115) wrote: “the existence of seemingly widely scattered populations of *V. nigrodorsalis* heighten the possibility that other species lie hidden under the cloak of that species name as currently circumscribed, and they may be unveiled after closer study of more specimens”.

Relationships and comparisons. *Ventrifossa nigrodorsalis* is closely related to *V. petersonii* (Alcock in Wood-Mason & Alcock, 1891), an Indian Ocean species also characterized by the presence of a prominent black spot on the first dorsal fin. The latter species differs most notably from *V. nigrodorsalis* in that an area of spinuleless scales is broadly developed posterior to the first dorsal-fin base (vs. absent or only narrowly developed). [Data for *V. petersonii* are based on 2 spec. collected from the Timor Sea (BSKU 16714 and BSKU 16715, 33.7–33.9 mm HL).]

***Ventrifossa rhipidodorsalis* Okamura, 1984**

[Japanese name: Homae-hige]

(Figs. 205C–D, 214–215; Appendix 3-13C)

Ventrifossa nigrodorsalis (not Gilbert & Hubbs 1920): Okamura 1982:147, 348, fig. 90 [first record from Japan; new Japanese name: “Homae-hige”; brief description; 2 spec. from Tosa Bay; photo based on BSKU 29495 (paratype of *V. rhipidodorsalis*)]; Ohta 1983: table A (listed; Suruga Bay).

Ventrifossa rhipidodorsalis Okamura, 1984a:205, fig. 146 (original description; holotype: BSKU 27695, from Okinawa Trough, 28°42.0'N, 127°09.0'E, in 500–535 m; 23 paratypes from Okinawa Trough and Tosa Bay, in 220–650 m); Okamura 1984b:93, pl. 81, fig. D (compiled); Okamura 1988:93, pl. 81, fig. D (compiled); Nakabo 1993:358 (in key; Japan); Okamura 1997:128, fig. 11 (compiled); Nakabo 2000:422 (in key; Japan); Shinohara *et al.* 2001:306 (listed; Tosa Bay); Nakabo 2002:422 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Chiou *et al.* 2004a:48, fig. 22, table 1 (brief description; 13 spec. from Nan-fang-ao; first record from Taiwan); Shinohara *et al.* 2005:418 (listed; Ryukyu Islands); Suetsugu & Ohta 2005: table 3 (listed; Enshu-nada); Shao *et al.* 2008b: table 2 (53 spec. listed from northeastern, southeastern, and southwestern Taiwan, and South China Sea); Furuhashi *et al.* 2010: table 2 (2 spec. listed from northern Okinawa Trough); Nakabo & Kai 2013:498 (in key; Japan); Iwamoto *et al.* 2015:115 [brief description; 82 spec. from northeastern Taiwan, Japan (including holotype), and Philippines]; Nakae *et al.* 2018:219 (1 spec. listed from Amami-oshima Island, Kagoshima Pref.); Motomura 2020:40 (listed; Japan).

Ventrifossa longibarbata (not Okamura 1984a): Okamura 1997:128, fig. 10 [in part, photo based on BSKU spec. (catalog no. unknown)].

Diagnosis. Second spinous ray of first dorsal fin finely serrated along its leading edge; large intense black spot on center of first dorsal fin; anal fin dusky, with narrow

FIGURE 214. Fresh specimens of *Ventrifossa rhipidodorsalis*. (A) BSKU 109208, 46.0 mm HL, 196+ mm TL, off Amami-oshima Island, Kagoshima Pref., Okinawa Trough, East China Sea, 607 m depth; (B) BSKU 45012, 48.4 mm HL, 276+ mm TL, Tosa Bay, Kochi Pref., Pacific, 500 m depth. Lateral views. [Photos: (A) N. Nakayama; (B) BSKU]

blackish margin anteriorly; front of premaxillary ascending process pale; oral cavity pale; median rostral ridge lacking narrow blackish streak; leading edges of snout, supranarial ridges, and suborbital shelves prominently marked with black; dorsum not strongly demarcated in color from rest of body; snout low, slightly protruding beyond upper jaw, preoral length 11–19% HL; no scute-like scale at snout tip; body scales covered with short, moderately erect, needle-like spinules in quincunx order; spinules on scales along second dorsal-fin base not enlarged; body scales moderately small, transverse scale rows below midbase of first dorsal fin 6.5–7.5, below second dorsal-fin origin 6.5–8; longitudinal scales 37–49; area of spinuleless scales posterior to first dorsal-fin base only narrowly developed; outer series of premaxillary teeth slightly enlarged; mandibular teeth arranged in narrow tapered band; suborbital shelf not greatly constricted anteriorly; inner gill rakers on first arch 15–17, outer gill rakers on second arch 14–17; orbit large, greatest diameter 29–35% HL; upper-jaw length 42–47% HL; interorbital space moderately narrow, width 20–27% HL; chin barbel moderately long, length 16–26% HL; pectoral-fin length 57–70% HL.

Material examined. 37 specimens. **Holotype of *Ventrifossa rhipidodorsalis*:** BSKU 27695 (62.2 mm HL, 328+ mm TL), west-northwest of Amami-oshima Island, Kagoshima Pref., Japan, Okinawa Trough, East China Sea, 28.7000°N, 127.1500°E, 500–535 m, F/V *Ryoan-maru*, No. 28, tr. 50, bottom trawl, coll. Y. Kinoshita and S. Hagino, 11 Mar. 1978. **Paratypes of *V. rhipidodorsalis*:** *BSKU 29571 (1, 57.2 mm HL, 315+ mm TL), northwest of Miyako-jima Island, Okinawa Trough, 25.7900°N, 124.3900°E, 600 m, F/V *Yuryo-maru*, No. 8, sta. 1-T13, bottom trawl, coll. T. Kitajima, 20 Sept. 1979; *BSKU 28599 (1, ca. 41 mm HL, 241+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.9500°N, 128.0000°E, 585–610 m, F/V *Ryoan-maru*, No. 28, tr. 76, bottom trawl, coll. Y. Kinoshita and S. Hagino, 20 Mar. 1978; BSKU 27184 (1, 51.7 mm HL, 269+ mm TL), BSKU 27190 (1, 32.8 mm HL, 187+ mm TL), BSKU 27191 (1, 40.7 mm HL, 225+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 29.4333°N, 127.5167°E, 570 m, F/V *Yuryo-maru*, No. 8, tr. 22, bottom trawl, coll. Y. Kinoshita and S. Hagino, 5 Feb. 1978; BSKU 27123 (1, 50.1 mm HL, 265+ mm TL), BSKU 27146 (1, 54.2 mm HL, 209+ mm TL), northwest of Amami-oshima

FIGURE 215. Holotype of *Ventrifossa rhipidodorsalis*. BSKU 27695, 62.2 mm HL, 328+ mm TL, west-northwest of Amami-oshima Island, Kagoshima Pref., Okinawa Trough, East China Sea, 500–535 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

Island, Okinawa Trough, 29.3833°N, 127.5150°E, 650 m, F/V *Yuryo-maru*, No. 8, tr. 21, bottom trawl, coll. Y. Kinoshita and S. Hagino, 5 Feb. 1978; *BSKU 27915 (1, 54.0 mm HL, 229+ mm TL), *BSKU 27916 (1, 46.6 mm HL, 246+ mm TL), BSKU 27917 (1, 30.2 mm HL, 171+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 28.7500°N, 127.1167°E, 520–542 m, F/V *Ryoan-maru*, No. 28, tr. 58, bottom trawl, coll. Y. Kinoshita and S. Hagino, 16 Mar. 1978; BSKU 27660 (1, 58.6 mm HL, 313+ mm TL), BSKU 27661 (1, 54.1 mm HL, 226+ mm TL), BSKU 27663 (1, 51.7 mm HL, 264+ mm TL), west of Okinawa-jima Island, Okinawa Trough, 28.7333°N, 127.0167°E, 610–640 m, F/V *Ryoan-maru*, No. 28, tr. 49, bottom trawl, coll. Y. Kinoshita and S. Hagino, 11 Mar. 1978; BSKU 27694 (1, 66.6 mm HL, 354+ mm TL), BSKU 27696 (1, 47.7 mm HL, 262+ mm TL), *BSKU 27698 (1, 34.3 mm HL, 191+ mm TL), *BSKU 29697 (1, 34.6 mm HL, 201+ mm TL), collected with holotype; BSKU 27858 (1, 42.6 mm HL, 172+ mm TL), BSKU 27861 (1, 58.8 mm HL, 268+ mm TL), *BSKU 27862 (1, 46.3 mm HL, 259+ mm TL), BSKU 27865 (1, 52.8 mm HL, 262+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 28.5833°N, 127.0000°E, 600 m, F/V *Ryoan-maru*, No. 28, tr. 57, bottom trawl, coll. Y. Kinoshita and S. Hagino, 16 Mar. 1978; BSKU 30498 (1, 30.8 mm HL, 156+ mm TL), Tosa Bay, 32.9667°N, 133.5333°E, 605 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, bottom trawl, coll. O. Okamura *et al.*, 21 Dec. 1979; BSKU 29495 (1, 28.6 mm HL, 168 mm TL), Tosa Bay, 32.9667°N, 133.5333°E, 605 m, F/Vs *Shinsei-maru*, No. 53 and *Kyoyo-maru*, No. 2, cr. 4-Ky, tr. 19, bottom trawl, coll. O. Okamura *et al.*, 21 Dec. 1979. **Non-types:**

Japan: BSKU 109208 (1, 46.0 mm HL, 196+ mm TL), off Amami, East China Sea, 28.5679°N, 127.0421°E, 607 m, T/V *Nagasaki-maru*, cr. N365, sta. J6, tr. 1, 3-m ORE beam trawl, coll. N. Nakayama, 16 Nov. 2012; BSKU 101640 (1, 48.9 mm HL, 282+ mm TL), BSKU 101641 (1, 48.7 mm HL, 270+ mm TL), off Muroto, Tosa Bay, 360–435 m, bottom trawl, 9 Jun. 1968; BSKU 45012 (1, 48.4 mm HL, 276+ mm TL), BSKU 45013 (1, 44.9 mm HL, 247+ mm TL), Tosa Bay, 500 m, FRV *Kotaka-maru*, otter trawl, 23 Aug. 1988; BSKU 48169 (1, 61.1 mm HL, 343+ mm TL), BSKU 99329 (1, 51.7 mm HL, 293+ mm TL), BSKU 99330 (1, 38.6 mm HL, 212+ mm TL), BSKU 99331 (1, 43.0 mm HL, 255+ mm TL), BSKU 99332 (1, 54.4 mm HL, 313+ mm TL), BSKU 99334 (1, 54.7 mm HL, 308+ mm TL), BSKU 99335 (1, 63.4 mm HL, 336+ mm TL), BSKU 99338 (1, 48.6 mm HL, 269+ mm TL), off Kochi, 600 m, bottom trawl, 24 Mar. 1989.

Counts and measurements. Based on 30 specimens (29–66.1 mm HL, 158+–353+ mm TL). Counts: first dorsal-fin rays II,10–12; pectoral-fin rays i19–i25; pelvic-fin rays 8–10; gill rakers on first arch (outer/inner) 9–16/15–17, on second arch 14–17/14–18; longitudinal scales 37–49; transverse scale rows below first dorsal-fin origin 7.5–11.5, below first dorsal-fin midbase 6–7.5, below second dorsal-fin origin 6.5–8.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 26–31 (34–43); orbit diameter 29–35 (38–46); postorbital length 41–48 (54–64); postrostral length 72–79; orbit–preopercle distance 39–44 (52–60); suborbital width 11–14 (14–19); upper-jaw length 42–47 (56–62); length of rictus 36–40 (48–54); length of premaxillary

tooth band 31–35 (40–47); preoral length 11–19 (15–25); distance between tip and lateral angle of snout 13–19 (18–26); snout width 23–31 (30–42); internasal width 18–22 (23–30); interorbital width 20–27 (27–35); occipital width 12–16 (16–21); body width over pectoral-fin bases 37–65 (50–84); body depth at first dorsal-fin origin 76–101 (100–130); body depth at anal-fin origin 64–86 (86–114); prepelvic length 99–117 (127–156); preanus length 106–128 (135–170); preanal length 130–149 (166–198); isthmus–pelvic distance 42–55 (57–73); isthmus–anus distance 51–63 (68–83); isthmus–anal distance 69–87 (93–115); pelvic–anal distance 29–43 (40–58); anus–anal distance 18–30 (25–40); pelvic-fin length 36–47 (45–62); pectoral-fin length 57–70 (75–93); predorsal length 107–119 (141–157); height of first dorsal fin 71–84 (95–111); length of first dorsal-fin base 25–33 (34–44); interdorsal length 43–64 (58–85); length of gill slit 22–29 (30–38); length of posterior nostril 3–5 (3–7); barbel length 16–26 (21–35).

Size. To about 35 cm TL (BSKU 27694, 353+ mm TL, paratype, Okinawa Trough, Japan).

Distribution. Distributed in the western Pacific from Japan to the Philippines (Chiou *et al.* 2004a; Shao *et al.* 2008a, 2008b; Iwamoto *et al.* 2015; this study); In Japanese waters, known from off the Pacific coasts northward to the Izu Peninsula (138.96°E) and the Okinawa Trough in the East China Sea (Appendix 3-13C). Depth range 315–1165 m. Uncommon.

Remarks. *Ventrifossa rhipidodorsalis* was originally described from the Okinawa Trough (type locality) and Tosa Bay by Okamura (1984a), to which the reader can refer for further morphological information. The two paratypes from Tosa Bay were initially reported as *V. nigrodorsalis* Gilbert & Hubbs, 1920 by Okamura (1982).

Relationships and comparisons. Among northwestern Pacific congeners, this species is most similar to *V. nigrodorsalis*, but readily differs from the latter in having a black anterodistal margin of the anal fin (vs. uniformly pale), higher counts of gill rakers (15–17 vs. 13–15 on the inner side of the first arch; 14–17 vs. 13–14 on the outer side of the second arch; and 14–18 vs. 12–13 on the inner side of the second arch), a larger mouth (upper-jaw length 42–47% HL vs. 38–40%), and a narrower interorbital space (20–27% HL vs. 27–29%).

Ventrifossa saikaiensis Okamura, 1984

[Japanese name: Saikai-hige]

(Figs. 205E–F, 216; Appendix 3-13D)

Ventrifossa saikaiensis Okamura, 1984a:209, fig. 147 (original description; holotype: BSKU 28004, from Okinawa Trough, 28°50.0'N, 127°14.0'E, in 700–740 m; 11 paratypes from Okinawa Trough, in 560–755 m; new Japanese name: “Saikai-hige”); Okamura 1988:453, pl. 373, fig. F (compiled);

Nakabo 1993:359 (in key; Japan); Nakabo 2000:423 (in key; Japan); Nakabo 2002:423 (in key; Japan); Yoda *et al.* 2002:11 (listed; East China and Yellow Seas); Chiou *et al.* 2004a:48, fig. 23, table 1 (brief description; 19 spec. from Da-xi; first record from Taiwan); Shinohara *et al.* 2005:418 (listed; Ryukyu Islands); Shao *et al.* 2008b: table 2 (46 spec. listed from northeastern and southwestern Taiwan, and South China Sea); Nakabo & Kai 2013:499 (in key; Japan); Iwamoto *et al.* 2015:116 (brief description; 64 spec. from northeastern Taiwan and East China Sea off Japan, including holotype); Motomura 2020:40 (listed; Japan).

Ventrifossa divergens (not Gilbert & Hubbs 1920): Chiou *et al.* 2004a:46, fig. 19, table 1 (brief description; 2 spec. from Dong-gang and Da-xi; first record from Taiwan) [fide Iwamoto *et al.* (2015)].

Diagnosis. Second spinous ray of first dorsal fin finely serrated along its leading edge; first dorsal and anal fins uniformly dusky; front of premaxillary ascending process lacking prominent black spot; oral cavity pale; median rostral ridge, leading edges of snout, supranarial ridges, and suborbital shelves not prominently blackish; body uniformly brown, except for dark abdomen; scale pockets narrowly outlined in dark; snout low, slightly protruding beyond upper jaw, preoral length 13–21% HL; no scute-like scale at snout tip; body scales covered with short, moderately erect, needle-like spinules in quincunx order; spinules on scales along second dorsal-fin base not enlarged; body scales moderately small, transverse scale rows below midbase of first dorsal fin 6–7.5, below second dorsal-fin origin 6–7.5; longitudinal scales 37–45; area of spinuleless scales posterior to first dorsal-fin base only narrowly developed; outer series of premaxillary teeth slightly enlarged; mandibular teeth arranged in narrow tapered band; suborbital shelf not greatly constricted anteriorly; inner gill rakers on first arch 15–19, outer gill rakers on second arch 15–17; orbit large, greatest diameter 28–38% HL; suborbital width 13–17% HL; upper-jaw length 45–52% HL; interorbital space broad, width 26–33% HL; chin barbel long, length 30–38% HL; pectoral-fin length 59–72% HL.

Material examined. 20 specimens. **Holotype of *Ventrifossa saikaiensis*:** BSKU 28004 (54.7 mm HL, 307+ mm TL), northwest of Amami-oshima Island, Kagoshima Pref., Japan, Okinawa Trough, East China Sea, 28.8333°N, 127.2333°E, 700–740 m, F/V *Ryoan-maru*, No. 28, tr. 59, bottom trawl, coll. Y. Kinoshita and S. Hagino, 16 Mar. 1978. **Paratypes of *V. saikaiensis*:** BSKU 26324 (1, 37.4 mm HL, 209+ mm TL), off Senkaku Islands, Okinawa Trough, 25.6283°N, 122.8933°E, 560–692 m, F/V *Yuryo-maru*, No. 8, tr. 3, bottom trawl, coll. T. Kitajima *et al.*, 14 Jan. 1978; *BSKU 28005 (1, 41.4 mm HL, 184+ mm TL), BSKU 28006 (1, 51.5 mm HL, 249+ mm TL), BSKU 28007 (1, 54.9 mm HL, 207+ mm TL), BSKU 28008 (1, 53.0 mm HL, 293+ mm TL), collected with holotype; *BSKU 27664 (1, 36.2 mm HL, 198+ mm

FIGURE 216. Holotype of *Ventrifossa saikaiensis*. BSKU 28004, 54.7 mm HL, 307+ mm TL, northwest of Amami-oshima Island, Kagoshima Pref., Okinawa Trough, East China Sea, 700–740 m depth. (A) Lateral view; (B) dorsal and (C) ventral views of the head and trunk. Preserved condition. [Photos: N. Nakayama]

TL), *BSKU 27665 (1, 42.1 mm HL, 200+ mm TL), west of Okinawa-jima Island, Okinawa Trough, 28.7333°N, 127.0167°E, 610–640 m, F/V *Ryoan-maru*, No. 28, tr. 49, bottom trawl, coll. Y. Kinoshita and S. Hagino, 11 Mar. 1978; BSKU 27576 (1, 57.2 mm HL, 287+ mm TL), BSKU 27577 (1, 50.4 mm HL, 222+ mm TL), *BSKU 27578 (1, 52.2 mm HL, 240+ mm TL), *BSKU 27579 (1, 52.6 mm HL, 294+ mm TL), northwest of Amami-oshima Island, Okinawa Trough, 28.7167°N, 127.2333°E, 750–755 m, F/V *Ryoan-maru*, No. 28, tr. 48, bottom trawl, coll. Y. Kinoshita and S. Hagino, 11 Mar. 1978. **Non-types: Japan:** BSKU 26325 (1, 53.7 mm HL, 280+ mm TL), off Senkaku Islands, Okinawa Trough, 25.6283°N, 122.8933°E, 560–692 m, F/V *Yuryo-maru*, No. 8, tr. 3, bottom trawl, coll. T. Kitajima *et al.*, 14 Jan. 1978; FFNU P-00738 (8, 24.7–47.7 mm HL, 155+–265+ mm TL), north of Hokutosekibi Bank, East China Sea, 26.4928°N, 125.1432°E, 675 m, T/V *Nagasaki-maru*, cr. N260, sta. S, 3-m ORE beam trawl, coll. J. Hashimoto, 21 Jun. 2008.

Counts and measurements. Based on 16 specimens (27.4–57.4 mm HL, 155+–307+ mm TL). Counts: first dorsal-fin rays II,9–11; pectoral-fin rays i18–i23; pelvic-fin rays 8–9; gill rakers on first arch (outer/inner) 10–14/15–19, on second arch 15–17/14–18; longitudinal scales 37–45; transverse scale rows below first dorsal-fin origin 7.5–10, below first dorsal-fin midbase 6–7.5, below second dorsal-fin origin 6–7.5.

The following measurements are in % of HL, followed by those in % of PRL in parentheses: snout length 27–33 (35–43); orbit diameter 28–38 (39–49); postorbital length 41–49 (54–63); postrostral length 72–80; orbit–preopercle distance 42–46 (54–63); suborbital width 13–17 (17–23); upper-jaw length 45–52 (59–69);

length of rictus 39–43 (51–58); length of premaxillary tooth band 33–39 (44–51); preoral length 13–21 (18–29); distance between tip and lateral angle of snout 16–22 (21–28); snout width 30–40 (39–52); internasal width 23–30 (30–39); interorbital width 26–33 (34–44); occipital width 15–18 (20–24); body width over pectoral-fin bases 39–60 (50–80); body depth at first dorsal-fin origin 82–97 (107–125); body depth at anal-fin origin 65–83 (86–106); prepelvic length 103–133 (134–171); preanus length 113–142 (146–178); preanal length 132–161 (169–203); isthmus–pelvic distance 39–48 (50–65); isthmus–anus distance 47–56 (61–74); isthmus–anal distance 70–77 (92–103); pelvic–anal distance 27–35 (35–47); anus–anal distance 18–25 (24–32); pelvic-fin length 37–40 (48–52); pectoral-fin length 59–72 (75–94); predorsal length 93–116 (121–155); length of first dorsal-fin base 25–32 (33–43); interdorsal length 45–63 (58–87); length of gill slit 26–35 (34–46); length of posterior nostril 2–6 (3–8); barbel length 30–38 (39–49).

Size. To about 31 cm TL (BSKU 28004, holotype).

Distribution. So far known only from the Okinawa Trough and the northern South China Sea at depths of 520–764 m (Chiou *et al.* 2004a; Shao *et al.* 2008b; this study; Appendix 3-13D). Rare in Japanese waters.

Remarks. For further morphological information see the original description given by Okamura (1984a). This species was originally described from the Okinawa Trough in the East China Sea, and additional specimens were subsequently recorded from Taiwan (Chiou *et al.* 2004a; Shao *et al.* 2008b; Iwamoto *et al.* 2015). Chiou *et al.* (2004a) first recorded *V. divergens* Gilbert & Hubbs, 1920 from Taiwanese waters based on two specimens obtained at fish markets in Dong-gang and Da-xi, but

their specimens were re-identified as *V. saikaiensis* by Iwamoto *et al.* (2015).

Relationships and comparisons. *Ventrifossa saikaiensis* closely resembles *V. divergens* distributed in the western Pacific from Taiwan to Indonesia. According to Iwamoto *et al.* (2015:117), the two species are readily distinguished by coloration: the body of *V. saikaiensis* is uniformly brownish, whereas in *V. divergens*, silvery reflections are present on the sides of the body. In Japanese waters, *V. saikaiensis* is most similar to *V. garmani* (Jordan & Gilbert in Jordan & Starks, 1904) in having a broad interorbital space (26–33% HL). However, it readily differs from the latter species in having needle-like spinules on the body scales (vs. broadly triangular), a wider snout (snout width 30–40% HL vs. 25–29%), and a deeper suborbital space (suborbital width 13–17% HL vs. 11–13%).

Funding

This study was financially supported by the JSPS KAKENHI Grant Numbers JP16H06896 and JP18K14509, and the Sasagawa Scientific Research Grant of the Japan Science Society.

Acknowledgments

I would like to express my deep appreciation to H. Endo (BSKU) for his expertise and supervision throughout this study. My sincere gratitude also goes to T. Iwamoto (CAS), the world's authority on macrourid taxonomy, who shared his knowledge on grenadiers. I am especially grateful to Y. Kai (FAKU), M. Motokawa and H. Nagamasu (Kyoto University Museum, Kyoto University, Kyoto), K. Sasaki (BSKU), and Y. Kondo (Department of Biology, Faculty of Science and Technology, Kochi University, Kochi) for their helpful advice and encouragement when most needed. The following researchers, museum specialists, and students variously supported this study, and without their kind assistance, this study could not have been completed: M. McGrouther, S. Reader, A. Hay and K. Graham (AMS); S.-P. Huang, P.-L. Lin and K.-T. Shao (ASIZP); T. Saruwatari (Atmosphere and Ocean Research Institute, the University of Tokyo, Kashiwa); J. MacLaine and O. Crimmen (BMNH); A. Suzumoto (BPBM); H. Endo, T. Yamakawa, R. Asaoka, M. Sato, K. Mizowaki, T. Yato, K. Sasaki and Y. Machida (BSKU); T. Iwamoto, D. Catania, J. Fong, M. Hoang and S.G. Poss (CAS); M. Miya (CBM); S. Tessier and K. Khidas (CMNFI); A. Graham, J. Pogonoski and P. Last (CSIRO); Y. Kai, T.P. Sato and T. Nakabo (FAKU); J. Hashimoto (FFNU); C. McMahan, S. Mochel and K. Swagel (FMNH); S. Kimura (FRLM); T. Naito, S. Ohashi, Y. Ohashi, F. Tanaka, S. Chiba, M. Miyamoto, T. Hayashibara and H. Honda

(FSFRL); H. Imamura, T. Kawai, F. Tashiro, M. Yabe, K. Nakaya and K. Amaoka (HUMZ); K. Uchikawa (Japan Sea National Fisheries Research Institute, Fisheries Research Agency of Japan, Niigata); H. Motomura, K. Fujiwara, Y. Fukui, G. Ogihara and M. Meguro (KAUM); Y. Yamamoto and T. Matsuzaki (Center for Advanced Marine Core Research, Kochi University, Kochi); A. Usui (Department of Geology, Faculty of Science and Technology, Kochi University, Kochi); M. Matsunuma and S. Watanabe (KUN); T. Ohkawa (Kochi Prefectural Fisheries Experimental Station, Susaki); Y. Hibino (Kitakyushu Museum of Natural History and Human History, Kitakyushu); H. Senou (KPM); W. Leo Smith (KU); T. Okamoto and T. Hikida (Division of Biological Science, Graduate School of Science, Kyoto University, Kyoto); K. Nishikawa (Department of Interdisciplinary Environment, Graduate School of Human and Environmental Studies, Kyoto University, Kyoto); M. Okamoto and H. Horikawa (Marine Fisheries Research and Development Center, Fisheries Research Agency of Japan, Yokohama); K.E. Hartel and C. Weisel (MCZ); R. Causse, C. Ferrara, Z. Gabsi and P. Pruvost (MNHN); S. Tomiyama (MSM); Y. Iwatsuki (MUFS); R.K. Hadiaty (MZB; deceased); Y.-C. Liao (National Museum of Marine Science and Technology, Keelung); K. Nashida, H. Sakaji and S. Yanagimoto (National Research Institute of Fisheries Science, Fisheries Research Agency of Japan, Yokohama); P. McMillan (NIWA); H.-C. Ho (NMMB); G. Shinohara, M. Nakae, K. Kuriwa, Y. Takata, Y. Inoue, T. Kubodera, H. Komatsu and K. Matsuura (NSMT); M. Hammer and G. Dally (NTM); W.-J. Chen, M.-Y. Lee, J.-N. Chen, M. Lin and S. Lavoue (NTUM); K. Miyamoto and H. Takaoka (OCF); K. Hatooka (OMNH); E. Katayama (Research Institute of Marine Invertebrates, Tokyo); K. Lahaise, M. van Oijen and R. de Ruiter (RMNH); A. Orlov (Russian Federal Research Institute of Fisheries and Oceanography, Moscow); J.-S. Zhong (SFU); K. Hoshino (SNFR); K. Shibukawa (SPMN); R. Misawa and D. Kitagawa (Tohoku National Fisheries Research Institute, Fisheries Research Agency of Japan, Hachinohe); A. Fukui, M. Takami and T. Tamai (Department of Fisheries, School of Marine Science and Technology, Tokai University, Shimizu); A.M. Prokofiev (Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Moscow); T. Okamoto (Tokyo); A. Teramura (Fisheries Laboratory, Graduate School of Agricultural and Life Sciences, the University of Tokyo, Hamamatsu); J.T. Williams, D.G. Smith, J. Finan, S.J. Raredon, A. Nonaka and G.D. Johnson (USNM); S.M. Morrison (WAM); E.D. Vasil'eva and I.A. Verigina (ZMMGU); I. Byrkjedal and G. Langhelle (ZMUB); P.R. Møller, T. Menne, W. Schwarzhans and J. Nielsen (ZMUC); and K. Sakamoto (ZMUT). Field surveys were supported by the FRVs *Kotaka-maru* (National Research Institute of Fisheries Science, Fisheries Research Agency of Japan, Yokohama) and *Wakataka-maru* (Tohoku National

Fisheries Research Institute, Fisheries Research Agency of Japan, Hachinohe); F/Vs *Kosei-maru*, *Tsukasa-maru*, and *Seiryō-maru* (Mimase fishing port, Kochi), *Maruko-maru* (Nomaïke fishing port, Minami Satsuma), and *Hinode-maru* (Heta fishing port, Heta); R/V *Tansei-maru* (Japan Agency for Marine-Earth Science and Technology, Yokosuka); T/V *Nagasaki-maru* (Nagasaki University, Nagasaki); and also students of Kochi University. My special thank goes to O. Okamura (deceased), a former Professor Emeritus of Kochi University, who established a large collection of grenadiers on which this study was largely based. I gratefully acknowledge H. Nagamasu (Kyoto University Museum) for his helpful nomenclatural advice. Finally, my deepest appreciation goes to G. Yearsley (Hobart), who thoroughly read an earlier draft of this paper and edited the English text; M. Matsunuma (KUN) provided a detailed review, catching errors and making suggestions for improvement; H. Endo (BSKU) and P. Konstantinidis (OS) also commented on the entire manuscript; all remaining errors and resistance to good advice are entirely my own responsibility.

References

Many references listed below were written in Japanese. If an English title was given in a paper, the title is cited exactly as given; if an English title was not given, a translated English title is provided here in square brackets.

- Akazaki, M. (1984) Aoshima fukin no kaisan gyorui [Marine fishes in the vicinity of Aoshima Island]. In: Miyazaki Linnean Society (Ed.), *Natural History of Aoshima*. Miyazaki Linnean Society, Miyazaki, pp. 149–266. [In Japanese.]
- Akihito, Sakamoto, K., Iwata, A. & Ikeda, Y. (1993) Cephalic sensory organs of the gobioid fishes. In: Nakabo, T. (Ed.), *Fishes of Japan with Pictorial Keys to the Species. 1st Edition*. Tokai University Press, Tokyo, pp. 1088–1116. [In Japanese.]
- Alcock, A. (1889) Natural history notes from H.M. Indian marine survey steamer ‘Investigator’, commander Alfred Carpenter, R.N., D.S.O., commanding.—No. 13. On the bathybial fishes of the Bay of Bengal and neighboring waters, obtained during the seasons 1885–1889. *Annals and Magazine of Natural History, Series 6*, 4, 376–399. <http://dx.doi.org/10.1080/00222938909460547>
- Alcock, A. (1890) Natural history notes from H.M. Indian marine survey steamer ‘Investigator’, Commander R.F. Hoskyn, R.N., commanding.—No. 18. On the bathybial fishes of the Arabian Sea, obtained during the season 1889–1890. *Annals and Magazine of Natural History, Series 6*, 6, 295–311. <http://dx.doi.org/10.1080/00222939008694038>
- Alcock, A. (1892) *Illustrations of the Zoology of H.M. Indian Marine Surveying Steamer Investigator, under the Command of Commander A. Carpenter, R.N., D.S.O., and of Commander R.F. Hoskyn, R.N. Fishes. Part I*. Royal Indian Museum, Calcutta, no pages, I–VII pls. <http://dx.doi.org/10.5962/bhl.title.64230>
- Alcock, A. (1894) Natural history notes from H.M. Indian marine survey steamer ‘Investigator’, Commander C.F. Oldham, R.N., commanding. Series II., No. 11. An account of a recent collection of bathybial fishes from the Bay of Bengal and from the Laccadive Sea. *Journal of the Asiatic Society of Bengal*, 63, 155–137, pls. VI–VII.
- Alcock, A. (1895) *Illustrations of the Zoology of Royal Indian Marine Surveying Steamer Investigator, under the Command of Commander A. Carpenter, R.N., D.S.O., of the Late Commander R.F. Hoskyn, R.N., and of Commander C.F. Oldham, R.N. Fishes. Part III*. Royal Indian Museum, Calcutta, no pages, XIV–XVI pls. <http://dx.doi.org/10.5962/bhl.title.64230>
- Alcock, A. (1899) *A Descriptive Catalogue of the Indian Deep-Sea Fishes in the Indian Museum. Being a Revised Account of the Deep-Sea Fishes Collected by the Royal Indian Marine Survey Ship Investigator*. Indian Museum, Calcutta, iii + 211 + viii pp. <http://dx.doi.org/10.5962/bhl.title.4684>
- Allain, V., Biseau, A. & Kergoat, B. (2003) Preliminary estimates of French deepwater fishery discards in the Northeast Atlantic Ocean. *Fisheries Research*, 60, 185–192. [http://dx.doi.org/10.1016/s0165-7836\(02\)00031-0](http://dx.doi.org/10.1016/s0165-7836(02)00031-0)
- Amante, C. & Eakins, B.W. (2009) ETOPO1 1 arc-minute global relief model: procedures, data sources and analysis. NOAA Technical Memorandum ESDIS NGDC-24. National Geophysical Data Center, NOAA, Boulder. <http://www.doi.org/10.7289/V5C8276M>
- Amaoka, K. (2007) Ōshū ni watatta dēderurain korekushon. 1. Gyorui [Döderlein’s Collections brought to Europe. 1. Fishes]. In: Fujita, T. & Namikawa, H. (Eds.), *Fauna Sagamiana*. Tokai University Press, Hadano, pp. 28–37. [In Japanese.]
- Amaoka, K. & Nakaya, K. (1983) Fishes. In: Amaoka, K., Nakaya, K., Yabu, H. & Yamamoto, H. (Eds.), *Fishes and Marine Algae of Northern Japan*. Kitanihon Kaiyo Center, Sapporo, pp. 5–175. [In Japanese.]
- Amaoka, K., Nakaya, K. & Yabe, M. (1995) *The Fishes of Northern Japan*. Kitanihon Kaiyo Center, Sapporo, 390 pp. [In Japanese.]
- Amaoka, K., Nakaya, K. & Yabe, M. (2011) *Fishes of Hokkaido*. Hokkaido Shimbun Press, Sapporo, 482 pp. [In Japanese.]
- Amaoka, K., Nakaya, K. & Yabe, M. (2020) *Pictorial Guide to the Fishes of Hokkaido*. Hokkaido Shimbun Press, Sapporo, 590 pp. [In Japanese.]
- Ambrose, D.A. (1996) Macrouridae: grenadiers. In: Moser, H.G. (Ed.), *The Early Stages of Fishes in the California Current Region. California Cooperative Oceanic Fisheries Investigations, Atlas No. 33*. Allen Press, Lawrence, pp. 483–499.

- Anderson, E.M. (1996) Notes on the genus *Hymenocephalus* Giglioli, 1884 (Teleostei: Macrouridae) in southern Africa. *Transactions of the Royal Society of South Africa*, 51, 299–232.
<http://dx.doi.org/10.1080/00359199609520608>
- Arai, T. & Iwamoto, T. (1979) A new species of the macrourid fish genus *Coelorinchus* from off Tasmania, New Zealand, and the Falkland Islands. *Japanese Journal of Ichthyology*, 26, 238–246.
- Arkhipkin, A.I., Laptikhovsky, V.V. & Brickle, P. (2010) An antipodal link between the North Pacific and South Atlantic Oceans? *Deep-Sea Research, Part I*, 57, 1009–1011.
<http://dx.doi.org/10.1016/j.dsr.2010.05.004>
- Arruda, L.M. (1997) Checklist of the marine fishes of the Azores. *Arquivos do Museu Bocage, Nova Série*, 3 (2), 13–164.
- Asano, N., Harada, K., Hujimoto, T. & Tange, M. (1952) Studies on the marine fauna in Ibaraki Prefecture—II. On the fish fauna. *Bulletin of the Fisheries Experimental Station of Ibaraki-ken*, 27, 87–97. [In Japanese with English abstract.]
- Bailly, N., Hureau, J.-C. & Pruvost, P. (1999) Catalogue critique des types de poissons du Muséum National d'Histoire Naturelle (et des Musées d'histoire naturelle en région) (suite). *Ordre des Gadiformes. Cybium*, 23, 219–245.
- Balanov, A.A. (2003) Additions to the ichthyofauna of the south western part of the Sea of Okhotsk. *Journal of Ichthyology*, 43, 483–485. [Originally published in Russian in *Voprosy Ikhtiologii*, 43, 565–567.]
- Balanov, A.A., Moku, M., Kawaguchi, K. & Shinohara, G. (2009) Fishes collected by commercial size midwater trawls from the Pacific coast off northern Japan. *National Museum of Nature and Science Monographs*, (39), 655–681.
- Barnard, K.H. (1925) Descriptions of new species of marine fishes from S. Africa. *Annals and Magazine of Natural History, Series 9*, 15, 498–504.
<http://dx.doi.org/10.1080/00222932508633240>
- Bean, T.H. (1884) Notes on some fishes collected by James G. Swan in Washington Territory, including a new species of *Macrurus*. *Proceedings of the United States National Museum*, 6, 362–364.
<http://dx.doi.org/10.5479/si.00963801.6-388.362>
- Bean, T.H. (1890) Scientific results of explorations by the U.S. Fish Commission steamer *Albatross*. No. XI. New fishes collected off the coast of Alaska and the adjacent region southward. *Proceedings of the United States National Museum*, 13, 37–45.
<http://dx.doi.org/10.5479/si.00963801.13-795.37>
- Bleeker, P. (1874) Typi nonnulli generici piscium neglecti. *Verslagen en Mededeelingen der Koninklijke Akademie van Wetenschappen, Afdeling Natuurkunde, Series 2*, 8, 367–371.
- Bloch, M.E. (1786) *Naturgeschichte der Ausländischen Fische. Vol. 2*. Schlesinger, Berlin, xii + 160 pp., CXLV–CLXXX pls.
- Boeseman, M. (1947) Revision of the fishes collected by Burger and von Siebold in Japan. *Zoologische Mededelingen, Leiden*, 28, 1–242, pls. I–V.
- Boeseman, M. (1997) Collectors and fish collections of the Rijksmuseum van Natuurlijke Historie in Leiden, Netherlands (1820–1980). In: Pietsch, T.W. & Anderson, W.D. Jr. (Eds.), *Collection Building in Ichthyology and Herpetology*. Allen Press, Lawrence, pp. 81–100.
- Bonaparte, C.L. (1831) Prospetto del sistema generale d'ittologia. In: Bonaparte, C.L. (1831–1832), *Saggio di una Distribuzione Metodica degli Animali Vertebrati*. Roma, pp. 89–123.
- Bonaparte, C.L. (1846) Catalogo metodico dei pesci europei. *Atti della Settima Adunanza degli Scienziati Italiani Tenuta in Napoli*, pt. 2, 1–95.
- Borets, L.A. (1986) Ichthyofauna of the Northwestern and Hawaiian Submarine Ridges. *Journal of Ichthyology*, 26 (2), 1–13. [Originally published in Russian in *Voprosy Ikhtiologii*, 26, 208–220.]
- Brauer, A. (1906) Die tiefsee-fische. I. Systematischer teil. *Wissenschaftliche Ergebnisse der Deutschen Tiefsee-Expedition auf dem Dampfer "Valdivia" 1898–99*, 15 (2), 1–432, pls. I–XVIII.
<http://dx.doi.org/10.5962/bhl.title.2171>
- Brickle, P. & Laptikhovsky, V. (2002) New records of deep-sea fishes from the waters around the Falkland Islands. *Journal of Fish Biology*, 60, 492–494.
<http://dx.doi.org/10.1006/jfbi.2001.1848>
- Brisson, M.J. (1760) *Ornithologie. Vol. 1*. Cl. Jean Baptiste Bauche, Paris, xxiv + 526 + lxxiii pp., XXXVII pls.
<http://dx.doi.org/10.5962/bhl.title.51902>
- Chen, T.-F. (1951) Check-list of the species of fishes known from Taiwan (Formosa). *Quarterly Journal of the Taiwan Museum*, 4, 181–210.
- Chiou, M.-L., Shao, K.-T. & Iwamoto, T. (2004a) A new species, *Caelorinchus sheni*, and 19 new records of grenadiers (Pisces: Gadiformes: Macrouridae) from Taiwan. *Zoological Studies*, 43, 35–50.
<http://dx.doi.org/10.1643/ci-03-128r1>
- Chiou, M.-L., Shao, K.-T. & Iwamoto, T. (2004b) New species of *Caelorinchus* (Macrouridae, Gadiformes, Teleostei) from Taiwan, with a redescription of *Caelorinchus brevirostris* Okamura. *Copeia*, 2004 (4), 298–304.
<http://dx.doi.org/10.1643/ci-03-128r1>
- Chu, Y.-T., Chan, C.-L. & Chen, C.-T. (1963) *Ichthyofauna of East China Sea*. Science Press, Beijing. xxviii + 642 pp. [In Chinese.]
- Chyung, M.-K. & Kim, K.-H. (1959) Thirteen unrecorded species of fish from Korean waters. *Korean Journal of Zoology*, 2, 1–10.
- Cocco, A. (1844) Intorno ad alcuni nuovi pesci del mare di Messina. *Giornale del Gabinetto Letterario di Messina*, anno 3, tomo 5, 1–10, figs. 1–4.
- Collett, R. (1896) Poissons provenant des campagnes du yacht "L'Hirondelle" (1885–1888). *Résultats des Campagnes Scientifiques Accomplies sur son Yacht par Albert Ier, Prince Souverain de Monaco*, fasc. 10, i–viii + 1–198, pls.

- Dalman J.W. (1826) Om Insekter inneslutne i Copal; jemte beskrifning på några dreibland förekommande nya släkten och arten. *Kungliga Svenska Vetenskapsakademiens Handlingar, Series 3*, 13, 375–410.
- Doflein, F. (1906) *Ostasienfahrt. Erlebnisse und Beobachtungen eines Naturforschers in China, Japan und Ceylon*. B.G. Teubner, Leipzig and Berlin, xiii + 511 pp.
- Dotsu, Y. & Nakamura, M. (1976) Tsushima no gyorui [Fishes of Tsushima Island]. In: Nagasaki Biological Society (Ed.), *Tsushima no Seibutsu [Fauna of Tsushima Island]*. Nagasaki Biological Society, Nagasaki, pp. 265–287. [In Japanese.]
- Dotsu, Y. & Tomiyama, I. (1967) The marine fishes from Saikai National Park of Japan. *Bulletin of the Faculty of Fisheries, Nagasaki University*, (23), 1–42. [In Japanese with English abstract.]
- Drazen, J.C. (2008) Energetics of grenadier fishes. In: Orlov, A.M. & Iwamoto, T. (Eds.), *Grenadiers of the World Ocean: Biology, Stock Assessment, and Fisheries*. American Fisheries Society Symposium 63. American Fisheries Society, Maryland, pp. 203–208.
- Dunlop, J.A. & Jekel, D. (2008) The oldest available fossil arachnid name. *Palaeodiversity*, 1, 87–92.
- Dunn, J.R. (1997) Charles Henry Gilbert (1859–1928): pioneer ichthyologist of the American West. In: Pietsch, T.W. & Anderson, W.D. Jr. (Eds.), *Collection Building in Ichthyology and Herpetology*. Allen Press, Lawrence, pp. 265–278.
- Endo, H. (1996) Sokodara-ka gyorui no shoki-seikatsushi ni kansuru kenkyū no genjō [Current progress in early life history of grenadiers]. *Tohoku Sokou Kenkyū [Tohoku Demersal Fish Research]*, 16, 26–34. [In Japanese.]
- Endo, H. (1997) Juveniles of roundhead grenadier *Odontomacrus murrayi* (Teleostei, Macrouridae) from the eastern Indian Ocean. *Memoirs of the Faculty of Science, Kochi University*, 18, 41–45.
- Endo, H. (2002) Phylogeny of the order Gadiformes (Teleostei, Paracanthopterygii). *Memoirs of the Graduate School of Fisheries Sciences, Hokkaido University*, 49, 75–149.
- Endo, H. (2014) Macrouridae. In: Okiyama, M. (Ed.), *An Atlas of Early Stage Fishes in Japan. 2nd Edition*. Tokai University Press, Hadano, pp. 411–420. [In Japanese.]
- Endo, H., Iwasaki, N., Machida, Y. & Iwai, M. (2000) Bathybenthic fauna of the Nankai Trough off Muroto observed using a deep seafloor observatory 1. Fishes. *JAMSTEC Journal of Deep Sea Research*, (17), 12–17. [In Japanese with English abstract.]
- Endo, H., Nakayama, N., Suetsugu, K. & Miyake, H. (2010) A larva of *Coryphaenoides pectoralis* (Gadiformes: Macrouridae) collected by deep-sea submersible from off Hokkaido, Japan. *Ichthyological Research*, 57, 272–277. <http://dx.doi.org/10.1007/s10228-010-0164-4>
- Endo, H. & Okamura, O. (1992) New records of the abyssal grenadiers *Coryphaenoides armatus* and *C. yaquinae* from the western North Pacific. *Japanese Journal of Ichthyology*, 38, 433–437.
- Endo, H., Tsutsui, D. & Amaoka, K. (1994) Range extensions of two deep-sea macrourids *Coryphaenoides filifer* and *Squalogadus modificatus* to the Sea of Okhotsk. *Japanese Journal of Ichthyology*, 41, 330–333.
- Endo, H., Yabe, M. & Amaoka, K. (1993) Occurrence of the macrourid alevins genera *Albatrossia* and *Coryphaenoides* in the northern North Pacific Ocean. *Japanese Journal of Ichthyology*, 40, 219–226.
- Eschmeyer, W.N. (1998) *Catalog of Fishes*. California Academy of Sciences, San Francisco, 2905 pp.
- Eschmeyer, W.N. & Fong, J.D. (2011) Pisces [In: Zhang, Z.-Q. (Ed.), *Animal Biodiversity: An Outline of Higher-Level Classification and Survey of Taxonomic Richness*]. *Zootaxa*, 3148, 26–38. <http://dx.doi.org/10.11646/zootaxa.3148.1.7>
- Fahay, M.P. & Markle, D.F. (1984) Gadiformes: development and relationships. In: Moser, H.G., Richards, W.J., Cohen, D.M., Fahay, P.M., Kendall, A.W. Jr. & Richardson, S.L. (Eds.), *Ontogeny and Systematics of Fishes Based on an International Symposium Dedicated to the Memory of Elbert Halvor Ahlstrom*. American Society of Ichthyologists and Herpetologists Special Publication Number 1. Allen Press, Lawrence, pp. 265–283.
- Filhol, H. (1884) Explorations sous-marines. Voyage du “Talisman”. *La Nature, Paris*, (559), 182–186.
- Fowler, H.W. (1925) New taxonomic names of West African marine fishes. *American Museum Novitates*, (162), 1–5.
- Fowler, H.W. (1936) The marine fishes of West Africa based on the collection of the American Museum Congo Expedition, 1909–1915. Part 1. *Bulletin of the American Museum of Natural History*, 70, 1–605.
- Franz, V. (1910) Die japanische knochenfische der sammlungen Haberer und Doflein (beiträge zur naturgeschichte Ostasiens). *Abhandlungen der Mathematisch-Physikalischen Klasse der K. Bayer, Akademie der Wissenschaften*, 4, 1–135, pls. I–XI. <http://dx.doi.org/10.5962/bhl.title.16274>
- Fraser-Brunner, A. (1935). New or rare fishes from the Irish Atlantic Slope. *Proceedings of the Royal Irish Academy, Section B, Biological, Geological, and Chemical Science*, 42, 319–326.
- Fricke, R. & Eschmeyer, W.N. (2020) A guide to fish collections in Eschmeyer’s Catalog of Fishes, online version. California Academy of Sciences, San Francisco. Available from: <http://researcharchive.calacademy.org/research/ichthyology/catalog/collections.asp> (accessed 2 January 2020)
- Fricke, R., Eschmeyer, W.N. & van der Laan, R. (2020) Eschmeyer’s Catalog of Fishes: genera, species, references, online version. California Academy of Sciences, San Francisco. Available from: <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp> (accessed 2 January 2020)
- Fujikura, K., Lindsay, D., Kitazato, H., Nishida, S. & Shirayama, Y. (2010) Marine biodiversity in Japanese waters. *PLoS*

ONE, 5 (8), e11836 (13 pp.).

<http://dx.doi.org/10.1371/journal.pone.0011836>

- Fujioka, Y. (1991) Yamaguchi-ken-san gyorui mokuroku [List of fishes from Yamaguchi Prefecture]. In: Fujioka Yutaka Kyōju Taikan Kinen Jigyō Kai [The Association to Commemorate the Retirement of Professor Yutaka Fujioka] (Ed.), *Yamaguchi no Sakana [Fishes of Yamaguchi]*. Fujioka Yutaka Kyōju Taikan Kinen Jigyō Kai [The Association to Commemorate the Retirement of Professor Yutaka Fujioka], Yamaguchi, pp. 106–153. [In Japanese.]
- Fujiwara, K., Tanoue, H., Mohori, M., Kamona, T., Hata, K., Okada, S., Nagai, S. & Motomura, H. (2018) Fishes of the Hibiki-nada Sea and Mi-shima Island, Yamaguchi Prefecture, Japan. *Journal of National Fisheries University*, 66, 47–80.
- Fukui, A., Takami, M., Tsuchiya, T., Sezaki, K., Igarashi, Y., Kinoshita, S. & Watabe, S. (2010) Pelagic eggs and larvae of *Coelorinchus kishinouyei* (Gadiformes: Macrouridae) collected from Suruga Bay, Japan. *Ichthyological Research*, 57, 169–179.
<http://dx.doi.org/10.1007/s10228-009-0144-8>
- Fukui, A. & Tsuchiya, T. (2005) Pelagic larvae of *Ventrifossa garmani* (Gadiformes: Macrouridae) from Suruga Bay and offshore waters of Japan. *Ichthyological Research*, 52, 311–315.
<http://dx.doi.org/10.1007/s10228-005-0285-3>
- Fukui, A., Tsuchiya, T., Sezaki, K. & Watabe, S. (2008) Pelagic eggs and larvae of *Coryphaenoides marginatus* (Gadiformes: Macrouridae) collected from Suruga Bay, Japan. *Ichthyological Research*, 55, 284–293.
<http://dx.doi.org/10.1007/s10228-007-0024-z>
- Fukui, Y., Matsunuma, M. & Motomura, H. (2015) A list of demersal fishes collected from off Kuro-shima Island in the Osumi Group, Kagoshima Prefecture, southern Japan, with record of *Hydrolagus mitsukurii* (Chimaeriformes: Chimaeridae). *Nature of Kagoshima*, 41, 177–186. [In Japanese.]
- Funabashi, M. (1998) Checklist of marine fishes from the coastal region of Ibaraki Prefecture. *Bulletin of Ibaraki Nature Museum*, (1), 75–96. [In Japanese with English abstract.]
- Furuhashi, N., Tsubaki, K., Mori, Y. & Hashimoto, J. (2010) Demersal fish assemblages from the continental shelf margin to the upper continental slope, southwest of Nagasaki, Japan. *Bulletin of the Faculty of Fisheries, Nagasaki University*, 91, 17–33. [In Japanese with English abstract.]
- Gage, J.D. & Tyler, P.A. (1992) *Deep-Sea Biology: A Natural History of Organisms at the Deep-Sea Floor*. Cambridge University Press, Cambridge, xiii + 504 pp.
<http://dx.doi.org/10.1017/CBO9781139163637>
- Garman, S. (1899) Reports on an exploration off the west coasts of Mexico, Central and South America, and off the Galapagos Islands, in charge of Alexander Agassiz, by the U.S. Fish Commission steamer “Albatross”, during 1891, Lieut. Commander Z.L. Tanner U.S.N., commanding. XXVI. Fishes. *Memoirs of the Museum of Comparative Zoology at Harvard College*, 24, 1–431, pls. I–LXXXV + A–N.
- Geistdoerfer, P. (1986) Macrouridae. In: Whitehead, P.J.P., Bauchot, M.-L., Hureau, J.-C., Nielsen, J. & Tortonese, E. (Eds.), *Fishes of the North-Eastern Atlantic and the Mediterranean. Volume II*. UNESCO, Paris, pp. 644–676.
- Geistdoerfer, P. (1990) Macrouridae. In: Quéro, J.C., Hureau, J.C., Karrer, C., Post, A. & Saldanha, L. (Eds.), *Clofeta. Check-List of the Fishes of the Eastern Tropical Atlantic*. Junta Nacional de Investigação Científica e Tecnológica, Lisbon, European Ichthyological Union, Paris, and United Nations Educational, Scientific and Cultural Organization, Paris, pp. 541–563.
- Giglioli, E.H. (1884) Esplorazione talassografica del Mediterraneo. In: Giglioli, E.H. & Issel, A. (Eds.), *Plagos, Saggio sulla Vita e sui Prodrutti del Mare*. Instituto de’ Sordo-muti, Genova, pp. 198–270.
<http://dx.doi.org/10.5962/bhl.title.42527>
- Giglioli, E.H. (1893) Di una nuova specie di Macruride appartenente alla fauna abissale del Mediterraneo. *Zoologischer Anzeiger*, 16, 343–345.
- Gilbert, C.H. (1890) A preliminary report on the fishes collected by the steamer *Albatross* on the Pacific coast of North America during the year 1889, with descriptions of twelve new genera and ninety-two new species. *Proceedings of the United States National Museum*, 13, 49–126.
<http://dx.doi.org/10.5479/si.00963801.13-797.49>
- Gilbert, C.H. (1891) Descriptions of thirty-four new species of fishes collected in 1888 and 1889, principally among the Santa Barbara Islands and in the Gulf of California. *Proceedings of the United States National Museum*, 14, 539–566.
<http://dx.doi.org/10.5479/si.00963801.14-880.539>
- Gilbert, C.H. (1896) The ichthyological collections of the steamer *Albatross* during the years 1890 and 1891. *Report of the United States Fish Commission*, 19, 393–476, pls. 20–35.
- Gilbert, C.H. (1905) The deep-sea fishes of the Hawaiian Islands. In: Jordan, D.S. & Evermann, B.W. (Eds.), *The Aquatic Resources of the Hawaiian Islands. Bulletin of the U.S. Fish Commission*, 23. Government Printing Office, Washington DC, pp. 573–713, figs. 230–276, pls. 66–101.
<http://dx.doi.org/10.5962/bhl.title.12624>
- Gilbert, C.H. (1915) Fishes collected by the United States Fisheries steamer “Albatross” in southern California 1904. *Proceedings of the United States National Museum*, 48, 305–380, pls. 14–22.
<https://doi.org/10.5479/si.00963801.48-2075.305>
- Gilbert, C.H. & Burke, C.V. (1912) Fishes from Bering Sea and Kamchatka. *Bulletin of the Bureau of Fisheries*, 30, 31–96.
- Gilbert, C.H. & Cramer, F. (1897) Report on the fishes dredged in deep water near the Hawaiian Islands, with descriptions and figures of twenty-three new species. *Proceedings of*

- the United States National Museum 19, 403–435, pls. XXXVI–XLVIII.
<http://dx.doi.org/10.5479/si.00963801.19-1114.403>
- Gilbert, C.H. & Hubbs, C.L. (1916) Report on the Japanese macrourid fishes collected by the United States fisheries steamer “Albatross” in 1906, with a synopsis of the genera. *Proceedings of the United States National Museum*, 51, 135–214, pls. 8–11.
<http://dx.doi.org/10.5479/si.00963801.51-2149.135>
- Gilbert, C.H. & Hubbs, C.L. (1917) Description of *Hymenoccephalus tenuis*, a new macrourid [sic] fish from the Hawaiian Islands. *Proceedings of the United States National Museum*, 54, 173–175.
<http://dx.doi.org/10.5479/si.00963801.54-2231.173>
- Gilbert, C.H. & Hubbs, C.L. (1920) The macrourid fishes of the Philippine Islands and the East Indies. *United States National Museum Bulletin*, 100 (vol. 1, pt. 7), 369–588.
<http://dx.doi.org/10.5962/bhl.title.13637>
- Gill, T.N. & Townsend, C.H. (1897) Diagnoses of new species of fishes found in Bering Sea. *Proceedings of the Biological Society of Washington*, 11, 231–234.
- Giorna, E.M. (1809) Suite et conclusion du mémoire. *Mémoires de l'Académie Impériale des Sciences, Littérature et Beaux-Arts, de Turin, Sér. Sciences Physiques et Mathématiques*, 16, 177–180, pls. I–II.
- Glare, P.G.W. (1968) *Oxford Latin Dictionary*. Oxford University Press, London, xxiii + 2126 pp.
- Goode, G.B. & Bean, T.H. (1883) Reports on the results of dredging under the supervision of Alexander Agassiz, on the east coast of the United States, during the summer of 1880, by the U.S. coast survey steamer “Blake”, Commander J.R. Bartlett, U.S.N., commanding. *Bulletin of the Museum of Comparative Zoology*, 10, 183–226.
- Goode, G.B. & Bean, T.H. (1885) Descriptions of new fishes obtained by the United States Fish Commission mainly from deep water off the Atlantic and Gulf coasts. *Proceedings of the United States National Museum*, 8, 589–605.
<http://dx.doi.org/10.5479/si.00963801.8-543.589>
- Goode, G.B. & Bean, T.H. (1896) Oceanic ichthyology, a treatise on the deep-sea and pelagic fishes of the world, based chiefly upon the collections made by the steamers *Blake*, *Albatross*, and *Fish Hawk* in the northwestern Atlantic, with an atlas containing 417 figures. *Memoirs of the Museum of Comparative Zoology at Harvard College*, 22, text: i–xxxv + 1–26 + 1–553; atlas: i–xxiii + 1–26, pls. I–CXXIII.
<http://dx.doi.org/10.5962/bhl.title.48521>
- Goto, T. (2000) Fauna of demersal fishes from off Iwate Prefecture, Pacific coast of northern Japan. *Bulletin of Iwate Prefectural Fisheries Technology Center*, 2, 1–13. [In Japanese with English abstract.]
- Grant, C.H.B. & Mackworth-Praed, C.W. (1956) Moehring's genera in his *Geslachten der Vogelen* (Avium Genera), Nozeman & Vosmaer Edition, Amsterdam, 1758. *Annals and Magazine of Natural History, Series 12*, 9, 774–778.
<http://dx.doi.org/10.1080/00222935608655893>
- Gray, J.E. (1854) *Catalogue of Fish Collected and Described by Laurence Theodore Gronow, Now in the British Museum*. British Museum, London, vii + 196 pp.
- Gunnerus, J.E. (1765) Efterretning om Berglaxen, en rar Norsk fisk, som kunde kaldes: *Coryphaenoides rupestris*. *Det Trondhiemske Selskabs Skrifter*, 3, 50–58, pl. III.
- Günther, A. (1862) *Catalogue in the Fishes of the British Museum. Vol. 4. Catalogue of the Acanthopterygii, Pharyngognathi and Anacanthini in the Collection of the British Museum*. British Museum, London, xxi + 534 pp.
<http://dx.doi.org/10.5962/bhl.title.8304>
- Günther, A. (1877) Preliminary notes on new fishes collected in Japan during the expedition of H.M.S. ‘Challenger’. *Annals and Magazine of Natural History, Series 4*, 20, 433–446.
<http://dx.doi.org/10.1080/00222937708682260>
- Günther, A. (1878) Preliminary notices of deep-sea fishes collected during the voyage of H.M.S. ‘Challenger’. *Annals and Magazine of Natural History, Series 5*, 2, 17–28.
<http://dx.doi.org/10.1080/00222937808682376>
- Günther, A. (1887) Report on the deep-sea fishes collected by H.M.S. *Challenger* during the years 1873–76. *Report on the Scientific Results of the Voyage of H.M.S. Challenger*, 22, i–ixv + 1–268.
<http://dx.doi.org/10.5962/bhl.title.15693>
- Hart, J.L. (1973) *Pacific Fishes of Canada. Bulletin, Fisheries Research Board of Canada, No. 180*. Fisheries Research Board of Canada, Ottawa, 740 pp.
- Headrich, R.L. & Merrett, N.R. (1988) Summary atlas of deep-living demersal fishes in the North Atlantic Basin. *Journal of Natural History*, 22, 1325–1362.
<http://dx.doi.org/10.1080/00222938800770811>
- Hector, J. (1875) Descriptions of five new species of fishes obtained in the New Zealand seas by H.M.S. “Challenger” Expedition July 1874. *Annals and Magazine of Natural History, Series 4*, 15, 78–82.
<http://dx.doi.org/10.1080/00222937508681027>
- Herring, P. (2002) *The Biology of the Deep Ocean*. Oxford University Press, Oxford, vi + 314 pp.
- Hikita, T. (1981) The fishes from eastern waters and rivers of the Shiretoko Peninsula in Hokkaido, Japan. *Scientific Reports of the Hokkaido Salmon Hatchery*, (35), 57–88. [In Japanese with English abstract.]
- Ho, H.-C. & Shao, K.-T. (2011) Annotated checklist and type catalog of fish genera and species described from Taiwan. *Zootaxa*, 2957, 1–74.
<http://dx.doi.org/10.11646/zootaxa.2957.1.1>
- Honma, Y., Mizusawa, R. & Okiyama, M. (1972) Further additions to “a list of the fishes collected in the Province of Echigo, including Sado Island” (IX). *Bulletin of the Biogeographical Society of Japan*, 28 (4), 47–57. [In Japanese with English abstract.]
- Honma, Y., Nakamura, Y., Imura, H., Shoji, T., Honma, R. & Inoue, N. (2008) Further additions to “a list of the fishes collected in the Province of Echigo including Sado Island” (XVIII). *Kashiwazaki City Museum (Bulletin)*, (22), 65–

74. [In Japanese with English abstract.]
- Hori, Y. (1996) Ibaraki-ken no gyorui · tōsokurui-sō oyobi shuyō-gyokaku-shurui nitsuite [Fishes and cephalopods of Ibaraki Prefecture, with comments on main commercial species]. *Bulletin of the Ibaraki Prefectural Fisheries Experimental Station*, 34, 1–52. [In Japanese.]
- Horikawa, H. (2009) Calendars of the distribution of demersal fishes in Tosa Bay. *Fisheries Biology and Oceanography in the Kuroshio*, (suppl. 1), i–xxi + 1–752.
- Hubbs, C.L. & Iwamoto, T. (1977) A new genus (*Mesobius*), and three new bathypelagic species of Macrouridae (Pisces, Gadiformes) from the Pacific Ocean. *Proceedings of the California Academy of Sciences*, 41, 233–251.
- ICZN (International Commission on Zoological Nomenclature) (1999) *International Code of Zoological Nomenclature. 4th Edition*. International Trust for Zoological Nomenclature, London, 306 pp.
<http://dx.doi.org/10.5962/bhl.title.50608>
- Ikeda, H. & Nakabo, T. (2015) *Fishes of the Pacific Coasts of Southern Japan*. Tokai University Press, Hadano, xxii + 597 pp. [In Japanese.]
- Ishikawa, C. & Matsuura, K. (1897) *Preliminary Catalogue of Fishes Including Dipnoi, Cyclostomi and Cephalochorda in the Collection of the Natural History Department, Imperial Museum*. Imperial Museum, Tokyo, 64 pp. [In Japanese with English preface.]
- Iwamoto, T. (1970) The R/V *Pillsbury* deep-sea biological expedition to the Gulf of Guinea, 1964–65. 19. Macrourid fishes of the Gulf of Guinea. *Studies in Tropical Oceanography*, 4, 316–431.
- Iwamoto, T. (1974) *Nezumia* (*Kuronezumia*) *bubonis*, a new subgenus and species of grenadier (Macrouridae: Pisces) from Hawaii and the western North Atlantic. *Proceedings of the California Academy of Sciences*, 39, 507–516.
- Iwamoto, T. (1978) Eastern Pacific macrourids of the genus *Coelorinchus* Giorna (Pisces: Gadiformes), with description of a new species from Chile. *Proceedings of the California Academy of Sciences*, 41, 307–337.
- Iwamoto, T. (1979) Eastern Pacific macrourine grenadiers with seven branchiostegal rays (Pisces: Macrouridae). *Proceedings of the California Academy of Sciences*, 42, 135–179.
- Iwamoto, T. (1982) *Ventrifossa johnborum*, a new grenadier from the western Pacific (Macrouridae: Pisces). *Australian Zoologist*, 21, 55–61.
<http://dx.doi.org/10.7882/az.1982.004>
- Iwamoto, T. (1986) Family no. 93: Macrouridae. In: Smith, M.M. & Heemstra, P.C. (Eds.), *Smiths' Sea Fishes*. Macmillan South Africa, Johannesburg, pp. 330–341.
<http://dx.doi.org/10.1007/978-3-642-82858-4>
- Iwamoto, T. (1990) Family Macrouridae. In: Cohen, D.M., Inada, T., Iwamoto, T. & Scialabba, N. (Eds.), *FAO Species Catalogue, Vol. 10. Gadiform Fishes of the World. An Annotated and Illustrated Catalogue of Cods, Hakes, Grenadiers and Other Gadiform Fishes Known to Date*. FAO, Rome, pp. 90–317.
- Iwamoto, T. (1997) *Trachonurus robinsi*, a new species of grenadier (Gadiformes, Macrouridae) from the Philippines. *Bulletin of Marine Science*, 60, 942–949.
- Iwamoto, T. (2008a) A note on the correct spelling of the names *Coelorinchus* (*Coelorinchus*) and *C. coelorhinchus* (*caelorhinchus*). In: Orlov, A.M. & Iwamoto, T. (Eds.), *Grenadiers of the World Ocean: Biology, Stock Assessment, and Fisheries. American Fisheries Society Symposium 63*. American Fisheries Society, Maryland, p. xi.
- Iwamoto, T. (2008b) A brief taxonomic history of grenadiers. In: Orlov, A.M. & Iwamoto, T. (Eds.), *Grenadiers of the World Ocean: Biology, Stock Assessment, and Fisheries. American Fisheries Society Symposium 63*. American Fisheries Society, Maryland, pp. 3–13.
- Iwamoto, T. & Anderson, M.E. (1994) Review of the grenadiers (Teleostei: Gadiformes) of southern Africa, with descriptions of four new species. *Ichthyological Bulletin of the J.L.B. Smith Institute of Ichthyology*, (61), 1–28.
- Iwamoto, T. & Anderson, M.E. (1999) New species of *Coelorinchus* (Macrouridae, Gadiformes, Teleostei) from the Indian Ocean. *Copeia*, 1999 (4), 1079–1083.
<http://dx.doi.org/10.2307/1447982>
- Iwamoto, T. & Arai, T. (1987) A new grenadier *Malacocephalus okamurai* (Pisces: Gadiformes: Macrouridae) from the western Atlantic. *Copeia*, 1987 (1), 204–208.
<http://dx.doi.org/10.2307/1446054>
- Iwamoto, T., Golani, D., Baranes, A. & Goren, M. (2006) Two new grenadiers (Teleostei, Gadiformes, Macrouridae) from the Seychelles and Mascarene Ridge, western Indian Ocean. *Proceedings of the California Academy of Sciences*, 57, 433–442.
- Iwamoto, T. & Graham, K.J. (2001) Grenadiers (families Bathygadidae and Macrouridae, Gadiformes, Pisces) of New South Wales, Australia. *Proceedings of the California Academy of Sciences*, 52, 407–509.
- Iwamoto, T. & Graham, K.J. (2008) Two new Australian grenadiers of the *Coelorinchus fasciatus* species group (Macrouridae: Gadiformes: Teleostei). *Proceedings of the California Academy of Sciences*, 59, 133–146.
- Iwamoto, T., Ho, H.-C. & Shao, K.-T. (2009) Description of a new *Coelorinchus* (Macrouridae, Gadiformes, Teleostei) from Taiwan, with notable new records of grenadiers from the South China Sea. *Zootaxa*, 2326, 39–50.
<https://doi.org/10.11646/zootaxa.2326.1.3>
- Iwamoto, T. & McCosker, J.E. (2001) Notes on Galápagos grenadiers (Pisces, Gadiformes, Macrouridae), with the description of a new species of *Coryphaenoides*. *Revista de Biología Tropical*, 49 (suppl. 1), 21–27.
- Iwamoto, T. & McMillan, P. (1997) A new grenadier, genus *Trachonurus*, from New Zealand and Australia (Pisces: Gadiformes: Macrouridae). *Memoirs of the Museum of Victoria*, 56, 255–259.
- Iwamoto, T., McMillan, P. & Shcherbachev, Y.N. (1999) A new grenadier, genus *Coelorinchus*, from Australia and New Zealand (Pisces, Gadiformes, Macrouridae). *New Zealand Journal of Marine and Freshwater Research*, 33, 49–54.

<http://dx.doi.org/10.1080/00288330.1999.9516855>

- Iwamoto, T. & Merrett, N.R. (1997) Pisces Gadiformes: taxonomy of grenadiers of the New Caledonian region, southwest Pacific. *Mémoires du Muséum National d'Histoire Naturelle*, 176, 473–570.
- Iwamoto, T., Nakayama, N., Shao, K.-T. & Ho, H.-C. (2015) Synopsis of the grenadier fishes (Gadiformes; Teleostei) of Taiwan. *Proceedings of the California Academy of Sciences*, 62, 31–126.
- Iwamoto, T. & Okamoto, M. (2015) A new grenadier fish of the genus *Lucigadus* (Macrouridae, Gadiformes, Teleostei) from the Emperor Seamounts, northwestern Pacific. *Proceedings of the California Academy of Sciences*, 62, 369–380.
- Iwamoto, T. & Orlov, A. (2006) *Paracetonus flagellicauda* (Koefoed, 1927) (Macrouridae, Gadiformes, Teleostei), new records from the Mid-Atlantic Ridge and Madagascar Plateau. *Proceedings of the California Academy of Sciences*, 57, 379–386.
- Iwamoto, T. & Orlov, A. (2008) First Atlantic record of *Asthenomacrus victoris* Sazonov & Shcherbachev (Macrouridae, Gadiformes, Teleostei). *Proceedings of the California Academy of Sciences*, 59, 125–131.
- Iwamoto, T. & Sazonov, Y.I. (1988) A review of the southeastern Pacific *Coryphaenoides* (sensu lato) (Pisces, Gadiformes, Macrouridae). *Proceedings of the California Academy of Sciences*, 45, 35–82.
- Iwamoto, T. & Sazonov, Y.I. (1994) Revision of the genus *Kumba* (Pisces, Gadiformes, Macrouridae), with the description of three new species. *Proceedings of the California Academy of Sciences*, 48, 221–237.
- Iwamoto, T., Shao, K.-T. & Ho, H.-C. (2011) Elevation of *Spicomacrus* (Gadiformes: Macrouridae) to generic status, with descriptions of two new species from the southwestern Pacific. *Bulletin of Marine Science*, 87, 513–530.
<http://dx.doi.org/10.5343/bms.2011.1004>
- Iwamoto, T. & Shcherbachev, Y.N. (1991) Macrourid fishes of the subgenus *Chalinura*, genus *Coryphaenoides*, from the Indian Ocean. *Proceedings of the California Academy of Sciences*, 47, 207–233.
- Iwamoto, T., Shcherbachev, Y.N. & Marquardt, B. (2004) Grenadiers (Gadiformes, Teleostei) of Walters Shoals, southwestern Indian Ocean, with description of a new “West-Wind Drift” species. *Proceedings of the California Academy of Sciences*, 55, 190–207.
- Iwamoto, T. & Stein, D.L. (1974) A systematic review of the rattail fishes (Macrouridae: Gadiformes) from Oregon and adjacent waters. *Occasional Papers of the California Academy of Sciences*, (111), 1–79.
- Iwamoto, T. & Ungaro, N. (2002) A new grenadier (Gadiformes, Macrouridae) from the Mediterranean. *Cybius*, 26, 27–32.
- Iwamoto, T. & Williams, A. (1999) Grenadiers (Pisces, Gadiformes) from the continental slope of western and northwestern Australia. *Proceedings of the California Academy of Sciences*, 51, 105–243.
- Iwatsubo, H. & Motomura, H. (2017) *Field Guide to Fishes of Kagoshima Bay in Southern Kyushu, Japan*. Kagoshima Museum of Aquatic Biodiversity, Kagoshima, and Kagoshima University Museum, Kagoshima, 304 pp. [In Japanese.]
- Iwatsuki, Y., Nagino, H., Tanaka, F., Wada, H., Tanahara, K., Wada, M., Tanaka, H., Hidaka, K. & Kimura, S. (2017) Annotated checklist of marine and freshwater fishes in the Hyuga Nada area, southern Japan. *Bulletin of the Graduate School of Bioresources Mie University*, (43), 27–55.
- Izuka, A. & Matsuura, K. (1920) *Catalogue of the Zoological Specimens Exhibited in the Natural History Department, Tōkyō Imperial Museum*. Tokyo Imperial Museum, Tokyo, 2 + 3 + 6 + iii + 192 + 25 + 23 pp., 5 pls. [In Japanese with English preface.]
- Jamieson, A.J., Fujii, T., Solan, M., Matsumoto, A.K., Bagley, P.M. & Priede, I.G. (2009) Liparid and macrourid fishes of the hadal zone: *in situ* observations of activity and feeding behaviour. *Proceedings of the Royal Society B*, 279, 1037–1045.
<http://dx.doi.org/10.1098/rspb.2008.1670>
- Jamieson, A.J., Priede, I.G. & Craig, J. (2012) Distinguishing between the abyssal macrourids *Coryphaenoides yaquinae* and *C. armatus* from *in situ* photography. *Deep-Sea Research, Part I*, 64, 78–85.
<http://dx.doi.org/10.1016/j.dsr.2012.02.001>
- Jeong, M.-K., Lee, S.-I. & Kim, J.-K. (2010) New record of Sagami grenadier, *Ventrifossa garmani* (Gadiformes: Macrouridae) from Korea. *Korean Journal of Systematic Zoology*, 26, 317–320.
<http://dx.doi.org/10.5635/kjsz.2010.26.3.317>
- Jordan, D.S. (1900) Notes on recent fish literature. *American Naturalist*, 34, 897–899.
<http://dx.doi.org/10.1086/277890>
- Jordan, D.S. & Evermann, B.W. (1898) The fishes of North and Middle America: a descriptive catalogue of the species of fish-like vertebrates found in the waters of North America north of the Isthmus of Panama. Part III. *Bulletin of the United States National Museum*, (47), i–xxiv + 2183a–3136.
<http://dx.doi.org/10.5962/bhl.title.39717>
- Jordan, D.S. & Gilbert, C.H. (1899). The fishes of Bering Sea. In: Jordan, D.S. (Ed.), *Fur Seals, and Fur-Seal Islands of the North Pacific Ocean. Part 3*. Government Printing Office, Washington DC, pp. 433–492, pls. LXII–LXXXV.
<http://dx.doi.org/10.5962/bhl.title.52024>
- Jordan, D.S. & Hubbs, C.L. (1925) Record of fishes obtained by David Starr Jordan in Japan, 1922. *Memoirs of the Carnegie Museum*, 10, 93–346, pls. V–XII.
- Jordan, D.S. & Snyder, J.O. (1900) A list of fishes collected in Japan by Keinosuke Otaki, and by the United States steamer *Albatross*, with descriptions of fourteen new species. *Proceedings of the United States National Museum*, 23, 335–380, pls. IX–XX.
<http://dx.doi.org/10.5479/si.00963801.23-1213.335>

- Jordan, D.S. & Snyder, J.O. (1901) A preliminary check list of the fishes of Japan. *Annotationes Zoologicae Japonenses*, 3, 31–159.
- Jordan, D.S. & Starks, E.C. (1904) List of fishes dredged by the steamer *Albatross* off the coast of Japan in the summer of 1900, with descriptions of new species and a review of the Japanese Macrouridae. *Bulletin of the U.S. Fish Commission*, 22, 577–630, pls. 1–8.
- Jordan, D.S., Tanaka, S. & Snyder, J.O. (1913) A catalogue of the fishes of Japan. *Journal of the College of Science, Imperial University of Tokyo*, 33, 1–497.
- Jordan, D.S. & Thompson, W.F. (1914) Record of the fishes obtained in Japan in 1911. *Memoirs of the Carnegie Museum*, 6, 205–313, pls. XXIV–XLII.
<http://dx.doi.org/10.5962/bhl.title.43189>
- Kamohara, T. (1931a) Kōchi-shi fukin no gyorui [Fishes collected in the vicinity of Kochi City]. *Dōbutsugaku Zasshi* [Zoological Magazine, Japan], 43, 79–95. [In Japanese.]
- Kamohara, T. (1931b) Kōchi-shi fukin no gyorui tsuiki (1) [Additions to the fishes collected in the vicinity of Kochi City (1)]. *Dōbutsugaku Zasshi* [Zoological Magazine, Japan], 43, 533–544. [In Japanese.]
- Kamohara, T. (1934a) Tosaoki no shinkaigyo ni tsuite [Deep-sea fishes of the Province of Tosa]. “*Shokubutsu oyobi Dōbutsu*” [Plants and Animals], 2, 1196–1202. [In Japanese.]
- Kamohara, T. (1934b) Kōchi-shi fukin no gyorui tsuiki (VI) [Supplementary notes on the fishes collected in the vicinity of Kochi City (VI)]. *Dōbutsugaku Zasshi* [Zoological Magazine, Japan], 46, 299–303. [In Japanese.]
- Kamohara, T. (1934c) Kōchi-shi fukin no gyorui tsuiki (VII) [Supplementary notes on the fishes collected in the vicinity of Kochi City (VII)]. *Dōbutsugaku Zasshi* [Zoological Magazine, Japan], 46, 457–463. [In Japanese.]
- Kamohara, T. (1936) Supplementary note on the fishes collected in the vicinity of Kōchi-shi (X). *Dōbutsugaku Zasshi* [Zoological Magazine, Japan], 48, 929–935. [In Japanese with English abstract.]
- Kamohara, T. (1937) On some rare and one new species of fishes from Japan. *Dōbutsugaku Zasshi* [Zoological Magazine, Japan], 49, 186–190. [In Japanese with English abstract.]
- Kamohara, T. (1938) *On the Offshore Bottom-Fishes of Prov. Tosa, Shikoku, Japan*. Maruzen Co., Ltd., Tokyo, 86 pp.
- Kamohara, T. (1950) *Description of the Fishes from the Provinces of Tosa and Kishu, Japan*. Kochi-Ken Bunkyo Kyōkai [Kochi Literature Education Association], Kochi, 288 + 5 + 48 + 26 pp. [In Japanese.]
- Kamohara, T. (1952) Revised descriptions of the offshore bottom-fishes of Prov. Tosa, Shikoku, Japan. *Reports of the Kochi University, Natural Science*, (3), 1–122.
- Kamohara, T. (1955) *Coloured Illustrations of the Fishes of Japan. 1st Edition*. Hoikusha Co., Ltd., Osaka, xi + 135 pp., 64 pls. [In Japanese.]
- Kamohara, T. (1958) A catalogue of fishes of Kochi Prefecture (Province Tosa), Japan. *Reports of the Usa Marine Biological Station*, 5, 1–76.
- Kamohara, T. (1961a) *Colored Illustrations of the Fishes of Japan. Revised Edition*. Hoikusha Co., Ltd., Osaka, xvii + 158 pp., 64 pls. [In Japanese.]
- Kamohara, T. (1961b) *Colored Illustrations of the Fishes of Japan. II*. Hoikusha Co., Ltd., Osaka, xi + 168 pp., 68 pls. [In Japanese.]
- Kamohara, T. (1961c) Notes on the type specimens of fishes in my laboratory. *Reports of the Usa Marine Biological Station*, 8, 1–9, pls. I–VII.
- Kamohara, T. (1964) Revised catalogue of fishes of Kōchi Prefecture, Japan. *Reports of the Usa Marine Biological Station*, 11, 1–99.
- Kamohara, T. (1967) *Fishes of Japan in Color*. Hoikusha Co., Ltd., Osaka, xi + 135 pp., 64 pls.
- Kamohara, T. & Okamura, O. (1985) *Colored Illustrations of the Marine Fishes of Japan. I*. Hoikusha Co., Ltd., Osaka, xvi + 212 pp., 72 pls. [In Japanese.]
- Kataoka, T. & Tomida, Y. (1981) *Fish Fauna of Mie Prefecture*. Mie Prefectural Museum, Tsu, 110 pp. [In Japanese with English summary.]
- Katayama, M. (1940) A catalogue of the fishes of Toyama Bay. *Toyama Hakubutu Gakkaishi* [Bulletin of the Natural History Society of Toyama], (3), 1–28. [In Japanese.]
- Katayama, M. (1942) A new macrouroid fish from the Japan Sea. *Dōbutsugaku Zasshi* [Zoological Magazine, Japan], 54, 332–334.
- Kawano, M., Miyake, H., Hoshino, N., Ito, K., Yamanaka, T., Komoto, R., Chubachi, T., Anzawa, W., Ikeda, S., Ookei, N., Kinoshita, H., Kodama, K., Tega, T., Yamasaki, A., Mori, T., Nagahama, T., Ootani, T., Yamada, H., Murayama, T., Ando, A., Kai, S., Doi, H., Sugiyama, H., Iida, S. & Funaki, S. (2014) List of the fishes in the Japan Sea. *Bulletin of Yamaguchi Prefectural Fisheries Research Center*, (11), 1–30. [In Japanese with English abstract.]
- Kikuchi, K. (1931) Toyama-wan-san gyorui mokuroku [List of fishes collected from Toyama Bay]. *Toyama Kyōiku* [Toyama Education], (206), 1–34. [In Japanese.]
- Kim, B.-J., Kim, I.-S., Nakaya, K., Yabe, M., Choi, Y. & Imamura, H. (2009a). Checklist of the fishes from Jeju Island, Korea. *Bulletin of Fisheries Sciences, Hokkaido University*, 59, 7–36.
- Kim, I.-S., Choi, Y., Lee, C.-L., Lee, Y.-J., Kim, B.-J. & Kim, J.-H. (2005) *Illustrated Book of Korean Fishes*. Kyohak Publishing Co., Ltd., Seoul, 615 pp. [In Korean.]
- Kim, S.-Y., Iwamoto, T. & Yabe, M. (2009b) Four new records of grenadiers (Macrouridae, Gadiformes, Teleostei) from Korea. *Korean Journal of Ichthyology*, 21, 106–117.
- King, N.J. & Priede, I.G. (2008). *Coryphaenoides armatus*, the Abyssal Grenadier: global distribution, abundance, and ecology as determined by baited landers. In: Orlov, A.M. & Iwamoto, T. (Eds.), *Grenadiers of the World Ocean: Biology, Stock Assessment, and Fisheries*. American Fisheries Society Symposium 63. American Fisheries Society, Maryland, pp. 139–161.
- Kishida, S. (1986) *Coelorinchus multispinulosus*. In: Okamura,

- O. (Ed.), *Fishes of the East China Sea and the Yellow Sea*. Seikai Regional Fisheries Research Laboratory, Nagasaki, p. 100. [In Japanese.]
- Kitagawa, D., Imamura, H., Goto, T., Ishito, Y., Fujiwara, K. & Ueda, Y. (2008) *Field Guide of the Fishes from the Tohoku District, North-Eastern Waters of Japan*. Tokai University Press, Hadano, xvii + 140 pp. [In Japanese.]
- Kobayashi, H. (1956) Atsumi-wan no gyorui fu sonota no suisan dōbutsu [Marine fishes and other aquatic animals of Atsumi Bay]. In: Aichi Prefecture (Ed.), *Mikawa-wan Shizen-kōen Chōsa Hōkokusho* [Survey Report in the Atsumi Bay Nature Park]. Aichi Prefecture, Aichi, pp. 62–77. [In Japanese.]
- Kobayashi, K. & Ikeda, T. (1962) Ichthyofauna of the marine waters adjacent to Shirikishinai in southern Hokkaido, Japan. *Fauna and Flora of the Sea around the Shirikishinai Marine Station*, (4), 1–21.
- Koefoed, E. (1927) Fishes from the sea-bottom from the “Michael Sars” North Atlantic Deep-Sea Expedition 1910. *Scientific Results of the Michael Sars North Atlantic Deep-Sea Expedition 1910*, 4, 1–148, pls. I–VI.
- Köhler, R. (1896) Résultats scientifiques de la Campagne du “Caudan” dans le Golfe de Gascogne. Aout-Septembre 1895. Poissons. *Annales de l'Université de Lyon*, 26, 475–526, pls. XXVI–XXVII.
<http://dx.doi.org/10.5962/bhl.title.58631>
- Kuroda, N. (1941) Suruga Shizuura fukin-san gyorui mokuroku tsuika (dai 6) [Additions to the fish fauna in the vicinity of Suruga and Shizuura (no. 6)]. *Shokubutsu oyobi Dōbutsu* [Plants and Animals], 9, 75–80. [In Japanese.]
- Kuroda, N. (1947) Additions to the catalogue of the fishes of Shizuura, Suruga and the vicinity, (VIII). *Seibutsu* [Biology], (suppl. 1), 25–31. [In Japanese.]
- Kuroda, N. (1951) A nominal list with distribution of the fishes of Suruga Bay, inclusive of the freshwater species found near coast (continued). *Japanese Journal of Ichthyology*, 1, 376–394. [In Japanese with English abstract.]
- Kuroda, N. (1952) Additions to the fishes of Suruga Bay (X), with corrections. *Dōbutsugaku Zasshi* [Zoological Magazine, Japan], 61, 169–177. [In Japanese with English summary.]
- Kuroda, N. (1962) On the life colours of some fishes—XV. *Japanese Journal of Ichthyology*, 10, 7–12. [In Japanese with English summary.]
- Kuroda, N. (1965) Additions and corrections to the list of fishes of Suruga Bay XVIII. *Dōbutsugaku Zasshi* [Zoological Magazine, Japan], 74, 88–90. [In Japanese with English abstract.]
- Kuroda, N. (1966) Additions and corrections to the list of fishes of Suruga Bay. XIX. *Dōbutsugaku Zasshi* [Zoological Magazine, Japan], 75, 27–29. [In Japanese with English abstract.]
- Kuroda, N. (1971) New additions and corrections to the list of fishes of Suruga Bay, Japan (no. 21). *Dōbutsugaku Zasshi* [Zoological Magazine, Japan], 80, 52–57. [In Japanese with English abstract.]
- Kyushin, K., Amaoka, K., Nakaya, K. & Ida, H. (1973) *Tōbu-Indoyō Andaman Shūhen Kaiiki no Gyorui* [Fishes in the Vicinity of Andaman, Eastern Indian Ocean]. Marine Fisheries Research and Development Center, Tokyo, 114 pp. [In Japanese.]
- Laptikhovsky, V., Gaither, M.R. & Black, A. (2013) A Pacific grenadier *Coryphaenoides acrolepis* in the south-west Atlantic and environmental changes in the Falkland deep seas. *Marine Biodiversity Records*, 6, 1–5.
<http://dx.doi.org/10.1017/S1755267213001061>
- Last, P.L., Pogonoski, J.J., Gledhill, D.C., White, W.T. & Walker, C.J. (2014) The deepwater demersal ichthyofauna of the western Coral Sea. *Zootaxa*, 1887: 191–224.
<https://doi.org/10.11646/zootaxa.3887.2.4>
- Lindberg, G.U. (1959) List of marine fauna of southern Sakhalin and south Kuril Islands. *Issledovaniia Dal'nevostochnykh Morei SSSR*, 6, 172–256. [In Russian.]
- Lindberg, G.U. & Legeza, M.I. (1969) *Fishes of the Sea of Japan and the Adjacent Areas of the Sea of Okhotsk and the Yellow Sea, Part 2. English Edition*. Israel Program for Scientific Translations, Jerusalem, 391 pp. [Originally published in Russian in 1965.]
- Lowe, R.T. (1843) Notices of fishes newly observed or discovered in Madeira during the years 1840, 1841, and 1842. *Proceedings of the Zoological Society of London*, 1843 (pt. 11), 81–95.
- Maeda, K. & Maruyama, S. (1991) Hokkaidō-san gyorui risuto [List of fishes of Hokkaido]. In: Nagasawa, K. & Torisawa, M. (Eds.), *Fishes and Marine Invertebrates of Hokkaido: Biology and Fisheries*. Kitanihon Kaiyo Center, Sapporo, pp. 360–377. [In Japanese.]
- Maeda, K. & Tsutsui, D. (2003) Hokkaidō-san gyorui risuto [List of fishes of Hokkaido]. In: Mizushima, T. & Torisawa, M. (Eds.), *Fisheries and Aquatic Life in Hokkaido*. Hokkaido Shimibun Press, Sapporo, pp. 481–504. [In Japanese.]
- Makushok, V.M. (1964) The specific identity of *Nematonurus longifilis* (Günther, 1877) and *N. clarki* (Jordan and Gilbert, 1898) and some remarks on age-dependent variations in Macrouridae (Pisces). *Trudy Instituta Okeanologii im. P.P. Shirshova*, 73, 139–162. [In Russian with English summary; English translation by Israel Program of Scientific Translation, Jerusalem, published in 1966.]
- Makushok, V.M. (1976) New species of rattails, *Coryphaenoides subserrolatus*, n. sp. (Macrouridae, Osteichthyes) collected to the south of New Zealand. *Trudy Instituta Okeanologii im. P.P. Shirshova*, 104, 144–155. [In Russian with English summary.]
- Marshall, N.B. (1964) Bathypelagic macrourid fishes. *Copeia*, 1964 (1), 86–93.
<http://dx.doi.org/10.2307/1440835>
- Marshall, N.B. (1965) Systematic and biological studies of the macrourid fishes (Anacanthini-Teleostii). *Deep-Sea Research and Oceanographic Abstracts*, 12, 299–322.
[http://dx.doi.org/10.1016/0011-7471\(65\)90004-5](http://dx.doi.org/10.1016/0011-7471(65)90004-5)
- Marshall, N.B. (1967) The olfactory organs of bathypelagic fishes. *Symposia of the Zoological Society of London*, 19,

- Marshall, N.B. (1973) Family Macrouridae [in part: *Hyomacrurus*, *Macrourus*, *Nematonurus*, *Chalinura*, *Lionurus*, *Echinomacrurus*, *Cetonurus*, *Paracetonurus*, *Kumba*, *Mataeocephalus*, *Trachonurus*, *Sphagemacrurus*, *Malacocephalus*, and *Ventrifossa*]. In: Cohen, D.M. (Ed.), *Fishes of the Western North Atlantic, Part 6*. Sears Foundation for Marine Research, Yale University, New Haven, pp. 564, 581–600, 613–623, 650–662.
- Marshall, N.B. (1979) *Development in Deep-Sea Biology*. Blandford Press, Poole, 566 pp.
- Marshall, N.B. & Iwamoto, T. (1973) Family Macrouridae [in part: *Cynomacrurus*, *Odontomacrurus*, *Coelorhynchus*, *Coryphaenoides*, *Hymenocephalus*, and *Nezumia*]. In: Cohen, D.M. (Ed.), *Fishes of the Western North Atlantic, Part 6*. Sears Foundation for Marine Research, Yale University, New Haven, pp. 496–515, 530–563, 565–580, 601–612, 624–649.
- Maruyama, K. (1971) A catalogue of fishes of Iwate Prefecture, Japan. *Bulletin of Iwate Prefectural Fisheries Experimental Station*, (1), 1–70. [In Japanese.]
- Masuda, H. & Kobayashi, Y. (1994) *Grand Atlas of Fish Life Modes. Color Variation in Japanese Fish*. Tokai University Press, Tokyo, 460 pp. [In Japanese.]
- Matsubara, K. (1936) On two new species of fishes found in Japan. *Annotationes Zoologicae Japonenses*, 15, 355–360.
- Matsubara, K. (1943) Ichthyological annotations from the depth of the Sea of Japan. VIII–IX. *Journal of the Sigenkagaku Kenkyusyo*, 1, 131–152.
- Matsubara, K. (1955) *Fish Morphology and Hierarchy*. Ishizaki Shoten, Tokyo, xi + v + 1950 + viii pp., 135 pls. [In Japanese.]
- Matsubara, K. (1965) Osteichthyes [sic] [in part: Macrouridae]. In: Okada, K., Uchida, K. & Uchida, T. (Eds.), *New Illustrated Encyclopedia of the Fauna of Japan*. Hokuryukan Publishing Co., Ltd., Tokyo, pp. 504–510. [In Japanese.]
- Matsubara, K. & Asano, N. (1943) On a new macrouroid fish, *Coelorhynchus kanetomoi*, obtained from Misaki. *Bulletin of the Biogeographical Society of Japan*, 13, 95–98.
- Matsubara, K., Okada, Y., Suzuki, K. & Hashimoto, T. (1951) Cyclostoma and Pisces. In: Mie-ken Seibutsu Chōsa Inkai [Committee of the Biological Survey of Mie Prefecture] (Ed.), *Mie-ken-san, Seibutsu Mokuroku [List of Organisms of Mie Prefecture]*, Mie Prefecture, Tsu, pp. 19–43. [In Japanese.]
- Matsumoto, H. (2005) Occurrence of fishes off Uiyagawa, Shimane Prefecture (I). *Report of Shimane Prefectural Fisheries Experimental Station*, (12), 79–86. [In Japanese.]
- Matsuura, K. (1997) Fish collection building in Japan, with comments on major Japanese ichthyologists. In: Pietsch, T.W. & Anderson, W.D. Jr. (Eds.), *Collection Building in Ichthyology and Herpetology*. Allen Press, Lawrence, pp. 171–182.
- Matsuura, K. & Kimura, S. (2007) New fishes of Japan: part 1. *Bulletin of the National Museum of Nature and Science, Series A*, (suppl. 1), i.
- Matsuura, K., Shinohara, G. & Nakae, M. (2009) Historical fish specimens collected from the Tohoku District by the Saito Ho-on Kai Museum of Natural History. *Bulletin of the National Museum of Nature and Science, Series A*, 35, 9–54.
- Maul, G.E. (1951) Monografia dos peixes do Museu Municipal do Funchal. Família Macrouridae e Merlucciidae. *Boletim do Museu Municipal do Funchal*, (5), 5–55.
- Maul, G.E. & Koefoed, E. (1950) On a new genus and species of macrourid fish, *Phalacromacrurus pantherinus*. *Annals and Magazine of Natural History, Series 12*, 3, 970–976. <http://dx.doi.org/10.1080/00222935008656101>
- McAllister, D.E. (1990) A list of the fishes of Canada. *Syllogeus*, (64), 1–310.
- McCann, C. & McKnight, D.G. (1980) The marine fauna of New Zealand: macrourid fishes (Pisces: Gadida). *New Zealand Oceanographic Institute Memoir*, 61, 1–91.
- McCulloch, A.R. (1907) The results of deep sea investigation in the Tasman Sea. II.—The expedition of the “Woy Woy.” 1. Fishes and crustaceans from eight hundred fathoms. *Records of the Australian Museum*, 6, 345–355, pls. LXIII–LXV. <http://dx.doi.org/10.3853/j.0067-1975.6.1907.1019>
- McMillan, P. & Iwamoto, T. (2009) Two new species of *Coelorhynchus* (Teleostei, Gadiformes, Macrouridae) from the Tasman Sea. *Proceedings of the California Academy of Sciences*, 60, 39–51.
- McMillan, P. & Iwamoto, T. (2014) Descriptions of four species of grenadier fishes of the genera *Hymenocephalus* and *Hymenogadus* (Teleostei, Gadiformes, Macrouridae) from the New Zealand region and Tasman Sea, including two new species of *Hymenocephalus*. *Zootaxa*, 3856, 117–134. <http://dx.doi.org/10.11646/zootaxa.3856.1.5>
- McMillan, P.J. & Paulin, C.D. (1993) Descriptions of nine new species of rattails of the genus *Caelorinchus* (Pisces, Macrouridae) from New Zealand. *Copeia*, 1993 (3), 819–840. <http://dx.doi.org/10.2307/1447247>
- Mecklenburg, C.W., Mecklenburg, T.A. & Thorsteinson, L.K. (2002) *Fishes of Alaska*. American Fisheries Society, Maryland, xxxvii + 1037 pp.
- Melo, M.R.S., Braga, A.C., Nunan, G.W.A. & Costa, R.A.S. (2010) On new collections of deep-sea Gadiformes (Actinopterygii: Teleostei) from the Brazilian continental slope, between 11° and 23° S. *Zootaxa*, 2433, 25–46. <https://doi.org/10.11646/zootaxa.2433.1.2>
- Merrett, N.R. (1989) The elusive macrourid alevin and its seeming lack of potential in contributing to interfamilial systematics. In: Cohen, D.M. (Ed.), *Papers on the Systematics of Gadiform Fishes*. Natural History Museum of Los Angeles County, Los Angeles. pp. 175–185.
- Merrett, N.R. (2006) Bathygadidae and Macrouridae: Grenadiers. In: Richards, W.J. (Ed.), *Early Stages of*

Atlantic Fishes. An Identification Guide for the Western Central North Atlantic, Vol. I. CRC Press, Boca Raton, pp. 595–616.

- Merrett, N.R. & Headrich, R.L. (1997) *Deep-Sea Demersal Fish and Fisheries*. Chapman & Hall, London, xii + 282 pp.
- Merrett, N.R. & Iwamoto, T. (2000) Pisces Gadiformes: grenadier fishes of the New Caledonian region, southwest Pacific Ocean. Taxonomy and distribution, with ecological notes. *Mémoires du Muséum National d'Histoire Naturelle*, 184, 723–781.
- Merrett, N.R., Sazonov, Y.I. & Shcherbachev, Y.N. (1983) A new genus and species of rattail fish (Macrouridae) from the eastern North Atlantic and eastern Indian Ocean, with notes on its ecology. *Journal of Fish Biology*, 22, 549–561.
<http://dx.doi.org/10.1111/j.1095-8649.1983.tb04214.x>
- Ministry of the Environment (Government of Japan) (2011) *Marine Biodiversity Conservation Strategy*. Japanese Ministry of the Environment, Tokyo, 51 + 23 pp.
- Miya, M., Yamaguchi, Y. & Okiyama, M. (1995) Midwater fishes off the Pacific coast of Boso Peninsula, central Japan: species composition, abundance, biomass, and zoogeographic affinities. *Japanese Journal of Ichthyology*, 42, 237–248.
- Miyazaki, Y., Teramura, A. & Senou, H. (2019) Preliminary report on bycatch fish species collected from the Tokyo Submarine Canyon, Japan. *ZooKeys*, 843, 117–128.
<http://dx.doi.org/10.3897/zookeys.843.32410>
- Moehring, P.H.G. (1752). *Avium Genera*, Aurica, 88 pp. [See also Moehring (1758).]
<http://dx.doi.org/10.5962/bhl.title.49203>
- Moehring, P.H.G. (1758) *Geslachten der Vögelen*. Pieter Meijer, Amsterdam, 28 pp. [Latin MS in 1752, 58 pp.]
<http://dx.doi.org/10.5962/bhl.title.13336>
- Moore, J.A., Hartel, K.E., Craddock, J.E. & Galbraith, J.K. (2003) An annotated list of deepwater fishes from off the New England region, with new area records. *Northeastern Naturalist*, 10, 159–248.
<http://dx.doi.org/10.2307/3858285>
- Mori, T. (1952) Check list of the fishes of Korea. *Memoirs of the Hyogo University of Agriculture*, 1, 1–228.
- Mori, T. (1956) Fishes of San-in District including Oki Islands and its adjacent waters (southern Japan Sea). *Memoirs of the Hyogo University of Agriculture, Biological Series*, 2, 1–62. [In Japanese.]
- Morita, T. (1999) Molecular phylogenetic relationships of the deep-sea fish genus *Coryphaenoides* (Gadiformes: Macrouridae) based on mitochondrial DNA. *Molecular Phylogenetics and Evolution*, 13, 447–454.
<http://dx.doi.org/10.1006/mpev.1999.0661>
- Motomura, H. (2020) *List of Japan's All Fish Species. Current Standard Japanese and Scientific Names of All Fish Species Recorded from Japanese Waters*. The Kagoshima University Museum, Kagoshima, 560 pp. [In Japanese.]
- Motomura, H., Kuriwa, K., Katayama, E., Senou, H., Ogihara, G., Meguro, M., Matsunuma, M., Takata, Y., Yoshida, T., Yamashita, M., Kimura, S., Endo, H., Murase, A., Iwatsuki, Y., Sakurai, Y., Harazaki, S., Hidaka, K., Izumi, H. & Matsuura, K. (2010) Annotated checklist of marine and estuarine fishes of Yaku-shima Island, Kagoshima, southern Japan. In: Motomura, H. & Matsuura, K. (Eds.), *Fishes of Yaku-shima Island*. National Museum of Nature and Science, Tokyo, pp. 65–248.
- Mundy, B.C. (2005) Checklist of the fishes of the Hawaiian Archipelago. *Bishop Museum Bulletins in Zoology*, (6), 1–703.
- Nagasaki Museum of History and Culture (2015). Visual encyclopedia. Japan in Edo period as witnessed by Keiga Kawahara (I). Keiga's works in the Netherlands and Japan. Nagasaki Museum of History and Culture, Nagasaki. Available from: <http://www.nmhc.jp/keiga01/> (accessed 15 May 2020)
- Nakabo, T. (1993) Macrouridae. In: Nakabo, T. (Ed.), *Fishes of Japan with Pictorial Keys to the Species. 1st Edition*. Tokai University Press, Tokyo, pp. 353–371, 1276–1277. [In Japanese.]
- Nakabo, T. (2000) Macrouridae. In: Nakabo, T. (Ed.), *Fishes of Japan with Pictorial Keys to the Species. 2nd Edition*. Tokai University Press, Tokyo, pp. 417–435, 1494. [In Japanese.]
- Nakabo, T. (2002) Macrouridae. In: Nakabo, T. (Ed.), *Fishes of Japan with Pictorial Keys to the Species. English Edition*. Tokai University Press, Tokyo, pp. 417–435, 1491–1492. [In Japanese.]
- Nakabo, T. & Hirashima, Y. (2015) *Scientific Names of Japanese Fishes: Etymology*. Tokai University Press, Hadano, xv + 372 pp. [In Japanese.]
- Nakabo, T. & Kai, Y. (2013) Macrouridae. In: Nakabo, T. (Ed.), *Fishes of Japan with Pictorial Keys to the Species. 3rd Edition*. Tokai University Press, Hadano, pp. 493–512, 1872–1876. [In Japanese.]
- Nakae, M., Motomura, H., Hagiwara, K., Senou, H., Koeda, K., Yoshida, T., Tashiro, S., Jeong, B., Hata, H., Fukui, Y., Fujiwara, K., Yamakawa, T., Aizawa, M., Shinohara, G. & Matsuura, K. (2018) An annotated checklist of fishes of Amami-oshima Island, the Ryukyu Islands, Japan. *Memoirs of the National Museum of Nature and Science, Tokyo*, 52, 205–361.
- Nakajima, T. (1975) A list of the fishes from the coastal waters of Aichi Prefecture, Honshyu, Japan. *Bulletin of the Biogeographical Society of Japan*, 30, 43–56. [In Japanese with English abstract.]
- Nakajima, T. (2003) *Records of Fishes from Coastal and Offshore Waters of Aichi Prefecture, Honshu, Japan*. Published by author, Toyohashi, 198 pp., 79 pls. [In Japanese.]
- Nakayama, N. (2016) Commercial trawl landings by local fishermen: an important source of scientific specimens for deep-sea fish taxonomy in Japan. In: Nakayama, N. (Ed.), *Proceedings of the International Symposium on Interactions of Human, Culture and Nature Explored with University Museum Collections*. The Kyoto University Museum, Kyoto, pp. 50–56.

- Nakayama, N. (2019) Family Macrouridae. In: Koeda, K. & Ho, H.-C. (Eds.), *Fishes of Southern Taiwan*. National Museum of Marine Biology and Aquarium, Pingtung, pp. 353–365.
- Nakayama, N. & Endo, H. (2012) A new grenadier of the genus *Nezumia* (Pisces: Gadiformes: Macrouridae) from southern Japan. *Zootaxa*, 3410, 61–68.
<http://dx.doi.org/10.11646/zootaxa.3410.1.5>
- Nakayama, N. & Endo, H. (2015) Redescription of *Nezumia infranudis* (Gilbert & Hubbs, 1920), with the first record of the species from the eastern Indian Ocean (Actinopterygii: Gadiformes: Macrouridae). *Marine Biology Research*, 11, 1108–1115.
<http://dx.doi.org/10.1080/17451000.2015.1064962>
- Nakayama, N. & Endo, H. (2016a) A new species of the grenadier genus *Coryphaenoides* (Actinopterygii: Gadiformes: Macrouridae) from Japan and a range extension of *Coryphaenoides rudis* Günther 1878 in the northwestern Pacific. *Ichthyological Research*, 64, 1–12.
<http://dx.doi.org/10.1007/s10228-016-0524-9>
- Nakayama, N. & Endo, H. (2016b) *Mesovagus*, a replacement name for the grenadier genus *Mesobius* Hubbs and Iwamoto 1977 (Actinopterygii: Gadiformes: Macrouridae), a junior homonym of *Mesobius* Chamberlin 1951 (Chilopoda: Lithobiomorpha: Lithobiidae). *Ichthyological Research*, 64, 120–122.
<http://dx.doi.org/10.1007/s10228-016-0531-x>
- Nakayama, N. & Endo, H. (2017) A new species of the grenadier genus *Coelorinchus* (Actinopterygii: Gadiformes: Macrouridae) from the Timor Sea, eastern Indian Ocean. *Ichthyological Research*, 65:12–20.
<http://dx.doi.org/10.1007/s10228-017-0585-4>
- Nakayama, N., Endo, H. & Schwarzhans, W. (2015a) A new grenadier of the genus *Hymenocephalus* from Tosa Bay, southern Japan (Actinopterygii: Gadiformes: Macrouridae). *Ichthyological Research*, 62, 504–511.
<http://dx.doi.org/10.1007/s10228-015-0464-9>
- Nakayama, N., Endo, H. & Uchikawa, K. (2015b) First record of the midwater grenadier, *Odontomacrus murrayi* (Actinopterygii: Gadiformes: Macrouridae), from the northwestern Pacific off Japan. *Species Diversity*, 20, 135–139.
<http://dx.doi.org/10.12782/sd.20.2.135>
- Nakayama, N., Matsunuma, M. & Endo, H. (2015c) Redescription of *Coelorinchus tokiensis* (Steindachner and Döderlein 1887) (Actinopterygii: Gadiformes: Macrouridae), with comments on its synonymy. *Ichthyological Research*, 63, 247–259.
<http://dx.doi.org/10.1007/s10228-015-0493-4>
- Nakayama, N., Prokofiev, A.M. & Kawai, T. (2020) *Coelorinchus posteromaculatus* (Actinopterygii, Gadiformes, Macrouridae), a new species of grenadier from the eastern Indian Ocean. *Ichthyological Research*, 8 pp.
<https://doi.org/10.1007/s10228-020-00741-x>
- Nakayama, N., Takaoka, H. & Miyamoto, K. (2018) First record of the grenadier *Coelorinchus sheni* (Actinopterygii: Gadiformes: Macrouridae) from Japan. *Species Diversity*, 23, 121–127.
<http://dx.doi.org/10.12782/specdiv.23.121>
- Naturalis (2020) Image collection. Naturalis Biodiversity Center, Leiden. Available from: <http://biportal.naturalis.nl/result?theme=arts> (accessed 15 May 2020)
- Nelson, J.S. (2006) *Fishes of the World. 4th Edition*. John Wiley & Sons, Inc., New Jersey, xv + 601 pp.
- Nelson, J.S., Grande, T.C. & Wilson, M.V.H. (2016) *Fishes of the World. 5th Edition*. John Wiley & Sons, Inc., New Jersey, xli + 707 pp.
- Nishida, T., Matsunaga, A., Nishida, T., Sashima, K. & Nakazono, A. (2004) The list of the fishes in Tsuyazaki Town, Munakata County. *Scientific Bulletin of the Faculty of Agriculture, Kyushu University*, 59, 113–136. [In Japanese with English summary.]
- Nishimura, S. (1992) *Guide to Seashore Animals of Japan with Color Pictures and Keys. Vol. I*. Hoikusha Publishing Co., Ltd., Osaka, xxxv + 425 pp., 72 pls. [In Japanese.]
- Norman, J.R. (1939) Fishes. The John Murray Expedition 1933–34. *Scientific Reports, John Murray Expedition*, 7, 1–116.
- Nybelin, O. (1957). Deep-sea bottom fishes. *Reports of the Swedish Deep-Sea Expedition 1947–1948*, 2, 247–345, pls. I–VII.
- Ohta, S. (1983) Photographic census of large-sized benthic organisms in the bathyal zone of Suruga Bay, central Japan. *Bulletin of the Ocean Research Institute, the University of Tokyo*, (15), 1–244.
- Ohtomi, J. (2006) Population structure, condition factor and relative growth of the grenadier *Coelorinchus jordani* (Gadiformes, Macrouridae) in Kagoshima Bay, southern Japan. *Nippon Suisan Gakkaishi*, 72, 217–222. [In Japanese with English abstract.]
- Okada, Y. (1955) *Fishes of Japan. Illustrations and Descriptions of Fishes of Japan*. Maruzen Co., Ltd., Tokyo, 1 + 434 + 28 pp.
- Okada, Y. & Matsubara, K. (1938) *Keys to the Fishes and Fish-Like Animals of Japan Including Kuril Islands, Southern Sakhalin, Bonin Islands, Ryukyu Islands, Korea and Formosa*. Sanseido Co., Ltd., Tokyo, x1 + 584 pp., 113 pls. [In Japanese.]
- Okada, Y., Suzuki, K. & Mori, K. (1959) Kumano-nada engan no teisei gyorui [List of demersal fishes from the coastal waters of Kumano-nada]. *Kumano-nada Engan Kokuritsu-Kōen Chōsa Hōkoku* [Survey Report in the Kumano-nada National Park], 11, 73–84, pls. 1–2. [In Japanese.]
- Okada, Y., Uchida, K. & Matsubara, K. (1935) *Color Atlas of Fishes of Japan*. Sanseido Co., Ltd., Tokyo, 436 pp., 166 pls. [In Japanese.]
- Okamura, O. (1963a) Two new and one rare macrouroid fishes of the genera, *Coelorhynchus* and *Lionurus*, found in the Japanese waters. *Bulletin of the Misaki Marine Biological Institute, Kyoto University*, (4), 21–35.
- Okamura, O. (1963b) A new macrouroid fish found in the adjacent waters of Formosa. *Bulletin of the Misaki Marine*

Biological Institute, Kyoto University, (4), 37–42.

- Okamura, O. (1970a) *Fauna Japonica, Macrourina (Pisces)*. Academic Press of Japan, Tokyo, 216 pp., 64 pls.
- Okamura, O. (1970b) Studies on the macrouroid fishes of Japan: morphology, ecology and phylogeny. *Reports of the Usa Marine Biological Station*, 17, 1–179, pls. I–V.
- Okamura, O. (1982) *Malacocephalus*, *Ventrifossa*, *Nezumia*, *Coryphaenoides*, and *Coelorhynchus* [in part]. In: Okamura, O., Amaoka, K. & Mitani, F. (Eds.), *Fishes of the Kyushu-Palau Ridge and Tosa Bay*. Japan Fisheries Resource Conservation Association, Tokyo, pp. 144–181, 347–352, 354.
- Okamura, O. (1984a) *Hymenocephalus*, *Hymenogadus*, *Malacocephalus*, *Ventrifossa*, *Nezumia*, *Abyssicola*, and *Coelorinchus* [in part]. In: Okamura, O. & Kitajima, T. (Eds.), *Fishes of the Okinawa Trough and the Adjacent Waters I*. Japan Fisheries Resource Conservation Association, Tokyo, pp. 198–217, 220–221, 224–229, 357–364, 366.
- Okamura, O. (1984b) Macrouroidei. In: Masuda, H., Amaoka, K., Araga, C., Uyeno, T. & Yoshino, T. (Eds.), *The Fishes of the Japanese Archipelago. 1st Edition*. Tokai University Press, Tokyo, pp. 93–99, pls. 79–83. [In Japanese.]
- Okamura, O. (1985) *Fishes of the Okinawa Trough and the Adjacent Waters II*. Japan Fisheries Resource Conservation Association, Tokyo, pp. 417–781.
- Okamura, O. (1988) Macrouroidei. In: Masuda, H., Amaoka, K., Araga, C., Uyeno, T. & Yoshino, T. (Eds.), *The Fishes of the Japanese Archipelago. 2nd Edition*. Tokai University Press, Tokyo, pp. 93–99, 453, pls. 79–83, 344–373.
- Okamura, O. (1997) Macrouridae. In: Okamura, O. & Amaoka, K. (Eds.), *Sea Fishes of Japan*. Yama-Kei Publishers Co., Ltd., Tokyo, pp. 124–129. [In Japanese.]
- Okamura, O., Amaoka, K. & Mitani, F. (1982) *Fishes of the Kyushu-Palau Ridge and Tosa Bay*. Japan Fisheries Resource Conservation Association, Tokyo, 435 pp.
- Okamura, O. & Kitajima, T. (1984) *Fishes of the Okinawa Trough and the Adjacent Waters I*. Japan Fisheries Resource Conservation Association, Tokyo, 414 pp.
- Okamura, O. & Yatou, T. (1984) *Coelorinchus divergens* and *C. sparsilepis*. In: Okamura, O. & Kitajima, T. (Eds.), *Fishes of the Okinawa Trough and the Adjacent Waters I*. Japan Fisheries Resource Conservation Association, Tokyo, pp. 236–243.
- Orlov, A.M. & Tokranov, A.M. (2008) Some ecological and biological features of Giant and Popeye grenadiers in the Pacific waters off the northern Kuril Islands and southeastern Kamchatka. In: Orlov, A.M. & Iwamoto, T. (Eds.), *Grenadiers of the World Ocean: Biology, Stock Assessment, and Fisheries*. American Fisheries Society Symposium 63. American Fisheries Society, Maryland, pp. 225–260.
- Ozawa, T. (1983) Studies on the bottom fishes of continental slope off Makurazaki, southern Japan I—Faunal composition and variation of abundance—. *Bulletin of the Japanese Society of Fisheries Oceanography*, (44), 9–16.
- Quelch, J.J. (1884) Preliminary notice of new genera and species of ‘Challenger’ reef-corals. *Annals and Magazine of Natural History, Series 5*, 13, 292–297.
<http://dx.doi.org/10.1080/00222938409459237>
- Parin, N.V., Nesis, K.N., Sagaidachny, A.Y. & Shcherbachev, Y.N. (1993) Fauna of Walters Shoals, a seamount in the southwestern Indian Ocean. *Trudy Instituta Okeanologii im. P.P. Shirshova*, 128, 199–216. [In Russian with English summary.]
- Parr, A.E. (1946) The Macrouridae of the western North Atlantic and Central American seas. *Bulletin of the Bingham Oceanographic Collection, Yale University*, 10, 1–99.
- Pequeño, G. (1971) Sinopsis de Macrouriformes de Chile (Pisces, Teleostomi). *Boletim do Museu Nacional de Chile*, 32, 269–298.
- Radcliffe, L. (1912) Description of a new family, two new genera, and twenty-nine new species of anacanthine fishes from the Philippine Islands and contiguous waters. *Proceedings of the United States National Museum*, 43, 105–140, pls. 22–31.
<http://dx.doi.org/10.5479/si.00963801.43-1924.105>
- Randall, J.E. & Lim, K.K.P. (2000) A checklist of the fishes of the South China Sea. *The Raffles Bulletin of Zoology*, (suppl. 8), 569–667.
- Rass, T.S. (1955) Deep-water fishes of the Kurile-Kamchatka Trench. *Trudy Instituta Okeanologii im. P.P. Shirshova*, 12, 328–339. [In Russian.]
- Regan, C.T. (1921) New fishes from deep water off the coast of Natal. *Annals and Magazine of Natural History, Series 9*, 7, 412–420.
<http://dx.doi.org/10.1080/00222932108632540>
- Richardson, J. (1839) Account of an interesting collection of fish formed at Port Arthur in Van Diemen’s Land, by T.J. Lempriere, Esq. *Proceedings of the Zoological Society of London*, 1839 (pt. 7), 95–100.
- Risso, A. (1810) *Ichthyologie de Nice, ou Histoire Naturelle des Poissons du Département des Alpes Maritimes*. F. Schoell, Paris, i–xxxvi + 1–388 pp., 1–11 pls.
<http://dx.doi.org/10.5962/bhl.title.7052>
- Roa-Varón, A. & Ortí, G. (2009) Phylogenetic relationships among families of Gadiformes (Teleostei, Paracanthopterygii) based on nuclear and mitochondrial data. *Molecular Phylogenetics and Evolution*, 52, 688–704.
<http://dx.doi.org/10.1016/j.ympev.2009.03.020>
- Roberts, C.D. (1993) Comparative morphology of spined scales and their phylogenetic significance in the Teleostei. *Bulletin of Marine Science*, 52, 60–113.
- Roule, L. & Angel, F. (1933) Poissons provenant des campagnes du Prince Albert Ier de Monaco. *Résultats des Campagnes Scientifiques Accomplies sur son Yacht par Albert Ier Monaco*, fasc. 86, 1–115 (+ 1–4), pls. I–IV.
- Santos, R.S., Porteiro, F.M. & Barreiros, J.P. (1997) Marine fishes of the Azores: an annotated checklist and bibliography. *Arquipélago. Life and Marine Sciences (Bulletin of the University of Azores)* (suppl. 1), i–xxiii + 1–242.

- Saruwatari, T., Lopez, J.A. & Pietsch, T.W. (1997) Cyanine blue: a versatile and harmless stain for specimen observations. *Copeia*, 1997, 840–841.
<https://doi.org/10.2307/1447302>
- Sawada, Y. (1983) Macrouridae. In: Amaoka, K., Nakaya, K., Araya, H. & Yasui, T. (Eds.), *Fishes from the North-Eastern Sea of Japan and the Okhotsk Sea off Hokkaido*. Japan Fisheries Resource Conservation Association, Tokyo, pp. 106–113, 192–196, 248–251, 324–325.
- Sazonov, Y.I. (1985) Two new macrourid species (Gadiformes) from Northern Pacific Seamounts. *Journal of Ichthyology*, 25, 13–21. [Originally published in Russian in *Voprosy Ikhtiologii*, 25, 719–727.]
- Sazonov, Y.I. (1993) Discovery of a rare grenadier species, *Hymenocephalus papyraceus* (Gadiformes, Macrouridae), in the East China Sea. *Journal of Ichthyology*, 33, 117–121. [Originally published in Russian in *Voprosy Ikhtiologii*, 33, 446–448.]
- Sazonov, Y.I. (1994) Additions to the list of macrourids (Gadiformes, Bathygadidae, and Macrouridae) from the Northwest Pacific Ridge. *Journal of Ichthyology*, 34, 98–115. [Originally published in Russian in *Voprosy Ikhtiologii*, 34, 149–160.]
- Sazonov, Y.I. & Iwamoto, T. (1992) Grenadiers (Pisces, Gadiformes) of the Nazca and Sala y Gomez Ridges, southeastern Pacific. *Proceedings of the California Academy of Sciences*, 48, 27–95.
- Sazonov, Y.I. & Shcherbachev, Y.N. (1982) A preliminary review of grenadiers related to the genus *Cetonurus* Günther (Gadiformes, Macrouridae). Descriptions of new taxa related to the genera *Cetonurus* Günther and *Kumba* Marshall. *Journal of Ichthyology*, 22, 1–15. [Originally published in Russian in *Voprosy Ikhtiologii*, 22, 707–721.]
- Sazonov, Y.I. & Shcherbachev, Y.N. (1985) Preliminary review of grenadiers of the *Cetonurus* group (Gadiformes, Macrouridae). II. The genus *Cetonurus* Günther: taxonomic characters of the group. *Journal of Ichthyology*, 25, 12–26. [Originally published in Russian in *Voprosy Ikhtiologii*, 26, 179–195.]
- Sazonov, Y.I. & Shcherbachev, Y.N. (1997) The composition and some problems pertaining to the variability of species of subgenus *Sokodara* (genus *Ventrifossa*, Macrouridae). *Journal of Ichthyology*, 37, 529–533. [Originally published in Russian in *Voprosy Ikhtiologii*, 37, 543–547.]
- Schmidt, P.J. (1931) Fishes of Japan, collected in 1910. *Transactions of the Pacific Committee of the Academy of Sciences of the USSR*, 2, 1–176.
- Schwarzshans, W. (2014) Head and otolith morphology of the genera *Hymenocephalus*, *Hymenogadus* and *Spicomacrus* (Macrouridae), with the description of three new species. *Zootaxa*, 3888, 1–73.
<http://dx.doi.org/10.11646/zootaxa.3888.1.1>
- Senou, H. (2008) Vertebrata. In: Fujikura, K., Okutani, T. & Maruyama, T. (Eds.), *Deep-Sea Life—Biological Observations Using Research Submersibles*. Tokai University Press, Hadano, pp. 207–211, 356–382. [In Japanese.]
- Senou, H., Matsuura, K. & Shinohara, G. (2006) Checklist of fishes in the Sagami Sea with zoogeographical comments on shallow water fishes occurring along the coastlines under the influence of the Kuroshio Current. *Memoirs of the National Science Museum, Tokyo*, (41), 389–542.
- Shao, K.-T. (1993) Gadiformes. In: Shen, S.-C. (Ed.), *Fishes of Taiwan*. Department of Zoology, National Taiwan University, Taipei, pp. 165–173. [In Chinese.]
- Shao, K.-T., Ho, H.-C., Lin, P.-L., Lee, P.-F., Lee, M.-Y., Tsai, C.-Y., Liao, Y.-C., Lin, Y.-C., Chen, J.-P. & Yeh, H.-M. (2008a) A checklist of the fishes of southern Taiwan, northern South China Sea. *The Raffles Bulletin of Zoology*, (suppl. 19), 233–271.
- Shao, K.-T., Iwamoto, T., Ho, H.-C., Cheng, T.-Y. & Chen, C.-Y. (2008b) Species composition and distribution pattern of grenadiers (family [sic] Bathygadidae, Macrouridae, and Macrouridae [sic]) from Taiwan. In: Orlov, A.M. & Iwamoto, T. (Eds.), *Grenadiers of the World Ocean: Biology, Stock Assessment, and Fisheries*. American Fisheries Society Symposium 63. American Fisheries Society, Maryland, pp. 17–29.
- Shcherbachev, Y.N. (1987) Preliminary list of thalassobathyal fishes of the tropical and subtropical waters of the Indian Ocean. *Journal of Ichthyology*, 27, 37–46. [Originally published in Russian in *Voprosy Ikhtiologii*, 27, 3–11.]
- Shcherbachev, Y.N. & Iwamoto, T. (1995) Indian Ocean grenadiers of the subgenus *Coryphaenoides*, genus *Coryphaenoides* (Macrouridae, Gadiformes, Pisces). *California Academy of Sciences*, 48, 285–314.
- Shcherbachev, Y.N., Sazonov, Y.I. & Iwamoto, T. (1992) Synopsis of the grenadier genus *Kuronezumia* (Pisces: Gadiformes: Macrouridae), with description of a new species. *Proceedings of the California Academy of Sciences*, 48, 97–108.
- Sherborn, C.D. & Jentink, F.A. (1895) On the dates of the parts of Siebold's "Fauna Japonica" and Giebel's "Allgemeine Zoologie" (first edition). *Proceedings of the Zoological Society of London*, 1895, 149–150.
<http://dx.doi.org/10.1111/j.1469-7998.1895.tb07888.x>
- Shinohara, G., Endo, H. & Matsuura, K. (1996) Deep-water fishes collected from the Pacific coast of northern Honshu, Japan. *Memoirs of the National Science Museum, Tokyo*, (29), 153–185.
- Shinohara, G., Endo, H., Matsuura, K., Machida, Y. & Honda, H. (2001) Annotated checklist of the deepwater fishes from Tosa Bay, Japan. *National Science Museum Tokyo Monographs*, (20), 283–343.
- Shinohara, G., Kojima, S. & Fukumori, H. (2020) Deep-sea fishes from the Kurile-Kamchatka Trench collected by the R/V *Sonne* during the KuramBio II expedition in 2016. *Progress in Oceanography*, 183, 9 pp.
<https://doi.org/10.1016/j.pocean.2020.102275>
- Shinohara, G. & Matsuura, K. (1997) Annotated checklist of deep-water fishes from Suruga Bay, Japan. *National*

- Science Museum Tokyo Monographs*, (12), 269–318, pls. 1–2.
- Shinohara, G., Matsuura, K. & Shirai, S. (1998) Fishes of Tachibana Bay, Nagasaki, Japan. *Memoirs of the National Science Museum, Tokyo*, (30), 105–138.
- Shinohara, G., Nakae, M., Ueda, Y., Kojima, S. & Matsuura, K. (2014) Annotated checklist of deep-sea fishes of the Sea of Japan. *National Museum of Nature and Science Monographs*, (44), 225–291.
- Shinohara, G., Narimatsu, Y., Hattori, T., Ito, M., Takata, Y. & Matsuura, K. (2009) Annotated checklist of deep-sea fishes from the Pacific coast off Tohoku District, Japan. *National Museum of Nature and Science Monographs*, (39), 683–735.
- Shinohara, G., Sato, T., Aonuma, Y., Horikawa, H., Matsuura, K., Nakabo, T. & Sato, K. (2005) Annotated checklist of deep-sea fishes from the waters around the Ryukyu Islands, Japan. *National Science Museum Tokyo Monographs*, (29), 385–452.
- Shinohara, G., Shirai, S.M., Nazarkin, M.V. & Yabe, M. (2011) Preliminary list of the deep-sea fishes of the Sea of Japan. *Bulletin of the National Museum of Nature and Science, Series A*, 37, 35–62.
- Shinohara, G. & Williams, J.T. (2006) Historical Japanese fish specimens from the Sagami Sea in the National Museum of Natural History, Smithsonian Institution. *Memoirs of the National Science Museum, Tokyo*, (41), 543–568.
- Shiogaki, M. (1982) A catalogue of the fishes collected from the waters of Aomori Prefecture, Japan. *Bulletin of the Fisheries Experimental Station of Aomori Prefecture*, 1982, 1–36. [In Japanese.]
- Shiogaki, M. (1988) Zoogeography of the fishes of the coastal waters of Aomori Prefecture, northern Japan. *Memoirs of the National Science Museum, Tokyo*, (21), 153–162. [In Japanese with English summary.]
- Shiogaki, M. & Dotsu, Y. (1973) Fishes collected from the coastal waters of Nomozaki near Nagasaki. *Bulletin of the Faculty of Fisheries, Nagasaki University*, (35), 11–39. [In Japanese with English abstract.]
- Shiogaki, M., Ishito, Y., Nomura, Y. & Sugimoto, T. (2004) Revised catalogue of the fishes collected from the waters of Aomori Prefecture. *Bulletin of Aomori Prefectural Fisheries Research Center*, (4), 39–80. [In Japanese with English abstract.]
- Shiogaki, M., Nomura, Y. & Sugimoto, T. (1992) Further additions to the catalogue of fishes from the waters of Aomori Prefecture—I. *Scientific Reports of Aquaculture Center, Aomori Prefecture*, 7, 17–31. [In Japanese with English abstract.]
- Smith, H.M. & Pope, E.B. (1906) List of fishes collected in Japan in 1903, with descriptions of new genera and species. *Proceedings of the United States National Museum*, 31, 459–499.
<http://dx.doi.org/10.5479/si.00963801.31-1489.459>
- Smith, J.L.B. (1935) New and little known fishes from South Africa. *Records of the Albany Museum Grahamstown*, 4, 169–235, pls. XVIII–XXIII.
- Smith, J.L.B. (1953) *The Sea Fishes of Southern Africa, Revised Enlarged Edition*. Central News Agency, Ltd., South Africa, Cape Town, 564 pp., 107 pls.
- Smith, J.L.B. (1968) New and interesting fishes from deepish water off Durban, Natal and southern Mozambique. *Investigational Report, South African Association for Marine Biological Research, Oceanographic Research Institute*, (19), 1–30, pls. 1–6.
- Sonoyama, T., Ogimoto, K., Hori, S., Uchida, Y. & Kawano, M. (2020) An annotated checklist of marine fishes of the Sea of Japan off Yamaguchi Prefecture, Japan. *Bulletin of the Kagoshima University Museum*, (11), 1–152. [In Japanese with English abstract.]
- Steindachner, F. & Döderlein, L. (1887) Beiträge zur kenntniss der fische Japan's. (IV.). *Denkschriften der Mathematisch-Naturwissenschaftlichen Classe der Kaiserlichen Akademie der Wissenschaften in Wien*, 53, 257–296, pls. I–IV.
- Suetsugu, K. & Ohta, S. (2005) Day and nighttime changes in species composition of deep-sea demersal fishes. *Journal of Oceanography*, 61, 187–196.
<http://dx.doi.org/10.1007/s10872-005-0031-5>
- Suzuki, K. & Kataoka, T. (1997) *Marine Fishes of Mie Prefecture, Japan*. Toba Aquarium, Toba, 297 pp., 152 pls. [In Japanese.]
- Takata, J., Tokunaga, T., Machida, S., Tashiro, M. & Yamamoto, N. (1979) Chec [sic] list of fishes obtained with the small trawl net in the northwestern coastal water of Tsushima Island. *Bulletin of Nagasaki Prefectural Institute of Fisheries*, (5), 95–98. [In Japanese.]
- Takegawa, Y. & Morino, H. (1970) Fishes from Wakasa Bay, Japan Sea. *Publications of the Seto Marine Biological Laboratory*, 17, 373–392.
- Tamai, T., Ichikawa, H. & Arao, K. (2012) Landing fishes on Isshiki fishing port, Aichi Prefecture (part II). *Science Report of the Toyohashi Museum of Natural History*, (22), 33–40. [In Japanese.]
- Tanaka, S. (1927) Cyclostoma and Pisces. In: Uchida, K. (Ed.), *Figuraro de Japanaj Bestoj*. Hokuryukan Publishing Co., Ltd., Tokyo, pp. 254–481. [In Japanese.]
- Tanaka, S. (1951) *Dr. Tanaka's Japanese Fishes in Life Colours*. Kazamashobo Co., Ltd., Tokyo, 203 pp.
- Taranetz, A.J. (1937) Handbook for identification of fishes of Soviet Far East and adjacent waters, *Bulletin of the Pacific Scientific Institute of Fisheries and Oceanography*, 11, 1–200. [In Russian.]
- Tatara, K., Kitamori, R., Nagata, K., Mito, T., Hayashi, T. & Kudo, S. (1965) Setonaikai oyobi rinsetsu tairikudana ni okeru gyorui mokuroku fu nantai dōbutsu mokuroku [Checklist of fishes and mollusks of the Seto Inland Sea and adjacent waters]. *Publications of the Naikai Regional Fisheries Research Laboratory C*, (3), 77–123. [In

- Japanese.]
- Temminck, C.J. & Schlegel, H. (1846) Pisces. Fascicle 15. In: von Siebold, P.F. (Ed.), *Fauna Japonica*. Müller, Amsterdam, pp. 249–268.
- Thompson, W.F. (1916) Fishes collected by the United States Bureau of Fisheries steamer “Albatross” during 1888, between Montevideo, Uruguay, and Tome, Chile, on the voyage through the Straits of Magellan. *Proceedings of the United States National Museum*, 50, 401–476, pls. 2–6. <http://dx.doi.org/10.5479/si.00963801.50-2133.401>
- Tominaga, Y. & Uyeno, T. (1981) List of Japanese fishes. In: Yasuda, F., Takagi, K., Tominaga, Y., Uyeno, T., Abe, T., Ishiyama, R., Iwai, T., Ochiai, A., Kuronuma, K. & Nakamura, M. (Eds.), *Dictionary of Japanese Fish Names and Their Foreign Equivalents*. Sanseido Co., Ltd., Tokyo, pp. 437–574.
- Toponogov, V.N. & Kurennoi, A.A. (1986) Giant grenadier. In: Vinogradov, M.E., Parin, N.V. & Shuntov, V.P. (Eds.), *Biological Resources of the Pacific Ocean*. Nauka, Moscow, pp. 233–241. [In Russian.]
- Townsend, C.H. & Nichols, J.T. (1925) Deep sea fishes of the ‘Albatross’ Lower California Expedition. *Bulletin of the American Museum of Natural History*, 52, 1–20, pls. I–IV, map.
- Trunov, I.A. (1981) *Parakumba maculisquama* genus and species family Macrouridae from the South Atlantic. *Journal of Ichthyology*, 21, 27–34. [Originally published in Russian in *Voprosy Ikhtiologii*, 21, 28–36.]
- Tsuda, T. (1964) Toyama-wan-san gyorui bunpu mokuroku [Checklist of fishes of Toyama Bay]. *Habataki [Wingbeat]*, 2, 1–8. [In Japanese.]
- Tsuda, T. (1990) *Fishes of the Sea of Japan*. Katsura Shobo [Katsura Publishing], Toyama. [In Japanese.]
- Uchida, K., Doutsu, Y., Tafuku, M., Mito, T., Shojima, Y., Inoko, Y., Mito, S., Arima, K. & Furukawa, T. (1957) *Tsuyazaki suisan-shi (I) [Fishes of Tsuyazaki (I)]*. Fisheries Research Center, Faculty of Agriculture, Kyushu University, Tsuyazaki, 31 pp. [In Japanese.]
- Uchinomi, F. (1951) Nanbu-oki sokobiki gyorui mokuroku [List of deep demersal fishes collected from off Nanbu]. *Nanki Seibutsu [Nankai Biology]*, 3, 69–76, 88. [In Japanese.]
- Ueno, T. (1965) Hokkaidō kinkai no sakana (1) [Marine fishes of Hokkaido]. *Hokusuishi Geppō [Monthly Report of the Hokkaido Fisheries Experimental Station]*, 22, 13–20. [In Japanese.]
- Ueno, T. (1971) List of the marine fishes from the waters of Hokkaido and its adjacent regions. *Scientific Reports of the Hokkaido Fisheries Experimental Station*, (3), 61–102. [In Japanese with English abstract.]
- Vaillant, L.L. (1888) *Expéditions Scientifiques du “Travailleur” et du “Talisman” Pendant les Années 1880, 1881, 1882, 1883. Poissons*. G. Masson, Paris, 406 pp., 28 pls. <http://dx.doi.org/10.5962/bhl.title.98313>
- Weber, M. (1913) *Die Fische der Siboga-Expedition*. E.J. Brill, Leiden, xii + 710 pp., 12 pls. <http://dx.doi.org/10.5962/bhl.title.35825>
- Weber, M. & de Beaufort, L.F. (1929) *The Fishes of the Indo-Australian Archipelago. V. Anacanthini, Allotriognathi, Heterostomata, Berycomorphi, Percomorphi: Families Kuhliidae, Apogonidae, Plesiopidae, Pseudoplesiopidae, Priacanthidae, Centropomidae*. E.J. Brill Ltd., Leiden, xiv + 458 pp.
- Wessel, P., Smith, W.H.F., Scharroo, R., Luis, J. & Wobbe, F. (2013) Generic Mapping Tools: improved version released. *Eos, Transactions, American Geophysical Union*, 94, 409–410. <https://doi.org/10.1002/2013EO450001>
- Wheeler, A.C. & O’Riordan, C.E. (1969) Type material of fishes from the ‘Challenger’ Expedition in the Natural Museum of Ireland. *Proceedings of the Royal Irish Academy, Section B, Biological, Geological, and Chemical Science*, 68, 89–100.
- Whitley, G.P. (1931) New names for Australian fishes. *Australian Zoologist*, 6, 310–334, pls. XXV–XXVII.
- Wikimedia Commons (2020) Kawahara Collection at Naturalis Biodiversity Center. Wikimedia Foundation, San Francisco. https://commons.wikimedia.org/wiki/Category:Kawahara_Collection_at_Naturalis_Biodiversity_Center?uselang=nl (accessed 15 May 2020)
- Wilson, R.R. Jr. (1994) Interrelationships of the subgenera of *Coryphaenoides* (Gadiformes: Macrouridae): comparison of protein electrophoresis and peptide mapping. *Copeia*, 1994 (1), 42–50. <http://dx.doi.org/10.2307/1446669>
- Wilson, R.R. Jr. (2001) A new species of *Nezumia* (Gadiformes: Macrouridae) from Fieberling Guyot, eastern North Pacific Ocean. *Revista de Biología Tropical*, 49 (suppl. 1), 29–37.
- Wilson, R.R. Jr. & Attia, P. (2003) Interrelationships of the subgenera of *Coryphaenoides* (Teleostei: Gadiformes: Macrouridae): synthesis of allozyme, peptide mapping, and DNA sequence data. *Molecular Phylogenetics and Evolution*, 27, 343–347. [http://dx.doi.org/10.1016/S1055-7903\(02\)00419-0](http://dx.doi.org/10.1016/S1055-7903(02)00419-0)
- Wilson, R.R. Jr., Siebenaller, J.F. & Davis, B.J. (1991) Phylogenetic analysis of species of three subgenera of *Coryphaenoides* (Teleostei: Macrouridae) by peptide mapping of homologs of LDH-A₄. *Biochemical Systematics and Ecology*, 19, 227–287. [http://dx.doi.org/10.1016/0305-1978\(91\)90015-R](http://dx.doi.org/10.1016/0305-1978(91)90015-R)
- Wilson, R.R. Jr. & Waples, R.S. (1983) Distribution, morphology, and biochemical genetics of *Coryphaenoides armatus* and *C. yaquinae* (Pisces: Macrouridae) in the central and eastern North Pacific. *Deep-Sea Research, Part A*, 30, 1127–1145. [http://dx.doi.org/10.1016/0198-0149\(83\)90092-4](http://dx.doi.org/10.1016/0198-0149(83)90092-4)
- Wilson, R.R. Jr. & Waples, R.S. (1984) Electrophoretic and biometric variability in the abyssal grenadier *Coryphaenoides armatus* of the western North Atlantic, eastern South Pacific and eastern North Pacific Oceans. *Marine Biology*, 80, 227–237. <http://dx.doi.org/10.1007/BF00392817>

- Wood-Mason, J. & Alcock, A. (1891) Natural history notes from H.M. Indian marine survey steamer 'Investigator', commander R.F. Hoskyn, R.N., commanding. Series II., no. 1. On the results of deep-sea dredging during the season 1890–91. *Annals and Magazine of Natural History, Series 6*, 8, 16–34, 119–138, pls. VII–VIII.
<http://dx.doi.org/10.1080/00222939109460385>
- Yamada, U. (1986) *Coelorinchus jordani*. In: Okamura, O. (Ed.), *Fishes of the East China Sea and the Yellow Sea*. Seikai Regional Fisheries Research Laboratory, Nagasaki, pp. 101. [In Japanese.]
- Yamada, U., Shirai, S., Irie, T., Tokimura, M., Deng, S., Zheng, Y., Li, C., Kim, Y.U. & Kim, Y.S. (1995) *Names and Illustrations of Fishes from the East China Sea and the Yellow Sea*. Overseas Fishery Cooperation Foundation, Tokyo, xi + 288 pp.
- Yamada, U., Tokimura, M., Horikawa, H. & Nakabo, T. (2007) *Fishes and Fisheries of the East China and Yellow Seas*. Tokai University Press, Hadano. xxiii + 1262 pp.
- Yamaguchi, T. & Machida, Y. (2003) Fish specimens collected in Japan by Ph. F. von Siebold and H. Bürger and now held by the Nationaal Natuurhistorisch Museum in Leiden and other two museums. *CALANUS, Bulletin of the Aitsu Marine Station, Kumamoto University*, (special no. 4), 87–337, pls. 1–87.
- Yamashita, H. (1959) Saga-ken Genkai engan gyorui mokuroku [List of shore fishes of Genkai, Saga Prefecture]. Published by author, Genkai, 1–22 pp. [In Japanese.]
- Yanai, T. (1950) The fishes of the San-in District. *Dōbutsugaku Zasshi* [Zoological Magazine, Japan], 59, 17–22. [In Japanese.]
- Yatou, T. (1982) *Hymenocephalus* and *Coelorhynchus* [in part]. In: Okamura, O., Amaoka, K. & Mitani, F. (Eds.), *Fishes of the Kyushu-Palau Ridge and Tosa Bay*. Japan Fisheries Resource Conservation Association, Tokyo, pp. 140–143, 170–171, 180–181, 346–347, 353.
- Yatou, T. (1984) *Coryphaenoides* and *Coelorinchus* [in part]. In: Okamura, O. & Kitajima, T. (Eds.), *Fishes of the Okinawa Trough and the Adjacent Waters I*. Japan Fisheries Resource Conservation Association, Tokyo, pp. 218–223, 228–235, 224–245, 363–370.
- Yoda, M., Tokimura, M., Horikawa, H. & Yamada, U. (2002) *A Catalogue of Fishes from the East China and Yellow Seas with Their Local Names*. Seikai National Fisheries Research Institute, Fisheries Research Agency, Nagasaki, 41 pp. [In Japanese.]
- Youn, C.-H. (2002) *Fishes of Korea, with Pictorial Key and Systematic List*. Academy Publications Committee, Seoul, 747 pp. [In Korean.]
- Zama, A. (2001) *Fish Fauna of Miyagi Prefecture, Japan*. Published by author, Ishimaki, 154 pp. [In Japanese.]
- Zugmayer, E. (1911) Poissons provenant des campagnes du yacht *Princesse-Alice* (1901–1910). *Résultats des Campagnes Scientifiques Accomplies sur Son Yacht par Albert 1er Prince Souverain de Monaco*, fasc. 35, 1–174, pls. I–VI.
<http://dx.doi.org/10.5962/bhl.title.55724>

APPENDIX 1. Additional material examined

The following specimens were examined in the course of this study and used for generating distribution maps in Appendix 3; meristic and morphometric data were not obtained for these specimens.

Coelorinchus anatirostris. 210 specimens. BSKU 32470 (1), northwest of Miyako-jima Island, Okinawa Trough, 25.8000°N, 124.4250°E, 400–420 m, 14 Sept. 1979; BSKU 32515 (1), northwest of Miyako-jima Island, Okinawa Trough, 25.7917°N, 124.5117°E, 500–550 m, 10 Oct. 1979; BSKU 26603 (1), BSKU 26604 (1), BSKU 26605 (1), off Miyako-jima Island, Okinawa Trough, 26.3883°N, 125.0667°E, 680–770 m, 24 Jan. 1978; BSKU 27379 (1), BSKU 27380 (1), BSKU 27381 (1), west of Okinawa-jima Island, Okinawa Trough, 26.7833°N, 125.6167°E, 330–365 m, 2 Mar. 1978; BSKU 34350 (1), northwest of Kume-jima Island, Okinawa Trough, 27.0083°N, 125.8417°E, 410–420 m, 10 Dec. 1979; BSKU 33084 (1), north-northeast of Miyako-jima Island, Okinawa Trough, 25.9833°N, 125.8500°E, 430 m, 10 Oct. 1979; BSKU 27413 (1), northwest of Okinawa-jima Island, Okinawa Trough, 27.4000°N, 126.1333°E, 340–440 m, 6 Mar. 1978; BSKU 27473 (1), west of Okinawa-jima Island, Okinawa Trough, 27.7767°N, 126.2833°E, 490 m, 9 Mar. 1978; KPM-NI 6314 (1), KPM-NI 6315 (1), ca. 120 km south of Jeju Island, East China Sea, 32.2783°N, 126.6633°E, 110 m, 29 Apr. 1999; SNFR 11081 (1), East China Sea, 28.3750°N, 126.9683°E, 378–382 m, 23 Oct. 1990; BSKU 27723 (1), BSKU 27724 (1), north-northwest of Okinawa Island, Okinawa Trough, 28.5000°N, 127.0000°E, 225–235 m, 12 Mar. 1978; BSKU 27868 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.5833°N, 127.0000°E, 600 m, 16 Mar. 1978; BSKU 27269 (1), BSKU 27270 (1), BSKU 27271 (1), BSKU 27272 (1), BSKU 27273 (1), BSKU 27274 (1), BSKU 27275 (1), BSKU 27276 (1), BSKU 27277 (1), BSKU 27278 (1), BSKU 27281 (1), BSKU 27282 (1), BSKU 27283 (1), BSKU 27284 (1), west-northwest of Amami-oshima Island, Okinawa Trough, 28.8033°N, 127.0617°E, 425 m, 6 Feb. 1978; BSKU 27928 (1), BSKU 27929 (1), BSKU 27930 (1), BSKU 27931 (1), BSKU 27933 (1), BSKU 27934 (1), BSKU 27935 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.7500°N, 127.1167°E, 520–542 m, 16 Mar. 1978; BSKU 27688 (1), BSKU 27689 (1), BSKU 27690 (1), BSKU 27691 (1), BSKU 27692 (1), BSKU 27708 (1), BSKU 27709 (1), west-northwest of Amami-oshima Island, Okinawa Trough, 28.7000°N, 127.1500°E, 500–535 m, 11 Mar. 1978; BSKU 34670 (1), BSKU 34671 (1), BSKU 34672 (1), BSKU 34673 (1), BSKU 34674 (1), BSKU 34675 (1), BSKU 34676 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.9983°N, 127.1550°E, 325 m, 17 Dec. 1979; BSKU 29838 (1), BSKU 34697 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.0667°N, 127.2250°E, 420–450 m, 17 Dec. 1979; BSKU 28271 (1),

BSKU 28272 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.3167°N, 127.3167°E, 255–260 m, 17 Mar. 1978; BSKU 28199 (1), BSKU 28202 (1), BSKU 28203 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.3000°N, 127.3333°E, 410–415 m, 17 Mar. 1978; BSKU 28219 (1), BSKU 28220 (1), BSKU 28222 (1), BSKU 28223 (1), BSKU 28225 (1), BSKU 28226 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.2667°N, 127.3500°E, 353–365 m, 17 Mar. 1978; SNFR 11292 (3), East China Sea, 29.3850°N, 127.3883°E, 338–339 m, 24 Oct. 1990; BSKU 34743 (1), BSKU 34744 (1), BSKU 34745 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.3833°N, 127.4250°E, 450–475 m, 18 Dec. 1979; SNFR 12507 (2), East China Sea, 29.3900°N, 127.4300°E, 410–416 m, 24 Oct. 1990; BSKU 27212 (1), BSKU 27213 (1), BSKU 27214 (1), BSKU 27216 (1), BSKU 27217 (1), BSKU 27218 (1), BSKU 27224 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.4333°N, 127.5167°E, 570 m, 5 Feb. 1978; BSKU 28311 (1), BSKU 28312 (1), BSKU 28313 (1), BSKU 28314 (1), BSKU 28315 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7167°N, 127.6167°E, 410–423 m, 18 Mar. 1978; BSKU 28633 (1), BSKU 28634 (1), BSKU 28635 (1), BSKU 28636 (1), BSKU 28637 (1), BSKU 28638 (1), BSKU 28639 (1), BSKU 28640 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.8000°N, 127.7000°E, 335–345 m, 21 Mar. 1978; BSKU 28654 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.9000°N, 127.7167°E, 250 m, 21 Mar. 1978; BSKU 27027 (1), BSKU 27029 (1), BSKU 27030 (1), BSKU 27031 (1), BSKU 27033 (1), BSKU 27034 (1), BSKU 27035 (1), BSKU 27036 (1), BSKU 27037 (1), BSKU 27038 (1), BSKU 27039 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.9267°N, 127.7483°E, 360 m, 4 Feb. 1978; BSKU 29843 (1), BSKU 33580 (1), northwest of Amami-oshima Island, Okinawa Trough, 30.0115°N, 127.8387°E, 400–420 m, 26 Oct. 1979; SNFR 14547 (1), East China Sea, 30.8043°N, 127.9207°E, 201–204 m, 18 Sept. 2009; BSKU 33766 (1), BSKU 33767 (1), BSKU 33768 (1), BSKU 33769 (1), BSKU 33770 (1), BSKU 34037 (1), BSKU 34039 (1), BSKU 34040 (1), BSKU 34041 (1), west of Yaku-shima Island, Okinawa Trough, 30.4433°N, 128.0177°E, 410–430 m, 28 Oct. 1979; SNFR 14252 (1), East China Sea, 30.8047°N, 128.0190°E, 380–409 m, 11 Aug. 2008; SNFR 14233 (1), SNFR 14234 (1), SNFR 14235 (2), East China Sea, 30.2230°N, 128.0288°E, 397–400 m, 9 Aug. 2008; SNFR 14248 (1), SNFR 14249 (3), East China Sea, 30.8050°N, 128.2205°E, 440–509 m, 6 Aug. 2008; SNFR 16334 (1), East China Sea, 30.4243°N, 128.2365°E, 500–512 m, 7 Nov. 2003; NSMT-P 67568 (1), west of Yaku-shima Island, East China Sea, 30.4243°N, 128.2365°E, 500–512 m, 6 Nov. 2003; NSMT-P 65654 (2), west of Yaku-shima Island, East China Sea, 30.4467°N, 128.2367°E, 499–500 m, 10 Nov. 2002; NSMT-P 102087 (1), west of Yaku-shima Island, Okinawa Trough,

30.4467°N, 128.2367°E, 499–500 m, 10 Nov. 2002; SNFR 13868 (2), East China Sea, 31.1358°N, 128.4412°E, 433–519 m, 30 Jul. 2008; FFNU P-00714 (3), west of Amakusa Knoll, East China Sea, 32.1593°N, 129.4715°E, 503 m, 12 Mar. 2008; FFNU P-00743 (3), west of Amakusa Knoll, East China Sea, 32.1625°N, 129.4838°E, 487 m, 17 Jul. 2008; BSKU 106775 (1), BSKU 106776 (1), BSKU 106778 (1), BSKU 106791 (1), BSKU 106792 (1), BSKU 106793 (1), BSKU 106795 (1), southeast of Shimokoshiki-jima Island, East China Sea, 31.5560°N, 129.8847°E, 415 m, 24 Apr. 2012; BSKU 106810 (1), BSKU 106812 (1), BSKU 106813 (1), southeast of Shimokoshiki-jima Island, East China Sea, 31.5652°N, 129.8878°E, 407 m, 24 Apr. 2012; BSKU 106772 (1), southeast of Shimokoshiki-jima Island, East China Sea, 31.5658°N, 129.8915°E, 380 m, 24 Apr. 2012; BSKU 58325 (1), Mimase fish market, 8 Dec. 2000; BSKU 39955 (1), Mimase fish market, 31 Mar. 1984; BSKU 36799 (1), Mimase fish market, 11 Mar. 1982; FAKU 34718 (1), Mimase fish market, 26–31 Mar. 1962; BSKU 1818 (1), Mimase fish market, Mar. 1952; BSKU 343 (1), Mimase fish market, 11 Apr. 1951; BSKU 4384 (1), Mimase fish market, 19 Dec. 1954; BSKU 1770 (1), Mimase fish market, 22 Mar. 1952; BSKU 4334 (1), Mimase fish market, 30 Nov. 1954; BSKU 38 (1), BSKU 39 (1), BSKU 40 (1), BSKU 43 (1), Mimase fish market, 4 Mar. 1951; BSKU 40215 (1), BSKU 40216 (1), Tosa Bay, 400 m, 25 Jul. 1984; BSKU 40330 (1), BSKU 40331 (1), BSKU 40332 (1), BSKU 40333 (1), BSKU 40334 (1), Tosa Bay, 400 m, 9 Oct. 1984; BSKU 77983 (1), Tosa Bay, 33.2000°N, 133.6317°E, 370–380 m, 8 Sept. 1998; BSKU 97959 (1), BSKU 97960 (1), Tosa Bay, 400 m, 9 May 2000; BSKU 13303 (1), ca. 10–15 km west of Cape Murotomisaki, Tosa Bay, 360–435 m, 8–9 Jun. 1968; FAKU 20924 (1), Owase fish market, Kumano-nada, 5 Nov. 1953; FAKU 27512 (1), FAKU 27516 (1), FAKU 27520 (1), FAKU 27550 (1), FAKU 27554 (1), FAKU 27555 (1), FAKU 27662 (1), FAKU 27663 (1), FAKU 27664 (1), FAKU 27665 (1), FAKU 27666 (1), FAKU 27667 (1), FAKU 27668 (1), Owase fish market, Kumano-nada, 25 Feb.–10 Mar. 1957; FAKU 23806 (1), Owase fish market, Kumano-nada, 20 Oct. 1954; FAKU 24917 (1), FAKU 24923 (1), Owase fish market, Kumano-nada, 25 Oct. 1954; FAKU 29108 (1), FAKU 29111 (1), FAKU 29113 (1), Owase fish market, Kumano-nada, May 1960; FRLM 34140 (1), FRLM 34332 (1), off Owase, Kumano-nada, 30 Jan. 2008; FAKU 26942 (1), FAKU 26943 (1), Miya fish market, Enshu-nada, 29–31 Dec. 1956; NSMT-P 101930 (1), southwest of Cape Omaezaki, Enshu-nada, 34.3816°N, 138.0010°E, 475–488 m, 10 Dec. 1995; BSKU 110011 (1), BSKU 110013 (1), Suruga Bay, 34.7080°N, 138.4587°E, 200–450 m, 23 Apr. 2013; BSKU 110429 (1), Suruga Bay, 34.7619°N, 138.4811°E, 196–387 m, 23 Apr. 2013; MSM uncat. (1), Suruga Bay, 21 Feb. 2012; BSKU 110425 (1), Suruga Bay, 34.7651°N, 138.4880°E, 200–450 m, 23 Apr. 2013; NSMT-P 101996

(4), west of Cape Hagachizaki, Suruga Bay, 34.6817°N, 138.6817°E, 418–457 m, 11 Nov. 1983; FAKU M7409 (1), Heta fish market, Suruga Bay, 150 ftn (275 m), 23 Jul. 1938.

Coelorinchus cingulatus. 2 specimens: FMNH 120864 (1), Nan-fang-ao fish market, Yilan, 7 Sept. 2009; BSKU uncat. (1), Mimase fish market, date unknown.

Coelorinchus formosanus. 10 specimens: BSKU 26581 (1), north of Miyako-jima Island, Okinawa Trough, 26.5200°N, 125.0417°E, 255–260 m, 23 Jan. 1978; BSKU 34669 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.9983°N, 127.1550°E, 325 m, 17 Dec. 1979; BSKU 99306 (1), Tosa Bay, 200 m, 11 Sept. 2007; FMNH 120861 (5 of 6), Dong-gang fish market, Pingtung, 12 Sept. 2009; KAUM-I. 44656 (1), off Kaohsiung, Yilan, 400 m, 1 Jul. 2011; MUFS 39535 (1), Dong-gang, Pingtung, 27 Feb. 1973.

Coelorinchus gilberti. 69 specimens: NSMT-P 93747 (1), Tosa Bay, 750 m, 25 Jun. 2000; BSKU 69282 (1), Tosa Bay, 33.1417°N, 133.6433°E, 582–597 m, 16 Jun. 1999; BSKU 98935 (1), Tosa Bay, 33.0452°N, 133.6448°E, 733–740 m, 25 Jan. 1989; SNFR 18053 (1), off Kochi, 21 Jun. 1988; BSKU 45009 (1), Tosa Bay, 500 m, 23 Aug. 1988; NSMT-P 93136 (1), off Oh-sima Island, Izu Islands, date unknown; FAKU 35863 (1), Sagami Bay, Apr. 1962; FMNH 57268 (1, ex. CM6207a), Misaki, Sagami Bay, date unknown; NSMT-P 101994 (1), southeast of Miyake-jima Island, Shichito-Iojima Ridge, 34.1287°N, 139.6413°E, 484–565 m, 19 May 1989; NSMT-P 97213 (2), east of Boso Peninsula, 35.1050°N, 140.8367°E, 627–673 m, 25 Apr. 1995; ZMUT 30601 (1), ZMUT 30606 (1), Choshi fish market, 10 Dec. 1934; HUMZ 94369 (1), Ishinomaki fish market, Miyagi, 18 Dec. 1981; BSKU 97949 (1), BSKU 97950 (1), BSKU 97951 (1), BSKU 97952 (1), off Onahama, 450 m, 16 Oct. 1997; BSKU 101979 (1), BSKU 101980 (1), BSKU 101981 (1), BSKU 101983 (1), BSKU 101984 (1), BSKU 101985 (1), BSKU 101986 (1), BSKU 101987 (1), off Iwaki, 36.9162°N, 141.5995°E, 615–633 m, 4 Nov. 1995; NSMT-P 102628 (1), NSMT-P 102629 (1), off Iwaki, 36.9690°N, 141.6327°E, 554–558 m, 28 Oct. 2006; NSMT-P 47403 (1), off Sendai, 600 m, 2 Nov. 1995; HUMZ 72718 (1), off Fukushima, 37.2233°N, 141.9333°E, 850–930 m, 19 Jan. 1978; NSMT-P 102792 (1), off Minami-soma, 37.6430°N, 142.0185°E, 503–511 m, 31 Oct. 2006; NSMT-P 101998 (1), off Minami-soma, 37.5367°N, 142.0550°E, 538–561 m, 1 Dec. 1981; NSMT-P 49054 (1), southeast of Oshika Peninsula, 37.9367°N, 142.0683°E, 596–626 m, 2 Nov. 1995; BSKU 99134 (1), southeast of Kinkazan, 37.9368°N, 142.0683°E, 596–626 m, 2 Nov. 1995; HUMZ 77534 (1), HUMZ 77535 (1), 10 km off Higashidori, Shimokita Peninsula, 520 m, 27 Sept. 1978; HUMZ 77536 (1), HUMZ 77537 (1), 15 km off Higashidori, Shimokita Peninsula, 640 m, 26 Sept. 1978; HUMZ 72687 (1), off Fukushima, 38.8900°N, 141.9333°E, 850–930 m, 19 Jan. 1978; HUMZ 112989 (1), off Hachinohe, 40.7450°N, 142.0583°E, 525 m, 8

Mar. 1988; FRLM 37475 (1), off Miyagi, 38.4062°N, 142.0633°E, 448 m, 27 Oct. 2010; BSKU 116164 (1), off Hachinohe, 40.4818°N, 142.2587°E, 557–595 m, 6 Nov. 1995; NSMT-P 102115 (1), off Kamaishi, 39.2133°N, 142.2667°E, 572–573 m, 14 Aug. 1985; BSKU 58334 (1), BSKU 58335 (1), BSKU 58336 (1), BSKU 58337 (1), off Kamaishi, 450 m, 17 Oct. 1999; HUMZ 124729 (1), HUMZ 6933 (1), Miyako, Iwate, Mar. 1935; HUMZ 18062 (1), Sirikisinai, Hokkaido, 5 Feb. 1955; FAKU M9315 (1), off Muroran, Mar. 1938; HUMZ 87121 (1), HUMZ 87122 (1), HUMZ 87123 (1), HUMZ 87124 (1), HUMZ 87125 (1), off Muroran, 3 Mar. 1980; HUMZ 6934 (1), Urakawa, Hokkaido, date unknown; BSKU 114866 (1), BSKU 114867 (1), BSKU 114868 (1), BSKU 114869 (1), BSKU 114870 (1), BSKU 114871 (1), BSKU 114873 (1), BSKU 114914 (1), east-southeast of Cape Erimomisaki, 41.8170°N, 143.7875°E, 576 m, 6 Jun. 2014; FAKU 17006 (1), off Kushiro, Jun. 1951; HUMZ 124728 (1), off Kushiro, 400 m, 20 May 1960; HUMZ 34043 (1), off Kushiro, 600 m, 17 Jul. 1974; BSKU 26102 (paratype of *Coelorhynchus longicephalus*), Kyushu-Palau Ridge, 27.9167°N, 134.6500°E, 700 m, 20 Feb. 1978.

***Coelorinchus hige*.** 19 specimens: BSKU 106835 (1), BSKU 106836 (1), BSKU 106837 (1), BSKU 106838 (1), BSKU 106839 (1), southeast of Shimokoshiki-jima Island, East China Sea, 31.5675°N, 129.8930°E, 393 m, 24 Apr. 2012; BSKU 13305 (1), BSKU 13306 (1), BSKU 13307 (1), BSKU 13308 (1), BSKU 13309 (1), BSKU 13310 (1), ca. 10–15 km west of Cape Murotomisaki, Tosa Bay, 360–435 m, 8–9 Jun. 1968; FAKU 24921 (1), Owase fish market, Kumano-nada, 25 Oct. 1954; FRLM 34770 (1), FRLM 34771 (1), off Owase, Kumano-nada, 24 Oct. 2008; NSMT-P 120920 (1), southwest of Cape Daiozaki, Kumano-nada, 34.0983°N, 136.6800°E, 422–425 m, 27 May 1986; NSMT-P 48607 (1), NSMT-P 120919 (1), off Kumomizaki, Suruga Bay, 34.7717°N, 138.7133°E, 275–500 m, 23 Oct. 1994; MSM 13-14 (2), southwest of Toi, Suruga Bay, 34.8791°N, 138.7159°E, 392–510 m, 21 Feb. 2012.

***Coelorinchus hubbsi*.** 174 specimens: BSKU 13236 (1), BSKU 13237 (1), BSKU 13238 (1), ca. 15 km east of Cape Ashizurimisaki, Tosa Bay, 420–555 m, 2–6 Jun. 1968; BSKU 41689 (1), BSKU 41690 (1), off Okitsu, Tosa Bay, 250–260 m, 18 Apr. 1985; BSKU 97964 (1), BSKU 97965 (1), BSKU 97966 (1), BSKU 97971 (1), BSKU 97972 (1), BSKU 97973 (1), BSKU 97976 (1), BSKU 97977 (1), Tosa Bay, 33.1412°N, 133.5388°E, 295–336 m, 18 Jul. 1988; KAUM-I. 43937 (1), Mimase fish market, 350 m, 12 Dec. 2011; BSKU 29955 (1), Mimase fish market, 1 Mar. 1980; BSKU 38605 (1), Mimase fish market, 1 Mar. 1983; BSKU 85969 (1), Mimase fish market, 10 Dec. 1998; BSKU 58481 (1), Mimase fish market, 10 Dec. 2001; BSKU 77501 (1), Mimase fish market, 11 Dec. 2005; BSKU 52827 (1), Mimase fish market, 11 Oct. 2000; BSKU 58137 (1), Mimase fish market, 12 Dec. 2001; BSKU 74816 (1),

Mimase fish market, 12 Dec. 2004; BSKU 76125 (1), Mimase fish market, 12 Oct. 2004; BSKU 72287 (1), Mimase fish market, 13 Apr. 2004; BSKU 82841 (1), Mimase fish market, 14 Dec. 1996; BSKU 53057 (1), Mimase fish market, 14 Dec. 2000; BSKU 70086 (1), Mimase fish market, 15 Jan. 2004; BSKU 89688 (1), Mimase fish market, 15 Jan. 2007; BSKU 69150 (1), Mimase fish market, 15 Mar. 2004; BSKU 53200 (1), Mimase fish market, 16 Nov. 2000; BSKU 54616 (1), BSKU 54617 (1), Mimase fish market, 19 Dec. 1998; BSKU 77823 (1), Mimase fish market, 19 Dec. 2004; BSKU 69237 (1), BSKU 69238 (1), BSKU 69239 (1), Mimase fish market, 19 Nov. 2003; BSKU 72781 (1), BSKU 72792 (1), BSKU 72794 (1), Mimase fish market, 20 Feb. 2004; BSKU 36880 (1), Mimase fish market, 20 Mar. 1982; BSKU 69579 (1), Mimase fish market, 21 Dec. 2003; BSKU 69785 (1), Mimase fish market, 22 Dec. 2003; BSKU 53762 (1), Mimase fish market, 22 Mar. 2001; BSKU 99252 (1), Mimase fish market, 23 Apr. 2007; BSKU 71540 (1), Mimase fish market, 25 Mar. 2004; BSKU 29990 (1), Mimase fish market, 26 Jan. 1980; BSKU 82935 (1), Mimase fish market, 26 Nov. 1996; BSKU 66799 (1), Mimase fish market, 27 Feb. 2002; BSKU 78386 (1), Mimase fish market, 27 Feb. 2004; BSKU 76104 (1), Mimase fish market, 27 Feb. 2005; BSKU 35559 (1), Mimase fish market, 27 Mar. 1980; BSKU 38323 (1), Mimase fish market, 3 Dec. 1982; BSKU 35465 (1), Mimase fish market, 3 Mar. 1981; BSKU 65043 (1), BSKU 65044 (1), Mimase fish market, 30 Jan. 2003; BSKU 52718 (1), Mimase fish market, 30 Nov. 2000; BSKU 91968 (1), Mimase fish market, 31 Oct. 2007; BSKU 89931 (1), Mimase fish market, 4 Dec. 2006; BSKU 90970 (1), Mimase fish market, 6 Apr. 2007; BSKU 76148 (1), BSKU 91634 (1), Mimase fish market, 6 Jan. 2005; BSKU 64309 (1), Mimase fish market, 7 Feb. 2003; BSKU 76117 (1), Mimase fish market, 7 Feb. 2005; MUFS 5756 (1), Mimase fish market, 7 Jan. 1976; BSKU 30179 (1), BSKU 30180 (1), Mimase fish market, 8 Dec. 1979; BSKU 69978 (1), Mimase fish market, 8 Feb. 2002; BSKU 56379 (1), Mimase fish market, 8 Nov. 2001; BSKU 77644 (1), Mimase fish market, 8 Nov. 2005; BSKU 63386 (1), Mimase fish market, 9 Jan. 2003; BSKU 75661 (1), Mimase fish market, 9 Nov. 2004; BSKU 77504 (1), BSKU 77505 (1), Mimase fish market, 9 Nov. 2005; BSKU 99304 (1), Mimase fish market, 11 Apr. 2007; BSKU 99299 (1), BSKU 99301 (1), BSKU 99303 (1), Mimase fish market, 8 Apr. 2007; BSKU 2436 (1), Mimase fish market, 1952; BSKU 10036 (1), BSKU 10037 (1), BSKU 10038 (1), Mimase fish market, 10 Feb. 1961; BSKU 75 (1), BSKU 76 (1), Mimase fish market, 11 Mar. 1951; BSKU 24 (1), Mimase fish market, 19 Feb. 1951; BSKU 9682 (1), Mimase fish market, 20 Feb. 1949; BSKU 1772 (1), Mimase fish market, 22 Mar. 1952; BSKU 210 (1), Mimase fish market, 3 Dec. 1950; BSKU 169 (1), Mimase fish market, 3 Feb. 1951; BSKU 9901 (1), Mimase fish market, 1955; BSKU 4412 (1), Mimase

fish market, 26 Dec. 1954; BSKU 4571 (1), Mimase fish market, 30 Apr. 1955; BSKU 94809 (1), Mimase fish market, 28 Nov. 2007; BSKU 95084 (1), Mimase fish market, 30 Nov. 2007; BSKU 92270 (1), BSKU 99374 (1), BSKU 99375 (1), Mimase fish market, 9 Nov. 2007; BSKU 92365 (1), Mimase fish market, 16 Nov. 2007; BSKU 13913 (1), BSKU 13914 (1), Mimase fish market, Jan. 1968; BSKU 13878 (1), Mimase fish market, Mar. 1968; KAUM-I. 12908 (1), KAUM-I. 12917 (1), Mimase fish market, 9 May 2008; BSKU 42764 (1), off Okitsu, Tosa Bay, 240–250 m, 13 Mar. 1986; BSKU 41722 (1), BSKU 41723 (1), BSKU 41724 (1), off Okitsu, Tosa Bay, 250 m, 24 Apr. 1985; BSKU 74622 (1), Tosa Bay, 300 m, 11 Jun. 1997; BSKU 97979 (1), BSKU 97980 (1), BSKU 97982 (1), BSKU 97983 (1), BSKU 97984 (1), BSKU 97986 (1), BSKU 97987 (1), BSKU 97989 (1), BSKU 97990 (1), Tosa Bay, 300 m, 16 Jul. 1997; BSKU 42506 (1), BSKU 42507 (1), BSKU 42508 (1), Tosa Bay, 14 Jan. 1986; BSKU 69311 (1), BSKU 69312 (1), Tosa Bay, 33.1800°N, 133.6217°E, 412–423 m, 8 Sept. 1998; BSKU 77893 (1), Tosa Bay, 33.2083°N, 133.6433°E, 364–382 m, 27 Sept. 1999; BSKU 13295 (1), BSKU 13296 (1), ca. 10–15 km west of Cape Murotomisaki, Tosa Bay, 360–435 m, 8–9 Jun. 1968; BSKU 42518 (1), off Kannoura, 240 m, 16 Jan. 1986; FAKU 27560 (1), FAKU 27561 (1), FAKU 27562 (1), FAKU 27563 (1), FAKU 27565 (1), FAKU 27566 (1), FAKU 27567 (1), FAKU 27568 (1), FAKU 27569 (1), FAKU 27570 (1), FAKU 27571 (1), FAKU 27572 (1), FAKU 27573 (1), Owase fish market, Kumano-nada, 25 Feb.–10 Mar. 1957; BSKU 99454 (1), BSKU 99455 (1), BSKU 99456 (1), BSKU 99457 (1), Owase fish market, Kumano-nada, 13 Mar. 1983; FRLM 8636 (1), Owase fish market, 16 Mar. 1967; FRLM 33596 (1), off Owase, Kumano-nada, 240 m, 6 Nov. 2007; FRLM 34981 (1), off Owase, Kumano-nada, 24 Oct. 2008; FRLM 33280 (1), off Owase, Kumano-nada, 17 May 2007; FRLM 34297 (1), FRLM 34460 (1), off Owase, Kumano-nada, 18 Feb. 2008; FRLM 34335 (1), off Owase, Kumano-nada, 30 Jan. 2008; FRLM 33911 (1), FRLM 33912 (1), FRLM 33913 (1), FRLM 33914 (1), FRLM 33915 (1), FRLM 33916 (1), off Owase, Kumano-nada, 16 Nov. 2007; FRLM 30764 (1), FRLM 30765 (1), off Owase, Kumano-nada, 2 Feb. 2004; FRLM 34358 (1), off Owase, Kumano-nada, date unknown; FRLM 8586 (1), Kumano-nada, date unknown; KPM-NI 24981 (1), eastern part of Suruga Bay, Nov. 2009; NSMT-P 101995 (1), southeast of Miyake-jima Island, Shichito-Iojima Ridge, 34.1132°N, 139.5845°E, 297–303 m, 19 May 1989; NSMT-P 99675 (1), off Choshi, Oct. 2009–11 May 1979; ZMUT 30612 (1), ZMUT 30613 (1), ZMUT 30614 (1), Choshi, 10 Dec. 1934; ZMUT 30611 (1), Choshi, 27 Jan. 1935; NSMT-P 65730 (1), off Hitachi, 36.5118°N, 141.0147°E, 309–321 m, 26 Oct. 2002; BSKU 19804 (1), BSKU 19805 (1), east of Hachijo Island, Shichito-Iojima Ridge, 33.1267°N, 139.9750°E, 430 m, 5 Jul. 1967; BSKU 19312 (1), BSKU 19313 (1), BSKU 19314 (1), BSKU 19315 (1), BSKU

19316 (1), east of Hachijo Island, Shichito-Iojima Ridge, 33.1367°N, 139.9900°E, 440 m, 8 Dec. 1962.

Coelorrinchus japonicus. 263 specimens: BSKU 28693 (1), BSKU 28694 (1), off Senkaku Islands, Okinawa Trough, 25.6283°N, 122.8933°E, 560–692 m, 14 Jan. 1978; BSKU 29636 (1), northwest of Miyako-jima Island, Okinawa Trough, 25.7900°N, 124.3333°E, 600 m, 20 Sept. 1979; BSKU 32580 (1), BSKU 32582 (1), northwest of Miyako-jima Island, Okinawa Trough, 25.7900°N, 124.3900°E, 600 m, 20 Sept. 1979; BSKU 26497 (1), northwest of Miyako-jima Island, Okinawa Trough, 25.2850°N, 124.7800°E, 910–990 m, 22 Jan. 1978; BSKU 27870 (1), BSKU 27871 (1), BSKU 27872 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.5833°N, 127.0000°E, 600 m, 16 Mar. 1978; BSKU 27676 (1), west of Okinawa-jima Island, Okinawa Trough, 28.7333°N, 127.0167°E, 610–640 m, 11 Mar. 1978; SNFR 16706 (1), East China Sea, 28.6768°N, 127.0703°E, 599–601 m, 28 Jun. 2000; BSKU 27936 (1), BSKU 27937 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.7500°N, 127.1167°E, 520–542 m, 16 Mar. 1978; BSKU 27583 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.7167°N, 127.2333°E, 750–755 m, 11 Mar. 1978; BSKU 27986 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.8333°N, 127.2333°E, 700–740 m, 16 Mar. 1978; BSKU 27110 (1), BSKU 27112 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.3833°N, 127.5150°E, 650 m, 5 Feb. 1978; BSKU 27219 (1), BSKU 27220 (1), BSKU 27221 (1), BSKU 27222 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.4333°N, 127.5167°E, 570 m, 5 Feb. 1978; BSKU 28367 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7333°N, 127.8000°E, 495–510 m, 19 Mar. 1978; BSKU 28439 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7667°N, 127.9833°E, 713–770 m, 20 Mar. 1978; BSKU 28587 (1), BSKU 28588 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.9500°N, 128.0000°E, 585–610 m, 20 Mar. 1978; BSKU 29886 (1), west of Yaku-shima Island, Okinawa Trough, 30.2083°N, 128.2088°E, 600–620 m, 26 Oct. 1979; NSMT-P 65669 (2), NSMT-P 65672 (1), south of Danjo Islands, East China Sea, 31.4217°N, 128.4017°E, 461–500 m, 8 Nov. 2002; SNFR 13867 (1), East China Sea, 31.1358°N, 128.4412°E, 433–519 m, 30 Jul. 2008; FFNU P-00671 (11), northwest of Daiichi-Danjo Knoll, East China Sea, 31.6917°N, 128.5828°E, 506 m, 9 May 2007; BSKU 109018 (1), west of Okikawayama Bank, 32.1503°N, 129.0072°E, 499–504 m, 19 Nov. 2012; FFNU P-00771 (1), east of Okihirazi Bank, East China Sea, 32.1635°N, 129.4663°E, 494 m, 3 Dec. 2008; FFNU P-00713 (2), west of Amakusa Knoll, East China Sea, 32.1593°N, 129.4715°E, 503 m, 12 Mar. 2008; FFNU P-00780 (2), southeast of Okihirazi Bank, East China Sea, 32.1020°N, 129.4767°E, 583 m, 7 Mar. 2009; BSKU 45521 (1), off Tosa-Shimizu, 31 Aug. 1988; BSKU 43972 (1), off Kamikawaguchi, Tosa Bay, 26 Aug. 1987;

BSKU 23210 (1), Tosa Bay, 32.9233°N, 133.4850°E, 489 m, 30 Jan. 1975; BSKU 29504 (1), Tosa Bay, 32.9167°N, 133.5500°E, 700 m, 21 Dec. 1979; BSKU 41028 (1), Mimase fish market, 13 Mar. 1985; BSKU 11716 (1), BSKU 11717 (1), BSKU 11718 (1), BSKU 11719 (1), Mimase fish market, Nov. 1965; FAKU 21770 (1), FAKU 21776 (1), Mimase fish market, 10–20 Apr. 1954; BSKU 42 (1), Mimase fish market, 22–27 Feb. 1951; BSKU 1820 (1), Mimase fish market, Mar. 1952; BSKU 1771 (1), Mimase fish market, 22 Mar. 1952; BSKU 5054 (1), Mimase fish market, 27 Dec. 1955; BSKU 6926 (1), Mimase fish market, 27 Oct. 1956; BSKU 4363 (1), Mimase fish market, 5 Dec. 1954; BSKU 12924 (1), Tosa Bay, 350–450 m, 14 May 1968; BSKU 65949 (1), BSKU 69818 (1), Tosa Bay, 400 m, 9 May 2000; BSKU 69307 (1), BSKU 69309 (1), BSKU 69310 (1), Tosa Bay, 33.1800°N, 133.6217°E, 412–423 m, 8 Sept. 1998; BSKU 71582 (1), BSKU 71584 (1), BSKU 71589 (1), Tosa Bay, 400 m, 9 May 2000; BSKU 70894 (1), BSKU 70895 (1), BSKU 70896 (1), Tosa Bay, 33.2250°N, 133.6933°E, 440–460 m, 24 Aug. 2000; BSKU 29937 (1), BSKU 32425 (1), BSKU 32427 (1), BSKU 32428 (1), BSKU 32429 (1), off Nahari, Tosa Bay, 33.2792°N, 133.8947°E, 450–470 m, 17 Nov. 1979; FAKU 27622 (1), FAKU 27648 (1), FAKU 27652 (1), FAKU 27653 (1), FAKU 27658 (1), FAKU 27702 (1), FAKU 27703 (1), FAKU 27704 (1), FAKU 27705 (1), FAKU 27707 (1), FAKU 27710 (1), FAKU 27711 (1), FAKU 27712 (1), FAKU 27714 (1), FAKU 27717 (1), FAKU 27719 (1), FAKU 27721 (1), Owase fish market, Kumano-nada, 25 Feb.–10 Mar. 1957; FAKU 21515 (1), Owase fish market, Kumano-nada, 15 Feb. 1954; FRLM 8667 (1), Owase fish market, 17 Dec. 1958; FRLM 34333 (1), FRLM 34334 (1), off Owase, Kumano-nada, 30 Jan. 2008; FRLM 34359 (1), off Owase, Kumano-nada, date unknown; FRLM 3567 (1), FRLM 3568 (1), FRLM 3569 (1), FRLM 3570 (1), FRLM 3571 (1), FRLM 3573 (1), FRLM 3574 (1), FRLM 3575 (1), FRLM 3576 (1), off Mikizaki, Kumano-nada, 18 May 1982; HUMZ 97114 (1), off Kii-Nagashima, Mie, 34.0527°N, 136.5328°E, 500 m, 6 Nov. 1982; FRLM 24168 (1), off Mie, Kumano-nada, 34.1090°N, 136.5808°E, 13 Oct. 1998; FRLM 2528 (1), FRLM 2529 (1), FRLM 2530 (1), FRLM 2532 (1), FRLM 2533 (1), FRLM 2534 (1), FRLM 2536 (1), FRLM 2537 (1), FRLM 2538 (1), FRLM 2539 (1), off Mie, Kumano-nada, 16 Apr. 1980; FRLM 3581 (1), off Mie, Kumano-nada, 18 May 1982; NSMT-P 101350 (1), southwest of Cape Daiozaki, Kumano-nada, 34.0836°N, 136.6554°E, 495–515 m, 12 Dec. 1995; NSMT-P 101997 (1), southwest of Cape Daiozaki, Kumano-nada, 34.0983°N, 136.6800°E, 422–425 m, 27 May 1986; FRLM 27656 (1), FRLM 27658 (1), FRLM 27660 (1), FRLM 27911 (1), FRLM 27912 (1), FRLM 27913 (1), FRLM 27914 (1), FRLM 27915 (1), FRLM 27916 (1), FRLM 27917 (1), FRLM 27918 (1), FRLM 27919 (1), FRLM 27920 (1), FRLM 27921 (1), FRLM 27922 (1), FRLM 27923 (1), FRLM 27924 (1), off Mie, Kumano-nada, 34.0848°N, 136.7193°E, 4 Jul. 2001; FRLM 3737 (1), FRLM 3738 (1), FRLM 3739 (1), FRLM 3740 (1), FRLM 3741 (1), FRLM 3742 (1), FRLM 3744 (1), FRLM 3745 (1), FRLM 3746 (1), FRLM 3747 (1), FRLM 3748 (1), FRLM 3749 (1), FRLM 3750 (1), FRLM 3751 (1), FRLM 3752 (1), FRLM 3753 (1), FRLM 3754 (1), FRLM 3755 (1), FRLM 3756 (1), FRLM 3757 (1), FRLM 3758 (1), FRLM 3759 (1), FRLM 3760 (1), FRLM 3761 (1), FRLM 3762 (1), FRLM 3763 (1), FRLM 3765 (1), FRLM 3766 (1), FRLM 3767 (1), FRLM 3768 (1), FRLM 3769 (1), FRLM 3770 (1), FRLM 3771 (1), FRLM 3778 (1), off Goza, Kumano-nada, 13 Oct. 1982; FRLM 3843 (1), FRLM 3844 (1), FRLM 3845 (1), FRLM 3846 (1), FRLM 3847 (1), FRLM 3848 (1), FRLM 3849 (1), FRLM 3850 (1), FRLM 3851 (1), FRLM 3852 (1), FRLM 3853 (1), FRLM 3854 (1), FRLM 3855 (1), FRLM 3856 (1), off Nakiri, Kumano-nada, 19 Nov. 1982; FRLM 3456 (1), FRLM 3457 (1), FRLM 3458 (1), FRLM 3459 (1), FRLM 3460 (1), FRLM 3461 (1), FRLM 3462 (1), off Shima Peninsula, Kumano-nada, 16 Feb. 1982; FRLM 3564 (1), FRLM 3565 (1), off Shima Peninsula, Kumano-nada, 29 Mar. 1982; FRLM 3343 (1), off Cape Daiozaki, Kumano-nada, 17 Dec. 1981; FRLM 3857 (1), off Cape Daiozaki, Kumano-nada, 19 Nov. 1982; FAKU 26944 (1), Miya fish market, Enshu-nada, 29–31 Dec. 1956; FAKU 28389 (1), FAKU 28390 (1), FAKU 28391 (1), Miya fish market, Enshu-nada, 5–7 Feb. 1959; ZMUT 33682 (1), Maisaka, Shizuoka, date unknown; NSMT-P 101359 (1), southeast of Oigawa River, Suruga Bay, 34.6790°N, 138.3580°E, 375–457 m, 26 Feb. 1986; BSKU 110078 (1), BSKU 110080 (1), Suruga Bay, 34.7481°N, 138.4652°E, 243–424 m, 23 Apr. 2013; KPM-NI 18043 (1), off Fuji River, Suruga Bay, 300 m, Nov. 2006; NSMT-P 101999 (1), Suruga Bay, 1 May 1973; NSMT-P 102356 (7), NSMT-P 114950 (1), west of Cape Hagachizaki, Suruga Bay, 34.6817°N, 138.6817°E, 418–457 m, 11 Nov. 1983; NSMT-P 102114 (1), west of Matsuzaki, Suruga Bay, 34.7717°N, 138.7000°E, 370–372 m, 3 Jul. 1980; NSMT-P 96041 (1), southwest of Heta, Suruga Bay, 34.9550°N, 138.7300°E, 340–740 m, 6 Oct. 1993; NSMT-P 7806 (1), off Heta, Suruga Bay, 18 Feb. 1969; NSMT-P 103025 (1), off Heta, Suruga Bay, 11 Dec. 2001; NSMT-P 93071 (1), off Heta, Suruga Bay, 15 Mar. 2000; NSMT-P 48540 (1), northwest of Heta, Suruga Bay, 34.9933°N, 138.7450°E, 310–680 m, 25 Oct. 1993; FAKU M6856 (1), FAKU M6860 (1), Heta fish market, Suruga Bay, 150 ftm (275 m), 23 Jul. 1938; FAKU 22841 (1), Heta fish market, Suruga Bay, 200–250 ftm (366–458 m), 24–25 Jun. 1939; FAKU M6974 (1), Heta fish market, Suruga Bay, 200–290 ftm (366–531 m), 24–25 Jun. 1939; KPM-NI 8635 (1), off Cape Irozaki, Izu Peninsula, 150–220 m, 22 Apr. 2001; NSMT-P 66214 (2), off Odawara, Sagami Bay, 35.2513°N, 139.2057°E, 223–229 m, 8 Nov. 2002; FAKU 34262 (1), off Oiso, Sagami Bay, Apr. 1969; NSMT-P 66216 (2), off Hiratsuka, Sagami Bay, 35.2427°N, 139.3588°E, 400–409 m, 13 Dec. 2002; FMNH 89118

(1), Sagami Bay, 22 Jul. 1904; FMNH 91771 (1), Sagami Bay, 9 Apr. 1903; FMNH 91706 (5), Misaki, Sagami Bay, 23 Dec. 1901; FMNH uncat. (1, ex. CM unknown), Misaki, Sagami Bay, 1922; NSMT-P 79318 (2), off Hitachi, 36.5350°N, 141.0683°E, 401–404 m, 18 Apr. 1997; BSKU 97933 (1), BSKU 97934 (1), BSKU 97935 (1), BSKU 97936 (1), BSKU 97937 (1), BSKU 97938 (1), BSKU 97939 (1), BSKU 97940 (1), BSKU 97941 (1), BSKU 97942 (1), BSKU 97943 (1), BSKU 97944 (1), BSKU 97945 (1), BSKU 97946 (1), BSKU 97947 (1), BSKU 97948 (1), off Mito, 350 m, 8 Oct. 1997.

Coelorinchus jordani. 258 specimens: BSKU 27501 (1), BSKU 27502 (1), BSKU 27503 (1), west of Okinawa-jima Island, Okinawa Trough, 28.1000°N, 126.3833°E, 143 m, 9 Mar. 1978; BSKU 34395 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.5583°N, 126.5233°E, 320–325 m, 16 Dec. 1979; SNFR 14069 (1), SNFR 14070 (4), East China Sea, 28.4578°N, 126.8727°E, 234–240 m, 6 Jun. 2008; SNFR 17190 (4), East China Sea, 28.3183°N, 126.8967°E, 314–315 m, 23 Oct. 1990; SNFR 11080 (1), East China Sea, 28.3750°N, 126.9683°E, 378–382 m, 23 Oct. 1990; BSKU 27726 (1), north-northwest of Okinawa Island, Okinawa Trough, 28.5000°N, 127.0000°E, 225–235 m, 12 Mar. 1978; SNFR 15388 (4), East China Sea, 28.8067°N, 127.0167°E, 343–355 m, 11 Jul. 1988; BSKU 28182 (1), BSKU 28183 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.1667°N, 127.0167°E, 160 m, 17 Mar. 1978; SNFR 11199 (2), East China Sea, 28.5233°N, 127.0250°E, 550 m, 23 Oct. 1990; BSKU 27739 (1), BSKU 27740 (1), west of Amami-oshima Island, Okinawa Trough, 28.4333°N, 127.1833°E, 300–355 m, 12 Mar. 1978; BSKU 28267 (1), BSKU 28268 (1), BSKU 28269 (1), BSKU 28270 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.3167°N, 127.3167°E, 255–260 m, 17 Mar. 1978; BSKU 33630 (1), BSKU 33632 (1), BSKU 33633 (1), BSKU 33634 (1), northwest of Amami-oshima Island, Okinawa Trough, 30.0773°N, 127.6788°E, 220–245 m, 27 Oct. 1979; BSKU 28659 (1), BSKU 28660 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.9000°N, 127.7167°E, 250 m, 21 Mar. 1978; BSKU 29834 (1), BSKU 33829 (1), northwest of Amami-oshima Island, Okinawa Trough, 30.4815°N, 127.8432°E, 320–360 m, 28 Oct. 1979; SNFR 14800 (1), SNFR 14801 (1), East China Sea, 30.2953°N, 127.8872°E, 329–345 m, 13 Sept. 2009; SNFR 16343 (2), East China Sea, 30.4833°N, 127.9482°E, 399 m, 7 Nov. 2003; BSKU 109025 (1), BSKU 109026 (1), BSKU 109028 (1), east of Kasayama Bank, 32.3884°N, 129.0505°E, 304–312 m, 19 Nov. 2012; BSKU 109055 (1), BSKU 109056 (1), west of Sanposone, 32.2872°N, 129.0732°E, 351–389 m, 19 Nov. 2012; FFNU P-00820 (4), southeast of Fukue-jima Island, East China Sea, 32.5077°N, 129.1522°E, 231 m, 21 Oct. 2009; FFNU P-00835 (10), southeast of Fukue-jima Island, East China Sea, 32.5007°N, 129.1543°E, 236 m, 23 Nov. 2009; FFNU P-00687 (10), west of Hirazi Bank, East China

Sea, 32.2398°N, 129.3057°E, 298 m, 2 Aug. 2007; KAUM-I. 6638 (1), off Nomaie, Kagoshima, 300 m, 25 Sept. 2007; KAUM-I. 6643 (1), off Nomaie, Kagoshima, 300 m, 27 Sept. 2007; KAUM-I. 6640 (1), off Nomaie, Kagoshima, 300 m, 30 Sept. 2007; KAUM-I. 32247 (1), off Nomaie, Kagoshima, 31.5000°N, 129.8833°E, 370–400 m, 10 Sept. 2010; BSKU 106789 (1), BSKU 106790 (1), southeast of Shimokoshiki-jima Island, East China Sea, 31.5560°N, 129.8847°E, 415 m, 24 Apr. 2012; BSKU 106816 (1), BSKU 106817 (1), BSKU 106818 (1), BSKU 106819 (1), southeast of Shimokoshiki-jima Island, East China Sea, 31.5652°N, 129.8878°E, 407 m, 24 Apr. 2012; BSKU 106769 (1), southeast of Shimokoshiki-jima Island, East China Sea, 31.5658°N, 129.8915°E, 380 m, 24 Apr. 2012; KAUM-I. 36195 (1), KAUM-I. 36196 (1), Kagoshima Bay, 18 Dec. 2010; KAUM-I. 43715 (1), KAUM-I. 43716 (1), KAUM-I. 43717 (1), KAUM-I. 43718 (1), KAUM-I. 43719 (1), KAUM-I. 43720 (1), KAUM-I. 43721 (1), KAUM-I. 43722 (1), KAUM-I. 43723 (1), KAUM-I. 43724 (1), KAUM-I. 43725 (1), KAUM-I. 43726 (1), KAUM-I. 43727 (1), KAUM-I. 43728 (1), KAUM-I. 43729 (1), KAUM-I. 43730 (1), KAUM-I. 43731 (1), KAUM-I. 43732 (1), KAUM-I. 43733 (1), KAUM-I. 43734 (1), KAUM-I. 43735 (1), KAUM-I. 43737 (1), KAUM-I. 43738 (1), KAUM-I. 43739 (1), KAUM-I. 43741 (1), KAUM-I. 43742 (1), off Sakura-jima Island, Kagoshima Bay, 31.6167°N, 130.6167°E, 200 m, 4 Sept. 1972; KAUM-I. 5174 (1), KAUM-I. 5180 (1), KAUM-I. 6009 (1), KAUM-I. 6010 (1), KAUM-I. 6011 (1), KAUM-I. 6012 (1), KAUM-I. 6013 (1), KAUM-I. 6014 (1), KAUM-I. 6015 (1), KAUM-I. 6016 (1), KAUM-I. 6017 (1), KAUM-I. 6018 (1), KAUM-I. 6019 (1), KAUM-I. 6020 (1), KAUM-I. 6021 (1), KAUM-I. 6022 (1), KAUM-I. 6023 (1), KAUM-I. 6024 (1), KAUM-I. 6025 (1), KAUM-I. 6026 (1), west of Sakura-jima Island, Kagoshima Bay, 31.6243°N, 130.6185°E, 140 m, 28 Jul. 2007; KAUM-I. 6980 (1), KAUM-I. 6981 (1), west of Sakura-jima Island, Kagoshima Bay, 31.6243°N, 130.6185°E, 140 m, 28 Oct. 2007; ZMUT 20157 (1), Kagoshima, date unknown; KAUM-I. 5876 (1), KAUM-I. 6061 (1), KAUM-I. 6074 (1), KAUM-I. 6075 (1), off Tarumizu, Kagoshima Bay, 31.4485°N, 130.6293°E, 227–227 m, 20 Aug. 2007; KAUM-I. 298 (1), off Tarumizu, Kagoshima Bay, 31.4688°N, 130.6303°E, 220 m, 20 Apr. 2006; KAUM-I. 38771 (1), south of Okiko-jima Island, Kagoshima Bay, 31.5333°N, 130.6333°E, 190 m, 26 Feb. 2011; KAUM-I. 11946 (1), off Tarumizu, Kagoshima Bay, Jul. 2008; BSKU 47209 (1), off Ibusuki, Kagoshima Bay, 190 m, 24 May 1989; BSKU 37575 (1), BSKU 37576 (1), BSKU 37577 (1), BSKU 37578 (1), BSKU 37580 (1), off Furue, Kagoshima Bay, 13 Oct. 1982; KAUM-I. 42791 (1), KAUM-I. 42792 (1), KAUM-I. 42793 (1), KAUM-I. 42794 (1), KAUM-I. 42795 (1), KAUM-I. 42796 (1), KAUM-I. 42797 (1), KAUM-I. 42798 (1), KAUM-I. 42799 (1), KAUM-I. 42800 (1), KAUM-I. 42801 (1),

KAUM-I. 42802 (1), KAUM-I. 42803 (1), KAUM-I. 42804 (1), KAUM-I. 42805 (1), KAUM-I. 42806 (1), KAUM-I. 42807 (1), KAUM-I. 42808 (1), KAUM-I. 42809 (1), KAUM-I. 42810 (1), KAUM-I. 42811 (1), KAUM-I. 42812 (1), KAUM-I. 42813 (1), KAUM-I. 42814 (1), KAUM-I. 42815 (1), KAUM-I. 42816 (1), KAUM-I. 42817 (1), KAUM-I. 42818 (1), KAUM-I. 42819 (1), KAUM-I. 42820 (1), KAUM-I. 42821 (1), off Furue, Kagoshima, 4 Sept. 1972; MUFS 4211 (1), MUFS 4212 (1), MUFS 4214 (1), Aoshima, Miyazaki, 300 m, 17 Apr. 1975; BSKU 41293 (1), Mimase fish market, 1 Apr. 1985; BSKU 52933 (1), BSKU 53429 (1), Mimase fish market, 11 Dec. 2000; BSKU 51771 (1), Mimase fish market, 18 Nov. 1999; BSKU 69481 (1), Mimase fish market, 24 Dec. 2003; BSKU 68130 (1), Mimase fish market, 26 Feb. 1974; BSKU 54046 (1), BSKU 54047 (1), BSKU 54048 (1), Mimase fish market, 4 Sept. 1999; BSKU 58381 (1), BSKU 58382 (1), Mimase fish market, 6 Dec. 2001; BSKU 35444 (1), Mimase fish market, 8 Jan. 1981; BSKU 74975 (1), Mimase fish market, 8 Nov. 2004; BSKU 36774 (1), Mimase fish market, 9 Mar. 1982; BSKU 99356 (1), Mimase fish market, 11 Jan. 2007; FAKU 33887 (1), FAKU 33888 (1), FAKU 33889 (1), Mimase fish market, Dec. 1960–Jan. 1961; FAKU 29398 (1), Mimase fish market, 24 Dec. 1959; BSKU 10033 (1), Mimase fish market, 10 Feb. 1961; BSKU 877 (1), BSKU 878 (1), Mimase fish market, 16 Jan. 1951; BSKU 5 (1), Mimase fish market, 19 Feb. 1951; BSKU 105 (1), Mimase fish market, 5 Feb. 1951; BSKU 99373 (1), Mimase fish market, 11 Jan. 2007; BSKU 99381 (1), BSKU 99382 (1), Mimase fish market, 9 Nov. 2007; BSKU 9902 (1), BSKU 9904 (1), Kochi, 1955; BSKU 82146 (1), Tosa Bay, 250 m, 12 Dec. 1995; ZMUT 47190 (1), Tosa Bay, date unknown; FRLM 34785 (1), off Owase, Kumano-nada, 27 Nov. 2008; FRLM 40505 (1), off Owase, Kumano-nada, 8 Dec. 2011; FRLM 34164 (1), FRLM 34165 (1), FRLM 34166 (1), off Owase, Kumano-nada, 4 Feb. 2008; FRLM 34260 (1), FRLM 34261 (1), FRLM 34262 (1), FRLM 34263 (1), FRLM 34264 (1), FRLM 34265 (1), FRLM 34282 (1), off Owase, Kumano-nada, 14 Jan. 2008; FRLM 34461 (1), off Owase, Kumano-nada, 18 Feb. 2008; FRLM 33332 (1), FRLM 33333 (1), FRLM 33334 (1), FRLM 33335 (1), FRLM 33336 (1), off Owase, Kumano-nada, 29 May 2007; FRLM 34426 (1), FRLM 34427 (1), off Owase, Kumano-nada, 7 Feb. 2008; FRLM 34211 (1), FRLM 34212 (1), off Owase, Kumano-nada, 7 Jan. 2008; FRLM 34360 (1), FRLM 34361 (1), off Owase, Kumano-nada, date unknown; BSKU 110038 (1), BSKU 110039 (1), BSKU 110040 (1), BSKU 110041 (1), BSKU 110042 (1), BSKU 110051 (1), Suruga Bay, 34.7127°N, 138.4553°E, 262–434 m, 23 Apr. 2013; BSKU 110025 (1), Suruga Bay, 34.7080°N, 138.4587°E, 200–450 m, 23 Apr. 2013; BSKU 110416 (1), Suruga Bay, 34.7481°N, 138.4652°E, 243–424 m, 23 Apr. 2013; BSKU 110427 (1), BSKU 110428 (1), Suruga Bay, 34.7619°N, 138.4811°E, 196–387 m, 23 Apr. 2013; BSKU 110142

(1), BSKU 110422 (1), BSKU 110436 (1), BSKU 110437 (1), BSKU 110438 (1), BSKU 110439 (1), BSKU 110440 (1), Suruga Bay, 34.7651°N, 138.4880°E, 200–450 m, 23 Apr. 2013; NSMT-P 46663 (1), off Tagoshima, Suruga Bay, 34.8233°N, 138.6833°E, 270–500 m, 23 Oct. 1994; KPM-NI 17957 (1), Suruga Bay, 400 m, 26 Nov. 2006; KPM-NI 10349 (1), eastern part of Suruga Bay, 110–450 m, 10 Dec. 1998; NSMT-P 46618 (1), off Kumomizaki, Suruga Bay, 34.7717°N, 138.7133°E, 275–500 m, 23 Oct. 1994; ZMUT 6844 (1), ZMUT 6845 (1), Tokyo, date unknown; BSKU 101997 (1), off Mito, 250 m, 9 Oct. 1997.

Coelorinchus kamoharai. 567 specimens: BSKU 26353 (1), north of Iriomote-jima Island, Okinawa Trough, 25.6783°N, 123.8783°E, 315–370 m, 20 Jan. 1978; BSKU 32469 (1), northwest of Miyako-jima Island, Okinawa Trough, 25.8000°N, 124.4250°E, 400–420 m, 14 Sept. 1979; BSKU 26582 (1), north of Miyako-jima Island, Okinawa Trough, 26.5200°N, 125.0417°E, 255–260 m, 23 Jan. 1978; BSKU 26640 (1), BSKU 26641 (1), BSKU 26642 (1), BSKU 26643 (1), BSKU 26644 (1), north of Miyako-jima Island, Okinawa Trough, 26.5533°N, 125.0817°E, 255–260 m, 25 Jan. 1978; BSKU 33040 (1), BSKU 33041 (1), west-northwest of Okinawa-jima Island, Okinawa Trough, 26.9500°N, 125.7500°E, 335 m, 10 Oct. 1979; BSKU 27385 (1), BSKU 27386 (1), BSKU 27387 (1), west of Amami-oshima Island, Okinawa Trough, 27.0333°N, 125.7667°E, 302–309 m, 3 Mar. 1978; BSKU 27504 (1), BSKU 27505 (1), BSKU 27506 (1), BSKU 27507 (1), BSKU 27508 (1), BSKU 27509 (1), west of Okinawa-jima Island, Okinawa Trough, 28.1000°N, 126.3833°E, 143 m, 9 Mar. 1978; BSKU 34391 (1), BSKU 34392 (1), BSKU 34393 (1), BSKU 34394 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.5583°N, 126.5233°E, 320–325 m, 16 Dec. 1979; BSKU 27331 (1), BSKU 27332 (1), BSKU 27333 (1), BSKU 27334 (1), BSKU 27335 (1), BSKU 27336 (1), BSKU 27337 (1), BSKU 27338 (1), west of Amami-oshima Island, Okinawa Trough, 28.6183°N, 126.8200°E, 220 m, 6 Feb. 1978; BSKU 27358 (1), west of Amami-oshima Island, Okinawa Trough, 28.4533°N, 126.9400°E, 300 m, 7 Feb. 1978; SNFR 11074 (2), East China Sea, 28.3750°N, 126.9683°E, 378–382 m, 23 Oct. 1990; BSKU 27721 (1), BSKU 27722 (1), north-northwest of Okinawa Island, Okinawa Trough, 28.5000°N, 127.0000°E, 225–235 m, 12 Mar. 1978; BSKU 29764 (1), BSKU 34620 (1), BSKU 34621 (1), BSKU 34622 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.9333°N, 127.0167°E, 220 m, 12 Oct. 1979; BSKU 28185 (1), BSKU 28186 (1), BSKU 28187 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.1667°N, 127.0167°E, 160 m, 17 Mar. 1978; BSKU 27279 (1), BSKU 27280 (1), BSKU 27323 (1), west-northwest of Amami-oshima Island, Okinawa Trough, 28.8033°N, 127.0617°E, 425 m, 6 Feb. 1978; BSKU 34668 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.9983°N, 127.1550°E, 325 m, 17 Dec. 1979;

SNFR 16345 (2), East China Sea, 28.9902°N, 127.1813°E, 400 m, 9 Nov. 2003; BSKU 27741 (1), BSKU 27742 (1), BSKU 27743 (1), BSKU 27744 (1), BSKU 27745 (1), BSKU 27746 (1), west of Amami-oshima Island, Okinawa Trough, 28.4333°N, 127.1833°E, 300–355 m, 12 Mar. 1978; BSKU 27240 (1), BSKU 27241 (1), BSKU 27242 (1), BSKU 27243 (1), BSKU 27244 (1), BSKU 27245 (1), BSKU 27246 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.2867°N, 127.2900°E, 250 m, 5 Feb. 1978; BSKU 28201 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.3000°N, 127.3333°E, 410–415 m, 17 Mar. 1978; BSKU 29713 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.3383°N, 127.3417°E, 315–330 m, 18 Dec. 1979; BSKU 28227 (1), BSKU 28228 (1), BSKU 28229 (1), BSKU 28230 (1), BSKU 28231 (1), BSKU 28232 (1), BSKU 28233 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.2667°N, 127.3500°E, 353–365 m, 17 Mar. 1978; BSKU 28287 (1), BSKU 28288 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7500°N, 127.5000°E, 325–360 m, 19 Mar. 1978; BSKU 28325 (1), BSKU 28326 (1), BSKU 28327 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7167°N, 127.6167°E, 410–423 m, 18 Mar. 1978; BSKU 33597 (1), BSKU 33598 (1), BSKU 33599 (1), BSKU 33600 (1), BSKU 33601 (1), BSKU 33602 (1), BSKU 33603 (1), BSKU 33604 (1), BSKU 33605 (1), BSKU 33606 (1), BSKU 33607 (1), BSKU 33608 (1), BSKU 33609 (1), BSKU 33610 (1), BSKU 33611 (1), BSKU 33612 (1), BSKU 33613 (1), BSKU 33614 (1), BSKU 33615 (1), BSKU 33616 (1), BSKU 33617 (1), BSKU 33618 (1), BSKU 33619 (1), BSKU 33620 (1), BSKU 33621 (1), BSKU 33622 (1), BSKU 33623 (1), BSKU 33624 (1), BSKU 33625 (1), BSKU 33626 (1), BSKU 33627 (1), BSKU 33628 (1), BSKU 33629 (1), northwest of Amami-oshima Island, Okinawa Trough, 30.0773°N, 127.6788°E, 220–245 m, 27 Oct. 1979; BSKU 28623 (1), BSKU 28624 (1), BSKU 28625 (1), BSKU 28626 (1), BSKU 28627 (1), BSKU 28628 (1), BSKU 28629 (1), BSKU 28630 (1), BSKU 28631 (1), BSKU 28632 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.8000°N, 127.7000°E, 335–345 m, 21 Mar. 1978; BSKU 28655 (1), BSKU 28656 (1), BSKU 28657 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.9000°N, 127.7167°E, 250 m, 21 Mar. 1978; BSKU 33822 (1), BSKU 33823 (1), BSKU 33824 (1), BSKU 33825 (1), BSKU 33826 (1), BSKU 33827 (1), BSKU 33828 (1), BSKU 33830 (1), BSKU 33831 (1), BSKU 33832 (1), BSKU 33833 (1), BSKU 33834 (1), BSKU 33835 (1), northwest of Amami-oshima Island, Okinawa Trough, 30.4815°N, 127.8432°E, 320–360 m, 28 Oct. 1979; FFNU P-00556 (1 of 2), southwest of Daini-Danjo Knoll, East China Sea, 30.8375°N, 127.9982°E, 340 m, 9 Nov. 2004; BSKU 33865 (1), BSKU 33866 (1), BSKU 33867 (1), BSKU 33868 (1), BSKU 33869 (1), BSKU 33870 (1), BSKU 33871 (1), BSKU 33872 (1), BSKU 33873 (1), BSKU 33874 (1), BSKU 33875 (1), BSKU 33876 (1), BSKU 33877 (1), BSKU 33878 (1), BSKU 33879 (1), BSKU 33880 (1), BSKU 33881 (1), west-northwest of Yaku-shima Island, Okinawa Trough, 31.0113°N, 128.0908°E, 315–320 m, 30 Oct. 1979; FFNU P-00590 (1), southwest of Daini-Danjo Knoll, East China Sea, 30.9763°N, 128.1740°E, 337 m, 24 Apr. 2005; BSKU 22898 (1), south of Danjo Islands, Okinawa Trough, 31.2617°N, 128.3433°E, 364–369 m, 5 Aug. 1974; FFNU P-00685 (12 of 13), northwest of Daiichi-Danjo Knoll, East China Sea, 31.7132°N, 128.4012°E, 302 m, 1 Aug. 2007; FFNU P-00610 (1 of 2), south of Danjo Islands, East China Sea, 31.6513°N, 128.4370°E, 419 m, 9 Nov. 2005; FFNU P-00547 (9), southwest of Fukue-jima Island, East China Sea, 32.5462°N, 128.4698°E, 257 m, 8 Nov. 2004; SNFR 17678 (1), SNFR 17679 (1), East China Sea, 32.2865°N, 128.9592°E, 301–304 m, 10 Jun. 2011; BSKU 109037 (1), west of Sanpo-sone, 32.2872°N, 129.0732°E, 351–389 m, 19 Nov. 2012; KAUM-I. 6644 (1), off Nomaie, Kagoshima, 300 m, 28 Sept. 2007; KAUM-I. 6645 (1), off Nomaie, Kagoshima, 300 m, 29 Sept. 2007; KAUM-I. 32208 (1), KAUM-I. 32246 (1), KAUM-I. 32258 (1), KAUM-I. 32259 (1), KAUM-I. 32260 (1), KAUM-I. 32261 (1), KAUM-I. 32262 (1), KAUM-I. 32263 (1), KAUM-I. 32264 (1), KAUM-I. 32265 (1), off Nomaie, Kagoshima, 31.5000°N, 129.8833°E, 370–400 m, 10 Sept. 2010; BSKU 106805 (1), BSKU 106806 (1), southeast of Shimokoshiki-jima Island, East China Sea, 31.5560°N, 129.8847°E, 415 m, 24 Apr. 2012; BSKU 106765 (1), BSKU 106766 (1), southeast of Shimokoshiki-jima Island, East China Sea, 31.5658°N, 129.8915°E, 380 m, 24 Apr. 2012; BSKU 19075 (1), BSKU 19076 (1), off Satsuma Peninsula, Okinawa Trough, 31.2267°N, 129.9750°E, 310 m, 12 Feb. 1959; MUFS 13348 (1), MUFS 13349 (1), MUFS 13350 (1), Meitsu, Miyazaki, 1996; MUFS 4213 (1), MUFS 4233 (1), Aoshima, Miyazaki, 300 m, 17 Apr. 1975; MUFS 39533 (20), Aoshima, Miyazaki, 300–350 m, 19 Apr. 1987; BSKU 78863 (1), Saga fish market, 22 Jun. 2006; BSKU 69285 (1), BSKU 69286 (1), BSKU 69287 (1), Tosa Bay, 33.1450°N, 133.5150°E, 240–249 m, 20 Aug. 1997; BSKU 99448 (1), BSKU 99449 (1), Tosa Bay, 33.1412°N, 133.5388°E, 295–336 m, 18 Jul. 1988; BSKU 53789 (1), Mimase fish market, 1 Apr. 2001; BSKU 75904 (1), BSKU 75905 (1), Mimase fish market, 1 Dec. 2004; BSKU 29954 (1), BSKU 30253 (1), Mimase fish market, 1 Mar. 1980; BSKU 59293 (1), Mimase fish market, 1 Mar. 2002; BSKU 99264 (1), BSKU 99265 (1), BSKU 99266 (1), Mimase fish market, 1 Mar. 2006; BSKU 58482 (1), Mimase fish market, 10 Dec. 2001; BSKU 99305 (1), Mimase fish market, 11 Apr. 2007; BSKU 92022 (1), Mimase fish market, 11 Dec. 2006; BSKU 99358 (1), BSKU 99359 (1), BSKU 99360 (1), Mimase fish market, 11 Jan. 2007; BSKU 99287 (1), Mimase fish market, 11 Oct. 2007; BSKU 71390 (1), BSKU 71391 (1), BSKU 71392 (1), Mimase fish market, 12 Apr. 2004; BSKU 70634 (1), Mimase fish market, 12 Jan. 2004;

BSKU 88707 (1), BSKU 88744 (1), Mimase fish market, 12 Oct. 2006; BSKU 72286 (1), Mimase fish market, 13 Apr. 2004; BSKU 99231 (1), Mimase fish market, 13 Apr. 2005; BSKU 79480 (1), Mimase fish market, 13 Jan. 2003; BSKU 30181 (1), BSKU 30182 (1), BSKU 30183 (1), Mimase fish market, 15 Apr. 1978; BSKU 29059 (1), Mimase fish market, 15 Dec. 1978; BSKU 63546 (1), Mimase fish market, 15 Dec. 2002; BSKU 75649 (1), Mimase fish market, 15 Dec. 2004; BSKU 89691 (1), Mimase fish market, 15 Jan. 2007; BSKU 69152 (1), Mimase fish market, 15 Mar. 2004; BSKU 76123 (1), Mimase fish market, 15 Mar. 2005; BSKU 64840 (1), Mimase fish market, 16 Apr. 2003; BSKU 89633 (1), BSKU 89634 (1), Mimase fish market, 17 Nov. 2006; BSKU 99257 (1), BSKU 99258 (1), BSKU 99259 (1), Mimase fish market, 18 Jan. 2006; BSKU 30292 (1), BSKU 30293 (1), Mimase fish market, 19 Apr. 1980; BSKU 78030 (1), BSKU 78031 (1), BSKU 78032 (1), BSKU 99406 (1), BSKU 99407 (1), BSKU 99408 (1), BSKU 99409 (1), BSKU 99412 (1), BSKU 99413 (1), Mimase fish market, 19 Apr. 2006; BSKU 54618 (1), Mimase fish market, 19 Dec. 1998; BSKU 89761 (1), BSKU 89762 (1), BSKU 89763 (1), BSKU 89764 (1), Mimase fish market, 19 Jan. 2007; BSKU 69231 (1), BSKU 69232 (1), BSKU 69233 (1), Mimase fish market, 19 Nov. 2003; BSKU 78494 (1), BSKU 78495 (1), BSKU 78496 (1), BSKU 78497 (1), BSKU 78498 (1), Mimase fish market, 19 Oct. 1996; BSKU 89410 (1), BSKU 89411 (1), Mimase fish market, 19 Oct. 2006; BSKU 76103 (1), Mimase fish market, 2 Mar. 2005; BSKU 55693 (1), Mimase fish market, 20 Feb. 1997; BSKU 68921 (1), Mimase fish market, 20 Jan. 2004; BSKU 58315 (1), BSKU 74698 (1), Mimase fish market, 20 Nov. 1996; BSKU 53281 (1), Mimase fish market, 21 Apr. 1995; BSKU 70801 (1), BSKU 70802 (1), BSKU 70803 (1), BSKU 70804 (1), Mimase fish market, 21 Feb. 2004; BSKU 89467 (1), Mimase fish market, 21 Feb. 2007; BSKU 63296 (1), Mimase fish market, 22 Jan. 2003; BSKU 82826 (1), BSKU 82827 (1), Mimase fish market, 22 Nov. 1996; BSKU 78372 (1), Mimase fish market, 23 Apr. 2006; BSKU 99241 (1), BSKU 99242 (1), BSKU 99243 (1), BSKU 99244 (1), Mimase fish market, 23 Apr. 2007; BSKU 62499 (1), Mimase fish market, 23 Dec. 2002; BSKU 53123 (1), BSKU 54124 (1), Mimase fish market, 23 Nov. 2000; BSKU 52995 (1), BSKU 52996 (1), Mimase fish market, 23 Oct. 2000; BSKU 99479 (1), BSKU 99480 (1), BSKU 99481 (1), BSKU 99482 (1), Mimase fish market, 23 Oct. 2006; BSKU 78100 (1), BSKU 78101 (1), Mimase fish market, 24 Apr. 2006; BSKU 86078 (1), Mimase fish market, 24 Dec. 1998; BSKU 75992 (1), Mimase fish market, 24 Oct. 2004; BSKU 71094 (1), Mimase fish market, 25 Apr. 2004; BSKU 89323 (1), Mimase fish market, 25 Oct. 2005; BSKU 70020 (1), BSKU 70021 (1), BSKU 70023 (1), BSKU 70024 (1), Mimase fish market, 26 Feb. 2003; BSKU 88121 (1), Mimase fish market, 26 Mar. 2006; BSKU 91577 (1), Mimase fish market, 26 Mar. 2007; BSKU 82934 (1), Mimase fish market, 26 Nov. 1996; BSKU 89529 (1), BSKU 89530 (1), Mimase fish market, 26 Oct. 2006; BSKU 99227 (1), BSKU 99228 (1), BSKU 99229 (1), BSKU 99230 (1), Mimase fish market, 27 Apr. 2007; BSKU 75933 (1), BSKU 75934 (1), Mimase fish market, 27 Dec. 2004; BSKU 71715 (1), BSKU 78387 (1), Mimase fish market, 27 Feb. 2004; BSKU 76105 (1), Mimase fish market, 27 Feb. 2005; BSKU 52486 (1), Mimase fish market, 27 Mar. 2000; BSKU 55380 (1), Mimase fish market, 27 Oct. 2000; BSKU 89736 (1), Mimase fish market, 27 Oct. 2006; BSKU 49878 (1), Mimase fish market, 28 Nov. 1991; BSKU 36048 (1), Mimase fish market, 28 Oct. 1981; BSKU 74968 (1), Mimase fish market, 28 Oct. 2004; BSKU 90319 (1), Mimase fish market, 29 Jan. 1998; BSKU 90190 (1), Mimase fish market, 29 Jan. 2007; BSKU 58413 (1), Mimase fish market, 29 Nov. 2001; BSKU 90052 (1), BSKU 90053 (1), Mimase fish market, 29 Oct. 2006; BSKU 35613 (1), BSKU 35614 (1), Mimase fish market, 3 Mar. 1981; BSKU 79524 (1), Mimase fish market, 3 Mar. 2006; BSKU 91650 (1), Mimase fish market, 3 Nov. 2006; BSKU 35372 (4), Mimase fish market, 3 Oct. 1980; BSKU 89320 (1), Mimase fish market, 3 Oct. 2006; BSKU 76070 (1), Mimase fish market, 30 Mar. 2005; BSKU 53704 (1), BSKU 87366 (1), Mimase fish market, 30 Nov. 1995; BSKU 52719 (1), Mimase fish market, 30 Nov. 2000; BSKU 69375 (1), Mimase fish market, 4 Feb. 2004; BSKU 69023 (1), Mimase fish market, 5 Dec. 2003; BSKU 90346 (2), Mimase fish market, 5 Oct. 1996; BSKU 53816 (1), Mimase fish market, 6 Apr. 2001; BSKU 90222 (1), Mimase fish market, 6 Dec. 2006; BSKU 53205 (1), BSKU 53206 (1), BSKU 53207 (1), Mimase fish market, 6 Nov. 2000; BSKU 30158 (1), Mimase fish market, 7 Dec. 1979; BSKU 83003 (1), BSKU 83004 (1), BSKU 83005 (1), Mimase fish market, 7 Dec. 1995; MUFS 5758 (1), Mimase fish market, 7 Jan. 1976; BSKU 70130 (1), Mimase fish market, 7 Jan. 2004; BSKU 76120 (1), BSKU 79525 (1), BSKU 79527 (1), Mimase fish market, 7 Mar. 2005; BSKU 57640 (1), BSKU 57641 (1), Mimase fish market, 7 Nov. 2001; BSKU 35443 (1), Mimase fish market, 8 Jan. 1981; BSKU 85042 (1), Mimase fish market, 8 Mar. 1978; BSKU 74974 (1), Mimase fish market, 8 Nov. 2004; BSKU 89641 (1), BSKU 89642 (1), BSKU 89643 (1), Mimase fish market, 8 Nov. 2006; BSKU 84087 (1), Mimase fish market, 9 Dec. 1995; BSKU 69533 (1), BSKU 69534 (1), Mimase fish market, 9 Jan. 2004; BSKU 74525 (1), BSKU 76684 (1), Mimase fish market, 9 Mar. 2005; BSKU 11715 (1), Mimase fish market, Nov. 1965; FAKU 21782 (1), Mimase fish market, 10–20 Apr. 1954; BSKU 89059 (1), BSKU 89061 (1), BSKU 89062 (1), Mimase fish market, 14 Dec. 2006; BSKU 91841 (1), Mimase fish market, 17 Oct. 2007; BSKU 94806 (1), Mimase fish market, 3 Nov. 2006; BSKU 1941 (1), Mimase fish market, 3 Jun. 1952; BSKU 3592 (1), BSKU 3593 (1),

BSKU 3594 (1), Mimase fish market, 23 Mar. 1954; BSKU 9671 (1), BSKU 9674 (1), BSKU 9675 (1), BSKU 9676 (1), BSKU 9677 (1), BSKU 9679 (1), BSKU 9680 (1), Mimase fish market, 24 Dec. 1950–Jan. 1956; BSKU 6734 (1), BSKU 6736 (1), Mimase fish market, Dec. 1955–Jan. 1956; BSKU 23289 (1), BSKU 89721 (1), BSKU 89722 (1), BSKU 89723 (1), BSKU 89724 (1), BSKU 12786 (1), Mimase fish market, date unknown; BSKU 92888 (1), Mimase fish market, 17 Dec. 2007; BSKU 99377 (1), Mimase fish market, 19 Nov. 2007; BSKU 94219 (1), Mimase fish market, 2 Mar. 2008; BSKU 93113 (1), Mimase fish market, 27 Dec. 2007; BSKU 94204 (1), Mimase fish market, 28 Feb. 2008; KAUM-I. 12645 (1), KAUM-I. 12907 (1), KAUM-I. 12910 (1), Mimase fish market, 9 May 2008; BSKU 2437 (1), Urado fish market, 1952; BSKU 78088 (1), Tosa Bay, 150 m, 17 Nov. 2005; BSKU 78502 (1), BSKU 78503 (1), BSKU 78504 (1), BSKU 78505 (1), BSKU 78506 (1), BSKU 78507 (1), BSKU 78508 (1), BSKU 78509 (1), BSKU 78510 (1), BSKU 78511 (1), BSKU 78512 (1), BSKU 78513 (1), Tosa Bay, 200 m, 11 Oct. 2005; BSKU 85962 (1), BSKU 99418 (1), BSKU 99419 (1), Tosa Bay, 200 m, 11 Sept. 1998; BSKU 73666 (1), Tosa Bay, 200 m, 13 Sept. 2004; BSKU 70364 (1), Tosa Bay, 200 m, 18 Nov. 2003; BSKU 95439 (1), Tosa Bay, 200 m, 10 Oct. 2007; BSKU 99307 (1), BSKU 99308 (1), BSKU 99309 (1), BSKU 99310 (1), BSKU 99311 (1), BSKU 99312 (1), BSKU 99313 (1), BSKU 99314 (1), BSKU 99315 (1), BSKU 99316 (1), BSKU 99317 (1), Tosa Bay, 200 m, 18 Jul. 2007; BSKU uncat. (3), BSKU 22593 (1), BSKU 22594 (1), BSKU 22595 (1), BSKU 22596 (1), BSKU 22597 (1), BSKU 22598 (1), BSKU 22599 (1), BSKU 22600 (1), BSKU 22601 (1), BSKU 22602 (1), BSKU 22603 (1), BSKU 22604 (1), Tosa Bay, 23 Jul. 1974; BSKU 40350 (1), Tosa Bay, 9 Oct. 1984; BSKU 13297 (1), ca. 10–15 km west of Cape Murotomisaki, Tosa Bay, 360–435 m, 8–9 Jun. 1968; BSKU 99288 (1), BSKU 99289 (1), Mimase fish market, 29 Oct. 2007; BSKU 66867 (1), Owase fish market, Kumano-nada, 20 Oct. 2003; FAKU 1597 (1), Owase fish market, Kumano-nada, 6–9 Dec. 1935; FAKU 15148 (1), Owase fish market, Kumano-nada, 18–20 Jan. 1950; FAKU 21505 (1), FAKU 21506 (1), Owase fish market, Kumano-nada, 15 Feb. 1954; FAKU 24888 (1), FAKU 24889 (1), FAKU 24894 (1), FAKU 24895 (1), FAKU 24896 (1), FAKU 24901 (1), FAKU 24903 (1), FAKU 24905 (1), FAKU 24907 (1), FAKU 24908 (1), Owase fish market, Kumano-nada, 25 Oct. 1954; BSKU 99451 (1), BSKU 99452 (1), BSKU 99453 (1), Owase fish market, Kumano-nada, 13 Mar. 1983; ZMUT 23378 (1), Owase, Mie, date unknown; FRLM 8634 (1), Owase fish market, 16 Mar. 1967; FRLM 34167 (1), off Owase, Kumano-nada, 4 Feb. 2008; FRLM 34272 (1), FRLM 34273 (1), off Owase, Kumano-nada, 14 Jan. 2008; FRLM 34459 (1), off Owase, Kumano-nada, 18 Feb. 2008; FRLM 34425 (1), off Owase, Kumano-nada, 7 Feb. 2008; FRLM 33944 (1), off Owase,

Kumano-nada, 14 Jan. 2008; FRLM 33917 (1), FRLM 33918 (1), FRLM 33919 (1), FRLM 33920 (1), FRLM 33921 (1), FRLM 33922 (1), off Owase, Kumano-nada, 16 Nov. 2007; FRLM 30762 (1), FRLM 30763 (1), off Owase, Kumano-nada, 2 Feb. 2004; FAKU 4922 (1), Kumano-nada, 1938; FAKU 20021 (1), FAKU 20022 (1), off Cape Daiozaki, Kumano-nada, 18–20 Dec. 1952; FRLM 27237 (1), FRLM 27238 (1), off Kowaura, Kumano-nada, 165–309 m, 5 Dec. 2000; FRLM 27216 (1), off Kowaura, Kumano-nada, 165–309 m, 5 Nov. 2000; FAKU 17720 (1), FAKU 17722 (1), Miya fish market, Enshu-nada, 15 Mar. 1952; FAKU 26940 (1), FAKU 26941 (1), Miya fish market, Enshu-nada, 29–31 Dec. 1956; FAKU 28380 (1), FAKU 28381 (1), FAKU 28382 (1), FAKU 28383 (1), FAKU 28384 (1), FAKU 28385 (1), FAKU 28386 (1), FAKU 28387 (1), Miya fish market, Enshu-nada, 5–7 Feb. 1959; BSKU 110409 (1), Suruga Bay, 34.7481°N, 138.4652°E, 243–424 m, 23 Apr. 2013; BSKU 110431 (1), Suruga Bay, 34.7489°N, 138.4664°E, 200–450 m, 23 Apr. 2013; KPM-NI 18044 (1), off Fuji River, Suruga Bay, 300 m, Nov. 2006; KPM-NI 10317 (1), eastern part of Suruga Bay, 110–450 m, 10 Dec. 1998; KPM-NI 27420 (1), eastern part of Suruga Bay, 200–400 m, Mar. 2009; KPM-NI 24982 (1), eastern part of Suruga Bay, Nov. 2009; NSMT-P 9972 (14), off Heta, Suruga Bay, 28 Nov. 1969; BSKU 74211 (1), off Numazu, Suruga Bay, 16 Jan. 2005; FAKU 10035 (1), Heta fish market, Suruga Bay, 1938; FAKU 25286 (1), Heta fish market, Suruga Bay, 26 Mar. 1939; KPM-NI 8600 (1), off Cape Irozaki, Izu Peninsula, 150–220 m, 22 Apr. 2001; ZMUT 30608 (1), Choshi fish market, 20 Nov. 1934; NSMT-P 48952 (1), east of Oshika Peninsula, 38.4380°N, 141.9098°E, 249–253 m, 29 Oct. 1995; SNFR 11821 (2), Sea of Japan, 33.6833°N, 128.3500°E, 115 m, 5 Jul. 1988; BSKU 14207 (1), off east coast of Japan, date unknown.

Coelorrinchus kishinouyei. 56 specimens: FAKU 34462 (1), Owase fish market, Kumano-nada, 12–14 Nov. 1961; FAKU 2474 (1), FAKU 2475 (1), FAKU 2476 (1), FAKU 2478 (1), FAKU 2481 (1), FAKU 2484 (1), Owase fish market, Kumano-nada, 4–9 Jan. 1936; FAKU 27789 (1), FAKU 27790 (1), FAKU 27791 (1), FAKU 27792 (1), FAKU 27793 (1), FAKU 27794 (1), FAKU 27795 (1), FAKU 27796 (1), FAKU 27797 (1), FAKU 27798 (1), FAKU 27799 (1), FAKU 27800 (1), Owase fish market, Kumano-nada, 25 Feb.–10 Mar. 1957; FRLM 8635 (1), Owase fish market, 16 Mar. 1967; FRLM 34210 (1), off Owase, Kumano-nada, 7 Jan. 2008; FAKU 28368 (1), Miya fish market, Enshu-nada, 5–7 Feb. 1959; ZMUT 18525 (1), Maisaka, Shizuoka, date unknown; KPM-NI 14642 (1), Mochimune fish market, Shizuoka, 8 Sept. 1992; BSKU 110037 (1), Suruga Bay, 34.7127°N, 138.4553°E, 262–434 m, 23 Apr. 2013; BSKU 110014 (1), BSKU 110018 (1), BSKU 110019 (1), BSKU 110021 (1), BSKU 110022 (1), Suruga Bay, 34.7080°N, 138.4587°E, 200–450 m, 23 Apr. 2013; BSKU 110415 (1), Suruga Bay,

34.7481°N, 138.4652°E, 243–424 m, 23 Apr. 2013; KPM-NI 18045 (1), off Fuji River, Suruga Bay, 300 m, Nov. 2006; NSMT-P 101347 (1), Suruga Bay, date unknown; NSMT-P 102000 (4), Suruga Bay, 1 May 1973; FAKU M7530 (1), Heta fish market, Suruga Bay, 150 ftn (275 m), 22–24 Jun. 1938; FAKU M7405 (1), FAKU M7406 (1), Heta fish market, Suruga Bay, 150 ftn (275 m), 23 Jul. 1938; FAKU M7229 (1), Heta fish market, Suruga Bay, 26 Mar. 1939; FAKU 34646 (1), off Oiso, Sagami Bay, 12 Mar. 1962; KPM-NI 1066 (1), Sagami Bay, 1 Jul. 1968; FAKU 34239 (1), FAKU 34245 (1), FAKU 34247 (1), FAKU 34248 (1), FAKU 34249 (1), FAKU 34251 (1), FAKU 34254 (1), FAKU 34255 (1), off Kurihama, Tokyo Bay (questionable), Apr. 1969; FAKU 34252 (1), off Kurihama, Sagami Bay, Apr. 1969; ZMUT 6842 (1), Tokyo, date unknown; FMNH 120862 (2), Dong-gang fish market, Pingtung, 10 Sept. 2009; MUFS 2600 (1), Dong-gang, Pingtung, 27 Feb. 1973.

Coelorinchus leptorhinus. 1 specimen: ASIZP61244 (1), Dong-gang fish market, Pingtung, 1 May 2000.

Coelorinchus longissimus. 130 specimens: BSKU 29823 (1), BSKU 33183 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.3167°N, 127.3000°E, 260 m, 13 Oct. 1979; SNFR 13754 (1), East China Sea, 29.7933°N, 127.5915°E, 273–278 m, 8 Jun. 2008; BSKU 29832 (1), BSKU 29835 (1), northwest of Amami-oshima Island, Okinawa Trough, 30.4815°N, 127.8432°E, 320–360 m, 28 Oct. 1979; SNFR 14798 (1), SNFR 14799 (4), East China Sea, 30.2953°N, 127.8872°E, 329–345 m, 13 Sept. 2009; FFNU P-00556 (1 of 2), southwest of Daini-Danjo Knoll, East China Sea, 30.8375°N, 127.9982°E, 340 m, 9 Nov. 2004; SNFR 14929 (1), SNFR 14930 (1), East China Sea, 30.8718°N, 128.0312°E, 325–366 m, 19 Sept. 2009; FFNU P-00685 (1 of 13), northwest of Daiichi-Danjo Knoll, East China Sea, 31.7132°N, 128.4012°E, 302 m, 1 Aug. 2007; FFNU P-00610 (1 of 2), south of Danjo Islands, East China Sea, 31.6513°N, 128.4370°E, 419 m, 9 Nov. 2005; SNFR 17677 (1), East China Sea, 32.2865°N, 128.9592°E, 301–304 m, 10 Jun. 2011; BSKU 109029 (1), BSKU 109030 (1), BSKU 109041 (1), BSKU 109051 (1), east of Kasayama Bank, 32.3884°N, 129.0505°E, 304–312 m, 19 Nov. 2012; BSKU 109036 (1), west of Sanpo-sone, 32.2872°N, 129.0732°E, 351–389 m, 19 Nov. 2012; KAUM-I. 6637 (1), off Nomaie, Kagoshima, 300 m, 24 Sept. 2007; KAUM-I. 6642 (1), off Nomaie, Kagoshima, 300 m, 26 Sept. 2007; KAUM-I. 32207 (1), off Nomaie, Kagoshima, 31.5000°N, 129.8833°E, 370–400 m, 10 Sept. 2010; MUFS 13351 (1), Meitsu, Miyazaki, 1996; MUFS 80524 (1), off Mimitsu, Miyazaki, 12 Jul. 1983; BSKU 69288 (1), BSKU 69289 (1), BSKU 69290 (1), BSKU 69292 (1), BSKU 69293 (1), BSKU 69294 (1), Tosa Bay, 33.1450°N, 133.5150°E, 240–249 m, 20 Aug. 1997; KAUM-I. 43938 (1), Mimase fish market, 350 m, 12 Dec. 2011; BSKU 99261 (1), BSKU 99272 (1), Mimase fish market, 1 Mar. 2006; BSKU 88705 (1), BSKU 88706 (1), Mimase fish market,

12 Oct. 2006; BSKU 53030 (1), Mimase fish market, 13 Apr. 2000; BSKU 99232 (1), BSKU 99233 (1), Mimase fish market, 13 Apr. 2005; BSKU 89394 (1), Mimase fish market, 13 Nov. 2006; BSKU 75631 (1), Mimase fish market, 14 Jan. 1995; BSKU 90842 (1), Mimase fish market, 14 Jan. 2007; BSKU 76106 (1), Mimase fish market, 16 Mar. 2005; BSKU 89393 (1), Mimase fish market, 17 Nov. 2006; BSKU 99405 (1), Mimase fish market, 19 Apr. 2006; BSKU 89395 (1), BSKU 89412 (1), Mimase fish market, 19 Oct. 2006; BSKU 99416 (1), BSKU 99417 (1), Mimase fish market, 2 Mar. 2005; BSKU 57972 (1), BSKU 57973 (1), Mimase fish market, 21 Oct. 2001; BSKU 39772 (1), Mimase fish market, 23 Jan. 1984; BSKU 70022 (1), Mimase fish market, 26 Feb. 2003; BSKU 49873 (1), Mimase fish market, 28 Nov. 1991; BSKU 39694 (1), Mimase fish market, 29 Nov. 1983; BSKU 40859 (1), BSKU 40860 (1), Mimase fish market, 29 Nov. 1984; BSKU 91969 (1), Mimase fish market, 31 Oct. 2007; BSKU 90969 (1), Mimase fish market, 6 Apr. 2007; MUFS 5757 (1), Mimase fish market, 7 Jan. 1976; BSKU 40443 (1), BSKU 40444 (1), BSKU 40446 (1), BSKU 40449 (1), BSKU 40450 (1), BSKU 40452 (1), BSKU 40453 (1), BSKU 40454 (1), BSKU 40455 (1), BSKU 40456 (1), BSKU 40458 (1), BSKU 40459 (1), Mimase fish market, 8 Mar. 1985; BSKU 39080 (1), Mimase fish market, 9 Apr. 1983; BSKU 91599 (1), BSKU 91600 (1), Mimase fish market, 9 Nov. 2006; BSKU 99286 (1), Mimase fish market, 12 Oct. 2007; BSKU 99343 (1), BSKU 99344 (1), BSKU 99345 (1), BSKU 99347 (1), BSKU 99348 (1), Mimase fish market, 23 Apr. 2008; BSKU 99234 (1), BSKU 99236 (1), Mimase fish market, 31 Oct. 2007; FAKU 29088 (1), Mimase fish market, 20 Mar. 1960; BSKU 1819 (1), Mimase fish market, Mar. 1952; BSKU 1517 (1), Mimase fish market, 15 Dec. 1951; BSKU 160 (1), Mimase fish market, 21 Mar. 1951; BSKU 262 (1), Mimase fish market, 27 Dec. 1950; BSKU 211 (1), Mimase fish market, 3 Dec. 1950; BSKU 54 (1), BSKU 56 (1), Mimase fish market, 5 Mar. 1951; BSKU 6740 (1), BSKU 6741 (1), BSKU 6751 (1), BSKU 6752 (1), Mimase fish market, Dec. 1955–Jan. 1956; BSKU 13915 (1), BSKU 13916 (1), BSKU 13917 (1), Mimase fish market, Jan. 1968; BSKU 13879 (1), Mimase fish market, Mar. 1968; BSKU 94483 (1), Mimase fish market, 14 Apr. 2008; BSKU 92336 (1), Mimase fish market, 15 Nov. 2007; BSKU 92368 (1), Mimase fish market, 16 Nov. 2007; BSKU 93574 (1), Mimase fish market, 21 Dec. 2007; BSKU 93089 (1), Mimase fish market, 24 Dec. 2007; BSKU 78514 (1), Tosa Bay, 200 m, 11 Oct. 2005; BSKU 92543 (1), BSKU 92544 (1), Tosa Bay, 200 m, 23 Aug. 2007; BSKU 79380 (1), BSKU 79475 (1), BSKU 79476 (1), BSKU 79477 (1), Tosa Bay, 200 m, 4 Aug. 2006; BSKU 82147 (1), Tosa Bay, 250 m, 12 Dec. 1995; BSKU 66410 (1), BSKU 67814 (1), BSKU 67815 (1), BSKU 67816 (1), Tosa Bay, 250 m, 8 Oct. 2003; BSKU 66819 (1), Tosa Bay, 250 m, 21 May 2003; BSKU 99290 (1), BSKU 99291 (1),

Mimase fish market, 29 Oct. 2007; BSKU 42678 (1), off Okitsu, Tosa Bay, 170–180 m, 7 Jan. 1986; FAKU 34472 (1), FAKU 34473 (1), FAKU 34475 (1), FAKU 34476 (1), Owase fish market, Kumano-nada, 12–14 Nov. 1961; FRLM 33785 (1), FRLM 33786 (1), FRLM 33787 (1), off Owase, Kumano-nada, 21 Nov. 2007; BSKU 110015 (1), Suruga Bay, 34.7080°N, 138.4587°E, 200–450 m, 23 Apr. 2013; FMNH 76861 (1), Boso Peninsula, 21 Mar. 1906.

Coelorinchus macrochir. 109 specimens: BSKU 26987 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.6467°N, 127.9733°E, 830 m, 4 Feb. 1978; BSKU 7192 (1), BSKU 7819 (1), Mimase fish market, 10 Apr. 1957; FAKU 27831 (1), Owase fish market, Kumano-nada, 25 Feb.–10 Mar. 1957; FRLM 34768 (1), FRLM 34769 (1), off Owase, Kumano-nada, 24 Oct. 2008; FRLM 33337 (1), FRLM 33338 (1), off Owase, Kumano-nada, 29 May 2007; FAKU 5784 (1), Kumano-nada, 26 Jun. 1938; BSKU 110043 (1), BSKU 110054 (1), BSKU 110125 (1), Suruga Bay, 34.7127°N, 138.4553°E, 262–434 m, 23 Apr. 2013; BSKU 110410 (1), BSKU 110414 (1), Suruga Bay, 34.7481°N, 138.4652°E, 243–424 m, 23 Apr. 2013; BSKU 110418 (1), BSKU 110430 (1), Suruga Bay, 34.7619°N, 138.4811°E, 196–387 m, 23 Apr. 2013; KPM-NI 17958 (1), Suruga Bay, 400 m, 26 Nov. 2006; KPM-NI 26655 (1), KPM-NI 26656 (1), Suruga Bay, Jan. 2010; FAKU 34670 (1), FAKU 34671 (1), off Shishihama, Suruga Bay, 15–20 Feb. 1962; NSMT-P 101342 (5), west of Nii-jima Island, Shichito-Iojima Ridge, 34.4117°N, 139.0883°E, 669–718 m, 8 Dec. 1987; FAKU 28413 (1), off Choshi, 15 Feb. 1957; ZMUT 23320 (1), ZMUT 23321 (1), ZMUT 23322 (1), off Minato, Ibaraki, Mar. 1931; ZMUT 14364 (1), ZMUT 14365 (1), Kujihama, Ibaraki, date unknown; ZMUT 30620 (1), ZMUT 30621 (1), ZMUT 30622 (1), ZMUT 30623 (1), Choshi fish market, 27 Jan. 1935; ZMUT 37682 (1), ZMUT 37683 (1), ZMUT 26288 (1), ZMUT 26289 (1), Onahama, Fukushima, date unknown; NSMT-P 66522 (1), Onahama fish market, 28 Dec. 2002; NSMT-P 75074 (1), off Hitachi, 36.5147°N, 140.9927°E, 250 m, 18 Apr. 1996; NSMT-P 77149 (1), Shiogama fish market, 2 Jun. 1940; BSKU 101970 (21), off Hitachi, 36.6143°N, 141.0965°E, 244–262 m, 9 Nov. 1995; BSKU 19255 (1), off Hitachi, 36.6067°N, 141.1683°E, 285–295 m, 14 May 1962; BSKU 101973 (14), off Iwaki, 37.0725°N, 141.5110°E, 271–303 m, 4 Nov. 1995; NSMT-P 102779 (1), off Minami-soma, 37.6148°N, 141.5977°E, 213 m, 29 Oct. 2006; NSMT-P 47406 (1), off Sendai, 600 m, 2 Nov. 1995; ZMUT 14139 (1), Kesen, Iwate, 12 Jan. 1925; ZMUT 13861 (1), Kesen, Iwate, 5 Jan. 1925; NSMT-P 53597 (10), east of Oshika Peninsula, 38.4750°N, 141.9018°E, 250–253 m, 11 Apr. 1996; NSMT-P 91546 (1), east of Oshika Peninsula, 38.3583°N, 141.9400°E, 280 m, 23 Oct. 2007; BSKU 45442 (1), off Kinkazan, 38.3217°N, 142.1000°E, 470 m, 24 May 1988; BSKU 116165 (1), off Rikuzen-Takata, 39.0500°N, 142.1557°E, 395–400 m, 23 Oct. 1995; BSKU 101974 (1), BSKU 101975 (1), BSKU 101976

(1), BSKU 101977 (1), BSKU 101978 (1), northeast of Kamaishi, 39.3800°N, 142.1700°E, 260–279 m, 27 Oct. 1995; FAKU 29134 (1), FAKU 29135 (1), FAKU 29136 (1), FAKU 29137 (1), FAKU 29138 (1), FAKU 29139 (1), FAKU 29140 (1), off Hachinohe, 250 m, Dec. 1959; FAKU 6062 (1), off Kushiro, 1938.

Coelorinchus matsubara. 6 specimens: BSKU 32227 (1), BSKU 32228 (1), BSKU 32247 (1), BSKU 32250 (1), collected with holotype; BSKU 30602 (1), Kyushu-Palau Ridge, 28.0667°N, 134.6833°E, 523 m, 18 Dec. 1978; FAKU KP82 (1), Kyushu-Palau Ridge, 28.0825°N, 134.7258°E, Nov. 1979.

Coelorinchus multispinulosus. 342 specimens: BSKU 27783 (1), BSKU 27784 (1), west of Amami-oshima Island, Okinawa Trough, 28.2000°N, 126.9167°E, 160 m, 12 Mar. 1978; SNFR 13258 (8), East China Sea, 32.2667°N, 125.7500°E, 78 m, 20 Sept. 1993; SNFR 13301 (5), East China Sea, 32.2300°N, 126.1983°E, 100 m, 20 Sept. 1993; SNFR 10423 (1), East China Sea, 29.2467°N, 126.2600°E, 100–103 m, 11 Oct. 1990; KAUM-I. 40451 (1), KAUM-I. 40452 (1), KAUM-I. 40453 (1), KAUM-I. 40454 (1), KAUM-I. 40455 (1), East China Sea, 29.7555°N, 126.4048°E, 110 m, 10 Dec. 2012; SNFR 11346 (2), East China Sea, 30.4833°N, 126.4083°E, 81 m, 2 Nov. 1986; SNFR 14282 (4), East China Sea, 30.4415°N, 126.4257°E, 82 m, 19 Jun. 2008; SNFR 12956 (3), East China Sea, 30.2150°N, 126.4267°E, 97 m, 30 Oct. 1985; SNFR 12288 (10), East China Sea, 31.1400°N, 126.6507°E, 94–95 m, 20 Jun. 2008; KAUM-I. 35735 (1), KAUM-I. 35736 (1), KAUM-I. 35737 (1), KAUM-I. 35738 (1), East China Sea, 29.7555°N, 126.6667°E, 112 m, 9 Dec. 2010; BSKU 27771 (1), BSKU 27772 (1), BSKU 27773 (1), west of Amami-oshima Island, Okinawa Trough, 28.4167°N, 127.0000°E, 200 m, 12 Mar. 1978; KAUM-I. 40464 (1), KAUM-I. 40465 (1), KAUM-I. 40466 (1), East China Sea, 29.7555°N, 127.0535°E, 144 m, 11 Dec. 2010; BSKU uncat. (1), BSKU 22955 (1), BSKU 22956 (1), BSKU 22959 (1), BSKU 22960 (1), BSKU 22961 (1), BSKU 22962 (1), BSKU 22963 (1), BSKU 22964 (1), BSKU 22965 (1), BSKU 22966 (1), BSKU 22967 (1), BSKU 22968 (1), BSKU 22972 (1), southwest of Danjo Islands, Okinawa Trough, 31.6150°N, 127.1083°E, 110 m, 6 Aug. 1974; BSKU 20190 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.3333°N, 127.1117°E, 128 m, 5 Feb. 1972; KAUM-I. 20513 (1), KAUM-I. 20520 (1), East China Sea, 30.9467°N, 127.3188°E, 116 m, 7 Nov. 2008; KAUM-I. 7194 (1), East China Sea, 30.9467°N, 127.3443°E, 121 m, 8 Nov. 2007; SNFR 12153 (1), SNFR 12154 (1), East China Sea, 31.5948°N, 127.4170°E, 130–131 m, 13 Jun. 2008; BSKU 28667 (1), northwest of Amami-oshima Island, Okinawa Trough, 30.0833°N, 127.5333°E, 146–167 m, 21 Mar. 1978; SNFR 12202 (1), East China Sea, 31.4230°N, 127.5417°E, 129–130 m, 13 Jun. 2008; FRLM 22133 (1), FRLM 22153 (1), East China Sea, Jul. 1998; ZMUT 49892 (1), ZMUT 49893

(1), Fukue-jima Island, 10 Jun. 1953; KAUM-I. 358 (1), off Kasasa, Kagoshima, 31.4833°N, 130.0333°E, 145–150 m, 10 Jun. 2006; KAUM-I. 3577 (1), off Kasasa, Kagoshima, 31.5307°N, 130.0390°E, 140 m, 8 May 2007; KAUM-I. 41326 (1), KAUM-I. 41327 (1), KAUM-I. 41328 (1), KAUM-I. 41329 (1), KAUM-I. 41330 (1), KAUM-I. 41331 (1), KAUM-I. 41332 (1), KAUM-I. 41333 (1), KAUM-I. 41334 (1), KAUM-I. 41335 (1), KAUM-I. 41336 (1), KAUM-I. 41337 (1), KAUM-I. 41338 (1), KAUM-I. 41339 (1), KAUM-I. 41340 (1), 1 km southwest of Kawajiri fish market, 31.1667°N, 130.5333°E, 20 Jun. 2011; NSMT-P 69708 (1), east-southeast of Tanega-shima Island, 30.3417°N, 131.2910°E, 101 m, 22 May 2002; MUFS 13661 (1), Meitsu, Miyazaki, 16 Jul. 1997; MUFS 20723 (1), Meitsu, Miyazaki, 27 Oct. 2000; MUFS 13265 (1), Meitsu, Miyazaki, Apr. 1997; MUFS 16249 (1), Meitsu, Miyazaki, 1 Jul. 1998; MUFS 16095 (1), MUFS 16096 (1), MUFS 16097 (1), MUFS 16098 (1), MUFS 16099 (1), Meitsu, Miyazaki, 24 Jun. 1998; ZMUT 50678 (1), Totoro, Miyazaki, 300–400 m, 9 Sept. 1959; BSKU 69893 (1), Irino fish market, 1 Apr. 2004; BSKU 78587 (1), BSKU 78588 (1), BSKU 78589 (1), BSKU 78590 (1), BSKU 78591 (1), BSKU 78592 (1), BSKU 78593 (1), BSKU 78594 (1), BSKU 78595 (1), BSKU 78596 (1), BSKU 78598 (1), Irino fish market, 11 Jun. 2006; BSKU 75804 (1), Irino fish market, 17 Jun. 2004; BSKU 54612 (1), BSKU 54613 (1), BSKU 54614 (1), BSKU 54615 (1), Irino fish market, 20 Feb. 1997; BSKU 67353 (1), Irino fish market, 22 Jun. 2003; BSKU 51946 (1), Irino fish market, 25 Apr. 2000; BSKU 78499 (1), BSKU 78500 (1), BSKU 78501 (1), Irino fish market, 8 Aug. 2004; KAUM-I. 44132 (1), off Irino, Kochi, 15 Dec. 2011; BSKU 72666 (1), Saga fish market, 16 Jan. 2004; BSKU 99397 (1), BSKU 99399 (1), BSKU 99401 (1), BSKU 99402 (1), Saga fish market, 17 Jun. 2007; BSKU 56418 (1), BSKU 56448 (1), BSKU 56449 (1), Saga fish market, 21 Jun. 2001; BSKU 78847 (1), BSKU 78848 (1), BSKU 78849 (1), BSKU 78850 (1), BSKU 78851 (1), BSKU 78852 (1), BSKU 78853 (1), BSKU 78854 (1), BSKU 78855 (1), BSKU 78856 (1), BSKU 78857 (1), BSKU 78858 (1), BSKU 78859 (1), BSKU 78860 (1), BSKU 78861 (1), BSKU 78862 (1), BSKU 78864 (1), Saga fish market, 22 Jun. 2006; BSKU 88116 (1), Saga fish market, 24 Aug. 2006; BSKU 77444 (1), BSKU 77552 (1), BSKU 77553 (1), BSKU 79580 (1), BSKU 79581 (1), Saga fish market, 24 Nov. 2005; BSKU 90885 (1), Saga fish market, 27 May 2007; BSKU 78904 (1), BSKU 78905 (1), BSKU 78906 (1), BSKU 78907 (1), BSKU 78908 (1), BSKU 78909 (1), BSKU 78910 (1), BSKU 78911 (1), BSKU 78912 (1), BSKU 78913 (1), BSKU 78914 (1), BSKU 78916 (1), BSKU 78917 (1), BSKU 78918 (1), BSKU 78919 (1), BSKU 78920 (1), BSKU 78921 (1), BSKU 78922 (1), BSKU 78923 (1), BSKU 78924 (1), BSKU 78925 (1), BSKU 78926 (1), BSKU 78927 (1), BSKU 78928 (1), BSKU 78929 (1), BSKU 78930 (1), BSKU 78931 (1), BSKU 78932 (1), BSKU 78933 (1), BSKU 78934 (1), BSKU 78935 (1), BSKU 78936 (1), BSKU 78937 (1), BSKU 78938 (1), BSKU 78939 (1), BSKU 78940 (1), BSKU 78941 (1), BSKU 78942 (1), BSKU 78943 (1), BSKU 78944 (1), BSKU 78945 (1), BSKU 78946 (1), BSKU 78947 (1), BSKU 78948 (1), BSKU 78949 (1), BSKU 78950 (1), BSKU 78951 (1), BSKU 78952 (1), BSKU 78953 (1), BSKU 78954 (1), BSKU 78955 (1), BSKU 78956 (1), BSKU 78957 (1), BSKU 78958 (1), BSKU 78959 (1), BSKU 78960 (1), BSKU 78961 (1), BSKU 78962 (1), BSKU 78963 (1), Saga fish market, 29 Jun. 2006; BSKU 76096 (1), Saga fish market, 4 Jul. 2005; BSKU 99395 (1), BSKU 99396 (1), BSKU 99400 (1), Saga fish market, 8 Jun. 2007; BSKU 76131 (1), BSKU 76132 (1), BSKU 76133 (1), BSKU 76134 (1), BSKU 76135 (1), BSKU 76136 (1), BSKU 76137 (1), BSKU 76138 (1), BSKU 76139 (1), BSKU 76140 (1), BSKU 76141 (1), Saga fish market, 9 Mar. 2005; BSKU 78490 (1), BSKU 78491 (1), BSKU 78492 (1), BSKU 78493 (1), Mimase fish market, 1 Dec. 2005; BSKU 99363 (1), BSKU 99366 (1), BSKU 99367 (1), BSKU 99368 (1), BSKU 99369 (1), Mimase fish market, 11 Jan. 2007; BSKU 36192 (1), Mimase fish market, 12 Jan. 1982; BSKU 77968 (1), Mimase fish market, 13 Apr. 2006; BSKU 53037 (1), Mimase fish market, 13 Dec. 2000; BSKU 90929 (1), BSKU 90930 (1), BSKU 90931 (1), BSKU 90932 (1), Mimase fish market, 14 Mar. 2007; BSKU 69386 (1), Mimase fish market, 14 Oct. 2003; BSKU 78099 (1), Mimase fish market, 24 Apr. 2006; BSKU 35612 (1), Mimase fish market, 3 Mar. 1981; BSKU 69296 (1), Mimase fish market, 3 Nov. 2003; BSKU 36151 (1), Mimase fish market, 5 Dec. 1981; BSKU 30169 (1), Mimase fish market, 8 Dec. 1979; BSKU 78489 (1), Mimase fish market, 8 Dec. 2005; BSKU 70973 (1), Mimase fish market, 9 Nov. 2003; BSKU 11714 (1), Mimase fish market, Nov. 1965; BSKU 919 (1), BSKU 9472 (1), Mimase fish market, 16 Jan. 1951; BSKU 9678 (1), BSKU 9681 (1), Mimase fish market, 24 Dec. 1950; BSKU 296 (1), BSKU 297 (1), Mimase fish market, 24 Oct. 1949; BSKU 6738 (1), BSKU 6739 (1), BSKU 6749 (1), Mimase fish market, Dec. 1955–Jan. 1956; BSKU 13638 (1), Mimase fish market, Jan.–Mar. 1968; ZMUT 30616 (1), ZMUT 30617 (1), Mimase fish market, 6 Jan. 1935; BSKU 94827 (1), Mimase fish market, 9 Dec. 2007; BSKU 52420 (1), Kokadai fish market, 16 Apr. 1994; BSKU 78521 (1), Tosa Bay, 120 m, 6 Jun. 2006; BSKU 79478 (1), BSKU 79481 (1), Tosa Bay, 120 m, 4 Aug. 2006; ZMUT 23090 (1), Yuasa-machi, Wakayama, 23 Oct. 1930; BSKU 39310 (1), BSKU 39311 (1), BSKU 39312 (1), BSKU 39313 (1), BSKU 39315 (1), BSKU 39316 (1), Owase fish market, Kumano-nada, 12 Mar. 1983; FAKU 27725 (1), FAKU 27736 (1), FAKU 27737 (1), FAKU 27748 (1), FAKU 27749 (1), FAKU 27751 (1), Owase fish market, Kumano-nada, 25 Feb.–10 Mar. 1957; FRLM 32912 (1), off Owase, Kumano-nada, 15 May 2007; FRLM 32942 (1), FRLM 32943 (1), FRLM 32944

(1), FRLM 32945 (1), FRLM 32946 (1), FRLM 32947 (1), FRLM 32948 (1), off Mie, Kumano-nada, 15 May 2007; NSMT-P 103585 (1), NSMT-P 103586 (1), Isshiki fish market, 25 Aug. 2010; FMNH 57216 (1, ex. CM 6155a), Misaki, Sagami Bay, date unknown; ZMUT 5440 (1), Tokyo, date unknown; NSMT-P 73336 (1, ex. SHMNH-P 26727), southeast off Onahama, 1931–1935; FAKU 57536 (1), east of Tsushima Island, 34.5200°N, 129.7433°E, 104 m, 22 Jul. 1990; SNFR 11781 (1), Sea of Japan, 34.7217°N, 130.1350°E, 15 Apr. 1982; ZMUT 51407 (1), East China Sea or Fukuoka fish market, Oct. 1959; ZMUT 51009 (1), Fukuoka fish market, 14 Mar. 1960; FAKU 27000 (1), FAKU 27001 (1), FAKU 27002 (1), Shimonoseki fish market, Sea of Japan, 24 Jan. 1957; FAKU 58494 (1), northwest of Hamada, Shimane, Sea of Japan, 35.3050°N, 131.3017°E, 123 m, 8 Jul. 1990; BSKU 40497 (1), BSKU 40547 (1), Hamada fish market, Sea of Japan, 26 Nov. 1981; BSKU 40178 (1), Kazue fish market, Sea of Japan, 28 Dec. 1984; BSKU 99420 (1), BSKU 99421 (1), BSKU 99422 (1), BSKU 99423 (1), BSKU 99424 (1), BSKU 99425 (1), BSKU 99426 (1), BSKU 99427 (1), BSKU 99428 (1), BSKU 99429 (1), BSKU 99430 (1), BSKU 99431 (1), BSKU 99432 (1), BSKU 99433 (1), BSKU 99434 (1), BSKU 99435 (1), BSKU 99436 (1), BSKU 99437 (1), BSKU 99438 (1), BSKU 99439 (1), BSKU 99440 (1), BSKU 99441 (1), BSKU 99442 (1), BSKU 99443 (1), BSKU 99444 (1), BSKU 99445 (1), BSKU 99446 (1), BSKU 99447 (1), Kazue fish market, Sea of Japan, 29 Nov. 1984; FAKU 131454 (1), off Takeno, Sea of Japan, date unknown; ZMUT 24979 (1), Horie, Ehime, Nov. 1931.

Coelorinchus parallelus. 21 specimens: BSKU 26556 (1), off Miyako-jima Island, Okinawa Trough, 26.4633°N, 124.8483°E, 680–770 m, 23 Jan. 1978; BSKU 27114 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.3833°N, 127.5150°E, 650 m, 5 Feb. 1978; BSKU 27026 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.9267°N, 127.7483°E, 360 m, 4 Feb. 1978; BSKU 26951 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.6483°N, 127.9333°E, 750–760 m, 4 Feb. 1978; BSKU 26988 (1), BSKU 26990 (1), BSKU 26992 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.6467°N, 127.9733°E, 830 m, 4 Feb. 1978; BSKU 28438 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7667°N, 127.9833°E, 713–770 m, 20 Mar. 1978; BSKU 28586 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.9500°N, 128.0000°E, 585–610 m, 20 Mar. 1978; BSKU 28506 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7333°N, 128.0500°E, 815 m, 20 Mar. 1978; BSKU 28547 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.8167°N, 128.1000°E, 915–932 m, 20 Mar. 1978; BSKU 33465 (1), BSKU 33469 (1), BSKU 33473 (1), BSKU 33475 (1), BSKU 33477 (1), BSKU 33479 (1), west of Yaku-shima Island, Okinawa Trough, 30.2067°N, 128.2833°E, 780–810 m, 26 Oct. 1979; BSKU 26864

(1), north-northwest of Amami-oshima Island, Okinawa Trough, 29.7850°N, 128.4467°E, 1000 m, 3 Feb. 1978; FFNU P-00573 (2), southwest of Danjo Basin, East China Sea, 31.6793°N, 128.9423°E, 733 m, 24 Apr. 2005; FMNH 89209 (1), Misaki, Sagami Bay, Jan. 1904.

Coelorinchus smithi. 55 specimens: BSKU 32513 (1), BSKU 32514 (1), northwest of Miyako-jima Island, Okinawa Trough, 25.7917°N, 124.5117°E, 500–550 m, 10 Oct. 1979; BSKU 27474 (1), west of Okinawa-jima Island, Okinawa Trough, 27.7767°N, 126.2833°E, 490 m, 9 Mar. 1978; KPM-NI 28290 (1), KPM-NI 28291 (1), KPM-NI 28292 (1), KPM-NI 28293 (1), KPM-NI 28294 (1), ca. 22 km south-southeast of Aguni-jima Island, East China Sea, 26.4157°N, 127.3410°E, 734–751 m, 17 Dec. 2008; KPM-NI 28331 (1), KPM-NI 28332 (1), ca. 25 km southeast of Aguni-jima Island, East China Sea, 26.4120°N, 127.3947°E, 774–780 m, 17 Dec. 2008; BSKU 27867 (1), BSKU 27869 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.5833°N, 127.0000°E, 600 m, 16 Mar. 1978; BSKU 27672 (1), BSKU 27674 (1), west of Okinawa-jima Island, Okinawa Trough, 28.7333°N, 127.0167°E, 610–640 m, 11 Mar. 1978; SNFR 11219 (3), East China Sea, 28.5233°N, 127.0250°E, 550 m, 23 Oct. 1990; BSKU 27987 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.8333°N, 127.2333°E, 700–740 m, 16 Mar. 1978; BSKU 28200 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.3000°N, 127.3333°E, 410–415 m, 17 Mar. 1978; SNFR 14231 (1), East China Sea, 30.2077°N, 127.7222°E, 218–236 m, 7 Aug. 2008; FFNU P-00749 (1), north of Minamiensei Knoll, East China Sea, 29.6828°N, 127.7400°E, 596 m, 30 Nov. 2008; BSKU 28363 (1), BSKU 28364 (1), BSKU 28365 (1), BSKU 28366 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7333°N, 127.8000°E, 495–510 m, 19 Mar. 1978; BSKU 28437 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7667°N, 127.9833°E, 713–770 m, 20 Mar. 1978; SNFR 11351 (2), East China Sea, 31.2717°N, 128.4350°E, 479 m, 30 Sept. 1995; FFNU P-00639 (2), east of Okirirazi Bank, East China Sea, 32.1753°N, 129.4948°E, 475 m, 1 Aug. 2006; BSKU 30493 (1), Tosa Bay, 32.9667°N, 133.5333°E, 605 m, 21 Dec. 1979; BSKU 10042 (1), BSKU 10043 (1), BSKU 10044 (1), Mimase fish market, 10 Feb. 1961; BSKU 4381 (1), Mimase fish market, 19 Dec. 1954; BSKU 4413 (1), Mimase fish market, 26 Dec. 1954; BSKU 12819 (1), Mimase fish market, date unknown; BSKU 69819 (1), Tosa Bay, 400 m, 9 May 2000; BSKU 77918 (1), BSKU 97992 (1), south of Muroto, 33.0790°N, 134.1645°E, 414–513 m, 17 Nov. 2005; FAKU 34514 (1), FAKU 34516 (1), Owase fish market, Kumano-nada, 12–14 Nov. 1961; FAKU 27651 (1), Owase fish market, Kumano-nada, 25 Feb.–10 Mar. 1957; FAKU 24932 (1), FAKU 24932 (1), FAKU 24933 (1), FAKU 24933 (1), FAKU 24934 (1), FAKU 24934 (1), Owase fish market, Kumano-nada, 25 Oct. 1954; NSMT-P 102069 (1), southwest of Cape Daiozaki, Kumano-nada, 34.0983°N,

136.6800°E, 422–425 m, 27 May 1986; FAKU 17713 (1), Miya fish market, Enshu-nada, 15 Mar. 1952; FAKU 28392 (1), Miya fish market, Enshu-nada, 5–7 Feb. 1959; BSKU 110417 (1), Suruga Bay, 34.7619°N, 138.4811°E, 196–387 m, 23 Apr. 2013; MSM 13-17 (1), Suruga Bay, 34.5937°N, 138.7282°E, 420–600 m, 25 Apr. 2012; ZMUT 6846 (1), Tokyo, date unknown.

Coelorrinchus tokiensis. 3 specimens: BSKU 106814 (1), southeast of Shimokoshiki-jima Island, East China Sea, 31.5652°N, 129.8878°E, 407 m, 24 Apr. 2012; SNFR 18056 (1), Tosa Bay, 19 Aug. 1987; BSKU 19189 (1), south of Miura Peninsula, Sagami Bay, 35.1250°N, 139.5100°E, 700 m, 10 Mar. 1961.

Coryphaenoides acrolepis. 157 specimens: KAUM-I. 30356 (1), KAUM-I. 30357 (1), northeast of Senoumi, Suruga Bay, 34.7913°N, 138.5893°E, 1500–1700 m, 15 Apr. 2010; NSMT-P 102197 (1), NSMT-P 102198 (1), east of Kozu-shima Island, Shichito-Iojima Ridge, 34.2500°N, 138.7583°E, 1934–2070 m, 9 Dec. 1987; KPM-NI 13876 (1), off Minami-izu, Izu Peninsula, Dec. 2002; BSKU 14189 (1), BSKU 14190 (1), BSKU 14191 (1), BSKU 14193 (1), off Odawara, Sagami Bay, 35.2014°N, 139.2003°E, 670–750 m, 14 Apr. 1969; KPM-NI 17446 (1), off Hatsu-shima Island, Sagami Bay, 900 m, 30 Mar. 2003; NSMT-P 102199 (3), northeast of Miyake-jima Island, Sagami-nada, 34.8683°N, 139.4617°E, 1656–1708 m, 12 Dec. 1978; FAKU M9555 (1), off Kominato, 15 May 1944; NSMT-P 102193 (1), NSMT-P 102194 (2), southeast of Boso Peninsula, 34.9862°N, 140.8350°E, 1361–1390 m, 15 Sept. 1989; NSMT-P 97424 (2), east of Boso Peninsula, 35.1123°N, 140.9150°E, 1351–1454 m, 24 Apr. 1995; NSMT-P 61287 (1), off Iwaki, 36.9400°N, 141.7000°E, 967–1219 m, Nov. 1995; HUMZ 72633 (1), off Fukushima, 37.4417°N, 142.1583°E, 900 m, 20 Jan. 1979; HUMZ 72748 (1), HUMZ 72749 (1), off Fukushima, 37.5750°N, 142.2950°E, 810–980 m, 21 Jan. 1978; NSMT-P 105669 (1), NSMT-P 105670 (1), off Minami-soma, 37.6000°N, 142.3950°E, 900 m, 21 Jul. 2008; HUMZ 72658 (1), off Miyagi, 38.0333°N, 142.4833°E, 1100–1150 m, 7 Feb. 1978; NSMT-P 19005 (2), off Minami-soma, 37.5083°N, 142.5417°E, 1350 m, 22 Jun. 1978; HUMZ 72676 (1), HUMZ 72677 (1), HUMZ 72678 (1), off Miyagi, 38.7250°N, 142.3917°E, 1000–1100 m, 4 Feb. 1978; NSMT-P 115470 (1), off Kamaishi, 39.1050°N, 142.3967°E, 926–950 m, 10 Jul. 1984; NSMT-P 97661 (4), off Kamaishi, 39.1300°N, 142.3967°E, 984–988 m, 25 Sept. 1988; NSMT-P 97654 (1), off Kamaishi, 39.1663°N, 142.3983°E, 975–988 m, 27 May 1989; HUMZ 113010 (1), east of Misawa, Aomori, 40.7717°N, 142.3983°E, 1125 m, 7 Mar. 1988; HUMZ 78253 (1), off Aomori, 40.7867°N, 142.4917°E, 1255 m, 10 Sept. 1978; HUMZ 78167 (1), off Iwate, 39.4517°N, 142.5550°E, 1100 m, 22 Sept. 1978; HUMZ 78207 (1), off Iwate, 40.2850°N, 142.5767°E, 1140 m, 17 Sept. 1978; NSMT-P 102182 (25), NSMT-P 102195 (1), off Kamaishi, 39.0150°N, 142.6167°E, 1237–1255 m, 21

May 1987; NSMT-P 97662 (3), off Kamaishi, 39.0627°N, 142.6250°E, 1203–1213 m, 24 May 1988; BSKU 97744 (1), BSKU 97745 (1), BSKU 97746 (1), BSKU 97747 (1), off Kuji, 40.0613°N, 142.6825°E, 1005–1016 m, 18 Nov. 1993; NSMT-P 90625 (1), off Miyako, 1700 m, 8–9 Nov. 2007; NSMT-P 78031 (1), NSMT-P 78032 (1), off Miyako, 870 m, 6 Aug. 2007; HUMZ 78177 (1), off Iwate, 39.0917°N, 142.7383°E, 1300 m, 20 Sept. 1978; NSMT-P 114949 (6), southeast of Otsuchi, 1225–1249 m, 12 Aug. 1985; HUMZ 78122 (1), off Iwate, 39.1217°N, 142.7750°E, 1480–1485 m, 20 Sept. 1978; NSMT-P 97153 (1), off Kamaishi, 39.0150°N, 142.8117°E, 1237–1255 m, 21 May 1987; BSKU 96420 (1), BSKU 96422 (1), BSKU 96423 (1), BSKU 96424 (1), BSKU 96426 (1), BSKU 96427 (1), BSKU 96428 (1), BSKU 96429 (1), BSKU 96430 (1), BSKU 96431 (1), off Miyako, 39.5668°N, 142.9008°E, 1586–1617 m, 21 Oct. 2008; NSMT-P 102402 (5), off Kesennuma, 38.9550°N, 142.9483°E, 1430–1440 m, 11 Jul. 1984; NSMT-P 102058 (1), NSMT-P 102059 (1), NSMT-P 102060 (1), NSMT-P 102061 (1), NSMT-P 102062 (1), NSMT-P 102063 (6), off Kesennuma, 38.7000°N, 143.0267°E, 1668–1685 m, 12 Aug. 1985; BSKU 35677 (1), BSKU 35678 (1), BSKU 35679 (1), BSKU 35680 (1), BSKU 35686 (1), BSKU 35687 (1), off Kinkazan, 38.7333°N, 143.1850°E, 1950–1960 m, 25 Jul. 1981; BSKU 35673 (1), off Kuji, 40.3683°N, 143.3800°E, 1970–2000 m, 10 Jul. 1981; BSKU 97739 (1), BSKU 97740 (1), BSKU 97778 (1), southeast of Cape Erimomisaki, Japan Trench, 41.2353°N, 144.0357°E, 2977–3014 m, 17 Nov. 1993; HUMZ 34255 (1), off Cape Erimomisaki, Hokkaido, 700–780 m, 4 Nov. 1973; BSKU 114845 (1), east-southeast of Cape Erimomisaki, 41.8002°N, 143.8150°E, 648 m, 6 Jun. 2014; BSKU 114798 (1), BSKU 114799 (1), BSKU 114800 (1), BSKU 114801 (1), BSKU 114802 (1), BSKU 114803 (1), BSKU 114804 (1), BSKU 114805 (1), southeast of Tokachi, 42.3320°N, 143.9187°E, 733 m, 10 Jun. 2014; BSKU 114817 (1), southeast of Tokachi, 42.3317°N, 143.9585°E, 719 m, 10 Jun. 2014; NSMT-P 97663 (9), off Tokachi, 42.2617°N, 144.0873°E, 992–1042 m, 14 Jun. 1990; HUMZ 41810 (1), Kushiro, Hokkaido, 620 m, 6 Jul. 1974; HUMZ 6923 (1), Kushiro, Hokkaido, date unknown; BSKU 114808 (1), BSKU 114814 (1), BSKU 114860 (1), south of Kushiro, 42.6542°N, 144.3770°E, 780 m, 7 Jun. 2014; BSKU 114858 (1), south of Kushiro, 42.6320°N, 144.3803°E, 877 m, 8 Jun. 2014; BSKU 97727 (1), BSKU 97730 (1), BSKU 97731 (1), BSKU 97732 (1), south-southeast of Akkeshi, 42.6052°N, 144.9938°E, 1113–1161 m, 16 Nov. 1993; HUMZ 79709 (1), Kitami-Yamato Bank, Okhotsk Sea, 44.4583°N, 144.5000°E, 1300–1340 m, 24 Sept. 1978; HUMZ 124353 (1), HUMZ 124354 (1), Kitami-Yamato Bank, Okhotsk Sea, 2 Jun. 1992; HUMZ 124286 (1), HUMZ 124287 (1), Kitami-Yamato Bank, Okhotsk Sea, 44.5700°N, 144.9983°E, 1300 m, 4 Jul. 1992.

Coryphaenoides cinereus. 289 specimens: BSKU

43472 (1), off Miyazaki, Hyuga-nada, 32.0650°N, 132.2950°E, 1463–1514 m, 3 Nov. 1986; MSM 13-12 (1), east of Shimizu, Suruga Bay, 35.0317°N, 138.6217°E, 1202–1402 m, 4 Jul. 2012; NSMT-P 93122 (1), off Oh-sima Island, Izu Islands, date unknown; NSMT-P 97217 (11), north of Miyake-jima Island, Sagami-nada, 34.9250°N, 139.3733°E, 1541–1630 m, 20 May 1999; NSMT-P 101929 (3), northeast of Miyake-jima Island, Sagami-nada, 34.8783°N, 139.4583°E, 1525–1685 m, 17 Dec. 1982; NSMT-P 94391 (1), Kashima-nada, 975–1020 m, 8 Aug. 1979; NSMT-P 94748 (1), off Kashima, 19 May 1987; NSMT-P 78115 (1), NSMT-P 78116 (1), NSMT-P 78117 (2), southeast off Onahama, 2000 m, 8 Aug. 2007; NSMT-P 18998 (1), NSMT-P 19001 (1), NSMT-P 19002 (1), off Ibaraki, 36.3083°N, 141.3600°E, 1100 m, 16 Mar. 1975; BSKU 97919 (1), BSKU 97920 (1), off Hitachi, 36.4205°N, 141.5420°E, 1845–1948 m, 10 Nov. 1995; BSKU 97822 (1), BSKU 97823 (1), BSKU 97824 (1), BSKU 97825 (1), BSKU 97826 (1), BSKU 97827 (1), BSKU 97828 (1), BSKU 97829 (1), BSKU 97830 (1), BSKU 97831 (1), BSKU 97832 (1), BSKU 97834 (1), BSKU 97835 (1), BSKU 97836 (1), BSKU 97837 (1), BSKU 97838 (1), BSKU 97839 (1), BSKU 97840 (1), off Kamaishi, 39.1444°N, 142.3748°E, 847–935 m, 10 Mar. 1989; NSMT-P 97659 (18), off Kamaishi, 39.1663°N, 142.3983°E, 975–988 m, 27 May 1989; NSMT-P 101603 (66), off Kamaishi, 39.1131°N, 142.4127°E, 1038–1055 m, 20 May 1987; NSMT-P 78017 (1), off Kinkazan, 2000 m, 6 Aug. 2007; BSKU 97849 (1), BSKU 97850 (1), BSKU 97851 (1), BSKU 97853 (1), BSKU 97854 (1), BSKU 97855 (1), BSKU 97856 (1), BSKU 97857 (1), BSKU 97858 (1), BSKU 97859 (1), BSKU 97860 (1), BSKU 97862 (1), BSKU 97863 (1), BSKU 97864 (1), BSKU 97865 (1), BSKU 97866 (1), BSKU 97867 (1), off Rikuzen-Takata, 39.0538°N, 142.6342°E, 1195–1202 m, 10 Mar. 1989; BSKU 97749 (1), off Kuji, 40.0613°N, 142.6825°E, 1005–1016 m, 18 Nov. 1993; BSKU 97725 (1), BSKU 97726 (1), BSKU 97769 (1), BSKU 97770 (1), BSKU 97781 (1), BSKU 97782 (1), off Miyako, 39.5747°N, 142.6852°E, 1132–1170 m, 13 Nov. 1993; BSKU 97916 (1), northeast of Miyako, 39.9633°N, 142.6987°E, 993–1007 m, 26 Oct. 1995; NSMT-P 97656 (2), off Otsuchi, 1997–2108 m, 31 May 1989; NSMT-P 94367 (1), southeast of Otsuchi, 1225–1249 m, 12 Aug. 1985; BSKU 14179 (1), BSKU 14180 (1), BSKU 14181 (1), BSKU 14182 (1), BSKU 14183 (1), off Kesennuma, 38.8506°N, 142.7675°E, 1270–1280 m, 12 Aug. 1967; NSMT-P 79349 (1), NSMT-P 79353 (1), off Kesennuma, 38.5367°N, 142.8000°E, 1502 m, 26 Oct. 2007; NSMT-P 101743 (1), off Kamaishi, 39.3971°N, 142.8647°E, 1650–1687 m, 8 Sept. 1996; BSKU 96499 (1), BSKU 96500 (1), BSKU 96501 (1), off Kinkazan, 38.4640°N, 143.0518°E, 1853–1906 m, 23 Oct. 2008; BSKU 35698 (1), BSKU 35699 (1), BSKU 35700 (1), BSKU 35701 (1), BSKU 35702 (1), BSKU 35703 (1), BSKU 35705 (1), BSKU 35707 (1), BSKU 35709 (1), BSKU 35711 (1), BSKU 35712 (1), BSKU 35717 (1), off Kesennuma, 38.7333°N, 143.1850°E, 1950–1960 m, 25 Jul. 1981; BSKU 97751 (1), BSKU 97752 (1), BSKU 97753 (1), BSKU 97754 (1), BSKU 97755 (1), BSKU 97756 (1), BSKU 97758 (1), BSKU 97759 (1), BSKU 97760 (1), BSKU 97761 (1), BSKU 97762 (1), BSKU 97764 (1), BSKU 97766 (1), BSKU 97767 (1), BSKU 97768 (1), south of Cape Erimomisaki, 41.0675°N, 143.3268°E, 2022–2027 m, 20 Oct. 2008; BSKU 97921 (1), off Hachinohe, 40.6018°N, 143.3585°E, 2014–2056 m, 10 Nov. 1995; BSKU 97741 (1), BSKU 97742 (1), BSKU 97779 (1), BSKU 97790 (1), off Kuji, 40.3727°N, 143.3827°E, 1923–1995 m, 18 Nov. 1993; BSKU 97735 (1), BSKU 97736 (1), BSKU 97737 (1), BSKU 97777 (1), southeast of Cape Erimomisaki, Japan Trench, 41.2353°N, 144.0357°E, 2977–3014 m, 17 Nov. 1993; HUMZ 90927 (1), Muroran, 560–580 m, 28 Apr. 1981; HUMZ 206293 (1), south of Tomakomai, Hokkaido, 42.2132°N, 141.6842°E, 800 m, 29 Sept. 2009; BSKU 96396 (1), BSKU 96397 (1), BSKU 96398 (1), BSKU 96399 (1), BSKU 96400 (1), BSKU 96402 (1), BSKU 96403 (1), BSKU 96404 (1), south of Cape Erimomisaki, 41.6675°N, 143.3268°E, 2022–2027 m, 20 Oct. 2008; BSKU 114912 (1), BSKU 114913 (1), east-southeast of Cape Erimomisaki, 41.8170°N, 143.7875°E, 576 m, 6 Jun. 2014; BSKU 114828 (1), BSKU 114829 (1), BSKU 114830 (1), BSKU 114831 (1), BSKU 114832 (1), BSKU 114833 (1), BSKU 114834 (1), BSKU 114835 (1), BSKU 114836 (1), BSKU 114837 (1), BSKU 114838 (1), BSKU 114839 (1), BSKU 114840 (1), BSKU 114841 (1), BSKU 114842 (1), BSKU 114843 (1), BSKU 114844 (1), east-southeast of Cape Erimomisaki, 41.8002°N, 143.8150°E, 648 m, 6 Jun. 2014; BSKU 114816 (1), BSKU 114824 (1), BSKU 114825 (1), BSKU 114826 (1), BSKU 114827 (1), southeast of Tokachi, 42.3317°N, 143.9585°E, 719 m, 10 Jun. 2014; HUMZ 34190 (1), off Kushiro, 450 m, 16 Jul. 1974; BSKU 115092 (1), south of Kushiro, 42.7440°N, 144.3763°E, 455 m, 8 Jun. 2014; BSKU 114810 (1), BSKU 114815 (1), south of Kushiro, 42.6542°N, 144.3770°E, 780 m, 7 Jun. 2014; BSKU 114806 (1), south of Kushiro, 42.6320°N, 144.3803°E, 877 m, 8 Jun. 2014; BSKU 115072 (1), south-southeast of Kushiro, 42.6615°N, 144.5527°E, 487 m, 7 Jun. 2014; HUMZ 34236 (1), HUMZ 34251 (1), HUMZ 34275 (1), Akkeshi, 10 Apr. 1974; BSKU 97772 (1), BSKU 97775 (1), BSKU 97787 (1), south-southeast of Akkeshi, 42.6052°N, 144.9938°E, 1113–1161 m, 16 Nov. 1993; BSKU 97734 (1), south-southeast of Akkeshi, 42.4608°N, 145.0665°E, 1938–1969 m, 16 Nov. 1993; BSKU 58780 (1), BSKU 58781 (1), BSKU 58782 (1), BSKU 58783 (1), BSKU 58784 (1), BSKU 58785 (1), BSKU 58786 (1), BSKU 58787 (1), BSKU 58788 (1), BSKU 58789 (1), BSKU 58790 (1), BSKU 58791 (1), BSKU 58792 (1), BSKU 58793 (1), BSKU 58794 (1), BSKU 58795 (1), BSKU 58796 (1), BSKU 58797 (1), BSKU 58798 (1), BSKU 58799 (1), BSKU 58800 (1), BSKU 58801 (1), BSKU 58802 (1), BSKU 58803 (1), BSKU 58804

(1), off Kuril Islands, 43.0630°N, 147.1152°E, 451–661 m, 21 Jun. 1996; HUMZ 49041 (1), off Okhotsk coast of Hokkaido, 45.8000°N, 143.8167°E, 350–605 m, 10 Oct. 1975; NSMT-P 90681 (1), Kitami-Yamato Bank, Okhotsk Sea, 2 Jun. 2008; HUMZ 44967 (1), Kitami-Yamato Bank, Okhotsk Sea, 44.4167°N, 144.2917°E, 385–400 m, 3 Jul. 1975; NSMT-P 78140 (3), NSMT-P 78141 (5), northwest of Shiretoko Peninsula, Okhotsk Sea, 44.3433°N, 144.9350°E, 1012–1033 m, 13 Sept. 2004.

***Coryphaenoides longifilis*.** 43 specimens: BSKU 43478 (1), BSKU 43479 (1), BSKU 43480 (1), BSKU 43482 (1), off Miyazaki, Hyuga-nada, 32.0650°N, 132.2950°E, 1463–1514 m, 3 Nov. 1986; BSKU 23050 (1), Tosa Bay, 33.0350°N, 133.7500°E, 945 m, 26 Jan. 1975; FMNH 76687 (1), Izu, Sagami Bay, 12 Feb. 1907; NSMT-P 78114 (1), southeast off Onahama, 2000 m, 8 Aug. 2007; BSKU 14195 (1), off Hitachi, 36.3189°N, 141.6667°E, 1690 m, 3 Dec. 1967; HUMZ 111934 (1), northeast of Hachinohe, Aomori, 40.9383°N, 142.1067°E, 950 m, 6 Mar. 1988; BSKU 97797 (1), off Kesennuma, 38.9300°N, 142.3763°E, 976–1004 m, 25 Oct. 1995; NSMT-P 102405 (1), NSMT-P 102406 (1), NSMT-P 102407 (1), NSMT-P 102408 (1), off Kamaishi, 39.1050°N, 142.3967°E, 926–950 m, 10 Jul. 1984; HUMZ 111949 (1), east of Misawa, Aomori, 40.7717°N, 142.3983°E, 1125 m, 7 Mar. 1988; BSKU 45554 (1), BSKU 45555 (1), off Rikuzen-Takata, 39.0538°N, 142.6342°E, 1195–1202 m, 10 Mar. 1989; NSMT-P 102183 (1), NSMT-P 102184 (1), NSMT-P 102185 (3), NSMT-P 102186 (1), NSMT-P 102187 (6), off Kesennuma, 39.0017°N, 142.6383°E, 1246–1270 m, 12 Aug. 1985; BSKU 97812 (1), BSKU 97813 (1), BSKU 97814 (1), off Kuji, 40.0613°N, 142.6825°E, 1005–1016 m, 18 Nov. 1993; NSMT-P 90644 (1), off Miyako, 1000 m, 5 Nov. 2007; HUMZ 78120 (1), off Iwate, 39.1217°N, 142.7750°E, 1480–1485 m, 20 Sept. 1978; BSKU 96432 (1), off Miyako, 39.5668°N, 142.9008°E, 1586–1617 m, 21 Oct. 2008; NSMT-P 102403 (2), off Kesennuma, 38.9550°N, 142.9483°E, 1430–1440 m, 11 Jul. 1984; BSKU 35691 (1), BSKU 35693 (1), BSKU 35695 (1), BSKU 35696 (1), BSKU 35697 (1), off Kinkazan, 38.7333°N, 143.1850°E, 1950–1960 m, 25 Jul. 1981; BSKU 19470 (1), off Hachijo-jima Island, Shichito-Iojima Ridge, 32.5867°N, 140.1333°E, 1560–1600 m, 12 Dec. 1963.

***Coryphaenoides marginatus*.** 255 specimens: BSKU 27850 (1), BSKU 27851 (1), BSKU 27852 (1), BSKU 27853 (1), BSKU 27854 (1), BSKU 27855 (1), BSKU 27856 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.5833°N, 127.0000°E, 600 m, 16 Mar. 1978; BSKU 27667 (1), BSKU 27668 (1), BSKU 27669 (1), BSKU 27670 (1), BSKU 27671 (1), west of Okinawa-jima Island, Okinawa Trough, 28.7333°N, 127.0167°E, 610–640 m, 11 Mar. 1978; BSKU 109209 (1), west of Amami-oshima Island, Okinawa Trough, 28.5679°N, 127.0421°E, 607 m, 16 Nov. 2012; BSKU 27920 (1), BSKU 27921 (1), BSKU 27922 (1), BSKU 27923 (1), BSKU 27924 (1),

BSKU 27926 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.7500°N, 127.1167°E, 520–542 m, 16 Mar. 1978; BSKU 27702 (1), BSKU 27703 (1), BSKU 27704 (1), BSKU 27705 (1), BSKU 27706 (1), BSKU 27707 (1), west of Okinawa-jima Island, Okinawa Trough, 28.7000°N, 127.1500°E, 535–550 m, 11 Mar. 1978; FFNU P-00810 (5), west of Minamiensei Knoll, East China Sea, 28.4825°N, 127.1793°E, 1024 m, 18 Jun. 2009; BSKU 27568 (1), BSKU 27569 (1), BSKU 27570 (1), BSKU 27571 (1), BSKU 27572 (1), BSKU 27573 (1), BSKU 27574 (1), BSKU 27575 (1), west of Okinawa-jima Island, Okinawa Trough, 28.7167°N, 127.2333°E, 750–755 m, 11 Mar. 1978; BSKU 27991 (1), BSKU 27992 (1), BSKU 27993 (1), BSKU 27994 (1), BSKU 27995 (1), BSKU 27996 (1), BSKU 27997 (1), BSKU 27998 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.8333°N, 127.2333°E, 700–740 m, 16 Mar. 1978; NSMT-P 69399 (6), west-northwest of Amami-oshima Island, East China Sea, 29.0053°N, 127.2390°E, 503 m, 8 Nov. 2003; BSKU 28075 (1), BSKU 28077 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.8833°N, 127.3000°E, 820–830 m, 16 Mar. 1978; BSKU 33257 (1), west of Okinawa-jima Island, Okinawa Trough, 29.7167°N, 127.3667°E, 520 m, 14 Oct. 1979; SNFR 10123 (1), East China Sea, 29.3050°N, 127.4933°E, 649–663 m, 11 Sept. 1993; BSKU 27125 (1), BSKU 27126 (1), BSKU 27127 (1), BSKU 27129 (1), BSKU 27130 (1), BSKU 27141 (1), BSKU 27142 (1), BSKU 27143 (1), BSKU 27144 (1), BSKU 27145 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.3833°N, 127.5150°E, 650 m, 5 Feb. 1978; BSKU 27197 (1), BSKU 27198 (1), BSKU 27199 (1), BSKU 27200 (1), BSKU 27201 (1), BSKU 27202 (1), BSKU 27203 (1), BSKU 27204 (1), BSKU 27205 (1), BSKU 27206 (1), BSKU 27207 (1), BSKU 27208 (1), BSKU 27209 (1), BSKU 27210 (1), BSKU 27211 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.4333°N, 127.5167°E, 570 m, 5 Feb. 1978; FFNU P-00748 (1), FFNU P-00751 (1), north of Minamiensei Knoll, East China Sea, 29.6828°N, 127.7400°E, 596 m, 30 Nov. 2008; BSKU 26946 (1), BSKU 26947 (1), BSKU 26948 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.6483°N, 127.9333°E, 750–760 m, 4 Feb. 1978; BSKU 26984 (1), BSKU 26985 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.6467°N, 127.9733°E, 830 m, 4 Feb. 1978; BSKU 28435 (1), BSKU 28436 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7667°N, 127.9833°E, 713–770 m, 20 Mar. 1978; BSKU 28591 (1), BSKU 28592 (1), BSKU 28593 (1), BSKU 28594 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.9500°N, 128.0000°E, 585–610 m, 20 Mar. 1978; SNFR 16419 (1), East China Sea, 30.7225°N, 128.0068°E, 364–398 m, 11 Aug. 2008; BSKU 28496 (1), BSKU 28497 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7333°N, 128.0500°E, 815 m, 20 Mar. 1978; BSKU 28549 (1), BSKU 28550 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.8167°N,

128.1000°E, 915–932 m, 20 Mar. 1978; BSKU 34006 (1), BSKU 34007 (1), BSKU 34008 (1), west of Yaku-shima Island, Okinawa Trough, 30.2083°N, 128.1922°E, 600–620 m, 26 Oct. 1979; BSKU 33452 (1), BSKU 33453 (1), BSKU 33454 (1), BSKU 33455 (1), BSKU 33456 (1), BSKU 33457 (1), BSKU 33458 (1), BSKU 33459 (1), BSKU 33460 (1), BSKU 33461 (1), BSKU 33462 (1), west of Yaku-shima Island, Okinawa Trough, 30.2067°N, 128.2833°E, 780–810 m, 26 Oct. 1979; BSKU 26669 (1), BSKU 26705 (1), northwest of Amami-oshima Island, Okinawa Trough, 30.0300°N, 128.3650°E, 900 m, 3 Feb. 1978; BSKU 33933 (1), BSKU 33934 (1), BSKU 33935 (1), BSKU 33936 (1), west of Yaku-shima Island, Okinawa Trough, 31.1367°N, 128.3867°E, 505–515 m, 3 Nov. 1979; BSKU 29862 (1), west of Kagoshima, Okinawa Trough, 31.1389°N, 128.3889°E, 505–515 m, 3 Nov. 1979; NSMT-P 65668 (2), south of Danjo Islands, East China Sea, 31.4217°N, 128.4017°E, 461–500 m, 8 Nov. 2002; SNFR 13869 (1), SNFR 13870 (1), East China Sea, 31.1358°N, 128.4412°E, 433–519 m, 30 Jul. 2008; SNFR 11137 (1), East China Sea, 31.5667°N, 128.4667°E, 520–565 m, 18 Jul. 1988; SNFR 15577 (1), East China Sea, 31.7033°N, 128.7367°E, 710–735 m, 18 Jul. 1988; FFNU P-00784 (4), northeast of Daini-Danjo Knoll, East China Sea, 31.4045°N, 128.7570°E, 659 m, 8 Mar. 2009; FFNU P-00768 (4), southwest of Amakusa Knoll, East China Sea, 32.1023°N, 129.4827°E, 600 m, 17 Jul. 2008; BSKU 23285 (1), Mimase fish market, date unknown; BSKU 44418 (1), Tosa Bay, 600 m, 11 Jan. 1988; BSKU 44300 (1), Tosa Bay, 600 m, 17 Nov. 1987; BSKU 23130 (1), BSKU 23131 (1), Tosa Bay, 33.1500°N, 133.6600°E, 615 m, 25 Jan. 1975; BSKU 110073 (1), Suruga Bay, 34.7481°N, 138.4652°E, 243–424 m, 23 Apr. 2013; BSKU 110088 (1), Suruga Bay, 34.7489°N, 138.4664°E, 200–450 m, 23 Apr. 2013; NSMT-P 101348 (41), Suruga Bay, 350–500 m, date unknown; NSMT-P 101919 (1 of 15), west of Matsuzaki, Suruga Bay, 34.7567°N, 138.6650°E, 652–676 m, 17 May 1989; NSMT-P 101887 (9), west of Matsuzaki, Suruga Bay, 34.7522°N, 138.6777°E, 490–547 m, 18 May 1989; NSMT-P 101885 (15), NSMT-P 101345 (10), west of Cape Hagachizaki, Suruga Bay, 34.6817°N, 138.6817°E, 418–457 m, 11 Nov. 1983; NSMT-P 101917 (1), west of Toi, Suruga Bay, 34.9167°N, 138.7117°E, 516–630 m, 23 Nov. 1978; NSMT-P 101886 (2), west of Toi, Suruga Bay, 34.9050°N, 138.7133°E, 520–545 m, 6 Nov. 1978; NSMT-P 48542 (1), NSMT-P 48543 (1), northwest of Heta, Suruga Bay, 34.9933°N, 138.7450°E, 310–680 m, 25 Oct. 1993; FAKU M7511 (1), Heta fish market, Suruga Bay, 150 fth (275 m), 22–24 Jun. 1938; FAKU M6861 (1), FAKU M6862 (1), FAKU M6864 (1), FAKU M6867 (1), FAKU M6868 (1), FAKU M6869 (1), FAKU M6870 (1), FAKU M6871 (1), FAKU M6873 (1), FAKU M6878 (1), FAKU M6879 (1), FAKU M6881 (1), FAKU M6882 (1), FAKU M7388 (1), FAKU M7389 (1), FAKU M7390 (1), FAKU M7391 (1), Heta fish market, Suruga Bay, 150 fth (275 m), 23 Jul. 1938; FAKU M6948

(1), Heta fish market, Suruga Bay, 200–290 fth (366–531 m), 24–25 Jun. 1939; FAKU M7226 (1), FAKU M7269 (1), Heta fish market, Suruga Bay, 26 Mar. 1939; KPM-NI 650 (1), off Cape Osezaki, Suruga Bay, 23 Oct. 1967; FAKU 34307 (1), FAKU 34308 (1), FAKU 34311 (1), FAKU 34313 (1), FAKU 34315 (1), off Oiso, Sagami Bay, Apr. 1961; NSMT-P 65715 (1), off Hitachi, 36.4725°N, 141.0140°E, 367–417 m, 26 Oct. 2002; NSMT-P 65699 (5), off Hitachi, 36.4697°N, 141.0272°E, 444–471 m, 24 Oct. 2002.

Coryphaenoides microps. 11 specimens. KAUM-I. 44837 (1), Da-xi fish market, Yilan, 6 Jul. 2011; KAUM-I. 44661 (1), KAUM-I. 44662 (1), KAUM-I. 44663 (1), KAUM-I. 44664 (1), KAUM-I. 44665 (1), KAUM-I. 44666 (1), KAUM-I. 44667 (1), KAUM-I. 44668 (1), KAUM-I. 44669 (1), KAUM-I. 44670 (1), off Kaohsiung, 400 m, 1 Jul. 2011.

Coryphaenoides nasutus. 272 specimens: BSKU 26491 (1), BSKU 26492 (1), BSKU 26493 (1), BSKU 26494 (1), BSKU 26495 (1), northwest of Miyako-jima Island, Okinawa Trough, 25.2850°N, 124.7800°E, 910–990 m, 22 Jan. 1978; FFNU P-00812 (1), Iheya Knolls, East China Sea, 27.5683°N, 126.3050°E, 1194 m, 25 Jun. 2009; BSKU 28146 (1), BSKU 28147 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.0500°N, 127.2667°E, 908–915 m, 16 Mar. 1978; BSKU 28073 (1), BSKU 28074 (1), BSKU 28078 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.8833°N, 127.3000°E, 820–830 m, 16 Mar. 1978; BSKU 26986 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.6467°N, 127.9733°E, 830 m, 4 Feb. 1978; BSKU 28494 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7333°N, 128.0500°E, 815 m, 20 Mar. 1978; FFNU P-00761 (1), north of Kitaensei Knoll, East China Sea, 29.1912°N, 128.1080°E, 1141 m, 2 Dec. 2008; BSKU 26704 (1), northwest of Amami-oshima Island, Okinawa Trough, 30.0300°N, 128.3650°E, 900 m, 3 Feb. 1978; BSKU 26850 (1), BSKU 26851 (1), BSKU 26853 (1), BSKU 26854 (1), BSKU 26855 (1), BSKU 26856 (1), BSKU 26857 (1), north-northwest of Amami-oshima Island, Okinawa Trough, 29.7850°N, 128.4467°E, 1000 m, 3 Feb. 1978; NSMT-P 101920 (1), northwest of Amami-oshima Island, East China Sea, 28.5640°N, 129.1390°E, 946–957 m, 21 Jun. 1993; NSMT-P 101923 (5), off Miyazaki, Hyuga-nada, 31.8870°N, 131.8693°E, 779–803 m, 26 Jun. 1993; BSKU 49569 (1), off Miyazaki, Hyuga-nada, 32.3083°N, 132.1817°E, 1453–1481 m, 3 Apr. 1991; NSMT-P 97658 (1), off Miyazaki, Hyuga-nada, 32.5483°N, 132.4533°E, 1075–1094 m, 2 Nov. 1986; BSKU 30331 (1), BSKU 30332 (1), BSKU 30333 (1), BSKU 30334 (1), BSKU 30335 (1), BSKU 30336 (1), BSKU 30337 (1), BSKU 30338 (1), Tosa Bay, 32.9167°N, 133.5500°E, 700 m, 21 Dec. 1979; BSKU 22781 (1), BSKU 22782 (1), BSKU 22783 (1), Tosa Bay, 32.9433°N, 133.5517°E, 707–750 m, 24 Jul. 1974; BSKU 22772 (1), Tosa Bay, 32.9467°N, 133.5567°E, 714–735 m,

24 Jul. 1974; BSKU 44845 (1), BSKU 44846 (1), BSKU 44847 (1), Tosa Bay, 32.9886°N, 133.5978°E, 700–720 m, 21 Jun. 1988; BSKU 95205 (1), BSKU 95206 (1), BSKU 95208 (1), BSKU 95209 (1), BSKU 95210 (1), BSKU 95211 (1), BSKU 95212 (1), BSKU 95214 (1), BSKU 95216 (1), BSKU 95217 (1), BSKU 95218 (1), BSKU 95220 (1), BSKU 95222 (1), BSKU 95223 (1), BSKU 95224 (1), BSKU 95225 (1), BSKU 95227 (1), BSKU 95228 (1), Tosa Bay, 750 m, 13 Sept. 1988; BSKU 22811 (1), BSKU 22812 (1), BSKU 22813 (1), BSKU 22814 (1), BSKU 22815 (1), BSKU 22817 (1), BSKU 22818 (1), BSKU 22819 (1), BSKU 22820 (1), BSKU 22821 (1), BSKU 22822 (1), BSKU 22823 (1), BSKU 22824 (1), BSKU 22825 (1), BSKU 22826 (1), BSKU 22827 (1), BSKU 22828 (1), BSKU 22829 (1), BSKU 22830 (1), BSKU 22831 (1), BSKU 22832 (1), BSKU 22833 (1), BSKU 22834 (1), BSKU 22835 (1), BSKU 22836 (1), BSKU 22837 (1), BSKU 22838 (1), BSKU 22839 (1), BSKU 22840 (1), off Tosa Bay, 32.8967°N, 133.6217°E, 960–985 m, 24 Jul. 1974; BSKU 22795 (1), Tosa Bay, 32.8967°N, 133.6217°E, 960–985 m, 24 Jul. 1974; NSMT-P 55419 (1), NSMT-P 55421 (3), off Katsurahama, Tosa Bay, 33.0167°N, 133.6417°E, 823–856 m, 22 Aug. 1997; SNFR 17952 (1), Tosa Bay, 750–780 m, 24 Nov. 1987; BSKU 47160 (1), BSKU 47161 (1), BSKU 47162 (1), BSKU 47165 (1), BSKU 47167 (1), BSKU 47168 (1), off Tosa Bay, 32.8953°N, 133.6777°E, 996–1010 m, 8 Nov. 1989; NSMT-P 55401 (2), off Nangoku, Tosa Bay, 33.0933°N, 133.6900°E, 753–768 m, 12 Nov. 1997; BSKU 70272 (1), Tosa Bay, 33.0933°N, 133.6900°E, 753–768 m, 12 Nov. 1997; BSKU 23051 (1), BSKU 23052 (1), BSKU 23053 (1), BSKU 23149 (1), BSKU 23150 (1), BSKU 23151 (1), Tosa Bay, 33.0350°N, 133.7500°E, 945 m, 26 Jan. 1975; BSKU uncat. (5), BSKU 23182 (1), BSKU 23086 (1), BSKU 23087 (1), BSKU 23088 (1), off Kochi, 32.9800°N, 133.8550°E, 1043 m, 26 Jan. 1975; NSMT-P 101337 (1), southwest of Kii Peninsula, Kii Channel, 33.5013°N, 135.1420°E, 975–1112 m, 2 Sept. 1984; NSMT-P 99690 (2), southwest of Cape Daiozaki, Kumano-nada, 34.0283°N, 136.6500°E, 906–955 m, 28 May 1986; NSMT-P 101922 (1), south of Cape Daiozaki, Kumano-nada, 33.9093°N, 136.8382°E, 1165–1181 m, 21 Aug. 1988; KPM-NI 18271 (1), off Shima Peninsula, Kumano-nada, 34.0103°N, 136.8793°E, 781–818 m, 24 Nov. 2005; KPM-NI 18258 (1), off Shima Peninsula, Kumano-nada, 34.0005°N, 136.8937°E, 771–787 m, 24 Nov. 2005; KPM-NI 18233 (1), KPM-NI 18234 (1), off Shima Peninsula, Kumano-nada, 34.0083°N, 136.9071°E, 767–774 m, 1 May 2004; BSKU 98923 (1), Suruga Bay, 29 Oct. 1972; NSMT-P 101919 (14 of 15), west of Matsuzaki, Suruga Bay, 34.7567°N, 138.6650°E, 652–676 m, 17 May 1989; NSMT-P 101924 (5), south-southeast of Fuji River, Suruga Bay, 35.0537°N, 138.6708°E, 971–1000 m, 13 Apr. 1993; NSMT-P 101918 (1), west-northwest of Toi, Suruga Bay, 34.9367°N, 138.7117°E, 548–590 m, 14 Apr. 1981; NSMT-P 93131 (1), NSMT-P 93125 (1), NSMT-P 93126 (1), NSMT-P 93121 (1), off Oh-sima Island, Izu Islands, date unknown; FMNH 55375 (3, ex. CM 6107a–c), Sagami Bay, date unknown; BSKU 19777 (1), Suruga Bay, 35.1650°N, 139.4083°E, 525–540 m, 30 Jun. 1967; BSKU 14197 (1), BSKU 14198 (1), Sagami Bay, 12 Apr. 1969; BSKU 14199 (1), BSKU 14200 (1), BSKU 14201 (1), Sagami Bay, 14 Apr. 1969; BSKU 19324 (1), Suruga Bay, 35.1333°N, 139.4683°E, 735 m, 14 Mar. 1963; BSKU 20368 (1), BSKU 20369 (1), BSKU 20370 (1), Sagami Bay, 35.1900°N, 139.4800°E, 805 m, 6 Jul. 1966; BSKU 20366 (1), southwest of Miura Peninsula, Sagami-nada, 35.0817°N, 139.4950°E, 770 m, 17 Nov. 1961; FMNH 57384 (2, ex. CM 6322a–b), Misaki, Sagami Bay, date unknown; FMNH uncat. (1, ex. CM unknown), Misaki, Sagami Bay, 1922; BSKU 19680 (1), BSKU 19681 (1), west of Izu Islands, Shichito-Iojima Ridge, 34.3550°N, 139.6217°E, 920–955 m, 18 Feb. 1966; NSMT-P 94370 (2), east of Miyake-jima Island, Shichito-Iojima Ridge, 565–644 m, 19 May 1989; BSKU 19652 (1), east of Miyake-jima Island, Shichito-Iojima Ridge, 34.0667°N, 140.3583°E, 1940–1980 m, 10 Oct. 1965; NSMT-P 102376 (1), off Kujukuri, 20 May 1988; FAKU 8022 (1), FAKU 8025 (1), FAKU 8026 (1), FAKU 8027 (1), off Choshi, 10 Apr. 1939; FAKU 5172 (1), FAKU 5175 (1), FAKU 5176 (1), off Choshi, 20–26 Nov. 1937; BSKU 97817 (1), off Mito, 450 m, 9 Oct. 1997; NSMT-P 97655 (3), southeast off Onahama, 621–710 m, 5 Sept. 1989; BSKU 101988 (1), BSKU 101989 (1), BSKU 101990 (1), BSKU 101991 (1), BSKU 101992 (1), BSKU 101993 (1), BSKU 101994 (1), off Iwaki, 36.9162°N, 141.5995°E, 615–633 m, 4 Nov. 1995; FAKU 102252 (1), off Ukedo, 3 Mar. 1967; NSMT-P 61234 (1), off Iwaki, 36.9483°N, 141.6317°E, 600–603 m, 17 Apr. 1997; NSMT-P 99687 (2), off Minami-soma, 37.5367°N, 142.0550°E, 538–561 m, 1 Dec. 1981; NSMT-P 102088 (1), east-southeast of Oshika Peninsula, 38.0367°N, 142.2250°E, 829–831 m, 19 May 1995; NSMT-P 79369 (1), east of Oshika Peninsula, 38.3867°N, 142.1783°E, 656 m, 25 Oct. 2007; BSKU 101968 (18), southeast of Kuji, 39.9997°N, 142.3687°E, 996–1010 m, 2 Nov. 1995; BSKU 97869 (1), BSKU 97870 (1), BSKU 97871 (1), BSKU 97874 (1), BSKU 97875 (1), BSKU 97876 (1), BSKU 97877 (1), BSKU 97878 (1), BSKU 97879 (1), BSKU 97880 (1), BSKU 97881 (1), BSKU 97882 (1), BSKU 97883 (1), BSKU 97884 (1), BSKU 97885 (1), BSKU 97886 (1), BSKU 97889 (1), BSKU 97891 (1), BSKU 97892 (1), BSKU 97894 (1), BSKU 97895 (1), BSKU 97896 (1), BSKU 97897 (1), BSKU 97898 (1), BSKU 97899 (1), BSKU 97902 (1), BSKU 97904 (1), BSKU 97905 (1), off Kamaishi, 39.1444°N, 142.3748°E, 847–935 m, 10 Mar. 1989; NSMT-P 100441 (1), east of Otsuchi, 39.1050°N, 142.3967°E, 926–950 m, 10 Jul. 1984; NSMT-P 102401 (14), off Kamaishi, 39.1050°N, 142.3967°E, 926–950 m, 10 Jul. 1984; NSMT-P 97653 (1), NSMT-P 97660 (1), off Kamaishi, 39.1663°N, 142.3983°E, 975–988 m, 27 May 1989; NSMT-P 101921 (2), off Kamaishi, 39.0150°N,

142.6167°E, 1237–1255 m, 21 May 1987; FAKU 29133 (1), off Hachinohe, Dec. 1959; BSKU 99135 (1), off Hachinohe, 40.6018°N, 143.3585°E, 2014–2056 m, 13 Nov. 1995; FAKU M9310 (1), FAKU M9311 (1), FAKU M9313 (1), off Muroran, Mar. 1938.

Coryphaenoides pectoralis. 66 specimens: NSMT-P 19030 (1), Sagami Bay, 35.1350°N, 139.3467°E, 780 m, 30 Nov. 1975; HUMZ 72581 (1), HUMZ 72582 (1), off Kashima, Ibaraki, 900 m, 12 Jan. 1978; HUMZ 72887 (1), off Miyagi, 1000 m, date unknown; NSMT-P 77869 (1), NSMT-P 77870 (1), off Minami-soma, 37.6050°N, 142.5600°E, 1513–1515 m, 1 Nov. 2006; BSKU 97801 (1), off Kitakami River, 38.5318°N, 142.3370°E, 772–783 m, 28 Oct. 1995; HUMZ 72681 (1), off Miyagi, 38.7250°N, 142.3917°E, 1000–1100 m, 4 Feb. 1978; NSMT-P 102404 (4), off Kamaishi, 39.1050°N, 142.3967°E, 926–950 m, 10 Jul. 1984; NSMT-P 101880 (1), off Kamaishi, 39.1131°N, 142.4127°E, 1038–1055 m, 20 May 1987; NSMT-P 101881 (1), off Kamaishi, 39.0626°N, 142.6244°E, 1203–1213 m, 24 May 1988; NSMT-P 90643 (1), off Miyako, 1000 m, 5 Nov. 2007; HUMZ 78178 (1), off Iwate, 39.0917°N, 142.7383°E, 1300 m, 20 Sept. 1978; NSMT-P 79352 (1), off Kesennuma, 38.5367°N, 142.8000°E, 1502 m, 26 Oct. 2007; NSMT-P 58946 (1), off Kesennuma, 39.0033°N, 143.4950°E, 650–679 m, 30 Jul. 1996; NSMT-P 58944 (1), off Kesennuma, 39.0683°N, 143.5500°E, 550–579 m, 30 Jul. 1996; HUMZ 206292 (1), south of Tomakomai, Hokkaido, 42.2132°N, 141.6842°E, 800 m, 29 Sept. 2009; HUMZ 6928 (1), off Urakawa, Hokkaido, date unknown; HUMZ 6921 (1), off Cape Erimomisaki, Hokkaido, 1936; NSMT-P 77255 (1), south-southeast of Cape Erimomisaki, 41.5677°N, 143.4767°E, 827–889 m, 28 May 2010; HUMZ 69383 (1), off Hiroo, Hokkaido, 650 m, 13 Aug. 1977; BSKU 114904 (1), east-southeast of Cape Erimomisaki, 42.2820°N, 143.8012°E, 572 m, 9 Jun. 2014; HUMZ 67357 (1), northwestern Pacific, 41.8333°N, 143.8333°E, 770–830 m, 5 Jul. 1977; BSKU 114823 (1), southeast of Tokachi, 42.3317°N, 143.9585°E, 719 m, 10 Jun. 2014; BSKU 114859 (1), south of Kushiro, 42.6542°N, 144.3770°E, 780 m, 7 Jun. 2014; HUMZ 49043 (1), off Okhotsk coast of Hokkaido, 45.2500°N, 143.9667°E, 580–645 m, 7 Oct. 1975; NSMT-P 90974 (1), NSMT-P 90975 (1), NSMT-P 90976 (1), NSMT-P 90977 (1), NSMT-P 90978 (1), NSMT-P 90979 (1), NSMT-P 90980 (1), NSMT-P 90981 (1), Kitami-Yamato Bank, Okhotsk Sea, 2 Jun. 2008; HUMZ 67511 (1), HUMZ 67512 (1), HUMZ 67513 (1), HUMZ 67514 (1), off Okhotsk coast of Hokkaido, 45.0833°N, 144.2167°E, 450–475 m, 20 May 1977; HUMZ 44962 (1), HUMZ 44968 (1), Kitami-Yamato Bank, Okhotsk Sea, 44.4167°N, 144.2917°E, 385–400 m, 3 Jul. 1975; HUMZ 68543 (1), off Okhotsk coast of Hokkaido, 44.8833°N, 144.3167°E, 152–153 m, 2 Jul. 1977; HUMZ 92128 (1), HUMZ 92129 (1), Kitami-Yamato Bank, Okhotsk Sea, 44.7333°N, 144.3333°E, 449 m, 30 Jun. 1981; HUMZ 78605 (1), HUMZ 78604 (1), Kitami-Yamato Bank,

Okhotsk Sea, 19 Oct. 1978; HUMZ 124352 (1), Kitami-Yamato Bank, Okhotsk Sea, 44.5700°N, 144.9983°E, 1300 m, 4 Jul. 1992; HUMZ 78752 (1), HUMZ 79674 (1), HUMZ 79675 (1), HUMZ 79676 (1), Kitami-Yamato Bank, Okhotsk Sea, 44.1833°N, 145.0000°E, 470–520 m, 11 Sept. 1978; HUMZ 79661 (1), Kitami-Yamato Bank, Okhotsk Sea, 44.1867°N, 145.0000°E, 455–500 m, 14 Sept. 1978; HUMZ 78822 (1), HUMZ 78823 (1), Kitami-Yamato Bank, Okhotsk Sea, 44.4000°N, 145.0333°E, 500 m, 9 Oct. 1978; NSMT-P 58884 (3), off Arahama, Miyagi, 40.3033°N, 144.9567°E, 630–657 m, 16 Jul. 1996; NSMT-P 58885 (4), off Kuji, Japan Trench, 40.0317°N, 144.9733°E, 630–657 m, 15 Jul. 1996; NSMT-P 58945 (1), off Kuji, Japan Trench, 39.9900°N, 145.0417°E, 630–658 m, 16 Jul. 1996; NSMT-P 59299 (2), southeast of Cape Erimomisaki, Japan Trench, 41.0033°N, 145.7717°E, 630–657 m, 8 Jul. 1996.

Coryphaenoides yaquinae. 22 specimens: NSMT-P 78122 (1), off Kushiro, 5700 m, 1–2 Aug. 2007; NSMT-P 102117 (1), east of Miyake-jima Island, Japan Trench, 34.2543°N, 142.7752°E, 5491–5497 m, 3 Sept. 1987; BSKU 19998 (1), BSKU 20000 (1), east of Japan Trench, 33.9750°N, 142.8550°E, 5200 m, 4 Jun. 1970; BSKU 35718 (1), BSKU 35721 (1), off Kuji, Japan Trench, 40.1133°N, 143.9700°E, 4200–4220 m, 15 Jul. 1981; BSKU 49931 (1), Japan Trench, 38.7342°N, 144.1233°E, 5500 m, 18 Nov. 1991; NSMT-P 98009 (1), off Miyako, 38.7983°N, 144.1345°E, 7340–7433 m, 30 Sept. 2001; BSKU 35728 (1), Kesennuma, Japan Trench, 38.5550°N, 144.3233°E, 6380–6450 m, 31 Jul. 1981; BSKU 37298 (1), BSKU 37300 (1), east of Cape Inbozaki, east of Japan Trench, 35.8367°N, 145.2083°E, 4100 m, 4 Nov. 1981; BSKU 35731 (1), BSKU 35732 (1), east of Japan Trench, 38.5650°N, 145.2617°E, 5350–5370 m, 1 Aug. 1981; NSMT-P 98010 (1), NSMT-P 98011 (1), NSMT-P 98012 (1), NSMT-P 98013 (1), NSMT-P 98014 (1), NSMT-P 98015 (1), NSMT-P 98016 (1), NSMT-P 98017 (1), NSMT-P 98018 (1), southeast of Kushiro, Japan Trench, 41.4250°N, 146.3900°E, 5509–5611 m, 22 Sept. 2001.

Hymenocephalus lethoenemus. 479 specimens: BSKU 26329 (1), BSKU 26330 (1), off Senkaku Islands, Okinawa Trough, 25.6283°N, 122.8933°E, 560–692 m, 14 Jan. 1978; ZMUT 53490 (1), Ishigaki-jima Island, date unknown; BSKU 27863 (1), BSKU 27874 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.5833°N, 127.0000°E, 600 m, 16 Mar. 1978; BSKU 27655 (1), BSKU 27656 (1), west of Okinawa-jima Island, Okinawa Trough, 28.7333°N, 127.0167°E, 610–640 m, 11 Mar. 1978; BSKU 27907 (1), BSKU 27909 (1), BSKU 27910 (1), BSKU 27911 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.7500°N, 127.1167°E, 520–542 m, 16 Mar. 1978; BSKU 27710 (1), BSKU 27711 (1), west of Okinawa-jima Island, Okinawa Trough, 28.7000°N, 127.1500°E, 535–550 m, 11 Mar. 1978; BSKU 27990 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.8333°N, 127.2333°E, 700–740 m, 16 Mar. 1978; SNFR

13230 (2), East China Sea, 29.3050°N, 127.4933°E, 649–663 m, 11 Sept. 1993; BSKU 27119 (1), BSKU 27120 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.3833°N, 127.5150°E, 650 m, 5 Feb. 1978; BSKU 27176 (1), BSKU 27177 (1), BSKU 27178 (1), BSKU 27179 (1), BSKU 27180 (1), BSKU 27181 (1), BSKU 27182 (1), BSKU 27183 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.4333°N, 127.5167°E, 570 m, 5 Feb. 1978; SNFR 12493 (1), East China Sea, 29.6983°N, 127.7000°E, 558–559 m, 12 Sept. 1993; BSKU 28385 (1), BSKU 28386 (1), BSKU 28387 (1), BSKU 28389 (1), BSKU 28390 (1), BSKU 28392 (1), BSKU 28393 (1), BSKU 28394 (1), BSKU 28395 (1), BSKU 28398 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7333°N, 127.8000°E, 495–510 m, 19 Mar. 1978; BSKU 26936 (1), BSKU 26937 (1), BSKU 26938 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.6483°N, 127.9333°E, 750–760 m, 4 Feb. 1978; BSKU 28443 (1), BSKU 28444 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7667°N, 127.9833°E, 713–770 m, 20 Mar. 1978; BSKU 28602 (1), BSKU 28603 (1), BSKU 28604 (1), BSKU 28605 (1), BSKU 28606 (1), BSKU 28607 (1), BSKU 28608 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.9500°N, 128.0000°E, 585–610 m, 20 Mar. 1978; BSKU 28500 (1), BSKU 28501 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7333°N, 128.0500°E, 815 m, 20 Mar. 1978; BSKU 33529 (1), BSKU 33530 (1), BSKU 33531 (1), BSKU 33532 (1), BSKU 33533 (1), west of Yaku-shima Island, Okinawa Trough, 30.2083°N, 128.1922°E, 600–620 m, 26 Oct. 1979; BSKU 29790 (1), BSKU 29889 (1), west of Yaku-shima Island, Okinawa Trough, 30.2083°N, 128.2088°E, 600–620 m, 26 Oct. 1979; BSKU 33501 (1), BSKU 33503 (1), BSKU 33504 (1), BSKU 33505 (1), BSKU 33506 (1), BSKU 33507 (1), BSKU 33508 (1), BSKU 33510 (1), BSKU 33512 (1), BSKU 33513 (1), west of Yaku-shima Island, Okinawa Trough, 30.2067°N, 128.2833°E, 780–810 m, 26 Oct. 1979; BSKU 26862 (1), north-northwest of Amami-oshima Island, Okinawa Trough, 29.7850°N, 128.4467°E, 1000 m, 3 Feb. 1978; FFNU P-00767 (4), southwest of Amakusa Knoll, East China Sea, 32.1023°N, 129.4827°E, 600 m, 3 Dec. 2008; BSKU 13113 (1), BSKU 13114 (1), BSKU 13115 (1), BSKU 13116 (1), BSKU 13268 (1), BSKU 13269 (1), BSKU 13270 (1), BSKU 13271 (1), BSKU 13272 (1), east of Cape Ashizurimisaki, Tosa Bay, 420–555 m, 2 Jun. 1968; BSKU 23105 (1), Tosa Bay, 32.9233°N, 133.4850°E, 480 m, 25 Jun. 1970; BSKU 30311 (1), Tosa Bay, 32.9167°N, 133.5500°E, 700 m, 21 Dec. 1979; BSKU 22777 (1), Tosa Bay, 32.9433°N, 133.5517°E, 707–750 m, 24 Jul. 1974; BSKU 22762 (1), BSKU 22763 (1), BSKU 22764 (1), Tosa Bay, 32.9467°N, 133.5567°E, 714–735 m, 24 Jul. 1974; FAKU 28292 (1), FAKU 28293 (1), FAKU 28294 (1), FAKU 28295 (1), FAKU 28296 (1), FAKU 28298 (1), FAKU 28300 (1), FAKU 28303 (1), FAKU 28306 (1), FAKU 28307 (1), FAKU 28308 (1),

FAKU 28312 (1), FAKU 28314 (1), FAKU 28316 (1), FAKU 28322 (1), FAKU 28324 (1), FAKU 28329 (1), FAKU 28330 (1), FAKU 28333 (1), FAKU 28341 (1), FAKU 28342 (1), FAKU 28343 (1), Mimase fish market, 31 Dec. 1958; BSKU 13877 (1), BSKU 13903 (1), BSKU 13904 (1), BSKU 13905 (1), BSKU 13906 (1), BSKU 13907 (1), BSKU 13908 (1), BSKU 13909 (1), BSKU 13910 (1), BSKU 13911 (1), BSKU 13912 (1), Mimase fish market, date unknown; BSKU 44841 (1), BSKU 44842 (1), BSKU 44843 (1), Tosa Bay, 32.9886°N, 133.5978°E, 700–720 m, 21 Jun. 1988; BSKU 12941 (1), BSKU 12942 (1), BSKU 12943 (1), BSKU 12944 (1), BSKU 12945 (1), BSKU 12947 (1), BSKU 12948 (1), BSKU 12949 (1), BSKU 12950 (1), Tosa Bay, 350–450 m, 14 May 1968; BSKU 40347 (1), BSKU 40348 (1), BSKU 40393 (1), BSKU 40394 (1), Tosa Bay, 400 m, 9 Oct. 1984; BSKU 41914 (1), BSKU 41915 (3), Tosa Bay, 420–500 m, 6 Jun. 1985; BSKU 44763 (1), BSKU 44764 (1), BSKU 44765 (1), Tosa Bay, 600 m, 16 May 1988; BSKU 43794 (1), BSKU 43796 (1), Tosa Bay, 600 m, 20 May 1987; BSKU 45120 (1), Tosa Bay, 750 m, 13 Sept. 1988; BSKU 46154 (9), Tosa Bay, 33.0400°N, 133.6098°E, 675–708 m, 22 May 1989; BSKU 97381 (1), BSKU 97382 (1), BSKU 97383 (1), BSKU 97384 (1), BSKU 97385 (1), BSKU 97386 (1), BSKU 97387 (1), BSKU 97388 (1), BSKU 97389 (1), BSKU 97390 (1), BSKU 97391 (1), BSKU 97392 (1), BSKU 97393 (1), BSKU 97394 (1), BSKU 97395 (1), BSKU 97396 (1), BSKU 97397 (1), BSKU 97398 (1), BSKU 97399 (1), BSKU 97400 (1), BSKU 97401 (1), BSKU 97402 (1), BSKU 97403 (1), BSKU 97404 (1), BSKU 97405 (1), BSKU 97406 (1), BSKU 97407 (1), Tosa Bay, 33.1428°N, 133.6248°E, 450–490 m, 1 Aug. 1988; BSKU 97409 (1), Tosa Bay, 400 m, 9 May 2000; BSKU 71616 (1), BSKU 71617 (1), BSKU 71618 (1), Tosa Bay, 33.1950°N, 133.6783°E, 528–537 m, 13 May 1997; BSKU 78010 (1), BSKU 78011 (1), Tosa Bay, 33.2083°N, 133.6983°E, 526–539 m, 3 Mar. 1999; BSKU 69189 (1), Tosa Bay, 33.1083°N, 133.7083°E, 763–803 m, 2 Mar. 1999; BSKU 13342 (1), BSKU 13343 (1), BSKU 13344 (1), BSKU 13345 (1), BSKU 13346 (1), BSKU 13347 (1), BSKU 13349 (1), west of Muroto, Tosa Bay, 350–450 m, 14 May 1968; BSKU 77303 (1), BSKU 77304 (1), BSKU 77305 (1), BSKU 77306 (1), BSKU 77307 (1), BSKU 77308 (1), BSKU 77309 (1), BSKU 77310 (1), BSKU 77311 (1), BSKU 77312 (1), BSKU 77313 (1), BSKU 77314 (1), BSKU 77315 (1), BSKU 77316 (1), BSKU 77317 (1), BSKU 77318 (1), BSKU 77319 (1), BSKU 77320 (1), BSKU 77321 (1), BSKU 77322 (1), BSKU 77323 (1), BSKU 77324 (1), BSKU 77325 (1), BSKU 77326 (1), BSKU 77327 (1), BSKU 77328 (1), BSKU 77329 (1), BSKU 77330 (1), BSKU 77331 (1), BSKU 77332 (1), BSKU 77333 (1), BSKU 77334 (1), BSKU 77335 (1), BSKU 77336 (1), BSKU 77337 (1), BSKU 77338 (1), BSKU 77339 (1), BSKU 77340 (1), BSKU 77341 (1), BSKU 77342 (1), BSKU 77343 (1), BSKU 77344

(1), BSKU 77345 (1), BSKU 77346 (1), BSKU 77347 (1), BSKU 77348 (1), BSKU 77349 (1), BSKU 77350 (1), BSKU 77351 (1), BSKU 77352 (1), BSKU 77353 (1), BSKU 77354 (1), BSKU 77355 (1), BSKU 77356 (1), BSKU 77357 (1), BSKU 77358 (1), BSKU 77359 (1), BSKU 77360 (1), BSKU 77361 (1), BSKU 77362 (1), BSKU 77363 (1), BSKU 77364 (1), BSKU 77365 (1), BSKU 77366 (1), BSKU 77367 (1), south of Muroto, 33.0790°N, 134.1645°E, 414–513 m, 17 Nov. 2005; KPM-NI 28848 (1), KPM-NI 28849 (1), ca. 24 km west-southwest of Shirahama, Wakayama, 33.5795°N, 135.0995°E, 532 m, 15 Aug. 2010; NSMT-P 101354 (9), southwest of Cape Daiozaki, Kumano-nada, 34.0836°N, 136.6554°E, 495–515 m, 12 Dec. 1995; NSMT-P 94396 (1), south of Daiozaki, Kumano-nada, 422–425 m, 28 May 1986; NSMT-P 94412 (1), south of Daiozaki, Kumano-nada, 475–494 m, 25 May 1994; FAKU 28399 (1), Miya fish market, Enshu-nada, 5–7 Feb. 1959; NSMT-P 101931 (9), southwest of Cape Omaezaki, Enshu-nada, 34.3816°N, 138.0010°E, 475–488 m, 10 Dec. 1995; BSKU 110105 (1), Suruga Bay, 34.7489°N, 138.4664°E, 200–450 m, 23 Apr. 2013; BSKU 98916 (1), BSKU 98917 (1), BSKU 98918 (1), BSKU 98919 (1), BSKU 98920 (1), BSKU 98921 (1), BSKU 98922 (1), Suruga Bay, 29 Oct. 1972; KPM-NI 24987 (1), eastern part of Suruga Bay, Nov. 2009; FAKU 22837 (1), FAKU 22838 (1), Heta fish market, Suruga Bay, 200–250 fm (366–458 m), 24–25 Jun. 1939; BSKU 18886 (1), off Atami, Sagami Bay, 35.0767°N, 139.1233°E, 380–425 m, 4 Oct. 1957; BSKU 20306 (1), BSKU 20307 (1), BSKU 20308 (1), BSKU 20309 (1), BSKU 20310 (1), BSKU 20311 (1), BSKU 20313 (1), off Atami, Sagami Bay, 35.0783°N, 139.1467°E, 450 m, 24 Nov. 1972; NSMT-P 93137 (1), off Oshima Island, Izu Islands, date unknown; BSKU 20004 (1), BSKU 20005 (1), off Chigasaki, Sagami Bay, 35.1683°N, 139.4233°E, 700 m, 25 Jun. 1970; NSMT-P 18948 (1), Sagami Bay, 35.1750°N, 139.4883°E, 650 m, 29 Nov. 1974; NSMT-P 94411 (1), southeast of Oshima Island, Izu Islands, 621–710 m, 5 Sept. 1989; NSMT-P 94417 (21), east of Miyake-jima Island, Shichito-Iojima Ridge, 484–565 m, 19 May 1989; NSMT-P 94415 (7), east of Miyake-jima Island, Shichito-Iojima Ridge, 565–644 m, 19 May 1989; NSMT-P 96997 (1), east of Boso Peninsula, 35.1287°N, 140.8364°E, 498–531 m, 11 May 2004; BSKU 19156 (1), BSKU 19157 (1), BSKU 19158 (1), off Mito, 36.2717°N, 140.9467°E, 355–370 m, 25 Jul. 1960; NSMT-P 77292 (1), NSMT-P 77293 (3), off Hitachi, 36.5150°N, 141.0517°E, 407–413 m, 13 Nov. 2006; NSMT-P 94385 (18), off Hitachi, 390–400 m, 9 Aug. 1979; BSKU 18839 (1), off Hitachi, 36.6133°N, 141.1600°E, 300 m, 7 Nov. 1956; NSMT-P 78962 (1), NSMT-P 78963 (3), off Iwaki, 36.8885°N, 141.4562°E, 376 m, 2 Nov. 2007; NSMT-P 53607 (2), off Soma, 37.8505°N, 141.4760°E, 410–415 m, 15 Apr. 1996; NSMT-P 48955 (29), off Iwaki, 36.9965°N, 141.5627°E, 387–420 m, 4 Nov. 1995; BSKU 99010 (1), BSKU 99011 (1), BSKU 99012 (1), BSKU

99013 (1), BSKU 99014 (1), BSKU 99015 (1), BSKU 99016 (1), BSKU 99017 (1), BSKU 99018 (1), BSKU 99019 (1), BSKU 99020 (1), BSKU 99021 (1), BSKU 99022 (1), BSKU 99023 (1), BSKU 99024 (1), BSKU 99025 (1), BSKU 99026 (1), BSKU 99027 (1), BSKU 99028 (1), BSKU 99029 (1), BSKU 99030 (1), BSKU 99031 (1), BSKU 99032 (1), BSKU 99033 (1), BSKU 99034 (1), BSKU 99035 (1), BSKU 99036 (1), BSKU 99037 (1), BSKU 99038 (1), BSKU 99039 (1), BSKU 99040 (1), BSKU 99041 (1), BSKU 99042 (1), BSKU 99043 (1), BSKU 99044 (1), BSKU 99045 (1), BSKU 99046 (1), BSKU 99047 (1), BSKU 99048 (1), BSKU 99049 (1), BSKU 99050 (1), BSKU 99051 (1), BSKU 99052 (1), BSKU 99053 (1), BSKU 99054 (1), BSKU 99055 (1), BSKU 99056 (1), BSKU 99057 (1), BSKU 99058 (1), BSKU 99059 (1), BSKU 99060 (1), BSKU 99061 (1), BSKU 99062 (1), BSKU 99063 (1), BSKU 99064 (1), BSKU 99065 (1), BSKU 99066 (1), BSKU 99067 (1), BSKU 99068 (1), off Onahama, 36.9965°N, 141.5627°E, 387–420 m, 4 Nov. 1995; BSKU 97410 (1), BSKU 97411 (1), BSKU 97412 (1), BSKU 97413 (1), off Onahama, 450 m, 16 Oct. 1997; BSKU 32202 (1), Kyushu-Palau Ridge, 27.9178°N, 134.7460°E, 685–710 m, 19 Dec. 1979; BSKU 19477 (1), BSKU 19478 (1), BSKU 19479 (1), BSKU 19480 (1), BSKU 19481 (1), BSKU 19482 (1), BSKU 19483 (1), BSKU 19484 (1), BSKU 19485 (1), BSKU 19486 (1), BSKU 19487 (1), BSKU 19488 (1), BSKU 19489 (1), BSKU 19490 (1), BSKU 19491 (1), BSKU 19492 (1), off Hachijo-jima Island, Shichito-Iojima Ridge, 33.1083°N, 140.0800°E, 490–495 m, 12 Dec. 1963.

Hymenocephalus longibarbis. 34 specimens: BSKU 29754 (1), west of Okinawa-jima Island, Okinawa Trough, 25.9833°N, 125.8500°E, 430 m, 16 Oct. 1979; BSKU 27470 (1), west of Okinawa-jima Island, Okinawa Trough, 27.7767°N, 126.2833°E, 490 m, 9 Mar. 1978; BSKU 28234 (1), BSKU 28235 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.2667°N, 127.3500°E, 353–365 m, 17 Mar. 1978; BSKU 28290 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7500°N, 127.5000°E, 325–360 m, 19 Mar. 1978; FFNU P-00725 (9), east of Nakakasayama Bank, East China Sea, 32.2072°N, 128.9563°E, 385 m, 9 Jun. 2008; FFNU P-00803 (1), southeast of Okihirazi Bank, East China Sea, 32.1553°N, 129.4998°E, 494 m, 10 Jun. 2009; KAUM-I. 32268 (1), off Nomaie, Kagoshima, 31.5000°N, 129.8833°E, 370–400 m, 10 Sept. 2010; BSKU 13266 (1), ca. 15 km east of Cape Ashizurimisaki, Tosa Bay, 420–555 m, 2–6 Jun. 1968; BSKU 48162 (1), Mimase fish market, 400 m, 19 Mar. 1989; BSKU 70089 (1), Mimase fish market, 15 Jan. 2004; BSKU 77901 (1), Mimase fish market, 19 Dec. 2004; BSKU 42497 (1), Mimase fish market, 21 Dec. 1985; BSKU 76886 (1), Mimase fish market, 11 Jan. 2005; BSKU 14175 (1), Mimase fish market, Dec. 1960; BSKU 7039 (1), BSKU 7040 (1), Mimase fish market, Oct. 1956; BSKU 14173 (1), Mimase fish

market, Jan. 1968; BSKU 92326 (1), Mimase fish market, 14 Nov. 2007; BSKU 99260 (1), Mimase fish market, 2 Dec. 2007; BSKU 92267 (1), Mimase fish market, 9 Nov. 2007; BSKU 44188 (1), off Okino-shima Island, Kochi, 230 m, 5 Nov. 1987; BSKU 97408 (1), Tosa Bay, 400 m, 9 May 2000; BSKU 110100 (1), Suruga Bay, 34.7481°N, 138.4652°E, 243–424 m, 23 Apr. 2013; BSKU 110104 (1), Suruga Bay, 34.7489°N, 138.4664°E, 200–450 m, 23 Apr. 2013; KAUM-I. 44660 (1), off Kaohsiung, 400 m, 1 Jul. 2011.

***Hymenocephalus papyraceus*.** 7 specimens: BSKU uncat. (1), off Yoron-jima Island, 27.1133°N, 128.6983°E, 870–945 m, 24 Feb. 1973; BSKU 22761 (1), Tosa Bay, 32.9467°N, 133.5567°E, 714–735 m, 24 Jul. 1974; BSKU 44293 (1), BSKU 44294 (1), BSKU 44295 (1), Tosa Bay, 600 m, 17 Nov. 1987; BSKU 14205 (1), off east coast of Japan, date unknown; CBM-ZF 3092 (1), southern Japan, date unknown.

***Hymenocephalus striatissimus*.** 455 specimens: BSKU 26328 (1), off Senkaku Islands, Okinawa Trough, 25.6283°N, 122.8933°E, 560–692 m, 14 Jan. 1978; ZMUT 53496 (1), ZMUT 53501 (1), Ishigaki-jima Island, date unknown; BSKU 27412 (1), west of Okinawa-jima Island, Okinawa Trough, 27.4000°N, 126.1333°E, 340–440 m, 6 Mar. 1978; KPM-NI 28343 (1), ca. 9 km east of Aguni-jima Island, East China Sea, 26.5700°N, 127.3307°E, 583–589 m, 17 Dec. 2008; BSKU 27294 (1), BSKU 27295 (1), BSKU 27296 (1), BSKU 27297 (1), BSKU 27298 (1), BSKU 27299 (1), BSKU 27300 (1), BSKU 27301 (1), BSKU 27302 (1), BSKU 27303 (1), BSKU 27304 (1), BSKU 27305 (1), BSKU 27306 (1), BSKU 27307 (1), BSKU 27308 (1), BSKU 27309 (1), BSKU 27310 (1), BSKU 27311 (1), BSKU 27312 (1), BSKU 27313 (1), BSKU 27314 (1), BSKU 27315 (1), BSKU 27316 (1), BSKU 27317 (1), BSKU 27318 (1), BSKU 27319 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.8033°N, 127.0617°E, 425 m, 6 Feb. 1978; BSKU 27905 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.7500°N, 127.1167°E, 520–542 m, 16 Mar. 1978; SNFR 11250 (2), East China Sea, 29.3900°N, 127.4300°E, 410–416 m, 24 Oct. 1990; BSKU 27175 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.4333°N, 127.5167°E, 570 m, 5 Feb. 1978; BSKU 28388 (1), BSKU 28391 (1), BSKU 28396 (1), BSKU 28397 (1), BSKU 28399 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7333°N, 127.8000°E, 495–510 m, 19 Mar. 1978; FFNU P-00585 (5), southwest of Daini-Danjo Knoll, East China Sea, 31.1487°N, 128.2227°E, 345 m, 24 Apr. 2005; FFNU P-00578 (1), south of Danjo Islands, East China Sea, 31.6248°N, 128.4040°E, 354 m, 24 Apr. 2005; FFNU P-00796 (14 of 18), west of Daini-Danjo Knoll, East China Sea, 31.3645°N, 128.4457°E, 448 m, 8 Jun. 2009; BSKU 22858 (1), BSKU 22859 (1), off Kagoshima, East China Sea, 31.3050°N, 128.5900°E, 460–512 m, 5 Aug. 1974; BSKU 106842 (1), BSKU 106843 (1), southeast of

Shimokoshiki-jima Island, East China Sea, 31.5675°N, 129.8930°E, 393 m, 24 Apr. 2012; MUFS 8526 (22), Aoshima, Miyazaki, 300–350 m, 19 Apr. 1987; ZMUT 50680 (1), Totoro, Miyazaki, 300–400 m, 9 Sept. 1959; FAKU 102514 (16), off Yawatahama, Hyuga-nada, 11 Nov. 1980; KPM-NI 28795 (1), KPM-NI 28796 (1), KPM-NI 28797 (1), KPM-NI 28798 (1), ca. 24 km east-southeast of Fuka-shima Island, Miyazaki, Bungo Channel, 32.6207°N, 132.1515°E, 550–587 m, 16 Aug. 2010; BSKU 43409 (1), BSKU 43411 (1), BSKU 43413 (1), BSKU 43414 (1), BSKU 43415 (1), BSKU 43416 (1), BSKU 43417 (1), south of Okino-shima Island, 32.6400°N, 132.5367°E, 440–451 m, 1 Nov. 1986; BSKU 13231 (1), BSKU 13232 (1), BSKU 13233 (1), BSKU 13262 (1), BSKU 13263 (1), BSKU 13264 (1), BSKU 13265 (1), BSKU 13267 (1), ca. 15 km east of Cape Ashizurimisaki, Tosa Bay, 420–555 m, 2–6 Jun. 1968; BSKU 29515 (1), Tosa Bay, 32.9833°N, 133.4833°E, 453 m, 21 Dec. 1979; BSKU 30561 (1), BSKU 30562 (1), Tosa Bay, 33.0833°N, 133.4833°E, 310 m, 20 Dec. 1979; BSKU 23104 (1), BSKU 23205 (1), BSKU 23206 (1), Tosa Bay, 32.9233°N, 133.4850°E, 480 m, 3 Jan. 1975; BSKU 30261 (1), Mimase fish market, 350 m, 26 Jan. 1980; BSKU 75908 (1), Mimase fish market, 1 Dec. 2004; BSKU 9686 (1), BSKU 9687 (1), BSKU 9688 (1), Mimase fish market, 1 May 1949; BSKU 10039 (1), BSKU 10040 (1), Mimase fish market, 10 Feb. 1996; BSKU 55646 (1), Mimase fish market, 10 Nov. 1995; BSKU 62515 (1), BSKU 62516 (1), Mimase fish market, 11 Dec. 2002; BSKU 71393 (1), Mimase fish market, 12 Apr. 2004; BSKU 72285 (1), Mimase fish market, 13 Apr. 2004; BSKU 71692 (6), Mimase fish market, 13 Feb. 2004; BSKU 42766 (1), Mimase fish market, 13 Mar. 1986; BSKU 82836 (1), BSKU 82837 (1), BSKU 82838 (1), BSKU 82839 (1), Mimase fish market, 14 Dec. 1996; BSKU 87834 (1), Mimase fish market, 14 Dec. 2003; BSKU 84435 (1), Mimase fish market, 14 Feb. 1998; BSKU 71548 (1), Mimase fish market, 14 Mar. 2004; BSKU 72260 (1), Mimase fish market, 14 Nov. 2003; BSKU 75646 (1), Mimase fish market, 15 Dec. 2004; BSKU 69151 (1), Mimase fish market, 15 Mar. 2004; BSKU 85983 (1), Mimase fish market, 17 Dec. 1998; BSKU 76000 (1), Mimase fish market, 18 Jan. 2005; BSKU 87875 (1), BSKU 87876 (1), Mimase fish market, 18 Nov. 2005; BSKU 53119 (1), Mimase fish market, 19 Dec. 2000; BSKU 69236 (1), Mimase fish market, 19 Nov. 2003; BSKU 72774 (5), Mimase fish market, 20 Feb. 2004; BSKU 78084 (7), Mimase fish market, 20 Nov. 2005; BSKU 69591 (1), BSKU 69592 (1), BSKU 69593 (1), Mimase fish market, 21 Dec. 2003; BSKU 69802 (1), Mimase fish market, 22 Dec. 2003; BSKU 92087 (1), Mimase fish market, 23 Apr. 2007; BSKU 54132 (1), BSKU 54133 (1), BSKU 54136 (1), Mimase fish market, 23 Nov. 2000; BSKU 75641 (8), Mimase fish market, 23 Nov. 2004; BSKU 58646 (1), Mimase fish market, 24 Jan. 2002; BSKU 75901 (1), Mimase fish

market, 24 Nov. 2004; BSKU 76516 (1), Mimase fish market, 25 Nov. 2004; BSKU 70019 (1), Mimase fish market, 26 Feb. 2003; BSKU 89531 (1), Mimase fish market, 26 Mar. 2006; BSKU 91576 (1), Mimase fish market, 26 Mar. 2007; BSKU 82936 (1), BSKU 82937 (1), Mimase fish market, 26 Nov. 1996; BSKU 75929 (1), Mimase fish market, 27 Dec. 2004; BSKU 52586 (1), Mimase fish market, 27 Nov. 2000; BSKU 79887 (1), Mimase fish market, 28 Dec. 2005; BSKU 73880 (1), Mimase fish market, 3 Dec. 2004; BSKU 58652 (1), Mimase fish market, 30 Jan. 2002; BSKU 53702 (1), BSKU 53703 (1), Mimase fish market, 30 Nov. 1995; BSKU 70451 (1), BSKU 70452 (1), Mimase fish market, 4 Dec. 2003; BSKU 64821 (1), Mimase fish market, 6 Apr. 2003; BSKU 30177 (1), BSKU 30178 (1), Mimase fish market, 8 Dec. 1978; BSKU 58318 (1), BSKU 58319 (1), BSKU 58320 (1), BSKU 58321 (1), BSKU 58322 (1), BSKU 58323 (1), BSKU 58324 (1), Mimase fish market, 8 Dec. 2000; BSKU 51458 (1), Mimase fish market, 9 Apr. 2000; BSKU 69527 (1), Mimase fish market, 9 Jan. 2004; BSKU 70971 (1), Mimase fish market, 9 Nov. 2003; BSKU 8318 (1), Mimase fish market, 5 Jan. 1951; BSKU 89725 (1), BSKU 89727 (1), Mimase fish market, date unknown; BSKU 36798 (1), Mimase fish market, 11 Mar. 1982; BSKU 12782 (1), BSKU 12784 (1), BSKU 12787 (1), BSKU 12790 (1), BSKU 12794 (1), BSKU 12796 (1), BSKU 12798 (1), BSKU 12802 (1), Mimase fish market, date unknown; BSKU 95077 (1), BSKU 95080 (1), BSKU 95083 (1), Mimase fish market, 30 Nov. 2007; BSKU 95065 (5), Mimase fish market, 18 Jan. 2008; KAUM-I. 12930 (1), Mimase fish market, 9 May 2008; BSKU 30034 (1), off Sukumo, Kochi, 350 m, 1 May 1980; BSKU 12933 (1), BSKU 12935 (1), BSKU 12936 (1), BSKU 12937 (1), BSKU 12939 (1), BSKU 12940 (1), Tosa Bay, 350–450 m, 14 May 1968; BSKU 40338 (1), BSKU 40339 (1), BSKU 40340 (1), BSKU 40341 (1), BSKU 40342 (1), BSKU 40343 (1), BSKU 40345 (1), BSKU 40346 (1), Tosa Bay, 400 m, 9 Oct. 1984; BSKU 41907 (1), BSKU 41908 (1), BSKU 41910 (1), BSKU 41911 (1), BSKU 41912 (1), BSKU 41913 (1), Tosa Bay, 420–450 m, 6 Jun. 1985; BSKU 98925 (1), BSKU 98926 (1), BSKU 98927 (1), BSKU 98928 (1), BSKU 98929 (1), BSKU 98930 (1), Tosa Bay, 33.1428°N, 133.6248°E, 450–490 m, 1 Aug. 1988; BSKU 77980 (1), Tosa Bay, 33.2000°N, 133.6317°E, 370–380 m, 8 Sept. 1998; BSKU 77984 (1), Tosa Bay, 33.1800°N, 133.6350°E, 408–458 m, 29 Jun. 1998; BSKU 77900 (7), Tosa Bay, 33.2083°N, 133.6433°E, 364–382 m, 27 Sept. 1999; BSKU 70908 (1), Tosa Bay, 33.2250°N, 133.6933°E, 440–460 m, 24 Aug. 2000; BSKU 77292 (1), BSKU 77293 (1), BSKU 77295 (1), BSKU 77296 (1), BSKU 77297 (1), BSKU 77298 (1), BSKU 77299 (1), BSKU 77300 (1), BSKU 77301 (1), BSKU 77302 (1), south of Muroto, 33.0790°N, 134.1645°E, 414–513 m, 17 Nov. 2005; BSKU 38816 (1), Owase fish market, Kumano-nada, 12 Mar. 1983; BSKU 66859 (1), BSKU 66860 (1), Owase fish market, Kumano-nada, 16 Sept. 2003; FAKU 20950 (1), FAKU 20951 (1), Owase fish market, Kumano-nada, 5 Nov. 1953; FAKU 15146 (1), Owase fish market, Kumano-nada, 18–20 Jan. 1950; FAKU 24953 (1), FAKU 24955 (1), FAKU 24965 (1), FAKU 24968 (1), Owase fish market, Kumano-nada, 25 Oct. 1954; FRLM 34279 (1), FRLM 34280 (1), off Owase, Kumano-nada, 14 Jan. 2008; FRLM 34467 (1), FRLM 34468 (1), off Owase, Kumano-nada, 18 Feb. 2008; FRLM 34340 (1), FRLM 34341 (1), off Owase, Kumano-nada, 30 Jan. 2008; FRLM 33828 (1), off Owase, Kumano-nada, 31 Nov. 2007; FRLM 30751 (1), FRLM 30752 (1), FRLM 30766 (1), off Owase, Kumano-nada, 2 Feb. 2004; FAKU 4267 (1), FAKU 4268 (1), FAKU 4270 (1), FAKU 4271 (1), FAKU 4272 (1), FAKU 4276 (1), Kumano-nada, Jan. 1937; NSMT-P 101353 (2), southwest of Cape Daiozaki, Kumano-nada, 34.0836°N, 136.6554°E, 495–515 m, 12 Dec. 1995; NSMT-P 94605 (3), south of Daiozaki, Kumano-nada, 422–425 m, 28 May 1986; FAKU 17733 (1), Miya fish market, Enshu-nada, 15 Mar. 1952; FAKU 102507 (56), Maisaka fish market, Enshu-nada, Mar. 1941; BSKU 110044 (1), BSKU 110055 (1), BSKU 110056 (1), Suruga Bay, 34.7127°N, 138.4553°E, 262–434 m, 23 Apr. 2013; BSKU 110411 (1), BSKU 110412 (1), BSKU 110413 (1), Suruga Bay, 34.7481°N, 138.4652°E, 243–424 m, 23 Apr. 2013; BSKU 110102 (1), BSKU 110103 (1), Suruga Bay, 34.7489°N, 138.4664°E, 200–450 m, 23 Apr. 2013; BSKU 110423 (1), Suruga Bay, 34.7651°N, 138.4880°E, 200–450 m, 23 Apr. 2013; KPM-NI 24984 (1), KPM-NI 24985 (1), KPM-NI 24986 (1), eastern part of Suruga Bay, Nov. 2009; FAKU M7416 (1), FAKU M7417 (1), FAKU M7421 (1), FAKU M7424 (1), FAKU M7426 (1), FAKU M7429 (1), FAKU M7431 (1), FAKU M7439 (1), FAKU M7440 (1), FAKU M7442 (1), FAKU M7445 (1), FAKU M7446 (1), FAKU M7450 (1), FAKU M7453 (1), FAKU M7456 (1), FAKU M7460 (1), FAKU M7461 (1), FAKU M7462 (1), FAKU M7463 (1), FAKU M7466 (1), FAKU M7467 (1), FAKU M7468 (1), FAKU M7470 (1), FAKU M7471 (1), FAKU M7473 (1), FAKU M7474 (1), FAKU M7476 (1), FAKU M7478 (1), FAKU M7485 (1), FAKU M7488 (1), FAKU M7489 (1), FAKU M7490 (1), FAKU M7492 (1), FAKU M7494 (1), FAKU M7495 (1), FAKU M7497 (1), Heta fish market, Suruga Bay, 150 fms (275 m), 23 Jul. 1938; KPM-NI 8633 (1), off Cape Irozaki, Izu Peninsula, 150–220 m, 22 Apr. 2001; BSKU 101972 (2), off Hitachi, 36.6143°N, 141.0965°E, 244–262 m, 9 Nov. 1995; NSMT-P 48956 (5), off Iwaki, 37.0725°N, 141.5110°E, 271–303 m, 4 Nov. 1995; NSMT-P 97760 (1), off Tsushima Island, date unknown; ZMUT KP-417 (1), Kyushu-Palau Ridge, 27.9178°N, 134.7460°E, 685–710 m, 19 Dec. 1979; BSKU 99099 (1), off Hong Kong, South China Sea, 19.5500°N, 113.8667°E, 21 May 1990.

Hymenogadus gracilis. 57 specimens: BSKU 26598 (1), BSKU 26599 (1), off Senkaku Islands, Okinawa Trough, 26.3883°N, 125.0667°E, 295–385 m, 24 Jan. 1978;

NSMT-P 65633 (1), west of Yaku-shima Island, Okinawa Trough East China Sea, 30.4450°N, 127.8450°E, 299–302 m, 10 Nov. 2002; BSKU 80770 (1), Mimase fish market, 1 Dec. 1992; BSKU 52909 (1), Mimase fish market, 10 Dec. 2000; BSKU 70635 (1), Mimase fish market, 12 Jan. 2004; BSKU 75648 (1), Mimase fish market, 15 Dec. 2004; BSKU 44375 (1), Mimase fish market, 19 Dec. 1987; BSKU 49862 (1), Mimase fish market, 19 Nov. 1991; BSKU 4973 (1), Mimase fish market, 1955; BSKU 40056 (1), Mimase fish market, 26 Apr. 1984; BSKU 49879 (1), Mimase fish market, 28 Nov. 1991; BSKU 40739 (1), Mimase fish market, 30 Nov. 1984; BSKU 64822 (1), Mimase fish market, 6 Apr. 2003; BSKU 64307 (1), Mimase fish market, 7 Feb. 2003; BSKU 62251 (1), Mimase fish market, 9 Jan. 2003; BSKU 41720 (1), off Okitsu, Tosa Bay, 240 m, 24 Apr. 1985; BSKU 41687 (1), BSKU 41688 (1), off Okitsu, Tosa Bay, 250–260 m, 18 Apr. 1985; BSKU 3517 (1), BSKU 3518 (1), Urado fish market, 1953; BSKU 60757 (1), Tosa Bay, 250 m, 14 Nov. 1997; BSKU 44550 (1), Tosa Bay, 250 m, 19 Jan. 1987; BSKU 99069 (1), BSKU 99070 (1), Tosa Bay, 250 m, 23 Aug. 2004; BSKU 11392 (1), BSKU 11393 (1), Tosa Bay, date unknown; BSKU 22590 (1), BSKU 22591 (1), BSKU 22592 (1), Tosa Bay, 21 Jul. 1974; NSMT-P 59309 (1), Tosa Bay, 250 m, 25 Jan. 1998; SNFR 18033 (5), Tosa Bay, 11 Apr. 1988; FAKU M7517 (1), FAKU M7518 (1), FAKU M7519 (1), FAKU M7527 (1), Owase fish market, Kumano-nada, 4–9 Jan. 1936; MSM 13-15 (1), southwest of Toi, Suruga Bay, 34.8867°N, 138.7416°E, 182–359 m, 21 Feb. 2012.

Kumba japonica. 30 specimens: BSKU 27291 (1), west-northwest of Amami-oshima Island, Okinawa Trough, 28.8033°N, 127.0617°E, 425 m, 6 Feb. 1978; NSMT-P 120917 (2), off Miyazaki, Hyuga-nada, 484–565 m, 19 May 1989; BSKU 43317 (1), Mimase fish market, 18 Oct. 1986; BSKU 44190 (1), Mimase fish market, 7 Nov. 1987; BSKU 8319 (1), Mimase fish market, 5 Jan. 1959; BSKU 12959 (1), Tosa Bay, 350–450 m, 14 May 1968; NSMT-P 96430 (4), southeast of Miyake-jima Island, Shichito-Iojima Ridge, 34.1287°N, 139.6413°E, 484–565 m, 19 May 1989; NSMT-P 65441 (1), off Hitachi, 36.4722°N, 140.9705°E, 213–272 m, 24 Oct. 2002; NSMT-P 65731 (1), off Hitachi, 36.5118°N, 141.0147°E, 309–321 m, 26 Oct. 2002; NSMT-P 102728 (1), off Minami-soma, 37.6128°N, 141.7277°E, 309–313 m, 29 Oct. 2006; NSMT-P 92271 (1), off Miyagi, 25 Oct. 2008; NSMT-P 91535 (1), east of Oshika Peninsula, 38.3400°N, 142.0283°E, 351 m, 24 Oct. 2007; NSMT-P 65733 (1), east of Oshika Peninsula, 38.3345°N, 142.0718°E, 412–452 m, 28 Oct. 2002; NSMT-P 58943 (1), off Pacific coast of Tohoku, 39.0683°N, 143.5500°E, 550–579 m, 30 Jul. 1996; BSKU 26066 (1), Kyushu-Palau Ridge, 27.9167°N, 134.6500°E, 700 m, 20 Feb. 1978; BSKU 26029 (1), Kyushu-Palau Ridge, 27.9000°N, 134.6550°E, 650 m, 15 Feb. 1978; BSKU 29332 (1), BSKU 32215 (1), BSKU 32218 (1), BSKU 32221 (1), BSKU 32222 (1), Kyushu-

Palau Ridge, 28.1117°N, 134.6656°E, 550–600 m, 19 Dec. 1979; BSKU 29382 (1), Kyushu-Palau Ridge, 27.9186°N, 134.7467°E, 685–710 m, 19 Dec. 1979; BSKU 14174 (1), off Hachijo-jima Island, Shichito-Iojima Ridge, Apr. 1968; BSKU 19309 (1), BSKU 19310 (1), east of Hachijo Island, Shichito-Iojima Ridge, 33.1367°N, 139.9900°E, 440 m, 8 Dec. 1962; BSKU 19759 (1), east of Hachijo Island, Shichito-Iojima Ridge, 33.1617°N, 140.0000°E, 525 m, 13 Nov. 1966.

Lucigadus nigromarginatus. 5 specimens: KAUM-I. 44836 (1), Da-xi fish market, Yilan, 6 Jul. 2011; FMNH 120866 (1), Dong-gang fish market, Pingtung, 12 Sept. 2009; FRLM 32268 (1), Dong-gang, Pingtung, 22 May 2005; KAUM-I. 44652 (1), KAUM-I. 44653 (1), off Kaohsiung, 400 m, 1 Jul. 2011.

Malacocephalus nipponensis. 63 specimens: BSKU 29587 (1), northwest of Miyako-jima Island, Okinawa Trough, 25.7917°N, 124.5117°E, 500–550 m, 15 Sept. 1979; BSKU 32507 (1), northwest of Miyako-jima Island, Okinawa Trough, 25.7917°N, 124.5117°E, 500–550 m, 10 Oct. 1979; BSKU 33082 (1), BSKU 33083 (1), north-northeast of Miyako-jima Island, Okinawa Trough, 25.9833°N, 125.8500°E, 430 m, 10 Oct. 1979; BSKU 27469 (1), northwest of Okinawa-jima Island, Okinawa Trough, 27.7667°N, 126.2833°E, 490 m, 9 Mar. 1978; BSKU 27320 (1), BSKU 27321 (1), BSKU 27322 (1), west-northwest of Amami-oshima Island, Okinawa Trough, 28.8033°N, 127.0617°E, 425 m, 6 Feb. 1978; BSKU 27693 (1), west-northwest of Amami-oshima Island, Okinawa Trough, 28.7000°N, 127.1500°E, 500–535 m, 11 Mar. 1978; SNFR 16337 (1), East China Sea, 29.0053°N, 127.2390°E, 503 m, 9 Nov. 2003; BSKU 28204 (1), BSKU 28205 (1), BSKU 28206 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.3000°N, 127.3333°E, 410–415 m, 17 Mar. 1978; KPM-NI 6214 (1), KPM-NI 6215 (1), KPM-NI 6217 (1), KPM-NI 6219 (1), ca. 230 km west-northwest of Amami-oshima Island, East China Sea, 29.3850°N, 127.4050°E, 400–432 m, 3 Oct. 1997; BSKU 28316 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7167°N, 127.6167°E, 410–423 m, 18 Mar. 1978; MUFS 39534 (1), Totoro, Nobeoka, Miyazaki, 27 Oct. 1977; BSKU 40015 (1), off Okino-shima Island, Tosa Bay, 250 m, 18 Apr. 1984; BSKU 10027 (1), BSKU 10028 (1), BSKU 10030 (1), BSKU 10031 (1), BSKU 10032 (1), Mimase fish market, 10 Feb. 1961; BSKU 3560 (1), Mimase fish market, 14 Feb. 1954; BSKU 220 (1), Mimase fish market, 28 Jan. 1951; BSKU 45340 (1), Mimase fish market, 9 Nov. 1988; FAKU 21779 (1), Mimase fish market, 10–20 Apr. 1954; FAKU 29123 (1), FAKU 29124 (1), FAKU 29125 (1), FAKU 29126 (1), FAKU 29127 (1), FAKU 29128 (1), FAKU 29129 (1), Mimase fish market, 20 Mar. 1960; BSKU 13636 (1), Mimase fish market, Jan.–Mar. 1968; BSKU 1822 (1), Mimase fish market, Mar. 1952; BSKU 13318 (1), BSKU 13322 (1), BSKU 13324 (1), BSKU 13325 (1), BSKU 13341 (1), ca. 10–15 km west

of Cape Murotomisaki, Tosa Bay, 360–435 m, 8–9 Jun. 1968; FAKU 34498 (1), FAKU 34499 (1), FAKU 34500 (1), FAKU 34501 (1), FAKU 34502 (1), Owase fish market, Kumano-nada, 12–14 Nov. 1961; FAKU 27644 (1), FAKU 27645 (1), Owase fish market, Kumano-nada, 25 Feb.–10 Mar. 1957; FAKU 21528 (1), Owase fish market, Kumano-nada, 15 Feb. 1954; FAKU 23807 (1), Owase fish market, Kumano-nada, 20 Oct. 1954; FRLM 40506 (1), off Owase, Kumano-nada, 8 Dec. 2011; FRLM 34378 (1), off Owase, Kumano-nada, date unknown; FAKU 4196 (1), Kumano-nada, Jan. 1937; FRLM 8587 (1), FRLM 8588 (1), FRLM 8589 (1), Kumano-nada, date unknown; FAKU 34116 (1), FAKU 34118 (1), Miya fish market, Enshu-nada, Apr. 1961; FAKU 28398 (1), Miya fish market, Enshu-nada, 5–7 Feb. 1959; FAKU 8008 (1), off Choshi, 10 Apr. 1939; FAKU 5174 (1), off Choshi, 20–26 Nov. 1937.

Nezumia condylura. 251 specimens: BSKU 26288 (1), BSKU 26289 (1), northwest of Yonaguni-jima Island, Okinawa Trough, 24.6833°N, 122.5933°E, 510–615 m, 13 Jan. 1978; BSKU 26429 (1), north of Miyako-jima Island, Okinawa Trough, 26.3350°N, 124.8450°E, 1000–1140 m, 22 Jan. 1978; BSKU 26552 (1), BSKU 26553 (1), BSKU 26554 (1), off Miyako-jima Island, Okinawa Trough, 26.4633°N, 124.8483°E, 680–770 m, 23 Jan. 1978; BSKU 26601 (1), BSKU 26602 (1), off Miyako-jima Island, Okinawa Trough, 26.3883°N, 125.0667°E, 680–770 m, 24 Jan. 1978; BSKU 27857 (1), BSKU 27864 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.5833°N, 127.0000°E, 600 m, 16 Mar. 1978; BSKU 27902 (1), BSKU 27903 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.7500°N, 127.1167°E, 520–542 m, 16 Mar. 1978; BSKU 27581 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.7167°N, 127.2333°E, 750–755 m, 11 Mar. 1978; NSMT-P 102110 (1), west-northwest of Amami-oshima Island, East China Sea, 29.0053°N, 127.2390°E, 503 m, 8 Nov. 2003; BSKU 28082 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.8833°N, 127.3000°E, 820–830 m, 16 Mar. 1978; BSKU 27187 (1), BSKU 27195 (1), BSKU 27196 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.4333°N, 127.5167°E, 570 m, 5 Feb. 1978; BSKU 26939 (1), BSKU 26940 (1), BSKU 26941 (1), BSKU 26942 (1), BSKU 26943 (1), BSKU 26944 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.6483°N, 127.9333°E, 750–760 m, 4 Feb. 1978; BSKU 28445 (1), BSKU 28447 (1), BSKU 28448 (1), BSKU 28449 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7667°N, 127.9833°E, 713–770 m, 20 Mar. 1978; BSKU 28502 (1), BSKU 28503 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7333°N, 128.0500°E, 815 m, 20 Mar. 1978; BSKU 28551 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.8167°N, 128.1000°E, 915–932 m, 20 Mar. 1978; BSKU 29888 (1), west of Yaku-shima Island, Okinawa Trough, 30.2083°N, 128.2088°E, 600–620 m, 26 Oct. 1979; BSKU 33451 (1),

BSKU 33514 (1), BSKU 33515 (1), west of Yaku-shima Island, Okinawa Trough, 30.2067°N, 128.2833°E, 780–810 m, 26 Oct. 1979; FFNU P-00715 (2), west of Amakusa Knoll, East China Sea, 32.1593°N, 129.4715°E, 503 m, 12 Mar. 2008; NSMT-P 94557 (2), off Yaku-shima and Tanega-shima Islands, 660–711 m, 17 Feb. 1992; KPM-NI 28794 (1), ca. 24 km east-southeast of Fuka-shima Island, Miyazaki, Bungo Channel, 32.6207°N, 132.1515°E, 550–587 m, 16 Aug. 2010; BSKU 13110 (1), BSKU 13111 (1), BSKU 13112 (1), BSKU 13229 (1), BSKU 13230 (1), ca. 15 km east of Cape Ashizurimisaki, Tosa Bay, 420–555 m, 2–6 Jun. 1968; BSKU 13040 (1), BSKU 13041 (1), BSKU 13042 (1), BSKU 13043 (1), BSKU 13044 (1), BSKU 13045 (1), BSKU 13046 (1), BSKU 13047 (1), BSKU 13048 (1), BSKU 13050 (1), BSKU 13051 (1), BSKU 13053 (1), BSKU 13054 (1), BSKU 13056 (1), BSKU 13057 (1), BSKU 13058 (1), BSKU 13059 (1), BSKU 13060 (1), BSKU 13061 (1), BSKU 13062 (1), BSKU 13063 (1), BSKU 13064 (1), BSKU 13065 (1), BSKU 13066 (1), BSKU 13067 (1), BSKU 13069 (1), BSKU 13070 (1), BSKU 13071 (1), off Susaki, Tosa Bay, 430–460 m, 22 May 1968; BSKU 23208 (1), BSKU 23209 (1), Tosa Bay, 32.9233°N, 133.4850°E, 489 m, 30 Jan. 1975; BSKU 30499 (1), BSKU 30500 (1), BSKU 30502 (1), BSKU 30503 (1), BSKU 30504 (1), BSKU 30505 (1), BSKU 30506 (1), BSKU 30507 (1), BSKU 29498 (1), Tosa Bay, 32.9667°N, 133.5333°E, 605 m, 21 Dec. 1979; BSKU 30313 (1), BSKU 30314 (1), BSKU 30315 (1), BSKU 30316 (1), BSKU 30317 (1), BSKU 30318 (1), BSKU 30319 (1), BSKU 30320 (1), BSKU 30322 (1), BSKU 30323 (1), BSKU 30324 (1), BSKU 30325 (1), BSKU 30326 (1), BSKU 30327 (1), BSKU 30328 (1), BSKU 30329 (1), BSKU 30330 (1), Tosa Bay, 32.9167°N, 133.5500°E, 700 m, 21 Dec. 1979; BSKU 22778 (1), BSKU 22779 (1), Tosa Bay, 32.9433°N, 133.5517°E, 707–750 m, 24 Jul. 1974; ZMUT 30609 (1), Mimase fish market, 10 Jan. 1935; BSKU 22768 (1), BSKU 22769 (1), BSKU 22770 (1), Tosa Bay, 32.9467°N, 133.5750°E, 714–735 m, 24 Jul. 1974; BSKU 44832 (1), BSKU 44833 (1), BSKU 44834 (1), BSKU 44835 (1), BSKU 44836 (1), Tosa Bay, 32.9886°N, 133.5978°E, 700–720 m, 21 Jun. 1988; BSKU 12956 (1), BSKU 12957 (1), BSKU 12958 (1), BSKU 12918 (1), Tosa Bay, 350–450 m, 14 May 1968; BSKU 99123 (1), BSKU 99124 (1), BSKU 99125 (1), BSKU 99127 (1), BSKU 99128 (1), BSKU 99130 (1), Tosa Bay, 600 m, 17 Nov. 1987; BSKU uncat. (1), Tosa Bay, date unknown; BSKU 69222 (1), BSKU 69313 (1), BSKU 69314 (1), BSKU 69315 (1), Tosa Bay, 33.1800°N, 133.6217°E, 412–423 m, 8 Sept. 1998; BSKU 98940 (1), BSKU 98943 (1), BSKU 98944 (1), BSKU 98945 (1), BSKU 98946 (1), BSKU 98947 (1), BSKU 98948 (1), BSKU 98949 (1), BSKU 98950 (1), BSKU 98951 (1), BSKU 98952 (1), BSKU 99121 (1), BSKU 99122 (1), Tosa Bay, 33.1428°N, 133.6248°E, 450–490 m, 1 Aug. 1988; BSKU 99120 (1), Tosa Bay, 33.1233°N, 133.6400°E, 602–631 m, 10 Sept. 1998;

BSKU 69276 (1), Tosa Bay, 33.1417°N, 133.6433°E, 582–597 m, 16 Jun. 1999; BSKU 70892 (1), BSKU 70893 (1), Tosa Bay, 33.2250°N, 133.6933°E, 440–460 m, 24 Aug. 2000; BSKU 13327 (1), BSKU 13328 (1), BSKU 13329 (1), BSKU 13330 (1), BSKU 13335 (1), BSKU 13336 (1), BSKU 13337 (1), BSKU 13352 (1), BSKU 13353 (1), BSKU 13354 (1), ca. 10–15 km west of Cape Murotomisaki, Tosa Bay, 360–435 m, 8–9 Jun. 1968; BSKU 13475 (1), ca. 10–15 km west of Murotomisaki, Tosa Bay, 360–435 m, 8–9 Jun. 1968; BSKU 99105 (1), BSKU 99106 (1), BSKU 99107 (1), BSKU 99108 (1), BSKU 99109 (1), BSKU 99110 (1), south of Muroto, 33.0790°N, 134.1645°E, 414–513 m, 17 Nov. 2005; KPM-NI 28850 (1), ca. 24 km west-southwest of Shirahama, Wakayama, 33.5795°N, 135.0995°E, 532 m, 15 Aug. 2010; FAKU 1945 (1), FAKU 1948 (1), FAKU 1949 (1), FAKU 1973 (1), Owase fish market, Kumano-nada, 4–9 Jan. 1936; FAKU 4929 (1), Kumano-nada, 1938; KPM-NI 28842 (1), ca. 22 km south-southeast of Mie-shima Island, Kumano-nada, 34.0713°N, 136.6453°E, 606–669 m, 14 Aug. 2010; NSMT-P 101355 (2), southwest of Cape Daiozaki, Kumano-nada, 34.0836°N, 136.6554°E, 495–515 m, 12 Dec. 1995; NSMT-P 94610 (4), south of Daiozaki, Kumano-nada, 505–533 m, 20 Aug. 1988; NSMT-P 94616 (4), south of Daiozaki, Kumano-nada, 25 May 1994; FAKU 28404 (1), FAKU 28405 (1), FAKU 28406 (1), FAKU 28407 (1), Miya fish market, Enshu-nada, 5–7 Feb. 1959; BSKU 98914 (1), Suruga Bay, 29 Oct. 1972; NSMT-P 100454 (1), west-southwest of Matsuzaki, Suruga Bay, 34.7483°N, 138.6717°E, 570–597 m, 3 Jul. 1980; NSMT-P 101396 (11), west of Cape Hagachizaki, Suruga Bay, 34.6817°N, 138.6817°E, 418–457 m, 11 Nov. 1983; FAKU M6863 (1), FAKU M6865 (1), FAKU M6872 (1), FAKU M6880 (1), FAKU M7393 (1), FAKU M7397 (1), FAKU M7399 (1), FAKU M7403 (1), Heta fish market, Suruga Bay, 150 fthm (275 m), 23 Jul. 1938; FAKU M6943 (1), Heta fish market, Suruga Bay, 200–290 fthm (366–531 m), 24–25 Jun. 1939; NSMT-P 101336 (1), east of Izu Peninsula, Sagami-nada, 34.7683°N, 139.1417°E, 574–670 m, 9 Mar. 1978; BSKU 20247 (1), BSKU 20248 (1), west of Misaki, Sagami Bay, 35.1283°N, 139.4983°E, 450 m, 30 Jun. 1972; NSMT-P 97206 (1), east of Boso Peninsula, 35.1048°N, 140.8367°E, 637–673 m, 25 Apr. 1995; ZMUT 30610 (1), Choshi, 27 Jan. 1935; ZMUT 25607 (1), Onahama, Fukushima, 400–500 m, 2 May 1933; NSMT-P 77294 (1), off Hitachi, 36.5150°N, 141.0517°E, 407–413 m, 13 Nov. 2006; NSMT-P 77295 (3), off Hitachi, 36.5157°N, 141.0517°E, 407–413 m, 13 Nov. 2006; NSMT-P 48958 (1), off Hitachi, 36.5670°N, 141.1085°E, 412–421 m, 9 Nov. 1995; BSKU 99080 (1), BSKU 99081 (1), BSKU 99082 (1), BSKU 99083 (1), BSKU 99084 (1), BSKU 99085 (1), BSKU 99086 (1), BSKU 99087 (1), BSKU 99088 (1), off Mito, 550 m, 15 Oct. 1997; BSKU 101995 (1), BSKU 101996 (1), off Iwaki, 36.9162°N, 141.5995°E, 615–633 m, 4 Nov. 1995; NSMT-P 102637 (1), off Iwaki,

36.9690°N, 141.6327°E, 554–558 m, 28 Oct. 2006; BSKU 83761 (1), BSKU 83762 (1), off Fukushima, 450 m, 6 Oct. 1997; BSKU 56985 (1), BSKU 56986 (1), BSKU 56987 (1), BSKU 56988 (1), BSKU 56989 (1), BSKU 56990 (1), BSKU 56991 (1), BSKU 56992 (1), BSKU 56993 (1), BSKU 56994 (1), off Fukushima, 5 Oct. 1997; BSKU 99090 (1), BSKU 99091 (1), off Onahama, 450 m, 16 Oct. 1997; NSMT-P 48957 (1), southeast of Oshika Peninsula, 37.9715°N, 141.7822°E, 250–270 m, 2 Nov. 1995; NSMT-P 48968 (2), off Minami-soma, 37.4878°N, 141.8532°E, 411–419 m, 3 Nov. 1995; BSKU 83763 (1), BSKU 83764 (1), off Kinkazan, 450 m, Oct. 1997; BSKU 45443 (1), off Kinkazan, 38.4533°N, 142.1000°E, 522 m, 24 May 1988; BSKU 101969 (1), southeast of Kuji, 39.9997°N, 142.3687°E, 996–1010 m, 2 Nov. 1995; BSKU 30588 (1), Kyushu-Palau Ridge, 27.9000°N, 134.6500°E, 910 m, 18 Dec. 1979; BSKU 26027 (1), Kyushu-Palau Ridge, 27.9000°N, 134.6550°E, 650 m, 15 Feb. 1978; BSKU 26114 (1), Kyushu-Palau Ridge, 27.9033°N, 134.6583°E, 700 m, 20 Feb. 1978.

Nezumia proxima. 132 specimens: FFNU P-00813 (1), Iheya Knolls, East China Sea, 27.5683°N, 126.3050°E, 1194 m, 25 Jun. 2009; BSKU 28071 (1), BSKU 28072 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.8833°N, 127.3000°E, 820–830 m, 16 Mar. 1978; BSKU 26945 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.6483°N, 127.9333°E, 750–760 m, 4 Feb. 1978; BSKU 28504 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7333°N, 128.0500°E, 815 m, 20 Mar. 1978; BSKU 28552 (1), BSKU 28553 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.8167°N, 128.1000°E, 915–932 m, 20 Mar. 1978; BSKU 29792 (1), west of Yaku-shima Island, Okinawa Trough, 30.2083°N, 128.2088°E, 600–620 m, 26 Oct. 1979; BSKU 29884 (1), west of Yaku-shima Island, Okinawa Trough, 30.2067°N, 128.2833°E, 780–810 m, 26 Oct. 1979; BSKU 26858 (1), BSKU 26859 (1), BSKU 26860 (1), north-northwest of Amami-oshima Island, Okinawa Trough, 29.7850°N, 128.4467°E, 1000 m, 3 Feb. 1978; NSMT-P 100504 (1), off Miyazaki, Hyuga-nada, 31.8867°N, 131.8700°E, 779–803 m, 26 Jun. 1993; BSKU 29500 (1), Tosa Bay, 32.9667°N, 133.5333°E, 605 m, 21 Dec. 1979; BSKU 30321 (1), Tosa Bay, 32.9167°N, 133.5500°E, 700 m, 21 Dec. 1979; BSKU 99071 (1), Tosa Bay, 33.0867°N, 133.5500°E, 820 m, 22 May 1997; BSKU 23239 (1), off Kochi, 32.9433°N, 133.5517°E, 707–750 m, 24 Jul. 1974; BSKU 22780 (1), Tosa Bay, 32.9433°N, 133.5517°E, 707–750 m, 24 Jul. 1974; BSKU 22765 (1), BSKU 22766 (1), BSKU 22767 (1), Tosa Bay, 32.9467°N, 133.5750°E, 714–735 m, 24 Jul. 1974; BSKU 22793 (1), BSKU 22794 (1), Tosa Bay, 32.8967°N, 133.6217°E, 960–985 m, 24 Jul. 1974; NSMT-P 55418 (1), off Katsurahama, Tosa Bay, 33.0167°N, 133.6417°E, 823–856 m, 22 Aug. 1997; BSKU 98936 (1), Tosa Bay, 33.0452°N, 133.6448°E, 733–740 m, 25 Jan. 1989; SNFR 17973 (1), off Kochi, 700–720 m, 21 Jun. 1988; SNFR 17972 (3), Tosa Bay, 750–780 m,

24 Nov. 1987; BSKU 23144 (1), BSKU 23145 (1), BSKU 23148 (1), Tosa Bay, 33.0350°N, 133.7500°E, 945 m, 26 Jan. 1975; BSKU 23077 (1), BSKU 23078 (1), BSKU 23180 (1), off Kochi, 32.9800°N, 133.8550°E, 1043 m, 26 Jan. 1975; KPM-NI 28840 (1), KPM-NI 28841 (1), ca. 22 km south-southeast of Mie-shima Island, Kumano-nada, 34.0713°N, 136.6453°E, 606–669 m, 14 Aug. 2010; KPM-NI 18268 (1), KPM-NI 18269 (1), KPM-NI 18270 (1), KPM-NI 18272 (1), off Shima Peninsula, Kumano-nada, 34.0103°N, 136.8793°E, 781–818 m, 24 Nov. 2005; KPM-NI 18257 (1), off Shima Peninsula, Kumano-nada, 34.0005°N, 136.8937°E, 771–787 m, 24 Nov. 2005; KPM-NI 18242 (1), off Shima Peninsula, Kumano-nada, 34.0144°N, 136.9054°E, 767–771 m, 1 May 2004; KPM-NI 18235 (1), KPM-NI 18236 (1), off Shima Peninsula, Kumano-nada, 34.0083°N, 136.9071°E, 767–774 m, 1 May 2004; NSMT-P 94363 (1), off Shima, Kumano-nada, 1190–1348 m, 25 May 1994; NSMT-P 100737 (1), NSMT-P 101340 (1), south of Atsumi Peninsula, Enshu-nada, 34.2288°N, 137.3630°E, 993–1004 m, 24 May 1994; NSMT-P 101360 (1), southeast of Yaizu, Suruga Bay, 34.8139°N, 138.4303°E, 865–878 m, 22 May 1994; MSM uncat. (2), Suruga Bay, date unknown; NSMT-P 101341 (2), Suruga Bay, 1978; NSMT-P 101167 (1), west-southwest of Matsuzaki, Suruga Bay, 34.7483°N, 138.6717°E, 570–597 m, 3 Jul. 1980; NSMT-P 101356 (1), west of Toi, Suruga Bay, 34.9167°N, 138.7117°E, 516–630 m, 23 Nov. 1978; NSMT-P 93119 (1), NSMT-P 93120 (1), off Oh-sima Island, Izu Islands, date unknown; NSMT-P 100468 (1), Sagami Bay, 810–850 m, 15 May 1970; BSKU 19776 (1), Suruga Bay, 35.1650°N, 139.4083°E, 525–540 m, 30 Jun. 1967; BSKU 19763 (1), Sagami Bay, 35.1167°N, 139.4500°E, 730–790 m, 23 Nov. 1966; BSKU 20358 (1), Sagami Bay, 35.1683°N, 139.4800°E, 480 m, 26 Nov. 1972; BSKU 14185 (1), BSKU 14187 (1), off Hiratsuka, Sagami Bay, 35.2522°N, 139.5178°E, 510–600 m, 12 Apr. 1969; NSMT-P 99988 (1), east of Boso Peninsula, 35.0802°N, 140.7500°E, 638–704 m, 15 Sept. 1989; NSMT-P 97668 (9), east of Boso Peninsula, 35.1048°N, 140.8367°E, 627–673 m, 25 Apr. 1995; NSMT-P 94347 (1), off Hitachi, 690–705 m, 9 Aug. 1979; BSKU 101194 (1), BSKU 101195 (1), BSKU 101196 (1), BSKU 101197 (1), BSKU 101198 (1), BSKU 101199 (1), off Hitachi, 900 m, 10 Oct. 1997; BSKU 101207 (1), BSKU 101208 (1), BSKU 101209 (1), off Hitachi, 36.9162°N, 141.5995°E, 615–633 m, 4 Nov. 1995; BSKU 101200 (1), BSKU 101201 (1), BSKU 101202 (1), BSKU 101203 (1), BSKU 101204 (1), BSKU 101205 (1), BSKU 101206 (1), off Hitachi, 36.9583°N, 141.6910°E, 777–905 m, 5 Nov. 1995; BSKU 101600 (1), BSKU 101601 (1), off Fukushima, 750 m, 5 Oct. 1997; BSKU 56983 (1), BSKU 56984 (1), off Fukushima, 5 Oct. 1997; BSKU 99089 (1), off Onahama, 450 m, 16 Oct. 1997; BSKU 99079 (1), off Onahama, 900 m, 7 Oct. 1997; BSKU 101184 (1), BSKU 101185 (1), BSKU 101186 (1), BSKU 101187 (1), BSKU 101188 (1), BSKU 101189 (1), off Onahama, 550 m, 8

Oct. 1997; HUMZ 72598 (1), off Fukushima, 37.1833°N, 141.9500°E, 810–820 m, 18 Jan. 1978; BSKU 99136 (1), off Fukushima, 37.4058°N, 141.9782°E, 603–610 m, 3 Nov. 1995; BSKU 99119 (1), off Fukushima, 650 m, 5 Oct. 1997; NSMT-P 94345 (2), off Minami-soma, 37.5367°N, 142.0550°E, 538–561 m, 1 Dec. 1981; NSMT-P 100461 (1), off Kamaishi, 39.2133°N, 142.2667°E, 562–563 m, 14 Aug. 1985; BSKU 87494 (1), off Kesennuma, 38.9300°N, 142.3763°E, 976–1004 m, 28 Oct. 1997; NSMT-P 115471 (4), off Kamaishi, 39.1050°N, 142.3967°E, 926–950 m, 10 Jul. 1984; BSKU 87588 (1), BSKU 99137 (1), BSKU 99138 (1), BSKU 99139 (1), BSKU 99140 (1), BSKU 99141 (1), BSKU 99142 (1), BSKU 99143 (1), BSKU 99144 (1), off Kesennuma, 38.6975°N, 142.4210°E, 986–998 m, 28 Oct. 1995; BSKU 101190 (1), BSKU 101191 (1), BSKU 101192 (1), off Rikuzen-Takata, 900 m, 18 Oct. 1997; BSKU 99131 (1), southeast of Kuji, 40.0013°N, 142.5240°E, 806–822 m, 26 Oct. 1995; BSKU 99100 (1), off Miyako, 900 m, 20 Oct. 1997; BSKU 99132 (1), BSKU 99133 (1), off Miyako, 39.9633°N, 142.6882°E, 993–1007 m, 26 Oct. 1995; NSMT-P 94369 (1), southeast of Otsuchi, 562–563 m, 14 Aug. 1985; NSMT-P 94368 (1), southeast of Otsuchi, 984–988 m, 25 Sept. 1988; NSMT-P 100984 (1), off Yoron-jima Island, East China Sea, 27.1133°N, 128.6983°E, 870–945 m, 24 Feb. 1973.

Pseudonezumia cetonuropsis. 1 specimen: BSKU 47907 (1), Kumano-nada, 33.7085°N, 136.6092°E, 2034–2036 m, 31 Aug. 1990.

Pseudonezumia japonica. 10 specimens: BSKU 49566 (1), BSKU 49571 (1), BSKU 49572 (1), BSKU 49573 (1), BSKU 49574 (1), BSKU 49575 (1), BSKU 49690 (1), off Miyazaki, Hyuga-nada, 32.3083°N, 132.1817°E, 1453–1481 m, 3 Apr. 1991; BSKU 56962 (1), BSKU 56968 (1), BSKU 56970 (1), off Hitachi, 36.4205°N, 141.5420°E, 1845–1948 m, 10 Nov. 1995.

Spicomacrurus kuronumai. 73 specimens: BSKU 28317 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7167°N, 127.6167°E, 410–423 m, 19 Mar. 1978; BSKU 28369 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7333°N, 127.8000°E, 495–510 m, 19 Mar. 1978; SNFR 16342 (2), East China Sea, 30.4833°N, 127.9482°E, 399 m, 7 Nov. 2003; BSKU 22896 (1), off Kagoshima, East China Sea, 31.2617°N, 128.3433°E, 364–369 m, 5 Aug. 1974; BSKU 29812 (1), northwest of Amami-oshima Island, Okinawa Trough, 31.5583°N, 128.4000°E, 400–410 m, 4 Nov. 1979; FFNU P-00579 (1), south of Danjo Islands, East China Sea, 31.6248°N, 128.4040°E, 354 m, 24 Apr. 2005; SNFR 11327 (1), East China Sea, 31.2717°N, 128.4350°E, 479 m, 30 Sept. 1995; FFNU P-00776 (3), east of Hirazi Bank, East China Sea, 32.2485°N, 129.4720°E, 403 m, 7 Mar. 2009; KAUM-I. 32266 (1), off Nomaie, Kagoshima, 31.5000°N, 129.8833°E, 370–400 m, 10 Sept. 2010; BSKU 13234 (1), BSKU 13257 (1), BSKU 13258 (1), east of Cape Ashizurimisaki, Tosa Bay, 420–555 m, 2 Jun. 1968; BSKU 4962 (1), BSKU 4963 (1), BSKU 4965 (1),

BSKU 4966 (1), BSKU 4967 (1), Mimase fish market, 2 Jun. 1968; FAKU 21766 (1), Mimase fish market, 10–20 Apr. 1954; BSKU 12917 (1), Tosa Bay, 350–450 m, 2 Jun. 1968; BSKU 13355 (1), BSKU 13362 (1), BSKU 13364 (1), BSKU 13365 (1), BSKU 13367 (1), BSKU 13370 (1), BSKU 13371 (1), BSKU 13382 (1), BSKU 13384 (1), BSKU 13386 (1), BSKU 13388 (1), BSKU 13390 (1), BSKU 13391 (1), west of Muroto, Tosa Bay, 360–435 m, 8 Jun. 1968; FAKU 20933 (1), FAKU 20934 (1), FAKU 20935 (1), Owase fish market, Kumano-nada, 5 Nov. 1953; FAKU 34506 (1), FAKU 34508 (1), FAKU 34510 (1), Owase fish market, Kumano-nada, 12–14 Nov. 1961; FAKU 1913 (1), FAKU 1914 (1), FAKU 1915 (1), Owase fish market, Kumano-nada, Mar. 1976; FAKU 27832 (1), FAKU 27833 (1), FAKU 27834 (1), FAKU 27835 (1), FAKU 27836 (1), FAKU 27837 (1), FAKU 27838 (1), FAKU 27839 (1), FAKU 27840 (1), FAKU 27841 (1), FAKU 27842 (1), FAKU 27843 (1), FAKU 27845 (1), FAKU 27846 (1), FAKU 27847 (1), FAKU 27848 (1), Owase fish market, Kumano-nada, 25 Feb.–10 Mar. 1957; FAKU 23803 (1), FAKU 23804 (1), Owase fish market, Kumano-nada, 20 Oct. 1954; FAKU 24937 (1), FAKU 24940 (1), FAKU 24941 (1), FAKU 24942 (1), Owase fish market, Kumano-nada, 25 Oct. 1954; FAKU 17728 (1), Miya fish market, Enshu-nada, 15 Mar. 1952; FAKU 17305 (1), Miya fish market, Enshu-nada, 8–11 Jan. 1951; FAKU 28399 (1), FAKU 28400 (1), Miya fish market, Enshu-nada, 5–7 Feb. 1959; NSMT-P 48322 (1), west-northwest of Matsuzaki, Suruga Bay, 34.7745°N, 138.7140°E, 205–423 m, 5 Oct. 1995; MSM 13-18 (1), west of Cape Irozaki, Suruga Bay, 34.6166°N, 138.7293°E, 300–555 m, 25 Apr. 2012; NSMT-P 49960 (1), north-northwest of Heta, Suruga Bay, 35.0068°N, 138.7490°E, 320–680 m, 10 Nov. 1996.

***Trachonurus villosus*.** 1 specimen: BSKU 23038 (1), Tosa Bay, 33.0350°N, 133.7500°E, 945 m, 26 Jan. 1975.

***Ventrifossa garmani*.** 728 specimens: BSKU 26285 (1), BSKU 26286 (1), northwest of Yonaguni-jima Island, Okinawa Trough, 24.6833°N, 122.5933°E, 510–615 m, 13 Jan. 1978; BSKU 32576 (1), BSKU 32577 (1), northwest of Miyako-jima Island, Okinawa Trough, 25.7900°N, 124.3900°E, 600 m, 20 Sept. 1979; BSKU 32468 (1), northwest of Miyako-jima Island, Okinawa Trough, 25.8000°N, 124.4250°E, 400–420 m, 14 Sept. 1979; BSKU 29585 (1), northwest of Miyako-jima Island, Okinawa Trough, 25.7917°N, 124.5117°E, 500–550 m, 15 Sept. 1979; BSKU 32508 (1), BSKU 32509 (1), BSKU 32510 (1), BSKU 32511 (1), northwest of Miyako-jima Island, Okinawa Trough, 25.7917°N, 124.5117°E, 500–550 m, 10 Oct. 1979; BSKU 27471 (1), west of Okinawa-jima Island, Okinawa Trough, 27.7767°N, 126.2833°E, 490 m, 9 Mar. 1978; BSKU 27498 (1), BSKU 27499 (1), BSKU 27500 (1), west of Okinawa-jima Island, Okinawa Trough, 28.1000°N, 126.3833°E, 143 m, 9 Mar. 1978; SNFR 11077 (2), East China Sea, 28.3750°N, 126.9683°E, 378–382 m, 23 Oct. 1990; BSKU 27286 (1), BSKU 27288

(1), BSKU 27289 (1), BSKU 27292 (1), BSKU 27293 (1), west-northwest of Amami-oshima Island, Okinawa Trough, 28.8033°N, 127.0617°E, 425 m, 6 Feb. 1978; BSKU 27699 (1), west-northwest of Amami-oshima Island, Okinawa Trough, 28.7000°N, 127.1500°E, 500–535 m, 11 Mar. 1978; BSKU 27747 (1), BSKU 27748 (1), west of Amami-oshima Island, Okinawa Trough, 28.4333°N, 127.1833°E, 300–355 m, 12 Mar. 1978; BSKU 29836 (1), northwest of Amami Islands, Okinawa Trough, 29.0667°N, 127.2250°E, 420–450 m, 17 Dec. 1979; BSKU 34696 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.0667°N, 127.2250°E, 420–450 m, 17 Dec. 1979; BSKU 28207 (1), BSKU 28208 (1), BSKU 28209 (1), BSKU 28210 (1), BSKU 28211 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.3000°N, 127.3333°E, 410–415 m, 17 Mar. 1978; KPM-NI 6216 (1), KPM-NI 6218 (1), KPM-NI 6220 (1), ca. 230 km west-northwest of Amami-oshima Island, East China Sea, 29.3850°N, 127.4050°E, 400–432 m, 3 Oct. 1997; BSKU 34740 (1), BSKU 34741 (1), BSKU 34742 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.3833°N, 127.4250°E, 450–475 m, 18 Dec. 1979; BSKU 28291 (1), BSKU 28292 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7500°N, 127.5000°E, 325–360 m, 19 Mar. 1978; BSKU 27118 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.3833°N, 127.5150°E, 650 m, 5 Feb. 1978; BSKU 28318 (1), BSKU 28319 (1), BSKU 28320 (1), BSKU 28321 (1), BSKU 28322 (1), BSKU 28323 (1), BSKU 28324 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7167°N, 127.6167°E, 410–423 m, 18 Mar. 1978; SNFR 14227 (1), SNFR 14228 (1), SNFR 14229 (1), SNFR 14230 (5), East China Sea, 30.2077°N, 127.7222°E, 218–236 m, 7 Aug. 2008; BSKU 28372 (1), BSKU 28373 (1), BSKU 28383 (1), BSKU 28384 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7333°N, 127.8000°E, 495–510 m, 19 Mar. 1978; BSKU 33555 (1), BSKU 33557 (1), northwest of Amami-oshima Island, Okinawa Trough, 30.0115°N, 127.8387°E, 400–420 m, 26 Oct. 1979; SNFR 16341 (2), East China Sea, 30.4833°N, 127.9482°E, 399 m, 7 Nov. 2003; SNFR 14807 (1), SNFR 14808 (1), East China Sea, 30.3018°N, 128.0553°E, 389–401 m, 14 Sept. 2009; SNFR 16049 (2), East China Sea, 30.5870°N, 128.1007°E, 439–464 m, 17 Sept. 2009; NSMT-P 65638 (1), west of Yaku-shima Island, East China Sea, 30.4467°N, 128.2367°E, 499–500 m, 10 Nov. 2002; FFNU P-00612 (2), northwest of Fukue Basin, East China Sea, 32.4923°N, 128.3042°E, 364 m, 10 Nov. 2005; FFNU P-00616 (11), north of Fukue Basin, East China Sea, 32.4367°N, 128.3980°E, 396 m, 10 Nov. 2005; FFNU P-00538 (2 of 3), southwest of Daini-Danjo Knoll, East China Sea, 31.1543°N, 128.4013°E, 512 m, 6 Nov. 2004; NSMT-P 65663 (3), west-southwest of Shimokoshiki-jima Island, Okinawa Trough, 31.4217°N, 128.4017°E, 461–500 m, 8 Nov. 2002; FFNU P-00702 (2), south of Danjo Islands, East China Sea, 31.6925°N, 128.4790°E, 461 m, 14 Nov.

2007; FFNU P-00744 (14), south of Danjo Islands, East China Sea, 31.7162°N, 128.5135°E, 441 m, 17 Jul. 2008; BSKU 33928 (1), BSKU 33929 (1), BSKU 33930 (1), east of Danjo Islands, Okinawa Trough, 32.0033°N, 128.7050°E, 360–440 m, 2 Nov. 1979; FFNU P-00774 (6), east of Hirazi Bank, East China Sea, 32.2485°N, 129.4720°E, 403 m, 7 Mar. 2009; FFNU P-00746 (4), east of Hirazi Bank, East China Sea, 32.2538°N, 129.4927°E, 419 m, 19 Jul. 2008; KAUM-I. 6641 (1), off Nomaie, Kagoshima, 300 m, 27 Sept. 2007; KAUM-I. 32267 (1), off Nomaie, Kagoshima, 31.5000°N, 129.8833°E, 370–400 m, 10 Sept. 2010; BSKU 106801 (1), BSKU 106802 (1), southeast of Shimokoshiki-jima Island, East China Sea, 31.5560°N, 129.8847°E, 415 m, 24 Apr. 2012; BSKU 106762 (1), BSKU 106826 (1), BSKU 106827 (1), southeast of Shimokoshiki-jima Island, East China Sea, 31.5658°N, 129.8915°E, 380 m, 24 Apr. 2012; BSKU 106844 (1), southeast of Shimokoshiki-jima Island, East China Sea, 31.5675°N, 129.8930°E, 393 m, 24 Apr. 2012; MUFS 4216 (1), Aoshima, Miyazaki, 300 m, 17 Apr. 1975; ZMUT 50679 (1), Totoro, Miyazaki, 300–400 m, 9 Sept. 1959; FAKU uncat. (4), off Yawatahama, Hyuganada, 11 Nov. 1980; BSKU 58345 (1), Kamikawaguchi fish market, 12 Mar. 1999; BSKU 30381 (1), BSKU 30382 (1), BSKU 30383 (1), BSKU 30384 (1), Kyushu-Palau Ridge, 32.9667°N, 133.4333°E, 335 m, 21 Dec. 1979; BSKU 29488 (1), Tosa Bay, 32.9667°N, 133.4333°E, 335 m, 21 Dec. 1979; BSKU 30482 (1), Tosa Bay, 32.9833°N, 133.4833°E, 453 m, 21 Dec. 1979; BSKU 30559 (1), BSKU 30560 (1), Tosa Bay, 33.0833°N, 133.4833°E, 310 m, 20 Dec. 1979; ZMUT 23514 (1), ZMUT 23515 (1), Kochi, 16 Mar. 1931; ZMUT 23480 (1), ZMUT 23481 (1), Kochi, 19 Oct. 1930; ZMUT 23489 (1), Kochi, 8 Jun. 1929; BSKU 30159 (1), Mimase fish market, 1 Dec. 1979; BSKU 29953 (1), Mimase fish market, 1 Mar. 1980; BSKU 59292 (1), Mimase fish market, 1 Mar. 2002; BSKU 76654 (1), Mimase fish market, 10 Apr. 2005; BSKU 58507 (1), Mimase fish market, 10 Dec. 2001; BSKU 62506 (1), BSKU 62517 (1), BSKU 62518 (1), BSKU 62519 (1), BSKU 62527 (1), Mimase fish market, 11 Dec. 2002; BSKU 76885 (1), Mimase fish market, 11 Jan. 2005; BSKU 74817 (1), Mimase fish market, 12 Dec. 2004; BSKU 79597 (1), BSKU 79598 (1), BSKU 79599 (1), Mimase fish market, 12 Jan. 2006; BSKU 58313 (1), Mimase fish market, 13 Nov. 2000; BSKU 53055 (1), BSKU 53056 (1), Mimase fish market, 14 Dec. 2000; BSKU 99476 (1), BSKU 99477 (1), Mimase fish market, 14 Nov. 2003; BSKU 69389 (1), Mimase fish market, 14 Oct. 2003; BSKU 67780 (1), BSKU 67787 (1), Mimase fish market, 15 Dec. 2003; BSKU 70087 (1), Mimase fish market, 15 Jan. 2004; BSKU 71776 (1), BSKU 71777 (1), BSKU 71778 (1), BSKU 71779 (1), Mimase fish market, 15 Mar. 2004; BSKU 83050 (1), Mimase fish market, 16 Apr. 1996; BSKU 64841 (1), Mimase fish market, 16 Apr. 2003; BSKU 90767 (1), Mimase fish market, 18 Apr. 1996; BSKU 53084 (1), Mimase fish market, 18 Dec. 2000; BSKU 63396 (1), BSKU 63397 (1), BSKU 63398 (1), Mimase fish market, 18 Dec. 2002; BSKU 72500 (1), Mimase fish market, 18 Jan. 2004; BSKU 87871 (1), BSKU 87872 (1), BSKU 87873 (1), BSKU 87874 (1), Mimase fish market, 18 Nov. 2005; BSKU 78033 (1), Mimase fish market, 19 Apr. 2006; BSKU 77902 (1), Mimase fish market, 19 Dec. 2004; BSKU 62641 (1), Mimase fish market, 19 Feb. 2003; BSKU 79932 (1), Mimase fish market, 19 Feb. 2006; BSKU 59236 (1), Mimase fish market, 19 Nov. 2001; BSKU 69235 (1), Mimase fish market, 19 Nov. 2003; BSKU 69815 (1), BSKU 69816 (1), BSKU 69817 (1), Mimase fish market, 19 Oct. 2003; BSKU 84997 (1), Mimase fish market, 2 Apr. 1988; BSKU 58314 (1), Mimase fish market, 2 Apr. 1996; BSKU 72789 (1), BSKU 72790 (1), BSKU 72791 (1), Mimase fish market, 20 Feb. 2004; BSKU 58316 (2), BSKU 74299 (1), BSKU 78039 (1), Mimase fish market, 20 Nov. 1996; BSKU 78440 (1), BSKU 78441 (1), BSKU 78442 (1), BSKU 78443 (1), Mimase fish market, 20 Nov. 2005; BSKU 69590 (1), Mimase fish market, 21 Dec. 2003; BSKU 75747 (1), Mimase fish market, 21 Nov. 2004; BSKU 69797 (1), BSKU 69798 (1), BSKU 69799 (1), BSKU 69800 (1), BSKU 69801 (1), Mimase fish market, 22 Dec. 2003; BSKU 55448 (1), Mimase fish market, 22 Feb. 2001; BSKU 62271 (1), BSKU 63295 (1), Mimase fish market, 22 Jan. 2003; BSKU 54121 (1), BSKU 54122 (1), Mimase fish market, 23 Nov. 2000; BSKU 75642 (2), Mimase fish market, 23 Nov. 2004; BSKU 77815 (1), BSKU 77816 (1), Mimase fish market, 23 Nov. 2005; BSKU 99475 (1), Mimase fish market, 25 Apr. 2006; BSKU 59152 (1), Mimase fish market, 25 Nov. 2001; BSKU 35588 (1), Mimase fish market, 26 Apr. 1980; BSKU 70027 (1), BSKU 70028 (1), BSKU 70029 (1), BSKU 70030 (1), Mimase fish market, 26 Feb. 2003; BSKU 99467 (1), Mimase fish market, 26 Mar. 2005; BSKU 82938 (1), Mimase fish market, 26 Nov. 1996; BSKU 75923 (1), BSKU 75924 (1), Mimase fish market, 27 Dec. 2004; BSKU 54527 (1), BSKU 54528 (1), Mimase fish market, 27 Mar. 2000; BSKU 69573 (1), Mimase fish market, 28 Dec. 2003; BSKU 79886 (1), Mimase fish market, 28 Dec. 2005; BSKU 79584 (1), Mimase fish market, 28 Feb. 2005; BSKU 70805 (1), Mimase fish market, 28 Nov. 1997; BSKU 36046 (1), Mimase fish market, 28 Oct. 1981; BSKU 90188 (1), Mimase fish market, 29 Jan. 2007; BSKU 55730 (1), Mimase fish market, 29 Mar. 2001; BSKU 70453 (1), BSKU 70454 (1), BSKU 70455 (1), BSKU 70456 (1), BSKU 70457 (1), BSKU 70458 (1), Mimase fish market, 4 Dec. 2003; BSKU 69376 (1), Mimase fish market, 4 Feb. 2004; BSKU 69686 (1), BSKU 69687 (1), BSKU 69688 (1), BSKU 69689 (1), Mimase fish market, 5 Dec. 2003; BSKU 89735 (1), Mimase fish market, 5 Nov. 2006; BSKU 77806 (1), Mimase fish market, 6 Apr. 2006; BSKU 58383 (1), BSKU 58384 (1), BSKU 58385 (1), BSKU 58386 (1), Mimase fish market, 6 Dec. 2001;

BSKU 53204 (1), Mimase fish market, 6 Nov. 2000; BSKU 51934 (1), Mimase fish market, 7 Apr. 2000; BSKU 75944 (1), Mimase fish market, 7 Dec. 2004; BSKU 64308 (1), Mimase fish market, 7 Feb. 2003; BSKU 30250 (1), BSKU 30251 (1), BSKU 30252 (1), Mimase fish market, 7 Jan. 1980; BSKU 70131 (1), Mimase fish market, 7 Jan. 2004; BSKU 79914 (1), Mimase fish market, 7 Nov. 2006; BSKU 99285 (1), Mimase fish market, 8 Nov. 2000; BSKU 74971 (1), BSKU 74972 (1), BSKU 74973 (1), Mimase fish market, 8 Nov. 2004; BSKU 77645 (1), Mimase fish market, 8 Nov. 2005; BSKU 70965 (1), BSKU 70966 (1), BSKU 70967 (1), BSKU 70968 (1), BSKU 70969 (1), BSKU 70970 (1), Mimase fish market, 9 Nov. 2003; BSKU 75660 (1), Mimase fish market, 9 Nov. 2004; BSKU 77512 (1), BSKU 77513 (1), BSKU 77514 (1), Mimase fish market, 9 Nov. 2005; FAKU 21780 (1), FAKU 21786 (1), Mimase fish market, 10–20 Apr. 1954; BSKU 99219 (1), BSKU 99220 (1), BSKU 99221 (1), Mimase fish market, 14 Dec. 2006; BSKU 99256 (1), Mimase fish market, 18 Jan. 2006; BSKU 89112 (1), Mimase fish market, 20 Dec. 2006; BSKU 99245 (1), BSKU 99246 (1), Mimase fish market, 23 Apr. 2007; BSKU 99276 (1), Mimase fish market, 25 Apr. 2006; BSKU 99201 (1), BSKU 99202 (1), Mimase fish market, 26 Apr. 2006; FAKU 33962 (1), FAKU 33963 (1), Mimase fish market, Dec. 1960–Jan. 1961; BSKU 89726 (1), Mimase fish market, date unknown; BSKU 94811 (1), BSKU 94812 (1), BSKU 94813 (1), Mimase fish market, 28 Nov. 2007; BSKU 95086 (1), Mimase fish market, 30 Nov. 2007; BSKU 99351 (1), BSKU 99352 (1), Mimase fish market, 28 Jan. 2008; BSKU 95069 (1), Mimase fish market, 18 Jan. 2008; BSKU 99752 (1), Mimase fish market, 30 Nov. 2007; BSKU 99282 (1), Mimase fish market, 14 Nov. 2007; BSKU 92383 (1), Mimase fish market, 20 Nov. 2007; BSKU 99378 (1), BSKU 99379 (1), BSKU 99380 (1), Mimase fish market, 9 Nov. 2007; BSKU 42765 (1), off Okitsu, Tosa Bay, 240–250 m, 13 Mar. 1986; BSKU 30030 (1), off Sukumo, Kochi, 350 m, 1 Mar. 1980; BSKU 12921 (1), BSKU 12951 (1), BSKU 12952 (1), Tosa Bay, 350–450 m, 14 May 1968; BSKU 43833 (1), BSKU 43834 (1), BSKU 43835 (1), BSKU 43836 (1), BSKU 43837 (1), Tosa Bay, 400 m, 11 Jun. 1987; NSMT-P 94256 (1), Tosa Bay, 500 m, 25 Jun. 2000; BSKU 75041 (2), Tosa Bay, 33.2600°N, 133.6150°E, 144 m, 10 May 2000; BSKU 69316 (1), BSKU 69317 (1), BSKU 69318 (1), BSKU 69319 (1), BSKU 69320 (1), Tosa Bay, 33.1800°N, 133.6217°E, 412–423 m, 8 Sept. 1998; NSMT-P 57387 (1), NSMT-P 57388 (4), off Katsurahama, Tosa Bay, 33.1802°N, 133.6217°E, 412–423 m, 8 Sept. 1998; BSKU 78012 (1), BSKU 78013 (1), BSKU 78014 (1), BSKU 78015 (1), BSKU 78016 (1), BSKU 78017 (1), BSKU 78018 (1), BSKU 78019 (1), BSKU 78020 (1), BSKU 78021 (1), BSKU 78022 (1), BSKU 78023 (1), BSKU 78024 (1), BSKU 78025 (1), BSKU 78026 (1), BSKU 78027 (1), BSKU 78028 (1), BSKU 85961 (1), Tosa Bay, 33.1800°N, 133.6350°E, 408–458 m, 29 Jun. 1998; BSKU 77894 (1), BSKU 77895 (1), BSKU 77896 (1), BSKU 77897 (1), BSKU 77898 (1), Tosa Bay, 33.2083°N, 133.6433°E, 364–382 m, 27 Sept. 1999; BSKU 77995 (1), BSKU 77996 (1), BSKU 77997 (1), BSKU 77998 (1), BSKU 77999 (1), BSKU 78000 (1), BSKU 78001 (1), BSKU 78002 (1), BSKU 78003 (1), BSKU 78004 (1), BSKU 78005 (1), BSKU 78006 (1), Tosa Bay, 400 m, 9 May 2000; BSKU 70890 (1), BSKU 70891 (1), Tosa Bay, 33.2250°N, 133.6933°E, 440–460 m, 24 Aug. 2000; ZMUT 18988 (1), Kannoura, 3 Apr. 1929; FAKU 101162 (1), Owase fish market, Kumano-nada, 1960–1969; FAKU 34486 (1), FAKU 34487 (1), FAKU 34488 (1), FAKU 34490 (1), Owase fish market, Kumano-nada, 12–14 Nov. 1961; FAKU 1946 (1), FAKU 2482 (1), FAKU 2489 (1), FAKU 2490 (1), FAKU 2491 (1), FAKU 2493 (1), FAKU 2494 (1), Owase fish market, Kumano-nada, 4–9 Jan. 1936; FAKU 1600 (1), FAKU 1601 (1), FAKU 1602 (1), FAKU 1603 (1), FAKU 1604 (1), FAKU 1605 (1), FAKU 1607 (1), Owase fish market, Kumano-nada, 6–9 Dec. 1935; FAKU 27822 (1), FAKU 27823 (1), FAKU 27824 (1), FAKU 27825 (1), FAKU 27826 (1), FAKU 27827 (1), FAKU 27828 (1), Owase fish market, Kumano-nada, 25 Feb.–10 Mar. 1957; FAKU 24945 (1), FAKU 24946 (1), FAKU 24947 (1), FAKU 24948 (1), FAKU 24949 (1), FAKU 24952 (1), FAKU 24956 (1), FAKU 24962 (1), FAKU 24966 (1), FAKU 24967 (1), Owase fish market, Kumano-nada, 25 Oct. 1954; FAKU 29116 (1), FAKU 29117 (1), FAKU 29118 (1), FAKU 29119 (1), FAKU 29120 (1), Owase fish market, Kumano-nada, May 1960; BSKU 99073 (1), BSKU 99074 (1), BSKU 99075 (1), BSKU 99076 (1), BSKU 99077 (1), BSKU 99078 (1), Owase fish market, Kumano-nada, 13 Mar. 1983; FRLM 33597 (1), off Owase, Kumano-nada, 240 m, 6 Nov. 2007; FRLM 34993 (1), off Owase, Kumano-nada, 24 Oct. 2008; FRLM 34784 (1), off Owase, Kumano-nada, 27 Nov. 2008; FRLM 35070 (1), off Owase, Kumano-nada, 25 Feb. 2009; FRLM 34274 (1), FRLM 34275 (1), FRLM 34276 (1), FRLM 34277 (1), FRLM 34278 (1), off Owase, Kumano-nada, 14 Jan. 2008; FRLM 34462 (1), FRLM 34463 (1), FRLM 34464 (1), FRLM 34465 (1), FRLM 34466 (1), off Owase, Kumano-nada, 18 Feb. 2008; FRLM 33339 (1), FRLM 33340 (1), FRLM 33341 (1), FRLM 33342 (1), off Owase, Kumano-nada, 29 May 2007; FRLM 34336 (1), FRLM 34337 (1), FRLM 34338 (1), FRLM 34339 (1), off Owase, Kumano-nada, 30 Jan. 2008; FRLM 33827 (1), off Owase, Kumano-nada, 30 Nov. 2007; FRLM 34428 (1), off Owase, Kumano-nada, 7 Feb. 2008; FRLM 34213 (1), FRLM 34214 (1), off Owase, Kumano-nada, 7 Jan. 2008; FRLM 33923 (1), FRLM 33924 (1), FRLM 33925 (1), FRLM 33926 (1), FRLM 33927 (1), FRLM 33928 (1), FRLM 33929 (1), FRLM 33930 (1), FRLM 33931 (1), FRLM 33932 (1), FRLM 33933 (1), off Owase, Kumano-nada, 16 Nov. 2007; FRLM 30750 (1), off Owase, Kumano-nada, 2 Feb.

2004; FRLM 33848 (1), off Owase, Kumano-nada, 2007; FRLM 34362 (1), off Owase, Kumano-nada, date unknown; FAKU 4261 (1), FAKU 4262 (1), FAKU 4263 (1), FAKU 4264 (1), FAKU 4273 (1), FAKU 4274 (1), FAKU 4277 (1), Kumano-nada, Jan. 1937; FRLM 27235 (1), FRLM 27236 (1), off Kowaura, Kumano-nada, 165–309 m, 5 Dec. 2000; FRLM 27210 (1), off Kowaura, Kumano-nada, 165–309 m, 5 Nov. 2000; FRLM 8639 (1), Kumano-nada, date unknown; ZMUT 25344 (1), ZMUT 25345 (1), south of Shima, 31 Mar. 1933; NSMT-P 97155 (1), south of Cape Daiozaki, Kumano-nada, 34.0853°N, 136.8533°E, 474–475 m, 25 May 1994; KPM-NI 12767 (1), ca. 35–37 km off Cape Irakomisaki, Atsumi Peninsula, 34.2667°N, 137.2667°E, 370–380 m, 4 Aug. 2003; FAKU 106980 (1), Miya fish market, Enshu-nada, 400–800 m, 7 Apr. 1967; FAKU 26945 (1), FAKU 26946 (1), Miya fish market, Enshu-nada, 29–31 Dec. 1956; FAKU 28370 (1), FAKU 28371 (1), FAKU 28372 (1), FAKU 28374 (1), FAKU 28375 (1), FAKU 28377 (1), FAKU 28378 (1), FAKU 28379 (1), Miya fish market, Enshu-nada, 5–7 Feb. 1959; ZMUT 18590 (1), ZMUT 33685 (1), Maisaka, Shizuoka, date unknown; NSMT-P 101927 (2), southwest of Cape Omaezaki, Enshu-nada, 34.3816°N, 138.0010°E, 475–488 m, 10 Dec. 1995; BSKU 110046 (1), Suruga Bay, 34.7127°N, 138.4553°E, 262–434 m, 23 Apr. 2013; BSKU 110023 (1), BSKU 110024 (1), Suruga Bay, 34.7080°N, 138.4587°E, 200–450 m, 23 Apr. 2013; BSKU 110075 (1), Suruga Bay, 34.7481°N, 138.4652°E, 243–424 m, 23 Apr. 2013; BSKU 110432 (1), BSKU 110433 (1), Suruga Bay, 34.7489°N, 138.4664°E, 200–450 m, 23 Apr. 2013; BSKU 110141 (1), BSKU 110424 (3), Suruga Bay, 34.7651°N, 138.4880°E, 200–450 m, 23 Apr. 2013; KPM-NI 18046 (1), KPM-NI 18047 (1), off Fuji River, Suruga Bay, 300 m, Nov. 2006; KPM-NI 26263 (1), ca. 6 km off Cape Hagachizaki, Suruga Bay, 34.7450°N, 138.6831°E, 282 m, 24 Mar. 2010; BSKU 98915 (1), Heta fish market, Suruga Bay, 29 Oct. 1972; KPM-NI 17954 (1), KPM-NI 17955 (1), KPM-NI 17956 (1), Suruga Bay, 400 m, 26 Nov. 2006; KPM-NI 10314 (1), KPM-NI 10315 (1), KPM-NI 10316 (1), eastern part of Suruga Bay, 110–450 m, 10 Dec. 1998; KPM-NI 27423 (1), eastern part of Suruga Bay, 200–400 m, Mar. 2009; KPM-NI 24983 (1), eastern part of Suruga Bay, Nov. 2009; NSMT-P 93073 (1), off Heta, Suruga Bay, 15 Mar. 2000; NSMT-P 92858 (1), 13857.4 N, N34 57.4, E138 44.544.5 E, 34.9567°N, 138.7417°E, 380–980 m, 26 Oct. 1993; NSMT-P 47164 (1), off Osezaki, Suruga Bay, 1 Apr. 1967; FAKU M6827 (1), FAKU M6828 (1), FAKU M6829 (1), FAKU M6830 (1), FAKU M6832 (1), FAKU M6833 (1), FAKU M6834 (1), FAKU M6835 (1), FAKU M6836 (1), FAKU M6837 (1), FAKU M6838 (1), FAKU M6839 (1), FAKU M6840 (1), FAKU M6841 (1), FAKU M6842 (1), FAKU M6844 (1), FAKU M6845 (1), FAKU M6846 (1), FAKU M6846 (1), FAKU M6847 (1), FAKU M6848 (1), FAKU M6849 (1), FAKU M6850 (1), FAKU M6851 (1), FAKU M6852 (1), FAKU M6853 (1), FAKU M6854 (1), FAKU M6855 (1), FAKU M7058 (1), FAKU M7059 (1), FAKU M7060 (1), FAKU M7061 (1), FAKU M7062 (1), FAKU M7063 (1), FAKU M7064 (1), FAKU M7065 (1), Heta fish market, Suruga Bay, 150 ftm (275 m), 22–24 Jun. 1938; FAKU M7394 (1), FAKU M7396 (1), FAKU M7398 (1), FAKU M7401 (1), FAKU M7402 (1), FAKU M7404 (1), Heta fish market, Suruga Bay, 150 ftm (275 m), 23 Jul. 1938; FAKU 22842 (1), Heta fish market, Suruga Bay, 200–250 ftm (366–458 m), 24–25 Jun. 1939; FAKU 25690 (1), FAKU M7089 (1), FAKU M7090 (1), FAKU M7091 (1), FAKU M7092 (1), FAKU M7095 (1), FAKU M7096 (1), FAKU M7100 (1), FAKU M7101 (1), FAKU M7102 (1), FAKU M7103 (1), FAKU M7106 (1), FAKU M7107 (1), FAKU M7108 (1), FAKU M7109 (1), FAKU M7115 (1), FAKU M7116 (1), FAKU M7119 (1), FAKU M7121 (1), FAKU M7126 (1), FAKU M7136 (1), FAKU M7142 (1), FAKU M7145 (1), FAKU M7146 (1), FAKU M7148 (1), FAKU M7153 (1), Heta fish market, Suruga Bay, 26 Mar. 1939; BSKU 20316 (1), off Atami, Sagami Bay, 35.0783°N, 139.1467°E, 450 m, 24 Nov. 1972; FMNH 55537 (2), Odawara, Sagami Bay, date unknown; FAKU 34647 (1), off Oiso, Sagami Bay, 12 Mar. 1962; BSKU 19185 (1), south of Miura Peninsula, Sagami Bay, 35.1250°N, 139.5100°E, 700 m, 10 Mar. 1961; ZMUT 21030 (1), Misaki, Kanagawa, date unknown; FMNH 57387 (1, ex. CM 6325a), Misaki, Sagami Bay, date unknown; ZMUT 22678 (1), near Tokyo, date unknown; BSKU 19863 (1), off Boso Peninsula, 34.9183°N, 140.0450°E, 270–305 m, 30 Nov. 1968; FMNH 57245 (1, ex. CM 6184a), Boso Peninsula, 12 Apr. 1907; NSMT-P 19806 (20), off Choshi, 12 Apr. 1971; FAKU 8028 (1), FAKU 8031 (1), FAKU 8032 (1), FAKU 8033 (1), FAKU 8034 (1), FAKU 8035 (1), FAKU 8040 (1), FAKU 8041 (1), FAKU 8043 (1), off Choshi, 10 Apr. 1939; FAKU M7606 (1), off Choshi, 14–16 Jul. 1939; FAKU 5254 (1), FAKU 5265 (1), FAKU 5273 (1), FAKU 5276 (1), FAKU 5281 (1), FAKU 5285 (1), FAKU 5286 (1), FAKU 5287 (1), FAKU 5291 (1), FAKU 5292 (1), FAKU 5293 (1), FAKU 5294 (1), FAKU 5296 (1), FAKU 5297 (1), FAKU 5298 (1), FAKU 5299 (1), FAKU 5301 (1), FAKU 5302 (1), FAKU 5303 (1), FAKU 5304 (1), FAKU 5305 (1), FAKU 5306 (1), FAKU 5307 (1), FAKU 5308 (1), FAKU 5309 (1), FAKU 5311 (1), FAKU 5312 (1), FAKU 5313 (1), FAKU 5314 (1), FAKU 5315 (1), FAKU 5316 (1), FAKU 5317 (1), FAKU 5318 (1), FAKU 5320 (1), FAKU 5321 (1), FAKU 5322 (1), FAKU 5323 (1), FAKU 5324 (1), FAKU 5325 (1), FAKU 5326 (1), FAKU 5328 (1), FAKU 5329 (1), FAKU 5330 (1), FAKU 5331 (1), FAKU 5332 (1), FAKU 5333 (1), FAKU 5334 (1), FAKU 5335 (1), FAKU 5336 (1), FAKU 5337 (1), FAKU 5338 (1), FAKU 5339 (1), FAKU 5340 (1), FAKU 5344 (1), FAKU 5345 (1), FAKU 5346 (1), FAKU 5347 (1), FAKU 5348 (1), FAKU 5349 (1), FAKU 5350 (1), FAKU 5351 (1), FAKU 5352 (1), FAKU 5353 (1), off Choshi, 20–26 Nov. 1937; ZMUT 30605 (1), Choshi fish market, 27 Jan. 1935; ZMUT 37679 (1), ZMUT 37680 (1), ZMUT 37681 (1),

ZMUT 37678 (1), Onahama, Fukushima, date unknown; BSKU uncat. (12), off Mito, 250 m, 9 Oct. 1997; BSKU 101971 (10), off Hitachi, 36.6143°N, 141.0965°E, 244–262 m, 9 Nov. 1995; NSMT-P 48962 (1), off Hitachi, 36.5670°N, 141.1085°E, 412–421 m, 9 Nov. 1995; BSKU 84024 (1), BSKU 84025 (1), BSKU 84026 (1), BSKU 84027 (1), BSKU 84028 (1), BSKU 84029 (1), BSKU 84030 (1), BSKU 84031 (1), BSKU 84032 (1), BSKU 84033 (1), BSKU 84034 (1), BSKU 84035 (1), off Mito, 350 m, 8 Oct. 1997; NSMT-P 73334 (2, ex. SHMNH-P 18325), southeast off Onahama, 1931–1935; BSKU 101643 (1), off Rikuzen-Takata, 38.9957°N, 142.0527°E, 255–259 m, 23 Oct. 1995.

***Ventrifossa longibarbata*.** 77 specimens: BSKU 26287 (1), northwest of Yonaguni-jima Island, Okinawa Trough, 24.6833°N, 122.5933°E, 510–615 m, 13 Jan. 1978; BSKU 27472 (1), west of Okinawa-jima Island, Okinawa Trough, 27.7767°N, 126.2833°E, 490 m, 9 Mar. 1978; BSKU 27859 (1), BSKU 27860 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.5833°N, 127.0000°E, 600 m, 16 Mar. 1978; BSKU 27662 (1), west of Okinawa-jima Island, Okinawa Trough, 28.7333°N, 127.0167°E, 610–640 m, 11 Mar. 1978; BSKU 27912 (1), BSKU 27913 (1), BSKU 27914 (1), BSKU 27918 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.7500°N, 127.1167°E, 520–542 m, 16 Mar. 1978; BSKU 27700 (1), BSKU 27701 (1), BSKU 27712 (1), west of Okinawa-jima Island, Okinawa Trough, 28.7000°N, 127.1500°E, 535–550 m, 11 Mar. 1978; BSKU 27188 (1), BSKU 27189 (1), BSKU 27193 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.4333°N, 127.5167°E, 570 m, 5 Feb. 1978; BSKU 33256 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7167°N, 127.7000°E, 520 m, 14 Oct. 1979; BSKU 28374 (1), BSKU 28375 (1), BSKU 28376 (1), BSKU 28377 (1), BSKU 28378 (1), BSKU 28379 (1), BSKU 28380 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.7333°N, 127.8000°E, 495–510 m, 19 Mar. 1978; BSKU 28597 (1), BSKU 28598 (1), BSKU 28600 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.9500°N, 128.0000°E, 585–610 m, 20 Mar. 1978; BSKU 29891 (1), west of Yaku-shima Island, Okinawa Trough, 30.2083°N, 128.2088°E, 600–620 m, 26 Oct. 1979; FFNU P-00538 (1 of 3), southwest of Daini-Danjo Knoll, East China Sea, 31.1543°N, 128.4013°E, 512 m, 6 Nov. 2004; BSKU 33927 (1), east of Danjo Islands, Okinawa Trough, 32.0033°N, 128.7050°E, 360–440 m, 2 Nov. 1979; FFNU P-00716 (4), west of Amakusa Knoll, East China Sea, 32.1593°N, 129.4715°E, 503 m, 12 Mar. 2008; FFNU P-00742 (2), west of Amakusa Knoll, East China Sea, 32.1625°N, 129.4838°E, 487 m, 17 Jul. 2008; NSMT-P 95186 (1), off Kochi, date unknown; BSKU 12923 (1), Tosa Bay, 350–450 m, 14 May 1968; BSKU 78007 (1), Tosa Bay, 400 m, 9 May 2000; BSKU 101608 (1), BSKU 101610 (1), BSKU 101611 (1), BSKU 101612 (1), BSKU 101613 (1), BSKU 101614 (1), BSKU 101615 (1), BSKU 101616 (1), BSKU 101617 (1), BSKU

101621 (1), BSKU 101622 (1), BSKU 101624 (1), BSKU 101625 (1), BSKU 101626 (1), BSKU 101627 (1), BSKU 101628 (1), BSKU 101629 (1), BSKU 101630 (1), BSKU 101631 (1), BSKU 101632 (1), BSKU 101633 (1), Tosa Bay, 33.1428°N, 133.6248°E, 450–480 m, 1 Aug. 1988; BSKU 44421 (1), Tosa Bay, 600 m, 11 Jan. 1988; BSKU 101637 (1), BSKU 101638 (1), BSKU 101639 (1), south of Muroto, 33.0790°N, 134.1645°E, 414–513 m, 17 Nov. 2005; FAKU 4925 (1), Kumano-nada, 1938; NSMT-P 101351 (2), southwest of Cape Daiozaki, Kumano-nada, 34.0836°N, 136.6554°E, 495–515 m, 12 Dec. 1995; MSM 13-16 (1), Suruga Bay, 34.8544°N, 138.7162°E, 372–563 m, 21 Feb. 2012; NSMT-P 93123 (1), NSMT-P 93124 (1), off Oh-sima Island, Izu Islands, date unknown; BSKU 19186 (1), south of Miura Peninsula, Sagami Bay, 35.1250°N, 139.5100°E, 700 m, 10 Mar. 1961; ZMUT 30602 (1), ZMUT 30603 (1), Choshi fish market, 27 Jan. 1935; NSMT-P 120918 (2), NSMT-P 120921 (1), off Iwaki, 36.8885°N, 141.4562°E, 376 m, 2 Nov. 2007; NSMT-P 48961 (1), off Iwaki, 36.9965°N, 141.5627°E, 387–420 m, 4 Nov. 1995; BSKU 84049 (1), off Onahama, 350 m, 16 Oct. 1997.

***Ventrifossa macroptera*.** 136 specimens: BSKU 30585 (1), BSKU 30586 (1), BSKU 30587 (1), Kyushu-Palau Ridge, 27.9000°N, 134.6500°E, 910 m, 18 Dec. 1979; BSKU uncat. (48), BSKU 26030 (1), BSKU 26031 (1), BSKU 26032 (1), BSKU 26033 (1), BSKU 26034 (1), BSKU 26035 (1), BSKU 26036 (1), BSKU 26037 (1), BSKU 26038 (1), BSKU 26039 (1), BSKU 26040 (1), BSKU 26041 (1), BSKU 26042 (1), BSKU 26043 (1), BSKU 26044 (1), BSKU 26045 (1), BSKU 26046 (1), BSKU 26047 (1), BSKU 26048 (1), BSKU 26049 (1), BSKU 26050 (1), BSKU 26051 (1), BSKU 26052 (1), BSKU 26053 (1), BSKU 26054 (1), BSKU 26055 (1), BSKU 26056 (1), BSKU 26057 (1), BSKU 26058 (1), BSKU 26077 (1), BSKU 26078 (1), BSKU 26079 (1), BSKU 26080 (1), BSKU 26082 (1), BSKU 26083 (1), BSKU 26084 (1), BSKU 26085 (1), BSKU 26086 (1), BSKU 26087 (1), BSKU 26088 (1), BSKU 26090 (1), BSKU 26091 (1), BSKU 26092 (1), BSKU 26093 (1), BSKU 26095 (1), BSKU 26096 (1), BSKU 26097 (1), BSKU 26098 (1), BSKU 26099 (1), BSKU 26117 (1), BSKU 26118 (1), BSKU 26120 (1), BSKU 26122 (1), BSKU 26123 (1), BSKU 26125 (1), BSKU 26126 (1), BSKU 26127 (1), BSKU 26128 (1), BSKU 26129 (1), BSKU 26131 (1), BSKU 26132 (1), BSKU 26133 (1), BSKU 26134 (1), BSKU 26135 (1), BSKU 26136 (1), BSKU 26137 (1), BSKU 26138 (1), BSKU 26140 (1), BSKU 26143 (1), Kyushu-Palau Ridge, 27.9167°N, 134.6500°E, 700 m, 20 Feb. 1978; BSKU 26059 (1), Kyushu-Palau Ridge, 27.9000°N, 134.6550°E, 650 m, 15 Feb. 1978; BSKU 40982 (1), BSKU 40983 (1), Kyushu-Palau Ridge, 27.9100°N, 134.6650°E, 910 m, 18 Dec. 1979; BSKU 32224 (1), BSKU 32225 (1), Kyushu-Palau Ridge, 28.1117°N, 134.6656°E, 550–600 m, 19 Dec. 1979; BSKU 30471 (1), Kyushu-Palau Ridge, 27.9000°N,

134.6667°E, 697–700 m, 18 Dec. 1979; BSKU 40976 (1), BSKU 40977 (1), BSKU 40978 (1), BSKU 40979 (1), Kyushu-Palau Ridge, 27.9100°N, 134.6667°E, 697–1320 m, 18 Dec. 1979; BSKU 32190 (1), BSKU 32191 (1), BSKU 32201 (1), collected with holotype; BSKU 30433 (1), BSKU 30434 (1), BSKU 30437 (1), Kyushu-Palau Ridge, 26.7500°N, 135.3167°E, 336 m, 17 Dec. 1979.

***Ventrifossa misakia*.** 258 specimens: NSMT-P 94395 (14), off Miyazaki, Hyuga-nada, 484–565 m, 19 May 1989; BSKU 13225 (1), BSKU 13226 (1), BSKU 13227 (1), BSKU 13228 (1), BSKU 13274 (1), BSKU 13275 (1), ca. 15 km east of Cape Ashizurimisaki, Tosa Bay, 420–555 m, 2–6 Jun. 1968; BSKU 12953 (1), BSKU 12954 (1), BSKU 12955 (1), Tosa Bay, 350–450 m, 14 May 1968; BSKU 44129 (1), SNFR 17969 (2), Tosa Bay, 450–500 m, 5 Oct. 1987; BSKU 13002 (1), off Tei, Tosa Bay, 415–430 m, 22 May 1968; BSKU 13340 (1), ca. 10–15 km west of Cape Murotomisaki, Tosa Bay, 360–435 m, 8–9 Jun. 1968; BSKU 13476 (1), ca. 10–15 km west of Murotomisaki, Tosa Bay, 360–435 m, 8–9 Jun. 1968; BSKU 78800 (1), BSKU 78802 (1), BSKU 78803 (1), BSKU 78804 (1), BSKU 78805 (1), BSKU 78806 (1), BSKU 78808 (1), south of Muroto, 33.0790°N, 134.1645°E, 414–513 m, 17 Nov. 2005; FAKU 34492 (1), FAKU 34493 (1), FAKU 34495 (1), FAKU 34496 (1), Owase fish market, Kumano-nada, 12–14 Nov. 1961; NSMT-P 101352 (2), southwest of Cape Daiozaki, Kumano-nada, 34.0836°N, 136.6554°E, 495–515 m, 12 Dec. 1995; FAKU 34130 (1), FAKU 34131 (1), FAKU 34134 (1), FAKU 34135 (1), FAKU 34136 (1), FAKU 34137 (1), FAKU 34138 (1), FAKU 34139 (1), Miya fish market, Enshu-nada, Apr. 1961; FAKU 28393 (1), FAKU 28396 (1), FAKU 28397 (1), Miya fish market, Enshu-nada, 5–7 Feb. 1959; NSMT-P 101928 (2), southwest of Cape Omaezaki, Enshu-nada, 34.3816°N, 138.0010°E, 475–488 m, 10 Dec. 1995; MSM uncat. (2), Suruga Bay, 21 Feb. 2012; NSMT-P 49952 (1), east of Matsuzaki, Suruga Bay, 34.7578°N, 138.6895°E, 366–440 m, 12 Nov. 1996; OMNH-P 9348 (1), OMNH-P 9349 (1), off Cape Irozaki, Izu Peninsula, 13 May 1996; FMNH 57091 (2, ex. CM 6029a–b), Misaki, Sagami Bay, 23 Dec. 1901; FMNH uncat. (1), off Misaki, Sagami Bay, date unknown; ZMUT 4736 (1), Tokyo market, date unknown; NSMT-P 94435 (1), southeast of Taitozaki, Boso Peninsula, 422–425 m, 28 May 1986; FAKU 8002 (1), FAKU 8003 (1), FAKU 8004 (1), FAKU 8005 (1), FAKU 8006 (1), FAKU 8007 (1), off Choshi, 10 Apr. 1939; FAKU M7631 (1), FAKU M7632 (1), FAKU M7634 (1), FAKU M7635 (1), FAKU M7636 (1), FAKU M7637 (1), FAKU M7638 (1), FAKU M7639 (1), FAKU M7640 (1), FAKU M7641 (1), FAKU M7642 (1), FAKU M7643 (1), FAKU M7644 (1), FAKU M7645 (1), FAKU M7646 (1), FAKU M7647 (1), FAKU M7648 (1), FAKU M7649 (1), FAKU M7650 (1), FAKU M7651 (1), FAKU M7652 (1), FAKU M7653 (1), FAKU M7654 (1), FAKU M7655 (1), FAKU M7656 (1), off Choshi, 14–16 Jul. 1939; FAKU 5160 (1), FAKU 5168

(1), FAKU 5188 (1), FAKU 5222 (1), FAKU 5270 (1), FAKU 5275 (1), FAKU 5279 (1), FAKU 5295 (1), FAKU 5319 (1), off Choshi, 20–26 Nov. 1937; FAKU 4002 (1), FAKU M7508 (1), off Choshi, Mar. 1937; ZMUT 30604 (1), Choshi fish market, 27 Jan. 1935; NSMT-P 77875 (1), NSMT-P 77876 (1), NSMT-P 77877 (1), NSMT-P 77878 (1), off Hitachi, 36.5017°N, 141.0300°E, 378–385 m, 12 Nov. 2006; NSMT-P 77290 (1), NSMT-P 77291 (4), off Hitachi, 36.5150°N, 141.0517°E, 407–413 m, 13 Nov. 2006; BSKU uncat. (8), BSKU 83324 (1), BSKU 83325 (1), BSKU 83326 (1), BSKU 83327 (1), BSKU 83328 (1), BSKU 83329 (1), BSKU 83330 (1), BSKU 83331 (1), BSKU 83332 (1), BSKU 83333 (1), BSKU 83334 (1), BSKU 83335 (1), BSKU 83336 (1), BSKU 83337 (1), BSKU 83338 (1), off Hitachi, 36.5670°N, 141.1085°E, 412–421 m, 9 Nov. 1995; BSKU 83733 (1), BSKU 83735 (1), BSKU 83736 (1), BSKU 83737 (1), BSKU 83738 (1), BSKU 83739 (1), BSKU 83740 (1), BSKU 83742 (1), BSKU 83743 (1), BSKU 83745 (1), BSKU 83746 (1), BSKU 83748 (1), BSKU 83749 (1), BSKU 83750 (1), BSKU 83751 (1), BSKU 83752 (1), BSKU 83753 (1), off Mito, 450 m, 9 Oct. 1997; BSKU 83758 (1), BSKU 83759 (1), BSKU 83760 (1), off Mito, 350 m, 8 Oct. 1997; NSMT-P 78961 (2), off Iwaki, 36.8885°N, 141.4562°E, 376 m, 2 Nov. 2007; BSKU 83339 (1), BSKU 83413 (1), BSKU 83414 (1), BSKU 83415 (1), BSKU 83416 (1), BSKU 83417 (1), BSKU 83418 (1), BSKU 83419 (1), off Onahama, 36.9965°N, 141.5627°E, 387–420 m, 4 Nov. 1995; BSKU 83756 (1), off Onahama, 350 m, 6 Oct. 1997; BSKU uncat. (36), off Onahama, 350 m, 16 Oct. 1997; BSKU uncat. (46), off Onahama, 350 m, 17 Oct. 1997; NSMT-P 101751 (2), south-southeast of Oshika Peninsula, 37.9215°N, 141.7400°E, 300 m, 15 Apr. 1997; BSKU uncat. (2), off Fukushima, 37.4878°N, 141.8532°E, 411–419 m, 3 Nov. 1995; BSKU 116163 (1), off Minami-soma, 37.4878°N, 141.8532°E, 411–419 m, 3 Nov. 1995.

***Ventrifossa rhipidodorsalis*.** 10 specimens: BSKU 27919 (1), northwest of Amami-oshima Island, Okinawa Trough, 28.7500°N, 127.1167°E, 520–542 m, 16 Mar. 1978; BSKU 27186 (1), northwest of Amami-oshima Island, Okinawa Trough, 29.4333°N, 127.5167°E, 570 m, 5 Feb. 1978; NSMT-P 77233 (1), west of Yaku-shima Island, East China Sea, 30.4467°N, 128.2367°E, 499–500 m, 10 Nov. 2002; BSKU 1821 (1), Mimase fish market, Mar. 1952; BSKU 99333 (1), BSKU 99336 (1), BSKU 99339 (1), off Kochi, 600 m, 24 Mar. 1989; FAKU 132253 (1), off Maisaka, Enshu-nada, Nov. 1941; NSMT-P 75111 (1), off Heta, Suruga Bay, 34.9050°N, 138.7133°E, 520–545 m, 23 Nov. 1978; NSMT-P 46728 (1), 2.0 miles west-southwest of Heta, Suruga Bay, 34.9567°N, 138.7250°E, 315–500 m, 22 Oct. 1994.

APPENDIX 2. Comparative material examined

Asthenomacrus fragilis. 1 specimen. **Panama:** CAS 87796 (171+ mm TL), southeast of Punta Mala, eastern Pacific, 6.8833°N, 79.4500°W, 3193 m, 5 May 1967.

Coelorinchus acanthiger. 15 specimens. **South Africa:** FAKU 108103 (1, 109 mm HL, 390+ mm TL), FAKU 108112 (1, 102 mm HL, 392+ mm TL), FAKU 108120 (1, 100 mm HL, 366+ mm TL), FAKU 108122 (1, 75.9 mm HL, 264+ mm TL), off Cape Agulhas, 35.6917°S, 18.8817°E, 980 m, 22 Nov. 1971. **Western Australia:** FSFRL EB436 (1, 74.8 mm HL, 231+ mm TL), off Perth, 31.6550°S, 114.7333°E, 1035 m, 15 Nov. 1975; FSFRL ED885 (1, 87.3 mm HL, 320+ mm TL), off Gregory, 28.3867°S, 113.0300°E, 855 m, 11 Nov. 1975. **Great Australian Bight:** FSFRL EA878 (1, 123 mm HL, 443+ mm TL), FSFRL EB674 (1, 98.5 mm HL, 299+ mm TL), west of Coffin Bay, 34.7783°S, 132.8283°E, 920 m, 5 Dec. 1975. **Louisville Ridge:** NSMT-P 41338 (1, 109 mm HL, 361+ mm TL), NSMT-P 41339 (1, 98.1 mm HL, 360+ mm TL), 41.4670°S, 164.2000°W, 874 m, 4 Apr. 1985; NSMT-P 41340 (1, 93.8 mm HL, 336+ mm TL), NSMT-P 41341 (1, 100 mm HL, 376+ mm TL), 40.6670°S, 165.4000°W, 1026 m, 5 Apr. 1985; BSKU 47717 (1, 98.7 mm HL, 342+ mm TL), BSKU 47718 (1, 115 mm HL, 416+ mm TL), BSKU 47719 (1, 101 mm HL, 399+ mm TL), 39.1000°S, 167.4667°W, 978 m, 2 Mar. 1986.

Coelorinchus acutirostris. 2 specimens. **Holotype of Coelorhynchus acutirostris:** USNM 72947 (62.3 mm HL, 177+ mm TL), between Cebu and Bohol, Philippines, 10.1472°N, 123.8750°E, 291 m, *Albatross* sta. 5418, 12-ft Agassiz beam trawl, 25 Mar. 1909. **Philippines:** USNM 149284 (paratype of *Coelorhynchus argentatus*, 59.5 mm HL, 166+ mm TL), between Cebu and Bohol, 10.1700°N, 123.8880°E, 302 m, 25 Mar. 1909.

Coelorinchus amirantensis. 3 specimens. **Paratypes of Coelorinchus amirantensis:** CAS 223467 (3, 95.6–127 mm HL, 286+–373+ mm TL), Mascarene Ridge, Mascarene Ridge, 8.4400°S, 59.4833°E, 1260–1300 m, 2 Sept. 1977.

Coelorinchus anisacanthus. 4 specimens. **Holotype of Coelorinchus anisacanthus:** ZMMGU P.18251 (80.7 mm HL, 290+ mm TL), Koko Seamount, 35.3500°N, 171.3167°E, 550 m, R/V *Mys Yunony*, tr. 87, 7 Sept. 1979. **Emperor Seamounts:** ZMMGU P.15374 (3, 74.9–81.5 mm HL, 270–298+ mm TL).

Coelorinchus aratrum. 1 specimen. **Hawaii:** BPBM 24344 (72.8 mm HL, 241+ mm TL), Pailolo Channel, 21.0833°N, 156.3833°W, 360–384 m, 23 Nov. 1968.

Coelorinchus argentatus. 8 specimens. **Holotype of Coelorhynchus argentatus:** USNM 72949 (94.3 mm HL, 360+ mm TL), vicinity of Jolo, Sulu Sea, 6.1000°N, 120.9810°E, 582 m, *Albatross* sta. 5138, 12-ft Agassiz beam trawl, 14 Feb. 1908. **Paratypes of C. argentatus:** USNM 149283 (1, 58.5 mm HL, 207+ mm TL), northern Mindanao, Bohol Sea, 8.7583°N, 123.5620°E, 309 m, 9

Aug. 1908; USNM 149285 (1, 37.6 mm HL, 136+ mm TL), off Malavatuan Island, South China Sea, 13.9583°N, 120.3710°E, 214 m, 17 Jul. 1908. **Philippines:** ZMUC P372922 (1, 58.7 mm HL, 220 mm TL), ZMUC P372923 (1, 61.6 mm HL, 221+ mm TL), ZMUC P372924 (1, 62.9 mm HL, 230+ mm TL), ZMUC P372925 (1, 70.1 mm HL, 256+ mm TL), ZMUC P372926 (1, 70.5 mm HL, 252+ mm TL), west of Jolo, Sulu Sea, 27 Mar. 1914.

Coelorinchus braueri. 11 specimens. **Indonesia:** HUMZ 191067 (1, 81.0 mm HL, 297+ mm TL), Indian Ocean off Java, 8.0050°S, 111.2583°E, 680–706 m, 10 Sept. 2004; HUMZ 191505 (1, ca. 78 mm HL, 290+ mm TL), HUMZ 191506 (1, 77.9 mm HL, 263+ mm TL), HUMZ 191507 (1, 78.7 mm HL, 287 mm TL), Indian Ocean off Java, 8.0017°S, 109.9033°E, 850–916 m, 12 Sept. 2004. **Indian Ocean:** ZMUC P373070 (1, 86.1 mm HL, 305+ mm TL), ZMUC P373072 (1, 69.0 mm HL, 241+ mm TL), ZMUC P373075 (1, 81.6 mm HL, 301+ mm TL), R/V *Galathea*, sta. 197, 14 Feb. 1951; ZMUC P373118 (1, 76.1 mm HL, 287+ mm TL), ZMUC P373119 (1, 85.0 mm HL, 308 mm TL), ZMUC P373120 (1, 74.1 mm HL, 265+ mm TL), R/V *Galathea*, sta. 203, 21 Feb. 1951; ZMUC P373106 (1, 72.3 mm HL, 280+ mm TL), R/V *Galathea*, sta. 554, 21 Feb. 1951.

Coelorinchus celanostoma. 4 specimens. **Tasman Sea:** CAS uncat. (2, 146–179 mm HL, 483+–560+ mm TL), West Norfolk Ridge, 33.8207°S, 167.0558°E, 805–938 m, 27 May 2003; NSMT-P 101629 (1, 238 mm HL, 704+ mm TL), Norfolk Ridge, 33.7617°S, 167.0367°E, 805 m, 12 Jan. 1976; CAS uncat. (1, 117 mm HL, 371+ mm TL), West Norfolk Ridge, 33.6038°S, 167.7848°E, 926–969 m, 31 May 2003.

Coelorinchus charius. 19 specimens. **Holotype of Coelorinchus charius:** AMS I.31166-005 (80.6 mm HL, 267+ mm TL), off Shark Bay, 25.3183°S, 111.9433°E, 610–612 m, FRV *Southern Surveyor*, sta. SS1/91/31, coll. J. Paxton, 29 Jan. 1991. **Paratypes of C. charius:** AMS I.31166-010 (6, ex. AMS I.31166-005, 73.3–98.2 mm HL, 266+–373+ mm TL), collected with holotype; CSIRO H2573-16 (4, 78.6–97.9 mm HL, 287+–354+ mm TL), west of Steep Point, 26.2417°S, 112.0533°E, 690–691 m, 30 Jan. 1991; CSIRO H2557-05 (1, 62.5 mm HL, 186+ mm TL), west of Cape Farquhar, 23.7150°S, 112.5983°E, 620 m, 1991; CSIRO H2554-10 (1, 73.1 mm HL, 228+ mm TL), off Point Cloates, 22.9600°S, 113.2400°E, 482–554 m, 26 Jan. 1991; CSIRO H1492-14 (1, 51.3 mm HL, 170+ mm TL), North West Shelf, south of Barrow Island, 20.6800°S, 114.7217°E, 420 m, 1988; CSIRO H1514-34 (5, 58.5–89.5 mm HL, 185+–296+ mm TL), North West Shelf, 18.1467°S, 117.9083°E, 582 m, 12 Oct. 1988.

Coelorinchus commutabilis. 1 specimen. **Holotype of Coelorhynchus commutabilis:** USNM 72945 (93.7 mm HL, 314+ mm TL), Palawan Passage, 10.9625°N, 118.6380°E, 686 m, *Albatross* sta. 5348, 12-ft Tanner beam trawl, 27 Dec. 1908.

Coelorinchus cylindricus. 1 specimen. **Holotype**

of *Caelorinchus cylindricus*: MNHN 1996-364 (42.2 mm HL, 193 mm TL), Seamount B, southeast of New Caledonia, Norfolk Ridge, 24.8783°S, 168.3633°E, 550–920 m, *BERYX II*, sta. CP4, 14 Oct. 1992.

Coelorinchus denticulatus. 2 specimens. **Holotype of *Coelorhynchus denticulatus***: BMNH 1921.3.1.9 (64.3 mm HL, 258+ mm TL), off Umuoti River, Natal, 15 miles out, 219 m, coll. H.W. Bell Marley and R. Robinson, date unknown. **Holotype of *Coelorhynchus natalensis***: ANSP 53067 (46.3 mm HL, 201+ mm TL), southeast of Bluff, Durban, trawl, coll. H.W. Bell Marley, date unknown.

Coelorinchus doryssus. 4 specimens. **Hawaii**: BPBM 24340 (4, 84.0–121 mm HL, 266+–364+ mm TL), Pailolo Channel, 21.0833°N, 156.1833°W, 623–667 m, 23 Nov. 1968.

Coelorinchus flabellispinnis. 2 specimens. **Indonesia**: HUMZ 191022 (1, 164 mm HL, 552+ mm TL), Indian Ocean off Sumatra, 0.1205°N, 97.5687°E, 510–524 m, 4 Oct. 2004; HUMZ 191011 (1, 147 mm HL, 506 mm TL), Indian Ocean off Sumatra, 5.0055°N, 94.1200°E, 783–1047 m, 22 Oct. 2004.

Coelorinchus gaesorhynchus. 2 specimens. **Holotype of *Caelorinchus gaesorhynchus***: CSIRO H2548-13 (75.8 mm HL, 254+ mm TL), off North West Cape, 21.7450°S, 113.8717°E, 290–320 m, FRV *Southern Surveyor*, sta. SS1/91/08, 24 Jan. 1991. **Paratype of *C. gaesorhynchus***: CSIRO H2548-16 (70.0 mm HL, 236+ mm TL), collected with holotype.

Coelorinchus gladius. 2 specimens. **Syntype of *Coelorhynchus gladius***: USNM 47706 (45.0 mm HL, 162+ mm TL), off Oahu, 21.2000°N, 157.8170°W, 539 m, *Albatross* sta. 3472, beam trawl, 4 Dec. 1891. **Hawaii**: HUMZ 99792 (1, 57.7 mm HL, 199+ mm TL), north-northeast of Hilo, 19.9483°N, 155.0050°W, 379 m, 7 Feb. 1983.

Coelorinchus goobala. 24 specimens. **Holotype of *Caelorinchus goobala***: NTM S.12631-004 (85.1 mm HL, 270+ mm TL), off Rowley Shoals, North West Shelf, 17.6167°S, 118.6667°E, 400 m, coll. W. Houston, 4 Nov. 1985. **Paratypes of *C. goobala***: AMS I.22808-042 (8, 57.3–94.4 mm HL, 190+–354+ mm TL), 220 km north of Port Hedland, North West Shelf, 420 m; AMS I.23423-014 (1, 63.5 mm HL, 218+ mm TL), North West Shelf, 376 m; AMS I.23425-019 (1, 57.2 mm HL, 204+ mm TL), North West Shelf, 400 m, FRV *Soela*, sta. SO4/82/leg1; CAS 200229 (1, 61.4 mm HL, 209+ mm TL), North West Shelf, 400 m; CSIRO H1514-31 (6, 60.0–77.1 mm HL, 207+–289+ mm TL), North West Shelf, 18.1467°S, 117.9083°E, 582 m, 12 Oct. 1988; NTM S.12614-023 (2, 58.0–85.1 mm HL, 180+–307+ mm TL), off Rowley Shoals, North West Shelf, 17.5667°S, 118.6333°E, 410 m, 7 Nov. 1985; NTM S.12588-023 (1, 101 mm HL, 376+ mm TL), off Rowley Shoals, North West Shelf, 17.3667°S, 118.6333°E, 403 m, 2 Jan. 1985; NTM S.13115-02 (1, 57.7 mm HL, 184+ mm TL), north of Cape Leveque, 14.1167°S, 122.1000°E, 423 m, 24 Jan. 1990. **Timor Sea**:

NTM S.14378-013 (2, 54.7–58.4 mm HL, 170+–201+ mm TL), south-southwest of Cartier Reef, 13.1320°S, 13.1320°E, 420 m, 19 Jun. 1996.

Coelorinchus immaculatus. 1 specimen. **Holotype of *Caelorinchus immaculatus***: ZMMGU P.17692 (68.9 mm HL, 264+ mm TL), Sala y Gomez Ridge, 24.9617°S, 88.6017°W, 560 m, R/V *Prof. Shtokman*, cr. 18, sta. 1955/62, trap, 29–30 Apr. 1987.

Coelorinchus kaiyomaru. 11 specimens. **Holotype of *Caelorinchus kaiyomaru***: NSMT-P 18504 (78.0 mm HL, 351 mm TL), Chatham Rise, 44.5700°S, 177.9650°W, 1006–1050 m, FRV *Kaiyo-maru*, otter trawl, 20 Dec. 1977. **Paratypes of *C. kaiyomaru***: NSMT-P 18506 (1, 82.0 mm HL, 337+ mm TL), NSMT-P 69285 (1, ex. FSFRL EH222, 91.9 mm HL, 353+ mm TL), collected with holotype; NSMT-P 18508 (1, 72.7 mm HL, 300+ mm TL), South Tasman Rise, 47.2517°S, 148.5133°E, 949 m, 22 Dec. 1977; NSMT-P 69286 (1, ex. FSFRL EH616, 49.2 mm HL, 212 mm TL), NSMT-P 69287 (1, ex. FSFRL EH616, 61.9 mm HL, 237+ mm TL), Campbell Plateau, 48.4883°S, 173.0133°E, 874–896 m, 8 Jan. 1978. **Australia**: BSKU 108818 (1, 71.8 mm HL, 293 mm TL), BSKU 108819 (1, 71.8 mm HL, 301+ mm TL), off southeastern coast of Australia. **New Zealand**: BSKU 44452 (1, 72.0 mm HL, 296+ mm TL), BSKU 44453 (1, 67.3 mm HL, 275 mm TL), southeast of New Zealand, 48.8000°S, 179.2667°E, 723–729 m, 14 Nov. 1983; FAKU 104966 (1, 85.5 mm HL, 362 mm TL), Mernoo Bank, Chatham Rise, 44.7333°S, 175.7000°E, 16 Jul. 1968.

Coelorinchus kermadecus. 6 specimens. **Syntypes of *Coelorhynchus kermadecus***: BMNH 1887.12.7.60 (1, 161 mm HL, 549+ mm TL), BMNH 1887.12.7.61 (1, 84.7 mm HL, 253+ mm TL), BMNH 1887.12.7.62 (1, 56.8 mm HL, 176+ mm TL), west of Kermadec Islands, 29.7500°S, 178.1833°E, *Challenger* sta. 170A, trawl, 14 Jul. 1874. **New Zealand**: FSFRL ED249 (1, 116 mm HL, 404+ mm TL); FSFRL ED246 (1, 133 mm HL, 426+ mm TL); FSFRL ED245 (1, 105 mm HL, 295+ mm TL).

Coelorinchus labiatus. 2 specimens. **Mid-Atlantic Ridge**: ZMUB 19303 (counts and measurements taken on 2 of 13, 89.0–106 mm HL, 393+–414+ mm TL), 49.8667°N, 29.6333°W, 966–1019 m, 15 Jul. 2004.

Coelorinchus lasti. 2 specimens. **Holotype of *Caelorinchus lasti***: CSIRO H2615-02 (158 mm HL, 613+ mm TL), west of Mandurah, 32.5500°S, 114.4300°E, 1140 m, FRV *Southern Surveyor*, sta. SS1/91/83, 1991. **Paratype of *C. lasti***: CSIRO H1925-01 (139 mm HL, 456+ mm TL), west of Temma, 41.2500°S, 143.9667°E, 959–1021 m, 18 Feb. 1989.

Coelorinchus macrorhynchus. 3 specimens. **Holotype of *Coelorhynchus macrorhynchus***: USNM 72944 (181 mm HL, 551+ mm TL), Verde Island Passage, 13.5769°N, 121.1250°E, 329 m, *Albatross* sta. 5367, 12-ft Agassiz beam trawl, 22 Feb. 1909. **Indonesia**: CSIRO LM160 (1, 231 mm HL), CSIRO LM161 (1, 250 mm

HL).

***Coelorinchus mayiae*.** 11 specimens. **Holotype of *Caelorinchus mayiae*:** NTM S.13577-001 (52.6 mm HL, 222 mm TL), Arafura Sea, 9.0850°S, 133.6617°E, 165–176 m, coll. R. Williams, 20 Oct. 1992. **Paratypes of *C. mayiae*:** NTM S.13577-024 (6, 45.1–56.0 mm HL, 181–224+ mm TL), collected with holotype; NTM S.13580-001 (2, 54.5–71.4 mm HL, 169+–282 mm TL), Arafura Sea, 9.0783°S, 133.0783°E, 179–205 m, 20 Oct. 1992. **Arafura Sea:** NTM S.13578-001 (1 of 6, paratype of *Caelorinchus pardus*, 48.2 mm HL, 195+ mm TL), 8.9217°S, 133.6850°E, 179–187 m, 20 Oct. 1992; NTM S.13313-013 (counts and measurements taken on 1 of 3, 44.5 mm HL, 166 mm TL), 9.3000°S, 133.2000°E, 100–160 m, 6 Nov. 1992.

***Coelorinchus mediterraneus*.** 4 specimens. **Paratypes of *Coelorinchus mediterraneus*:** CAS 213887 (4, 49.9–61.8 mm HL, 167+–222+ mm TL), Adriatic Sea, Mediterranean, 41.7057°N, 17.5770°E, 1085–1159 m, Oct. 2000.

***Coelorinchus melanobranchus*.** 9 specimens. **Holotype of *Caelorinchus melanobranchus*:** MNHN 1994-911 (43.9 mm HL, 201 mm TL), off southern tip of New Caledonia, 22.9917°S, 167.3667°E, 490–515 m, *MUSORSTOM 4*, sta. CP216, 29 Sept. 1985. **Paratypes of *C. melanobranchus*:** CAS 86493 (1, 49.2 mm HL, 207 mm TL), off southern tip of New Caledonia, 22.9283°S, 167.2833°E, 485–520 m, 28 Sept. 1985; MNHN 1994-910 (1, 33.3 mm HL, 152 mm TL), off southern tip of New Caledonia, 22.8550°S, 167.2000°E, 405–530 m, 28 Sept. 1985. **Tasman Sea:** CAS 218440 (3, 67.2–76.7 mm HL, 319+–361+ mm TL), West Norfolk Ridge, 32.5928°S, 167.6468°E, 348–362 m, 30 May 2003; AMS I.42758-001 (1, 71.2 mm HL, 320+ mm TL), AMS I.42758-002 (1, 67.8 mm HL, 312+ mm TL), West Norfolk Ridge, 32.5928°S, 167.6469°E, 348–362 m, 30 May 2003; NSMT-P 121969 (1, 47.6 mm HL, 171+ mm TL), Derwent Hunter Guyot, 30.5817°S, 156.1550°E, 467–476 m, 13 Dec. 1976.

***Coelorinchus melanosagmatus*.** 19 specimens. **Holotype of *Caelorinchus melanosagmatus*:** CAS 50127 (53.9 mm HL, 171+ mm TL), Andaman Sea, 10.6500°N, 97.1000°E, 293 m, R/V *Anton Bruun*, cr. 1, sta. 22B, 24 Mar. 1963. **Paratypes of *C. melanosagmatus*:** AMS I.39209-001 (2, ex. CAS 50127, 50.0–53.8 mm HL, 157+–164+ mm TL), CAS 203714 (counts and measurements taken on 10 of 11, 38.2–60.1 mm HL, 121+–200+ mm TL, ex. CAS 50127), NTM S.14764-001 (2, ex. CAS 50127, 46.9–47.1 mm HL, 146+–154+ mm TL), collected with holotype; CAS 50165 (1, 54.5 mm HL, 190+ mm TL), off Dar es Salaam, 6.8000°S, 39.8500°E, 300–400 m, 20 Nov. 1964; CAS 50130 (counts and measurements taken on 6 of 8, 40.3–63.5 mm HL, 134+–212+ mm TL), Andaman Sea, 10.6500°N, 96.5833°E, 384 m, 24 Mar. 1963.

***Coelorinchus multifasciatus*.** 1 specimen. **Holotype of *Caelorinchus multifasciatus*:** ZMMGU P.18117 (32.3 mm HL, 134+ mm TL), Sala y Gomez Ridge, 25.1167°S,

99.6667°W, 439–500 m, R/V *Prof. Shtokman*, cr. 18, sta. 2023, bottom trawl, 8 May 1987.

***Coelorinchus mycterismus*.** 4 specimens. **New Zealand:** FSFRL EI555 (1, 110 mm HL, 352+ mm TL), FSFRL EB749 (1, 109 mm HL, 369+ mm TL), FSFRL ED137 (1, 111 mm HL, 407+ mm TL), FSFRL ED122 (1, 77.9 mm HL, 257+ mm TL).

***Coelorinchus obscuratus*.** 3 specimens. **Paratypes of *Coelorinchus obscuratus*:** CAS 90752 (1, 87.5 mm HL, 292+ mm TL), Norfolk Ridge, 25.5667°S, 167.4167°E, 1132–1160 m, 15 Nov. 1996; CAS 218477 (2, 64.6–75.3 mm HL, 211+–246+ mm TL), West Norfolk Ridge, 32.6065°S, 167.8432°E, 1331–1345 m, 30 May 2003.

***Coelorinchus occa*.** 3 specimens. **Caribbean Sea:** FMNH 113548 (1, 135 mm HL, 471+ mm TL), northwest of Honduras, 16.8000°N, 82.5500°W, 1006 m, 17 Sept. 1957. **Atlantic:** FMNH 49186 (1, 121 mm HL, 414+ mm TL), off Bermuda, 1908; FMNH 74632 (1, 121 mm HL, 427+ mm TL), no further data.

***Coelorinchus okamurai*.** 5 specimens. **Holotype of *Coelorinchus okamurai*:** MZB 23338 (ex. BSKU 16695, 121 mm HL, 457+ mm TL), south of Leti Islands, Timor Sea, 9.4500°S, 127.9767°E, 610–690 m, R/V *Hakuho-maru*, cr. KH-72-01, sta. 26, 3-m beam trawl, coll. O. Okamura, 18 Jun. 1972. **Paratypes of *C. okamurai*:** BSKU 16696 (1, 85.8 mm HL, 282+ mm TL), BSKU 16697 (1, 81.0 mm HL, 275+ mm TL), BSKU 16698 (1, 114 mm HL, 412+ mm TL), BSKU 16699 (1, 102 mm HL, 317+ mm TL), collected with holotype.

***Coelorinchus osipullus*.** 5 specimens. **Paratypes of *Coelorinchus osipullus*:** CAS 90984 (1, 116 mm HL, 396+ mm TL), Norfolk Ridge, 25.3833°S, 168.9498°E, 640–740 m, 12 Nov. 1996; CAS 90982 (1, 142 mm HL, 495 mm TL), Norfolk Ridge, 25.3322°S, 168.9427°E, 697–1340 m, 12 Nov. 1996. **Tasman Sea:** NSMT-P 101630 (1, 185 mm HL, 662+ mm TL), NSMT-P 101631 (1, 171 mm HL, 692+ mm TL), Norfolk Ridge, 29.8017°S, 168.0017°E, 611 m, 17 Jan. 1976. **New Zealand:** NSMT-P 33297 (1, 204 mm HL).

***Coelorinchus pardus*.** 22 specimens. **Holotype of *Caelorinchus pardus*:** NTM S.13551-001 (56.8 mm HL, 218+ mm TL), Arafura Sea, 9.6550°S, 133.0733°E, 123 m, 3 Oct. 1992. **Paratypes of *C. pardus*:** NTM S.13550-001 (5, 45.3–49.3 mm HL), Arafura Sea, 9.5117°S, 132.9383°E, 136 m, 6 Nov. 1990; NTM S.12903-011 (4, 47.6–58.7 mm HL, 171+–214+ mm TL), northeast of Cape Don, Arafura Sea, 9.3167°S, 133.2000°E, 152 m, 7 Nov. 1990; NTM S.13313-028 (2, 46.0–57.0 mm HL, 197+–220+ mm TL), Arafura Sea, 9.3000°S, 133.2000°E, 153 m, 6 Nov. 1990; CAS 200227 (4, 44.3–54.0 mm HL, 171+–197+ mm TL), Arafura Sea, 9.7050°S, 133.2867°E, 112–116 m, 3 Oct. 1992; AMS I.39216-001 (2, 41.1–43.0 mm HL, 168+–187+ mm TL), BSKU 85717 (1, ex. NTM S.13574-001, 49.2 mm HL, 193 mm TL), BSKU 85718 (1, ex. NTM S.13574-001, 117+ mm TL), Arafura Sea, 9.7067°S, 133.3883°E, 107 m, 3 Oct. 1992; NTM

S.13578-001 (counts and measurements taken on 2 of 6, 51.7–53.0 mm HL, 198+–212+ mm TL), Arafura Sea, 8.9217°S, 133.6850°E, 179–187 m, 20 Oct. 1992.

***Coelorinchus posteromaculatus*.** 9 specimens.

Holotype of *Coelorinchus posteromaculatus*: AMS I.44527-001 (66.4 mm HL, 211+ mm TL), south of Lombok Island, eastern Indian Ocean, 8.9700°S, 116.0800°E, 314 m, 1 Aug. 1981. **Paratypes of *C. posteromaculatus*:** AMS I.44527-002 (2, 57.8–65.3 mm HL, 165+–187+ mm TL), collected with holotype; BSKU 106747 (1, 63.3 mm HL, 196+ mm TL), Thailand, no further data; IORAS 001-022 (2, 62.3–70.0 mm HL, 203+–220+ mm TL), Andaman Sea, 11.7675°N, 96.8511°E, 315–323 m, 11 Mar. 1966; HUMZ 190722 (1, 58.2 mm HL, 197+ mm TL), HUMZ 190723 (1, 61.0 mm HL, 197+ mm TL), HUMZ 190724 (1, 59.1 mm HL, 192+ mm TL), west of Sumatra, 1.1228°S, 98.4590°E, 100–170 m, 3 Oct. 2004.

***Coelorinchus quincunciatus*.** 11 specimens.

Holotype of *Coelorhynchus quincunciatus*: USNM 78213 (66.5 mm HL, 223+ mm TL), vicinity southern Luzon, 13.4708°N, 121.0750°E, 329 m, *Albatross* sta. 5293, 12-ft Agassiz beam trawl, 23 Jul. 1908. **Paratypes of *C. quincunciatus*:** USNM 137824 (counts and measurements taken on 3 of 7, 41.5–46.1 mm HL, 145+–176 mm TL), between Samar and Masbate, 12.2097°N, 124.0470°E, 247 m, 13 Mar. 1909. **Philippines:** CAS 33797 (counts and measurements taken on 1 of 3, 72.9 mm HL, 261+ mm TL), Tayabas Bay, northwest of Marinduque Island, 13.5825°N, 121.8518°E, 165–196 m, 13 Sept. 1966; CAS 34450 (1, 51.1 mm HL, 185+ mm TL), Batangas Bay, 13.6916°N, 120.9746°E, 124–150 m, 25 Jul. 1966; CAS 34135 (counts and measurements taken on 2 of 3, 40.6–46.2 mm HL, 155+–180+ mm TL), Batangas Bay, 13.6916°N, 120.9746°E, 152–181 m, 25 Jul. 1966; CAS 34902 (1, 59.8 mm HL, 235+ mm TL), Batangas Bay, 13.6916°N, 120.9746°E, 219–245 m, 9 Aug. 1966; CAS 64570 (1, 55.7 mm HL, 207+ mm TL), Batangas Bay, 13.6916°N, 120.9746°E, 283–315 m, 18 Aug. 1966; CAS 34604 (1, 57.1 mm HL, 231 mm TL), Batangas Bay, 13.6916°N, 120.9746°E, 313–329 m, 2 Aug. 1966.

***Coelorinchus radcliffei*.** 1 specimen. **Holotype of *Coelorhynchus radcliffei*:** USNM 78224 (76.9 mm HL, 259+ mm TL), off northern Mindanao, Bohol Sea, 8.6072°N, 124.6020°E, 413 m, *Albatross* sta. 5503, 12-ft Tanner beam trawl, 4 Aug. 1909.

***Coelorinchus semaphoreus*.** 2 specimens. **Holotype of *Coelorinchus semaphoreus*:** MNHN 1996-956 (103 mm HL, 342+ mm TL), New Caledonia, 18.9500°S, 163.2100°E, 485 m, *MUSORSTOM 4*, sta. CP170, 17 Sept. 1985. **Paratype of *C. semaphoreus*:** CSIRO H1188-06 (1, 48.8 mm HL, 162+ mm TL), east of Flinders Reef, Coral Sea, 17.5200°S, 149.6700°E, 400–402 m, 3 Dec. 1985.

***Coelorinchus sereti*.** 7 specimens. **Holotype of *Coelorinchus sereti*:** MNHN 1994-925 (119 mm HL,

423+ mm TL), New Caledonia, 22.3108°S, 167.3883°E, 825 m, *BIOCAL*, sta. CP75, 4 Sept. 1985. **Paratypes of *C. sereti*:** MNHN 1994-924 (2, 107–117 mm HL, 381+–436+ mm TL), Chesterfield and Bellona Plateau, 19.8967°S, 158.6333°E, 412–430 m, 15 Oct. 1986; MNHN 1994-923 (1, 62.1 mm HL), MNHN 1994-926 (1, 124 mm HL, 437+ mm TL), Chesterfield and Bellona Plateau, 19.7067°S, 158.8467°E, 756–772 m, 21 Oct. 1986; CAS 86484 (1, 101 mm HL, 362+ mm TL), New Caledonia, 21.5635°S, 166.4512°E, 760–790 m, 12 Apr. 1987; CAS 86492 (1, 98.5 mm HL, 367+ mm TL), New Caledonia, 23.1163°S, 166.8533°E, 825 m, 29 Aug. 1985.

***Coelorinchus sexradiatus*.** 10 specimens.

Philippines: BSKU 15877 (1, 55.6 mm HL, 199+ mm TL), BSKU 15878 (1, 57.2 mm HL, 212+ mm TL), BSKU 15880 (1, 53.8 mm HL, 202+ mm TL), BSKU 15881 (1, 52.2 mm HL, 180+ mm TL), BSKU 15882 (1, 47.0 mm HL, 175+ mm TL), BSKU 15883 (1, 48.8 mm HL, 168+ mm TL), BSKU 15884 (1, 45.6 mm HL, 173+ mm TL), BSKU 15885 (1, 44.1 mm HL, 168+ mm TL), BSKU 15886 (1, 44.9 mm HL, 167+ mm TL), BSKU 15887 (1, 47.4 mm HL, 172+ mm TL), Sibutu Passage, Sulu Sea, 5.6817°N, 119.7717°E, 460–515 m, 10 Jun. 1972.

***Coelorinchus shcherbachevi*.** 2 specimens.

Holotype of *Coelorinchus shcherbachevi*: MNHN 1996-966 (ex. NMNZ P.27508, 130 mm HL, 453+ mm TL), Seamount K, Loyalty Ridge, 24.7425°S, 170.1167°E, 790–825 m, *BERYX 2*, sta. 9, 26 Oct. 1991. **Paratype of *C. shcherbachevi*:** CAS 90995 (123 mm HL, 440+ mm TL), Espoir K, Loyalty Ridge, 24.3000°S, 169.9500°E, 805–835 m, 8 Nov. 1996.

***Coelorinchus spilonotus*.** 2 specimens. **Paratypes of *Coelorinchus spilonotus*:** ZMMGU P.17707 (counts and measurements taken on 2 of 6, 49.4–58.5 mm HL, 188–199 mm TL), Sala y Gomez Ridge, 25.7667°S, 86.5500°W, 410 m, 31 Oct. 1979.

***Coelorinchus spinifer*.** 1 specimen. **South China Sea:** ASIZP 66748 (52.9 mm HL, 177 mm TL), 20.7333°N, 117.6500°E, 1098 m, 19 Aug. 2005.

***Coelorinchus supernasutus*.** 3 specimens. **New Zealand:** NSMT-P 41297 (1, 129 mm HL, 397+ mm TL), NSMT-P 41298 (1, 127 mm HL, 403+ mm TL), NSMT-P 41299 (1, 119 mm HL, 365+ mm TL), Three Kings Ridge, 30.2500°S, 172.9300°E, 737 m, 24 Apr. 1985.

***Coelorinchus thompsoni*.** 8 specimens. **Holotype of *Coelorhynchus thompsoni*:** USNM 78214 (1, 75.8 mm HL, 207+ mm TL), off Luzon, Balayan Bay, 13.7889°N, 120.7250°E, 329 m, *Albatross* sta. 5363, 12-ft Agassiz beam trawl, 20 Feb. 1909. **Paratypes of *C. thompsoni*:** USNM 149838 (1, 57.1 mm HL, 177+ mm TL), Balayan Bay and Verde Island Passage, 13.8125°N, 120.6980°E, 291 m, 21 Jan. 1908; USNM 148886 (counts and measurements taken on 1 of 2, 56.5 mm HL, 165+ mm TL), off northern Mindanao, Bohol Sea, 8.8000°N, 123.5200°E, 366 m, 9 Aug. 1909. **Philippines:** CAS 57426 (1, 62.5 mm HL, 200+ mm TL), east of Gaspar

Island, 13.2484°N, 121.8913°E, 134–155 m, 29 Sept. 1966; CAS 33787 (counts and measurements taken on 1 of 2, 67.1 mm HL, 193+ mm TL), Batangas Bay, 13.6916°N, 120.9746°E, 362–373 m, 6 Aug. 1966; CAS 64569 (1, 68.0 mm HL, 227+ mm TL), Balayan Bay, 13.8032°N, 120.7984°E, 238–254 m, 19 Jul. 1966; CAS 34551 (1, 66.6 mm HL, 226+ mm TL), Balayan Bay, 13.8032°N, 120.7984°E, 408–441 m, 28 Jul. 1966; CAS 34537 (1, 62.0 mm HL, 208+ mm TL), Balayan Bay, 13.8323°N, 120.7370°E, 216–236 m, 27 Jul. 1966.

***Coelorinchus thurla*.** 7 specimens. **Holotype of *Caelorinchus thurla*:** WAM P.21638-43 (48.7 mm HL, 207+ mm TL), North West Shelf, 13.7511°S, 123.5000°E, *Umitaka-maru*, date unknown. **Paratypes of *C. thurla*:** WAM P.21638-43 (in part, 4, 28.6–44.2 mm HL, 106+–186 mm TL), collected with holotype; WAM P.26270-016 (2, 47.5 mm HL, 198–203+ mm TL), North West Shelf, 15.5000°S, 120.9667°E, 280–320 m, 28 Jun. 1978.

***Coelorinchus trachycarus*.** 3 specimens. **Paratype of *Caelorinchus trachycarus*:** CAS 90983 (107 mm HL, 415+ mm TL), Norfolk Ridge, 24.1897°S, 167.7587°E, 1349–1350 m, 21 Nov. 1996. **New Zealand:** BMNH 1887.12.7.59 (1, 114 mm HL, 448+ mm TL), off East Cape, Gisborne, 37.5667°S, 179.3667°E, 10 Jul. 1874; FSFRL EH090 (1, 112 mm HL, 434+ mm TL).

***Coelorinchus trunovi*.** 8 specimens. **Indian Ocean:** ZMUC P373407 (1, 54.7 mm HL, 234 mm TL), ZMUC P373408 (1, 52.3 mm HL, 204+ mm TL), ZMUC P373409 (1, 91.1 mm HL, 365 mm TL), ZMUC P373410 (1, 111 mm HL, 385+ mm TL), R/V *Galathea*, sta. 196, 14 Feb. 1951; ZMUC P373112 (1, 76.5 mm HL, 271+ mm TL), ZMUC P373114 (1, 44.4 mm HL, 186 mm TL), R/V *Galathea*, sta. 197, 14 Feb. 1951; ZMUC P373066 (1, 106 mm HL, 341+ mm TL), R/V *Galathea*, sta. 202, 21 Feb. 1951; ZMUC P373117 (1, 123 mm HL, 432+ mm TL), R/V *Galathea*, sta. 203, 21 Feb. 1951.

***Coelorinchus velifer*.** 3 specimens. **Philippines:** BSKU 15876 (1, 52.1 mm HL, 188+ mm TL), BSKU 15879 (1, 50.4 mm HL, 200+ mm TL), BSKU 15897 (1, 37.1 mm HL, 155+ mm TL), Sibutu Passage, Sulu Sea, 5.6817°N, 119.7717°E, 460–515 m, 10 Jun. 1972.

***Coelorinchus weberi*.** 1 specimen. **Holotype of *Coelorhynchus weberi*:** USNM 78225 (88.0 mm HL, 305+ mm TL), off northern Luzon, South China Sea, 18.5708°N, 121.8540°E, 410 m, *Albatross* sta. D. 5325, 12-ft Tanner beam trawl and mud bag, 12 Nov. 1908.

***Coelorinchus yurii*.** 5 specimens. **Paratypes of *Coelorinchus yurii*:** CAS 223466 (3, 71.8–77.8 mm HL, 228+–256 mm TL), Saya de Malha Bank, Mascarene Ridge, 9.5450°S, 60.0333°E, 800 m, 26 Sept. 1977; CAS 66472 (2, 83.8–94.3 mm HL, 284+–316+ mm TL), off Saya de Malha Bank, Mascarene Ridge, 8.5333°S, 59.6833°E, 960–1130 m, 10 Jan. 1989.

***Coryphaenoides brevibarbis*.** 10 specimens. **Mid-Atlantic Ridge:** ZMUB 17797 (1, 73.5 mm HL), 53.0333°N, 33.6167°W, 2995–3071 m, 24 Jul. 2004.

Greenland: BSKU 81833 (1, 42.4 mm HL, 191+ mm TL), BSKU 81834 (1, 38.2 mm HL, 178+ mm TL), BSKU 81835 (1, 34.3 mm HL, 163+ mm TL), BSKU 81836 (1, 29.3 mm HL, 146+ mm TL), BSKU 81837 (1, 36.0 mm HL, 147+ mm TL), BSKU 81838 (1, 29.9 mm HL, 104+ mm TL), BSKU 81839 (1, 26.3 mm HL, 116+ mm TL), BSKU 81840 (1, 26.1 mm HL, 115+ mm TL), Davis Strait, 63.1840°N, 54.6808°W, 1495 m, 6 Aug. 1991; BSKU 80134 (1, 41.1 mm HL, 216 mm TL), Davis Strait, 63.1843°N, 54.6808°W, 1493 m, 6 Aug. 1991.

***Coryphaenoides fernandezianus*.** 3 specimens. **Holotype of *Macrurus fernandezianus*:** BMNH 1887.12.7.110 (52.8 mm HL, 286 mm TL), south of Juan Fernández Islands, 33.7000°S, 78.3000°W, 1375 m, *Challenger* sta. 300, trawl, 17 Dec. 1875. **Australia:** CSIRO H5313-11 (2, 45.0–48.6 mm HL, 204+–232+ mm TL), east of Flinders Island, 39.7833°S, 149.0833°E, 29 Apr. 2000.

***Coryphaenoides mediterraneus*.** 3 specimens. **Gulf of Mexico:** FMNH 64081 (1, 133 mm HL, 728 mm TL), southwest of New Orleans, 28.7833°N, 87.8333°W, 2103 m, 26 Mar. 1955. **Porcupine Seabight:** ZMUC P372980 (1, 90.7 mm HL, 437+ mm TL), southwest of Ireland, 51.3333°N, 12.5167°W, 24 Sept. 1983. **Greenland:** ZMUC P374426 (1, 114 mm HL, 544+ mm TL), Irminger Sea, 61.8333°N, 40.1500°W, 1455 m, 7 Jul. 1998.

***Coryphaenoides murrayi*.** 2 specimens. **Tasmania:** CSIRO H711-03 (1, 48.6 mm HL, 263+ mm TL), west of Granville Harbour, 41.7833°S, 144.3500°E, 1360 m, 20 Jul. 1986; CSIRO H1566-13 (1, 58.9 mm HL, 336 mm TL), south of King Island, 40.9833°S, 143.7000°E, 1253 m, 15 Sept. 1985.

***Coryphaenoides profundicola*.** 1 specimen. **Mid-Atlantic Ridge:** ZMUB 19503 (1, 197 mm HL, 1144 mm TL), 51.3333°N, 28.8667°W, 3505–3527 m, 16 Jul. 2004.

***Coryphaenoides tydemani*.** 3 specimens. **Indonesia:** BSKU 97929 (1, 49.7 mm HL, 248+ mm TL), BSKU 97930 (1, 60.4 mm HL, 294 mm TL), BSKU 97932 (1, 64.3 mm HL, 291+ mm TL), eastern Java Sea, 6.3183°S, 119.5817°E, 936–960 m, 12 Feb. 1985.

***Hymenocephalus aterrimus*.** 1 specimen. **Paratype of *Hymenocephalus aterrimus*:** FMNH 55952 (1, 34.1 mm HL, 126+ mm TL), Hawaiian Islands.

***Hymenocephalus longipes*.** 15 specimens. **Philippines:** BSKU 15519 (1), BSKU 15520 (1), BSKU 15521 (1), BSKU 15522 (1), BSKU 15523 (1), BSKU 15524 (1), BSKU 15525 (1), BSKU 15526 (1), BSKU 15527 (1), BSKU 15528 (1), BSKU 15529 (1), BSKU 15530 (1), BSKU 15531 (1), BSKU 94745 (3), BSKU 97414 (1), Sulu Sea, 8.2117°N, 117.9933°E, 285–306 m, 26 Jun. 1972.

***Kuronezumia bubonis*.** 3 specimens. **Western Australia:** NSMT-P 113029 (1, 116 mm HL, 639+ mm TL), northwest of Barrow Island, 20.0200°S, 114.3600°E, 805 m, 12 Nov. 1975. **New Zealand:** NSMT-P 42974 (1,

53.5 mm HL, 347+ mm TL), Mernoo Bank, Chatham Rise, 42.7500°S, 176.7670°E, 747–751 m, 24 Nov. 1983. **Suriname:** NSMT-P 42280 (1, 102 mm HL), off Paramaribo, 7.9167°N, 53.9833°W, 920 m, 15 Jun. 1981.

***Kuronezumia leonis*.** 3 specimens. **South Africa:** BSKU 48494 (1, 69.0 mm HL, 405+ mm TL), southeast of Port Elizabeth, 34.4433°S, 26.0133°E, 750 m, 3 Jun. 1990. **Louisville Ridge:** NSMT-P 41355 (1, 74.1 mm HL, 451+ mm TL), 41.4670°S, 164.2000°W, 874 m, 4 Apr. 1985; NSMT-P 41187 (1, 91.3 mm HL, 569+ mm TL), 40.9170°S, 165.1170°W, 857 m, 12 Apr. 1985.

***Nezumia africana*.** 11 specimens. **Holotype of *Ventrifossa africana*:** USNM 202771 (49.5 mm HL, 237+ mm TL), Gulf of Guinea, 4.4200°N, 9.4000°W, 641–733 m, R/V *Pillsbury*, cr. 6405, sta. 74, 4 Jun. 1964. **Angola:** CAS 223135 (1, 40.4 mm HL, 189+ mm TL), off Lobito, 12.3833°S, 13.2833°E, 729–733 m, 1 Apr. 2005; CAS 224397 (1, 49.7 mm HL, 255+ mm TL), off Lobito, 12.3833°S, 13.2833°E, 729–733 m, 1 Apr. 2005; CAS 224399 (1, 67.5 mm HL, 337+ mm TL), off Lobito, 11.9333°S, 13.3333°E, 652–658 m, 3 Apr. 2005; CAS 223136 (2, 39.5–43.3 mm HL, 169+–193+ mm TL), off Porto Amboim, 10.9333°S, 13.3500°E, 640–647 m, 5 Apr. 2005; CAS 223172 (1, 44.9 mm HL, 200+ mm TL), off Luanda, 8.4500°S, 12.7667°E, 704–716 m, 12 Apr. 2005; CAS 223139 (1, 45.2 mm HL, 248+ mm TL), off Ambriz, 7.7833°S, 12.5000°E, 749–758 m, 18 Apr. 2005; CAS 224398 (1, 57.3 mm HL, 263+ mm TL), off Santo António do Zaire, 6.6167°S, 11.4500°E, 654–655 m, 22 Apr. 2005. **Ghana:** CAS 233908 (1, 55.6 mm HL, 264+ mm TL), off Takoradi, 4.1307°N, 1.7603°W, 814–830 m, 4 May 2010. **Southeastern Atlantic:** CAS 224926 (1, 43.4 mm HL, 202+ mm TL), 719–730 m, 7 Mar. 2007.

***Nezumia aspidontata*.** 1 specimen. **Holotype of *Nezumia aspidontata*:** MNHN 1996-958 (34.2 mm HL, 193 mm TL), off New Caledonia, 19.6500°S, 158.8167°E, 700–720 m, 18 Oct. 1986.

***Nezumia atlantica*.** 18 specimens. **French Guiana:** NSMT-P 105011 (1, 84.3 mm HL), off Mana, 7.5833°N, 53.0500°W, 922 m, 3 Feb. 1980. **Suriname:** NSMT-P 42338 (1, 67.8 mm HL, 341 mm TL), off Paramaribo, 7.7830°N, 53.7170°W, 850 m, 19 Jul. 1981; BSKU 44463 (1, 55.7 mm HL, 271+ mm TL), off Paramaribo, 7.7833°N, 53.7167°W, 850 m, 19 Jul. 1991; BSKU 108867 (1, 70.7 mm HL, 369+ mm TL), BSKU 108868 (1, 61.0 mm HL, 284+ mm TL), off Paramaribo, 7.8200°N, 53.6833°W, 6 Dec. 1973; NSMT-P 40182 (1, 70.5 mm HL, 350+ mm TL), off Paramaribo, 7.8500°N, 54.3000°W, 800–820 m, 27 Apr. 1980; NSMT-P 42346 (1, 66.7 mm HL, 325+ mm TL), off Paramaribo, 7.8830°N, 54.4330°W, 823–825 m, 21 Jul. 1979; NSMT-P 42345 (1, 61.4 mm HL, 296+ mm TL), off Paramaribo, 7.8830°N, 54.2670°W, 810 m, 31 Jan. 1981; BSKU 44462 (1, 58.8 mm HL, 286+ mm TL), NSMT-P 42301 (1, 59.2 mm HL, 229+ mm TL), off Paramaribo, 7.8833°N, 54.2500°W, 810–820 m, 31 Jan. 1980; NSMT-P 31640 (4, 57.0–67.7 mm HL, 278+–375+

mm TL), off Paramaribo, 7.8900°N, 54.2500°W, 810–820 m, 31 Jan. 1980; NSMT-P 42303 (1, 71.3 mm HL, 360+ mm TL), off Paramaribo, 7.9167°N, 53.9833°W, 920 m, 15 Jun. 1981; NSMT-P 42339 (1, 70.1 mm HL, 343+ mm TL), off Paramaribo, 7.9500°N, 54.1330°W, 930 m, 15 Jun. 1981. **Caribbean Sea:** CAS 14541 (1, 78.5 mm HL, 417+ mm TL), southwest of Curaçao Island, 11.9000°N, 69.3000°W, 914 m, 4 Oct. 1963. **Gulf of Mexico:** CAS 14540 (1, 92.2 mm HL, 448+ mm TL), off Louisiana, 27.7000°N, 90.5417°W, 914 m, 22 Feb. 1964.

***Nezumia burragei*.** 1 specimen. **Holotype of *Macrourus burragei*:** USNM 51641 (44.2 mm HL, 254+ mm TL), off Oahu, 21.2666°N, 157.8700°W, 386 m, *Albatross* sta. 3810, 8-ft Tanner trawl, 27 Mar. 1902.

***Nezumia holocentra*.** 6 specimens. **Paratype of *Macrourus holocentrus*:** USNM 47723 (1, 39.6 mm HL, 171+ mm TL), off Hawaii, 21.2000°N, 157.6417°W, 686 m, 6 Dec. 1981. **Hawaii:** BPBM 24579 (5, 30.9–38.6 mm HL, 151+–181+ mm TL), 650 m, 10 Mar. 1971.

***Nezumia infranudis*.** 2 specimens. **Holotype of *Lionurus infranudis*:** USNM 82669 (40.2 mm HL, 202+ mm TL), off Borneo, Celebes Sea, 4.1139°N, 118.7889°E, 635 m, *Albatross* sta. 5586, 28 Sept. 1909. **Timor Sea:** BSKU 16686 (1, 32.9 mm HL, 174+ mm TL), 9.4500°S, 127.9767°E, 610–690 m, 18–19 Jan. 1972.

***Nezumia kensmithi*.** 14 specimens. **Northeastern Pacific:** NSMT-P 104501 (1, 87.8 mm HL, 357+ mm TL), no further data; NSMT-P 104502 (1, 78.1 mm HL, 404+ mm TL), NSMT-P 104901 (1, 84.9 mm HL, 375+ mm TL), NSMT-P 104902 (1, 84.6 mm HL, 467+ mm TL), NSMT-P 104905 (1, 72.1 mm HL, 411 mm TL), NSMT-P 104996 (1, 76.5 mm HL, 355+ mm TL), NSMT-P 105001 (1, 60.8 mm HL, 322+ mm TL), NSMT-P 105002 (1, 79.0 mm HL, 423+ mm TL), NSMT-P 105005 (1, 68.6 mm HL, 358+ mm TL), NSMT-P 105006 (1, 60.3 mm HL, 332+ mm TL), Fieberling Seamount, 32.4383°N, 127.7717°W, 13 Jan. 1979; FAKU 119097 (1, 61.3 mm HL, 302+ mm TL), FAKU 119099 (1, 70.3 mm HL, 258+ mm TL), FAKU 119100 (1, 80.3 mm HL, 398+ mm TL), FAKU 119101 (1, 67.1 mm HL, 329+ mm TL), Fieberling Seamount, 32.4400°N, 127.7400°W, 499–576 m, 12 Jan. 1979.

***Nezumia merretti*.** 3 specimens. **Holotype of *Nezumia merretti*:** CSIRO H2584-20 (54.3 mm HL, 244+ mm TL), off Western Australia, 18.0667°S, 112.7000°E, 853–854 m, 1 Feb. 1991. **Paratype of *N. merretti*:** CSIRO H1966-01 (1, 39.3 mm HL, 203+ mm TL), off Queensland, 18.9833°S, 150.5333°E, 879–886 m, 25 Nov. 1985. **Indonesia:** HUMZ 190198 (1, 36.7 mm HL), northwest of Sumatra Island, 5.4860°N, 93.7357°E, 568–590 m, 13 Oct. 2004.

***Nezumia namatahi*.** 4 specimens. **Tasman Sea:** AMS I.24507-009 (1, 57.1 mm HL, 360+ mm TL), off Kiama, 34.8833°S, 151.2333°E, 951–978 m, 24 Aug. 1983; AMS I.42743-007 (1, 47.5 mm HL, 282+ mm TL), Lord How Rise, 34.2028°S, 163.3561°E, 26 May 2003;

AMS I.29754-005 (1, 33.7 mm HL, 194+ mm TL), east of Seal Rocks, 32.5000°S, 153.0000°E, 1006–1052 m, 15 Aug. 1989; AMS I.24059-022 (1, 40.0 mm HL, 237+ mm TL), off Norah Head, 32.0050°S, 152.1500°E, 933–969 m, 23 Aug. 1983.

Nezumia stelgidolepis. 1 specimen. **Northeastern Pacific:** FAKU 119106 (1, 73.9 mm HL, 309+ mm TL), Dowd Seamount, 13.4967°N, 119.8883°W, 801–845 m, 10 Feb. 1979.

Pseudocetonurus septifer. 1 specimen. **Holotype of *Pseudocetonurus septifer*:** ZMMGU P.16011 (55.1 mm HL), Sala y Gomez Ridge, 25.3339°S, 93.5917°W, 850–860 m, date unknown.

Pseudocetonurus flagellicauda. 46 specimens. **Syntypes of *Macrurus flagellicauda*:** ZMUB 3383 (4, 55.1–67.8 mm HL, 295+–394+ mm TL), Mid-Atlantic Ridge, 45.4333°N, 25.4167°W, 3120 m, *Michael Sars*, sta. 88, 18 Jul. 1910. **Mid-Atlantic Ridge:** ZMUB 16356 (9, 37.5–58.3 mm HL, 205+–375+ mm TL), ZMUB 16357 (1, 40.8 mm HL, 231+ mm TL), 42.7667°N, 29.2667°W, 3005–3050 m, 11 Jul. 2004; ZMUB 16353 (1, 22.7 mm HL, 129+ mm TL), ZMUB 16354 (10, 31.5–59.7 mm HL, 179+–395 mm TL), 42.9167°N, 30.3333°W, 2660–2670 m, 7 Jul. 2004; ZMUB 16358 (1, 62.3 mm HL, 373+ mm TL), ZMUB 16359 (7, 37.4–61.6 mm HL, 270+–392 mm TL), ZMUB 16360 (counts and measurements taken on 6 of 11, ca. 33–61.4 mm HL, 205+–376+ mm TL), 42.9167°N, 28.1333°W, 2973–2979 m, 13 Jul. 2004. **Porcupine Seabight:** BMNH 2003.11.3.29–31 (3, 56.1–66.7 mm HL, 261+–331+ mm TL), 50.1717°N, 13.3717°W, 1 Apr. 1982; ZMH 115359 (1, 58.0 mm HL, 391+ mm TL), 50.2000°N, 13.6000°W, 3000 m, 6 May 1978; MNHN 2001-2728 (2, 50.4–57.8 mm HL, 277+–317+ mm TL), 50.2500°N, 13.2670°W, 2691 m, Jul. 1977. **Walters Shoals:** ZMMGU P.21618 (1, 45.9 mm HL, 282+ mm TL), 30.5000°S, 46.8833°E, 2580–2680 m, 26 Dec. 1988; ZMUB 16356 (1, 52.7 mm HL, 295+ mm TL).

Pseudocetonurus parvipes. 7 specimens. **Paratypes of *Macrourus parvipes*:** USNM 148896 (1, 31.9 mm HL, 212+ mm TL), Chenoki Point Makassar Strait, Celebes Island, Java Sea, 1.3167°S, 118.7167°E, 30 Dec. 1909; USNM 148893 (2, 37.3–52.2 mm HL, 179+–181+ mm TL), west of Binang Unauna Island, Togian Islands, Gulf of Tomini, 0.1833°S, 121.2667°E, 18 Nov. 1909; USNM 148894 (1, 39.6 mm HL, 273+ mm TL), northwest of Binang Unauna Island, Togian Islands, Gulf of Tomini, 0.1333°S, 121.3167°E, 18 Nov. 1909; USNM 148895 (1, 47.7 mm HL, 309+ mm TL), Gomomo Island, Piti Passage, Obi Islands, 1.9167°S, 127.7083°E, 3 Dec. 1909. **New Caledonia:** MNHN 1994-972 (1, 37.2 mm HL, 198+ mm TL), 22.1503°S, 167.5530°E, 2100 m, 4 Sept. 1985. **South China Sea:** BSKU 17717 (1, 47.3 mm HL, 272+ mm TL), off Ho Chi Minh, Vietnam, 9.2417°N, 109.6417°E, 1680–1720 m, 13 Jul. 1972.

Pseudocetonurus pusilla. 31 specimens. **Holotype of *Paracetonurus pusillus*:** ZMMGU P.15306 (33.4 mm

HL, 220+ mm TL), Ninety-Degree East Ridge, 11.4050°S, 88.8333°E, 1500–1600 m, R/V *Fiolent*, cr. 9, tr. 22, coll. Y.I. Sazonov, 13 Jan. 1977. **Paratypes of *P. pusillus*:** ZMMGU P.15307 (5, 32.3–34.7 mm HL, 167+–196+ mm TL), collected with holotype; ZMMGU P.15309 (1, 26.2 mm HL, 201+ mm TL), Ninety-Degree East Ridge, 11.2800°S, 87.8333°E, 1800–2000 m, 19 Jan. 1977; ZMMGU P.15308 (3, 30.8–35.8 mm HL, 155+–270+ mm TL), Ninety-Degree East Ridge, 11.2400°S, 88.7967°E, 1500–1600 m, 13 Jan. 1977; ZMMGU P.15311 (6, 26.0–32.7 mm HL, 151+–225 mm TL), Ninety-Degree East Ridge, 11.5167°S, 88.9233°E, 1600–1700 m, 18 Feb. 1979; ZMMGU P.15310 (1, 31.0 mm HL, 218+ mm TL), Mascarene Ridge, 12.7500°S, 61.1833°E, 1500 m, 31 Aug. 1977; ZMMGU P.14357 (1, 34.4 mm HL, 226+ mm TL), Broken Ridge, 25.8083°S, 100.1867°E, 1630–1740 m, 9 Oct. 1976. **Seychelles:** BMNH 1995.1.12.2 (1, 34.6 mm HL), BMNH 1996.8.5.1 (1, 33.1 mm HL), Saya de Malha Bank, 9.1694°S, 60.5194°E, 1520–1720 m, 9 Jan. 1989. **Ninety-Degree East Ridge:** ZMMGU P.15348 (2, 22.0–34.8 mm HL, 122+–207+ mm TL), 14.7067°S, 86.8233°E, 1760 m, 19 Mar. 1979; ZMMGU P.15312 (1, 29.4 mm HL, 161+ mm TL), west of Ninety-Degree East Ridge, 3.0500°S, 83.0417°E, 1680–1800 m, 9 Sept. 1979. **Western Australia:** CSIRO H2551-19 (1, 28.7 mm HL, 178+ mm TL), CSIRO H2551-20 (1, 30.2 mm HL, 185+ mm TL), west of North West Cape, 21.9667°S, 113.1333°E, 1460–1500 m, 25 Jan. 1991. **Loyalty Islands:** MNHN 1996-964 (1, 40.2 mm HL, 206+ mm TL), 21.0682°S, 167.0067°E, 1760–1870 m, 18 Apr. 1987. **South China Sea:** CAS 224176 (1, 27.6 mm HL, 173+ mm TL), south of Parcel Island, 15.6333°N, 111.9000°E, 1234–1264 m, 29 Feb. 1960; ASIZP 66424 (1, 30.1 mm HL, 221+ mm TL), 20.7333°N, 117.6500°E, 1098 m, 19 Aug. 2005; ASIZP 66945 (3, 20.5–32.9 mm HL, 135+–161+ mm TL), east of Dong-sha Atoll, 20.7451°N, 117.6538°E, 1098 m, 19 Aug. 2005.

Ventrifossa atherodon. 2 specimens. **Paratype of *Optonurus atherodon*:** FMNH 2552 (1, paratype, 40.3 mm HL, 171+ mm TL), Hawaiian Islands, 6 Dec. 1891. **Hawaii:** FMNH 55979 (1, 63.7 mm HL, 307+ mm TL).

Ventrifossa petersonii. 2 specimens. **Timor Sea:** BSKU 16714 (1, 33.9 mm HL, 184+ mm TL), BSKU 16715 (1, 33.7 mm HL, 181+ mm TL), 9.4500°S, 127.9767°E, 690–850 m, 18–19 Jun. 1972.

Ventrifossa sazonovi. 1 specimen. **Taiwan:** BSKU 98976 (1, 38.5 mm HL, 202+ mm TL), Dong-gang fish market, Pingtung, 16 Nov. 2007.

APPENDIX 3. Maps of the northwestern Pacific around Japan showing the distribution of each species of grenadier occurring in the area

Asthenomacrus victoris

Cetonurus globiceps

Coelorinchus anatirostris

Coelorinchus brevirostris

Coelorinchus cingulatus

Coelorinchus divergens

Coelorinchus formosanus

Coelorinchus fuscigulus

Coelorinchus gilberti

Coelorinchus hexafasciatus

Coelorinchus hige

Coelorinchus hubbsi

Coelorinchus japonicus

Coelorinchus jordani

Coelorinchus kamoharai

Coelorinchus kishinouyei

Coelorinchus lanceolatus sp. nov.

Coelorinchus leptorhinus

Coelorinchus longissimus

Coelorinchus macrochir

Coelorinchus matsubarai

Coelorinchus multispinulosus

Coelorinchus nox sp. nov.

Coelorinchus parallelus

Coelorinchus sheni

Coelorinchus smithi

Coelorinchus tokiensis

Coryphaenoides acrolepis

Coryphaenoides altipinnis

Coryphaenoides armatus

Coryphaenoides asper

Coryphaenoides cinereus

Coryphaenoides filifer

Coryphaenoides leptolepis

Coryphaenoides longifilis

Coryphaenoides marginatus

Coryphaenoides microps

Coryphaenoides nasutus

Coryphaenoides pectoralis

Coryphaenoides rudis

Coryphaenoides soyoae

Coryphaenoides yaquinae

Hymenocephalus hachijoensis

Hymenocephalus lethonemus

Hymenocephalus longibarbis

Hymenocephalus papyraceus

Hymenocephalus striatissimus

Hymenocephalus yamasakiorum

Hymenogadus gracilis

Kumba japonica

Kuronezumia dara

Kuronezumia endoi sp. nov.

Lucigadus nigromarginatus

Malacocephalus nipponensis

Nezumia condylura

Nezumia kamoharai

Nezumia proxima

Nezumia rara sp. nov.

Nezumia shinoharai

Nezumia spinosa

Nezumia tomiyamai

Odontomacrus murrayi

Pseudocetonurus sp. cf. septifer

Pseudonezumia cetonuropsis

Pseudonezumia japonica

Sphagemacrus pumiliceps

Spicomacrus kuronumai

Trachonurus villosus

Ventrifossa garmani

Ventrifossa johnboborum

Ventrifossa longibarbata

Ventrifossa macroptera

Ventrifossa misakia

Ventrifossa nigrodorsalis

Ventrifossa rhipidodorsalis

Ventrifossa saikaiensis