



## New combinations in the fern genus *Tectaria* (Tectariaceae) for the Flora of China

MAARTEN J.M. CHRISTENHUSZ

Botanical Garden and Herbarium, Finnish Museum of Natural History, University of Helsinki, 00014 Helsinki, Finland.  
phytotaxa@googlemail.com

### Abstract

For studies on the fern family Tectariaceae for the Flora of China, four species previously treated as *Ctenitopsis* need new combinations in *Tectaria*. The new combinations are proposed here and information on their types is provided.

### Introduction

The generic delimitation in Tectariaceae has been in considerable flux in the recent past. Molecular studies (e.g. Hasebe *et al.* 1995) have greatly increased the understanding of the placement of many genera and has redefined genera in Tectariaceae. Currently the family is considered to be composed of ca 10 genera (*Aenigmopteris*, *Arthropteris*, *Heterogonium*, *Hypoderris*, *Pleocnemia*, *Psammiosorus*, *Psomiocarpa*, *Pteridrys*, *Tectaria* and *Triplophyllum*). The genera *Ataxipteris*, *Ctenitis*, *Dryopsis* and *Lastreopsis*, previously associated with Tectariaceae, are now considered to belong to the Dryopteridaceae (Smith *et al.* 2006), and Liu *et al.* (2007) also placed *Pleocnemia* tentatively in that family. *Arthropteris* and *Psammiosorus* form a group sister to the rest of Tectariaceae (Hasebe *et al.* 1995), and for this reason are placed there (Liu *et al.* 2007), even though this makes the family difficult to define morphologically (Smith *et al.* 2006). Genera such as *Amphiblestra*, *Camptodium*, *Chlamydogramme*, *Cionidium*, *Ctenitopsis*, *Dictyoxiphium*, *Fadyenia*, *Hemigramma*, *Pleuroderris*, *Pseudotectaria* and *Quercifilix* are found to be embedded in *Tectaria* and are therefore included into that genus. Many combinations already exist to accommodate these generic changes, but for four of the Chinese species of *Ctenitopsis*, combinations are still needed. While revising the Tectariaceae for the *Flora of China*, I provide these combinations below with their type information.

### ***Tectaria acrocarpa* (Ching) Christenh., comb. nov.**

Basionym:—*Ctenitopsis acrocarpa* Ching in Ching & Wang (1981: 124).

Type:—CHINA: Yunnan: Jinping, S.K. Wu 3985 (holotype PE).

### ***Tectaria chinensis* (Ching & Chu H.Wang) Christenh., comb. nov.**

Basionym:—*Ctenitopsis chinensis* Ching & Wang (1981: 124).

Type:—CHINA: Yunnan: Hekou, Department of Biology of Yunnan University 512 (holotype PE)

### ***Tectaria sagenioides* (Mett.) Christenh., comb. nov.**

Basionym:—*Aspidium sagenioides* Mettenius (1858: 397).

Homotypic synonyms:—*Dryopteris sagenioides* (Mett.) Kuntze (1891: 813), *Ctenitopsis sagenioides* (Mett.)

Ching (1938: 312), *Ctenitis sagenioides* (Mett.) Copeland (1947: 124), *Heterogonium sagenioides* (Mett.) Holttum (1949: 161).

Type:—INDONESIA: Java: *Zollinger 1803* (lectotype Z, designated here; the holotype at B is no longer extant).

***Tectaria subsageniacea* (Christ) Christenh., comb. nov.**

Basionym:—*Aspidium subsageniaceum* Christ (1906: 240)

Homotypic synonyms:—*Ctenitopsis subsageniacea* (Christ) Ching (1938: 311), *Dryopteris subsageniacea* (Christ) Christensen (1913: 40), *Heterogonium subsageniaceum* (Christ) Holttum (1955: 273).

Type:—CHINA: Guizhou: Tien-Sen-Kiao, sud Tin-Fan, près rivière, nov. 1904, *Cavalerie 1916* (holotype P?, not found).

## References

- Ching, R.C. (1938) A revision of the Chinese and Sikkim Himalaya *Dryopteris* with reference to some species in the neighbouring regions. 7. *Ctenitopsis*. *Bulletin of the Fan Memorial Institute of Biology, Botany* 8: 304–321.
- Ching & Wang, C.H. (1964) Additional material for the Pteridophytioflora of Hainan. *Acta Phytotaxonomica Sinica* 9: 345–373.
- Ching & Wang, C.H. (1981) Breviarum plantarum novarum Aspidacearum Sinicarum. *Acta Phytotaxonomica Sinica* 19: 118–130.
- Christ, H. (1906) Filices Cavalerianae II. *Bulletin de l'Académie Internationale de Géographie Botanique* 16: 233–246.
- Christensen, C.F.A. (1913) *Index Filicum, Supplementum 1906–1912*. H. Hagerup, Copenhagen.
- Copeland, E.B. (1947) *Genera Filicum: The Genera of Ferns*. Chronica Botanica Co., Waltham, Massachusetts.
- Hasebe, M., Wolf, P.G., Pryer, K.M., Ueda, K., Ito, M., Sano, R., Gastony, G.J., Yokoyama, J., Manhart, J.R., Murakami, N., Crane, E.H., Haufler, C.H. & Hauk, W.D. (1995) Fern phylogeny based on *rbcL* nucleotide sequences. *American Fern Journal* 85: 134–181.
- Holttum, R.E. (1949) The fern genus *Heterogonium* Presl. *Sarawak Museum Journal* 5: 156–166.
- Holttum, R.E. (1955) Some additional species of *Heterogonium*. *Reinwardtia* 3: 269–274.
- Kuntze, O. (1891) *Revisio Generum Plantarum* vol. 2: 375–1011. A. Felix, Leipzig.
- Liu, H.M., Zhang, X.C., Chen, Z.D. Dong, S.Y. & Qiu, Y.L. (2007) Polyphyly of the fern family Tectariaceae sensu Ching: insights from cpDNA sequence data. *Science in China Series C: Life Sciences* 50: 789–798.
- Mettenius, G. (1858) Über einige FarnGattungen. IV. *Phegopteris* und *Aspidium*. *Abhandlungen herausgegeben von der Senckenbergischen Naturforschenden Gesellschaft* 2(4): 288–420.
- Smith, A.R., Pryer, K.M., Schuettpehl, E., Korall, P., Schneider, H. & Wolf, P.G. (2006) A classification for extant ferns. *Taxon* 55: 705–731.