



Correspondence

Notes on Early Land Plants Today. 11. *Microlejeunea bischlerae* (B.M.Thiers) comb. nov.

BARBARA M. THIERS¹, LARS SÖDERSTRÖM², ANDERS HAGBORG³ & MATT VON KONRAT³

¹William and Lynda Steere Herbarium, The New York Botanical Garden, Bronx, New York 10458-5126, USA; bthiers@nybg.org

²Department of Biology, Norwegian University of Science and Technology, N-7491 Trondheim, Norway; lars.soderstrom@bio.ntnu.no

³Department of Botany, The Field Museum, 1400 South Lake Shore Drive, Chicago, IL 60605–2496, USA; hagborg@pobox.com, mvonkonrat@fieldmuseum.org

In the past the genus *Microlejeunea* Stephani (1888) has often been regarded as a subgenus of *Lejeunea* Libert (1820). *Lejeunea bischlerae* Thiers (1997) was described using this concept. However, recent molecular studies (e.g. Wilson *et al.* 2004) resolve *Microlejeunea* as a sister of *Lejeunea*. If this concept is followed, a new combination is needed.

***Microlejeunea bischlerae* (B.M.Thiers) B.M.Thiers comb. nov.** Basionym:—*Lejeunea bischlerae* B.M.Thiers, *Cryptog. Bryol. Lichénol.* 18: 223, 1997 (Thiers 1997). Holotype:—AUSTRALIA, Queensland, Yungaburra, Gillies Road, near Mt. Nomico, 8 Sept. 1985, G. A. M. Scott s. n. (MUCV 6939; isotype in NY).

References

- Libert, M.-L. (1820) Sur un genre nouveau d'hépatiques, *Lejeunia*. *Annales générales des sciences physiques* 6: 372–374.
- Stephani, F. (1888) Hepaticae africanae. *Hedwigia* 27: 59–63.
- Thiers, B.M. (1997) *Lejeunea bischlerae*, a new species of *Lejeunea* subgenus *Microlejeunea* from Australia. *Cryptogamie, Bryologie et Lichénologie* 18: 223–226.
- Wilson, R., Gradstein, S.R., Heinrichs, J., Groth, H., Ilku-Borges, A.L. & Hartmann, F.A. (2004) Phylogeny of Lejeuneaceae: a cladistic analysis of chloroplast gene rbcL sequences and morphology with preliminary comments on the mitochondrial NAD4-2 spacer region. *Monographs in Systematic Botany from the Missouri Botanical Garden* 98: 189–202.