



<http://dx.doi.org/10.11646/phytotaxa.170.2.5>

***Rhaponticoides gokceoglu* (Asteraceae), a striking new species from Turkey**

İLKER ÇİNİLGEL¹, ÖZKAN EREN^{2*} & HAYRİ DUMAN³

¹ Akdeniz University, Akseki Vocational School, Program of Hunting and Wildlife, 07630, Akseki, Antalya, Turkey; email: icinbilgel@akdeniz.edu.tr

² Adnan Menderes University, Faculty of Arts and Sciences, Department of Biology, 09100, Aydin, Turkey; *corresponding author; email: ozkaneren@adu.edu.tr

³ Gazi University, Faculty of Arts and Sciences, Department of Biology, 06500, Teknikokullar, Ankara, Turkey; email: hduman@gazi.edu.tr

Abstract

Rhaponticoides gokceoglu is described as a species new to science. Its affinities within the genus are discussed. The new species is closely related to *R. wagenitziana* and *R. aytachii* from which we delimit it. *Rhaponticoides gokceoglu* is readily distinguished from those species by its entirely leafy stem, lacking a distinct appendage on phyllaries, ovate outer and ovate-oblong to oblong middle phyllaries with a very narrow scarious margin, and inner phyllaries cucullate at apex only. Habitat conditions and conservation status of *R. gokceoglu* are considered. Finally, a distribution map and a key to the pinkish-flowered species of *Rhaponticoides* in Turkey are provided.

Key words: Antalya, Compositae, taxonomy

Introduction

The genus *Rhaponticoides* Vaillant (1754: 165) was recently split from *Centaurea* Linnaeus (1753: 909). It occurs from Portugal and Morocco in the west to Mongolia in the east (Agababian 1997, Greuter 2003, Hellwig 2004, Greuter et al. 2005, Garcia-Jacas et al. 2006). If we exclude *R. ruthenica* (Lamarck 1785: 663) Agababian & Greuter (2003: 61) which occurs over large areas, most of the representatives of the genus are narrow endemics (Wagenitz 1975, 1986, Eren 2007, Puntillo & Peruzzi 2009, Tan et al. 2009, Doğu et al. 2009). In addition, there are only a few species such as *R. wagenitziana* (Bancheva & Kit Tan 2009: 52) Agababian & Greuter in Greuter & Raus (2009: 337) which display disjunct distributions.

Recently, six species of *Rhaponticoides* were recognised in Turkey (Eren 2007). Eren provided the first key, as well as a distribution map of all Turkish representatives of the genus. Since that publication, *R. aytachii* Bağcı, Doğu & Dinç in Doğu et al. (2009: 479) and *R. wagenitziana* were described (Tan et al. 2009, Greuter & Raus 2009). The latter is described as a species new to science on the basis of plants collected in Bulgaria and Turkey, both formerly identified as *R. amplifolia* (Boissier & Heldreich in Boissier 1856: 68) Agababian & Greuter (2003: 60) in standard Floras of those countries and subsequent publications. This statement was no longer supported by Tan et al. (2009) who recognized the Turkish and Bulgarian populations as a distinct species. As indicated by Tan et al. (2009), *R. amplifolia* became endemic to the Peloponnese in southern Greece.

In early July 2010 a very striking species of *Rhaponticoides*, distinct from any other Turkish member of this genus, was collected in Suçikan mevkii, above Başlar village that belongs to İbradı district in the province of Antalya. The few plants gathered in early July 2010 were complemented by collections conducted in June and July 2011 from the same site. The population of *Rhaponticoides* from the Suçikan mevkii turned out to represent a hitherto unknown species related to *R. wagenitziana* and *R. aytachii*. Hence, this species is described here as new to science, bringing a total number of the species of *Rhaponticoides* known from Turkey to eight. Seven of those species, namely, *R. gokceoglu* sp. nov., *R. aytachii*, *R. amasiensis* (Bornmüller 1906: 54) Agababian & Greuter (2003: 60), *R. pythiae* (Aznavour & Bornmüller in Bornmüller 1927: 48) Agababian & Greuter (2003: 61), *R. iconiensis* (Huber-Morath 1981: 77) Agababian & Greuter (2003: 60), *R. mykalea* (Huber-Morath 1979: 370) Agababian & Greuter (2003: 60), and *R. hierroi* Eren (2007: 15) are endemic to Turkey. *Rhaponticoides wagenitziana* is the only species which occurs both in Turkey and Bulgaria. The genus *Rhaponticoides* exhibits a high level of endemism in Turkey.

2. Terminal segments of basal leaves broadly ovate; stem arachnoid below; all phyllaries or at least the middle and inner ones distinctly cucullate *Rhaponticoides aytachii*
 – Terminal segments of basal leaves elliptic-lanceolate; stem puberulent below; phyllaries not cucullate *Rhaponticoides wagenitziana*

Specimens examined

***Rhaponticoides amplifolia*:**—GREECE. In montibus Kylleus faucibus Phlamburtya, T. *Orphanides* G 129 (E00394662, E00394663, type, photos!).

***Rhaponticoides aytachii*:**—TURKEY. C4 Karaman: Sariveliler, Dumlugöze köyü, bozulmuş *Quercus* ormanı, açıklıklar, 1000–1050 m, 20 June 2007, S. *Doğu* 1398, Bağcı & Dinç (GAZI, isotype).

***Rhaponticoides wagenitziana*:**—BULGARIA. Toundzha Hills, Municipality of Elhovo, north of the village of Golyam Dervent, 42°01'18.1" N, 26°43'00.0" E, 348 m, 26 July 2006, S. *Bancheva* & S. *Stoyanov* SB-07-06 (GAZI, isotype). TURKEY. A2 İstanbul: Aydos hill, on the slopes, 02 July 1974, *Tuzlaci* E 30115 (E00199159, E00474023, photos!).

Acknowledgements

We thank Dr. José L. Hierro (Argentina) for linguistic improvement of the text. Funding was provided by Akdeniz University Management Unit of Scientific Research Projects (project number 2008.03.0121.008).

References

- Agababian, M. (1997) *Centaurea* subg. *Centaurea* (Compositae): Delimitation and distribution of sections and subsections. *Lagascalia* 19: 889–902.
- Boissier, E. (1856) *Diagnoses plantarum orientalium novarum*, ser. 2, vol. 3. Lipsia [Leipzig], 177 pp.
- Bornmüller, J. (1906) *Centaurea amasiensis* Bornm. (sect. *Centaurium*), florae Anatoliae species indescripta nova. *Feddes Repertorium* 3: 54–55.
<http://dx.doi.org/10.1002/fedr.19060032906>
- Bornmüller, J. (1927) Über eine neue *Centaurea* (*C. pythiae* Aznav. et. Bornm.) der Flora Kleinasiens. *Mitteilungen des Thüringischen Botanischen Vereins, Neue Folge* 37: 48–49.
- Candolle, A.P. de (1838) *Prodromus systematis naturalis regni vegetabilis*, vol. 6. Treuttel & Würtz, Paris, 687 pp.
<http://dx.doi.org/10.5962/bhl.title.286>
- Doğu, S., Bağcı Y. & Dinç, M. (2009) *Rhaponticoides aytachii* sp. nov. (Asteraceae) from south Anatolia, Turkey. *Nordic Journal of Botany* 27: 479–482.
<http://dx.doi.org/10.1111/j.1756-1051.2009.00537.x>
- Eren, Ö. (2007) The genus *Rhaponticoides* Vaill. (Asteraceae) in Turkey: a new species and first key. *Plant Systematics and Evolution* 267: 13–23.
<http://dx.doi.org/10.1007/s00606-007-0576-4>
- Garcia-Jacas, N., Uysal, T., Romashchenko, K., Suárez-Santiago, V.N., Ertuğrul, K. & Susanna, A. (2006) *Centaurea* revisited: A molecular survey of the *Jacea* group. *Annals of Botany* 98: 741–753.
<http://dx.doi.org/10.1093/aob/mcl157>
- Greuter, W. (2003) The Euro+Med treatment of Cardueae (Compositae)—generic concepts and required new names. *Willdenowia* 33: 49–61.
- Greuter, W., Agababian, M. & Wagenitz, G. (2005) Vaillant on Compositae: Systematic impact and nomenclatural consequences. *Taxon* 54: 149–174.
<http://dx.doi.org/10.2307/25065315>
- Greuter, W. & Raus, T. (eds.) (2009) Med-Checklist Notulae 28. *Willdenowia* 39: 335–345.
<http://dx.doi.org/10.3372/wi.39.39211>
- Greuter, W., Wagenitz, G., Agababian, M. & Hellwig, F.H. (2001) Proposal to conserve the name *Centaurea* (Compositae) with a conserved type. *Taxon* 50: 1201–1205.

<http://dx.doi.org/10.2307/1224746>

Hellwig, F.H. (2004) Centaureinae (Asteraceae) in the Mediterranean—history of ecogeographical radiation. *Plant Systematics and Evolution* 246: 137–162.

<http://dx.doi.org/10.1007/s00606-004-0150-2>

Huber-Morath, A. (1979) Novitiae Florae Anatolicae 13. *Bauhinia* 6: 370–371.

Huber-Morath, A. (1981) Novitiae Florae Anatolicae 14. *Bauhinia* 7: 77–80.

IUCN (2012) *IUCN Red List Categories & Criteria*: version 3.1. IUCN Species Survival Commission. Gland, Switzerland & Cambridge, UK.

Lamarck, J.B. de (1785) *Encyclopédie méthodique: botanique*, vol. 1(2). Paris, pp. 345–752.

Linnaeus, C. (1753) *Species plantarum*. L.Salvius, Holmia [Stockholm], 1200 pp.

Puntillo, D. & Peruzzi, L. (2009) A new species of *Rhaponticoides* (Asteraceae) from Southern Italy. *Folia Geobotanica* 44: 191–197.
<http://dx.doi.org/10.1007/s12224-009-9037-x>

Tan, K., Bancheva, S., Vural, M. & Strid, A. (2009) *Centaurea wagenitziana* (Asteraceae: Centaureinae), a new species from the Eastern Balkans. *Phytologia Balcanica* 15 (1): 51–58.

Vaillant, S. (1754) Neue Kennzeichen dreyer Classen von Pflanzen mit zusammengesetzten Blumen, nämlich: der *Cynarocephalarum*, derer mit Artischockenhäuptern; *Corymbiferarum*, die zusammengesetzte, scheibenförmige Blumen tragen; *Cichoracearum*, wegwartartiger. *Der Königl. Akademie der Wissenschaften in Paris. Anatomische, Chymische und Botanische Abhandlungen* 5: 143–194, 333–377, 545–613, 697–750.

Wagenitz, G. (1975) *Centaurea* L. In: Davis, P.H. (ed.), *Flora of Turkey and the East Aegean Islands*, vol. 5. Edinburgh University Press, Edinburgh, pp. 465–585.

Wagenitz, G. (1986) *Centaurea* in South-West Asia: patterns of distribution and diversity. *Proceedings of the Royal Society of Edinburgh* 89B: 11–21.

<http://dx.doi.org/10.1017/S0269727000008861>