



<http://dx.doi.org/10.11646/phytotaxa.173.2.6>

***Porophyllum spathulatum* (Asteraceae: Tageteae), a new species from the southern Brazilian coast**

CAMILA REZENDO CARNEIRO¹, ANGELO ALBERTO SCHNEIDER² & MARA REJANE RITTER¹

¹ Programa de Pós-Graduação em Botânica, Universidade Federal do Rio Grande do Sul, Av. Bento Gonçalves, 9500, CEP 91501-970, Porto Alegre, Rio Grande do Sul, Brazil; email: milarezendo@yahoo.com.br

² Universidade Federal do Pampa, UNIPAMPA, Campus São Gabriel, Av. Antônio Trilha, 1847, São Gabriel, Rio Grande do Sul, Brazil.

Abstract

Porophyllum spathulatum is described and illustrated. The new species is distinguished by its decumbent habit, spatulate leaves and large heads bearing pale yellow florets. According to the collector's information given on the labels, the plant was successfully re-collected and is known to occur only in Balneário Hermenegildo, Santa Vitória do Palmar, Rio Grande do Sul, southernmost Brazil. Additionally, conservation status and ecological information about the new species are provided.

Resumo

Porophyllum spathulatum é descrito e ilustrado. A nova espécie distingue-se pelo hábito decumbente, folhas espatuladas e capítulos grandes com flores amarelo-claras. Baseado em dados de coleta constantes em etiquetas de exsicatas, a planta foi recoletada e sabe-se que ocorre apenas no Balneário Hermenegildo, Santa Vitória do Palmar, Rio Grande do Sul, extremo sul do Brasil. São fornecidos o estado de conservação e informações ecológicas sobre a nova espécie.

Key words: Compositae, dune vegetation, endemism, Heliantheae Alliance

Introduction

Porophyllum Guettard (1754: 377) belongs to tribe Tageteae Cassini (1819: 162), which is part of the “Heliantheae Alliance” (Panero 2007), a monophyletic group of 13 related tribes of Asteraceae. The genus has its center of distribution and probably center of origin in Mexico or Central America and comprises about 28 species distributed from southwestern USA to northern Argentina (Johnson 1969), in tropical and subtropical areas, in dry grasslands, vegetation dominated by subshrubs or shrubs, on rocky or sandy soils. *Porophyllum* is characterized by the presence of discoid heads, involucle uniserial of free (or almost free) phyllaries numbering 5–9, pappus of bristles and presence of translucent oil glands in leaves and phyllaries, which confers to these plants a strong and characteristic scent. Johnson (1969), in his monograph of *Porophyllum*, divided the genus in two sections: *P.* sect. *Hunteria* Candolle (1836: 649), restricted to North America and occurring from southwestern USA to arid and semi-arid regions of northern and central Mexico, and *P.* sect. *Porophyllum*, widespread from North to South America in dry or mesic areas.

During the taxonomic revision of the genus *Porophyllum* in southern Brazil, we concluded that we were in face of a new species represented by two herbarium sheets that did not match any previously known species. Further field work led to the discovery of a small population, in the same region where the available specimens were collected before. Morphological studies of the material available allowed the description of the new species *Porophyllum spathulatum*, and additional information about the new species is provided.

Taxonomic treatment

***Porophyllum spathulatum* C.R.Carneiro & A.A.Schneid., sp. nov. (Figs. 1 and 2)**

Inter species generis Porophylli habitu decumbenti, foliis spathulatis, capitulis ultra 50-floris, corolla albo-lutea, pedunculis apice dilatatis ab aliis recedens.

Acknowledgements

The first author thanks the PPG-Botânica/UFRGS for financial aid. The authors also would like to thank Gabriel Emiliano Ferreira for the map design, Márcia Vignoli da Silva for the species illustration, Ana Carolina Oliveira da Costa for helping with photos, Martin Grings for collection of *Porophyllum brevifolium*, Edson Luís de Carvalho Soares for plate preparation, Rafael Espindola Pereira Soares and Rafaela Carneiro Soares for field trips collaboration.

References

- Andrews, H.C. (1802) *Botanist's Repository, for New and Rare Plants* 3. Author, London, pl. 145–288.
<http://dx.doi.org/10.5962/bhl.title.51972>
- Arduino, P. (1764) *Animadversionum Botanicarum Specimen Alterum*. Typographia Sansoniana, Venezia, 42 pp.
- Baker, J.G. (1884) Compositae IV. Helenioideae. In: Martius, C.F.P. von (ed.) *Flora Brasiliensis* 6(3). Lipsiae, Leipzig, pp. 268–290.
- Boldrini, I.I., Ferreira, P.M.A., Andrade, B.O., Schneider, A.A., Setubal, R.B., Trevisan, R. & Freitas, E.M. (2010) *Bioma Pampa: diversidade florística e fisionômica*. Pallotti, Porto Alegre, 64 pp.
- Cabrera, A.L. (1974) Compuestas. Tribu VI. Helenieae. In: Burkart, A. (ed.) *Flora Ilustrada de Entre Ríos* 6. I.N.T.A., Buenos Aires, pp. 399–415.
- Cassini, A.H.G. de (1819) Sixième mémoire sur la famille des Synanthérées, contenant les caractères des tribus. *Journal de Physique, de Chimie, d'Histoire Naturelle et des Arts* 88: 150–163.
- Candolle, A.P. de (1836) *Prodromus Systematis Naturalis Regni Vegetabilis* 5. Treuttel & Würtz, Paris, 706 pp.
<http://dx.doi.org/10.5962/bhl.title.286>
- Gardner, G. (1848) Contributions towards a Flora of Brazil. *London Journal of Botany* 7: 395–425.
- Guettard, J.E. (1754) Huitième mémoire sur les glandes des plants. *Histoire de l'Academie Royale des Sciences. Avec les Mémoires de Mathematique & de Physique* 1750: 345–384.
- Hooker, W.J. (1836) Contributions towards a flora of South America and the islands of the Pacific. *Companion to the Botanical Magazine* 2: 41–52.
- IUCN (2013) *The IUCN Red List of Threatened Species*, version 2013.2. IUCN Red List Unit, Cambridge U.K. Available from: <http://www.iucnredlist.org> (accessed: 26 January 2014).
<http://dx.doi.org/10.5962/bhl.title.44776>
- Johnson, R.R. (1969) Monograph of the Plant Genus *Porophyllum* (Compositae-Helenieae). *The University of Kansas Science Bulletin* 48(7): 225–267.
- Lamarck, J.B. (1789) *Encyclopédie Méthodique, Botanique* 3(1). Panckoucke, Paris, 759 pp.
<http://dx.doi.org/10.5962/bhl.title.49178>
- Lessing, C.F. (1832) *Synopsis Generum Compositarum*. Duncker & Humblot, Berlin, 473 pp.
<http://dx.doi.org/10.5962/bhl.title.51470>
- Malme, G.O.A. (1899) Die Compositen der Ersten Regnell'schen Expedition. *Kongliga Svenska Vetenskapsakademien Handlingar (ser. 2)* 32(5): 1–90.
- Nakajima, J. (2013) *Porophyllum* in *Lista de Espécies da Flora do Brasil*. Jardim Botânico do Rio de Janeiro. Available from: <http://floradobrasil.jbrj.gov.br/jabot/floradobrasil/FB16259>. (accessed: 10 December 2013).
- Palisot de Beauvois, A.M.F.J. (1812) *Essai d'une Nouvelle Agrostographie*. Chez l'auteur, Paris, 182 pp.
<http://dx.doi.org/10.5962/bhl.title.474>
- Panero, J.L. (2007) Tribe Tageteae. In: Kadereit, J.W. & Jeffrey, C. (eds.) *The Families and Genera of Vascular Plants* 8. Springer, Berlin, pp. 420–431.
- Petenatti, E.M. & Espinar, L.A. (1997) Asteraceae, parte 6. Tribu VI. Helenieae. In: Hunziker, A.T. (ed.) *Flora Fanerogámica Argentina* 45. Programa Proflora (CONICET), Córdoba, pp. 1–34.
- Pfeiffer, H.H. (1937) *Androtrichum trigynum* (Spreng.) H. Pfeiff. Ein oft verkanntes, halophiles Sauergras Südamerikas. *Repertorium Specierum Novarum Regni Vegetabilis* 42: 10–14.
<http://dx.doi.org/10.1002/fedr.19370420105>
- Sprengel, C.P.J. (1824) *Systema Vegetabilium* (Ed. 16) 1. Sumtibus Librariae Dieterichiana, Gottingae, 992 pp.
<http://dx.doi.org/10.5962/bhl.title.822>
- Sprengel, C.P.J. (1826) *Systema Vegetabilium* (Ed. 16) 3. Sumtibus Librariae Dieterichiana, Gottingae, 936 pp.
<http://dx.doi.org/10.5962/bhl.title.822>
- Willdenow, C.L. von (1806) *Species Plantarum* (Ed. 4) 4(2). G.C. Nauk, Berlin, pp. 633–1157.