



Taxonomic revision of Chenopodiaceae in Nepal

ALEXANDER P. SUKHORUKOV^{1*} & MARIA KUSHUNINA²

¹ Department of Higher Plants, Biological Faculty, Lomonosov Moscow State University, 119234, Moscow, Russia; suchor@mail.ru

² Department of Plant Physiology, Biological Faculty, Lomonosov Moscow State University, 119234, Moscow, Russia;
mkushunina@gmail.com

* author for correspondence

Abstract

A taxonomic revision of the family Chenopodiaceae in Nepal with new diagnostic keys, descriptions and detailed distribution patterns is here presented for the first time. 24 species from 13 genera are reported. The apophytes in Himalaya are found in both ruderal and segetal plant communities in contrast to Chenopodiaceae in temperate parts of Eurasia which prefer ruderal places only. The Himalaya and West Tibet are considered to be the regions with the greatest taxonomic diversity of native *Dysphania* in Eurasia. *Chenopodium perttii* is described as new for science, and its differences from other related taxa are described, including its unique minute-papillate (almost smooth when dry) pericarp. *C. bengalense* seems to be an older name for the taxa known as *C. giganteum* or *C. album* subsp. *amaranticolor*. The lectotypes of *Chenopodium giganteum*, *C. album* subsp. *yunnanense*, *C. aristatum* f. *muticum* (here synonymized with *Dysphania kitiae*) and *Microgynoecium tibeticum* are designated. Both lectotype and epitype for *Atriplex bengalensis* (basionym of *C. bengalense*), *C. giganteum* and the re-instated species *C. pallidum* are designated.

Key words: *Chenopodiaceae*, *Dysphania*, distribution, ecology, Himalaya, Nepal, new species, typification, taxonomic revision.

Introduction

The family Chenopodiaceae Vent. comprises about 100 genera and approximately 1600 species distributed worldwide (Sukhorukov 2014) and it is a sister group to Amaranthaceae according to molecular investigations (Kadereit *et al.* 2003). Most of Chenopodiaceae are widespread in the semiarid and arid regions of the World. The Himalaya, especially the northern territories including NW India and Pakistan, is rich in Chenopodiaceae (Rechinger 1997, Klimeš & Dickoré 2005), but the number of the representatives decreases towards the south-east territories. In Bhutan, only eight species in the family were reported (Long 1984).

The revision of Chenopodiaceae in Nepal has not been carried out before now. In the first comprehensive account of the Eastern Himalaya (Bhutan, parts of India and Nepal) based on expeditions lasting several years, only two representatives [*Chenopodium album* Linnaeus (1753: 219) and *Dysphania ambrosioides* (Linnaeus 1753: 219) Mosyakin & Clemants (2002: 382) (sub *C. ambrosioides*)] were recorded (Hara 1966). The treatment of the entire family (14 species from 7 genera) was provided for Mustang Province in the mountainous Central Nepal (Yonekura 2008), as well as revised contributions relating to the complicated Himalayan and Tibetan species of the genera *Axyris* Linnaeus (1753: 979) (Sukhorukov 2011) and *Dysphania* Brown (1810: 411) (Sukhorukov 2012a, 2012b, 2014; Uotila 2013). The aim of the present study is a revision of the family in Nepal using recent and improved taxonomy and detailed distribution patterns.

Material and Methods

The field investigations were carried out by the first author during 2005–2013 in many provinces of Nepal in all major parts of the country (West, Central and Eastern Nepal). The herbarium material is preserved in the herbaria B, BM,

(Press *et al.* 2000). This remarkable taxon with its triangular leaf shape is distributed in North India, northern Pakistan and Afghanistan (Uotila 1997), but it does not apparently occur in north-eastern Indian states close to the Nepali border. The most predictable taxa are *Dysphania tibetica* (Uotila 2013) and *Corispermum nanum* Sukhor. & Zhang (Sukhorukov *et al.* 2014: 84) found in West Tibet near Nepali territory. The Terai region of Nepal may include some *Chenopodium* species recently described (Pandeya & Pandeya 2003) and accepted in Paul's (2012) treatment of Indian Chenopodiaceae, but the composition of this genus in tropical part of India has been very poorly studied so far and requires additional investigation.

Acknowledgements

We thank D. Iamonico (editor of Caryophyllales section of “Phytotaxa”), P. Uotila (Botanical Museum, University of Helsinki), G. Harper (Royal Botanic Garden Edinburgh), I. Belyaeva (Royal Botanic Gardens, Kew), J. Wajer (Natural History Museum, London) and A. I. Solopov (Moscow Lomonosov State University) for discussion and suggestions regarding some parts of the article. The investigation was carried out with the support of grants from the Russian Fund for Basic Research (taxonomic part: projects 14-04-00136-a; 14-04-01852-a) and Russian Scientific Foundation (seed research: 14-50-00029).

References

- Aellen, P. (1929a) Beitrag zur Systematik der *Chenopodium*-Arten Amerikas, vorwiegend auf Grund der Sammlung des United States National Museum in Washington I, II. *Feddes Repertorium* 26: 31–64, 119–160.
<http://dx.doi.org/10.1002/fedr.19290260108>
- Aellen, P. (1929b) *Chenopodium amaranticolor* Coste & Reynier, *Ch. purpurascens* “Jacquin”, *Ch. giganteum* Don, *Ch. quinoa* Willd., *Ch. moquinianum* Aellen und *Ch. reynieri* Ludwig & Aellen. *Berichte der Schweizerischen Botanischen Gesellschaft* 38: 1–19.
- Aellen, P. & Iljin, M.M. (1936) *Chenopodium*. In: Shishkin, B.K. (Ed.) *Flora of USSR*, vol. 6. Izdatelstvo Akademii Nauk, Moscow–Leningrad, pp. 41–73. [in Russian]
- Aellen, P. (1960–61) Chenopodiaceae Vent. In: Hegi, G. (Ed.) *Illustrierte Flora von Mitteleuropa*. ed. 2. vol. 3(2). Parey. Berlin & Hamburg, pp. 533–762.
- Akhani, H., Edwards, G. & Roalson E.H. (2007) Diversification of the Old World Salsoleae s.l. (Chenopodiaceae): Molecular phylogenetic analysis of nuclear and chloroplast data sets and a revised classification. *International Journal of Plant Sciences* 168: 931–956.
<http://dx.doi.org/10.1086/518263>
- Allioni, C. (1766) Stirpium aliquot descriptions cum duorum novorum generum constitutione. *Mélanges de philosophie et de mathématique de la société Royale de Turin* 3: 176–184.
- Ascherson, P.F.A. (1864) *Flora der Provinz Brandenburg, der Altmark und des Herzogthums Magdeburg*. vol. 1. Verlag von Hirschwald, Berlin, 143 pp.
- Bamber, C.J. (1916) *Plants of the Punjab. A descriptive key to the flora of the Punjab, North-West frontier province and Kashmir*. Government Printing, Lahore, 652 pp.
- Bentham, G. & Hooker, J.D. (1880) *Genera Plantarum*, vol. 3(1). Reeve & Co, London, 459 pp.
- Bhargava, A., Shukla, S. & Ohri, D. (2007) Genome size variation in some cultivated and wild species of *Chenopodium* (Chenopodiaceae). *Caryologia* 60 (3): 245–250.
<http://dx.doi.org/10.1080/00087114.2007.10797943>
- Brenan, J.P.M. (1954) Chenopodiaceae. In: Turrill, W.B. & Milne-Redhead, E. (Eds.) *Flora of Tropical East Africa. Chenopodiaceae*. vol. 7. Crown Agents, London, pp. 1–26.
- Britton, N. & Brown, A. (1913) *An Illustrated flora of the Northern United States, Canada and the British possessions*. 2nd ed. vol. 2. New-York, 735 pp.
- Brown, R. (1810) *Prodromus florae Novae Hollandiae et insulae Van-Diemen, exhibens characteres plantarum quas annis 1802-1805 per oras utriusque insulae collegit et descripsit*. Taylor & socii, Londini, 590 pp.
<http://dx.doi.org/10.5962/bhl.title.52309>
- Bryant, G.J. (2013) *Emergence of British power in India, 1600-1784: A grand strategic interpretation*. The Boydell Press, New-York, 350 pp.
- Bunge, A.A. (1876) *Chenopodium bryoniifolium* [description]. *Delectus seminum, quae Hortus botanicus imperialis Petropolitanus pro*

- mutua commutatione offert.* Hortus Botanicus, St.-Petersburg, without pagination.
- Chu (Zhu), G.-L. (1987) *Archiatriplex*, a new Chenopodiaceous genus from China. *Journal of the Arnold Arboretum* 68: 461–469.
- Colla, A. (1836) *Herbarium Pedemontanum*. Regio typographeo, Turin, 602 pp.
- Dobeš, C. & Hahn, B. (1997) Chromosomenzahlen zur Gefäßpflanzen-Flora Österreichs. *Linzer Biologische Beiträge* 29(1): 5–43.
- Don, D., Hamilton, F. & Wallich, N. (1825) *Prodromus florae Nepalensis, sive enumeration vegetabilium, quae in itinere per Nepalam proprie dictam et regions conterminas, ann. 1802–1803*. Wilks, London, 256 pp.
- Dumortier, B.-C. (1827) *Florula Belgica*. J. Casterman, Tournai, 172 pp.
- Freitag, H. & Rilke, S. (1997) *Salsola* [s.l.]. In: Rechinger, K.H. (Ed.) *Flora Iranica*. vol. 172. Akademische Druck- und Verlagsanstalt, Graz, pp. 154–255.
- Fu, J.Q. (2000) *Flora Loess-Plateaus Sinicae*. vol. 1. Science Press, Beijing, 648 pp.
- Fuentes-Bazan, S., Mansion, G. & Borsch, T. (2012a) Towards a species level tree of the globally diverse genus *Chenopodium*. *Molecular Phylogenetics & Evolution* 62: 359–374.
<http://dx.doi.org/10.1016/j.ympev.2011.10.006>
- Fuentes-Bazan, S., Uotila, P. & Borsch, T. (2012b) A novel phylogeny-based generic classification for *Chenopodium* sensu lato, and a tribal rearrangement of Chenopodioideae (Chenopodiaceae). *Willdenowia* 42: 5–24.
<http://dx.doi.org/10.3372/wi42.42101>
- Grozeva, N. & Stoeva, M. (2006) Mediterranean chromosome number reports 16 (1466–1472). *Flora Mediterranea* 16: 400–408.
- Grubov, V.I. (1961) Due species novae e familia Chenopodiaceae. *Botanicheskiye materialy gerbariya Botanicheskogo Instituta im. V.L. Komarova* 21: 125–129.
- Gueldenstaedt, A.I. (1772) *Krascheninnikovia*, novum plantarum genus. *Novi Commentarii Academiae Scientiarum Imperialis Petropolitanae* 16: 548–560.
- Gutermann, W. (2011) Notulae nomenclaturales 41–45 (Neue Namen bei *Cruciata* und *Kali* sowie einige kleinere Korrekturen). *Phytom (Horn)* 51: 95–102.
- Handel-Mazzetti, H. (1929) *Symbolae sinicae. Botanische Ergebnisse der Expedition der Akademie der Wissenschaften in Wien nach Südwest-China 1914–1918*, vol. 7. Springer, Vienna, 730 pp.
- Hara, H. (1966) *The Flora of Eastern Himalaya: Results of the Botanical Expedition to Eastern Himalaya organized by the University of Tokyo 1960 and 1963*, vol. 1. The University of Tokyo Press, Tokyo, 744 pp.
- Heklau, H. & Röser, M. (2008) Delineation, taxonomy and phylogenetic relationships of the genus *Krascheninnikovia* (Amaranthaceae subtribe Axyridinae). *Taxon* 57: 563–576.
- Herder, F. (1889) Plantae Raddeanae Apetalae. Chenopodeae et Amaranthaceae. *Acta Horti Petropolitani* 10: 583–627.
- Hooker, J.D. (1890) *The Flora of British India, Chenopodiaceae to Orchidaceae*. vol. 5. Reeve & Co., London, 910 pp.
- Iamonico, D. (2010) Confirmation of the occurrence of *Chenopodium strictum* subsp. *strictum* (Amaranthaceae s. l.) in Italy. *Phytom (Horn, Austria)* 49(2): 235–240.
- Iamonico, D. (2011) *Dysphania anthelmintica* (Amaranthaceae), new to the non-native flora of Italy, and taxonomic considerations on the related species.
<http://dx.doi.org/10.2478/v10028-011-0002-x>
- Iamonico, D. & Jarvis, C.E. (2012) Lectotypification of two Linnaean names in *Chenopodium* L. (Amaranthaceae). *Taxon* 61: 864–865.
- IUCN (2010) *The IUCN red list of threatened species*, version 2010.4. IUCN Red List Unit, Cambridge, UK. Available from: <http://www.iucnredlist.org/> (accessed 20 January 2014).
- Jacquin, N.J. (1776) *Hortus botanicus Vindobonensis*, vol. 3. Typ. Joseph Michael, Vienna, 52 pp.
- Jafri, S.M.H. & Rateeb, F.B. (1978) Chenopodiaceae. In: Jafri, S.M.H. & El-Gadi, A. (Eds.) *Flora of Libya*. vol. 58. Al Faateh University, Tripoli, 109 pp.
- Jarvis, C. (2007) *Order out of Chaos*. Linnaean plant names and their types. The Linnaean Society of London & Natural History Museum, London, 1017 pp.
- Kadereit, G., Borsch, T., Weising, K. & Freitag, H. (2003) Phylogeny of Amaranthaceae and Chenopodiaceae and the evolution of C4 photosynthesis. *International Journal of Plant Sciences* 164: 959–986.
<http://dx.doi.org/10.1086/378649>
- Kadereit, G. & Freitag, H. (2011) Molecular phylogeny of Camphorosmeae (Camphorosmoideae, Chenopodiaceae): Implications for biogeography, evolution of C4-photosynthesis and taxonomy. *Taxon* 60: 51–78.
- Kadereit, G., Zacharias, E., Mavrodiev, E. & Sukhorukov, A.P. (2010) Molecular phylogeny of Atripliceae (Chenopodioideae, Chenopodiaceae): Implications for systematics, biogeography, flower and fruit evolution, and the origin of C₄ photosynthesis. *American Journal of Botany* 97: 1664–1687.
<http://dx.doi.org/10.3732/ajb.1000169>
- Karelin, G.S. & Kirilov, I.P. (1841) *Enumeratio plantarum anno 1840 in regionibus altaicis et confinibus collectarum*. *Bulletin de la CHENOPODIACEAE*

- Société Imperiale des Naturalistes de Moscou 4: 703–870.
- Klimeš, L. & Dickoré, B. (2005) A contribution to the vascular plant flora of Lower Ladakh (Jammu & Kashmir, India). *Willdenowia* 35: 125–153.
<http://dx.doi.org/10.3372/wi.35.35110>
- Lamarck, J.B. (1783) *Encyclopédie méthodique Botanique*. vol. 1. Paris, Liege, 752 pp.
- Léveillé, H. (1913) Decades plantarum novarum. CIV–CV. *Repertorium specierum novarum regni vegetabilis (Feddes Repertorium)* 11: 548–552.
<http://dx.doi.org/10.1002/fedr.19130113404>
- Linnaeus, C. (1753) *Species Plantarum*. vol. 1–2. Salvius, Stockholm, 1200 pp.
- Linnaeus, C. (1782) *Supplementum plantarum systematis vegetabilium editionis decimae tertiae, generum plantarum editiones sextae, et specierum plantarum editionis secundae*. Imp. Orphanotrophei, Braunschweig, 468 pp.
<http://dx.doi.org/10.5962/bhl.title.555>
- Lomonosova, M.N. (2006) Chromosome numbers of some Chenopodiaceae representatives of the flora of Russia. *Botanicheskiy Zhurnal* 91: 1757–1759.
- Lomonosova, M.N. & Krasnikov, A.A. (1993) Chromosome numbers in some members of the Chenopodiaceae. *Botanicheskiy Zhurnal* 78: 158–159.
- Lomonosova, M.N., Krasnikova, S.A., Krasnikov, A.A., Ebel, A.L. & Rudaja N.A. (2003) Chromosome numbers of the family members Chenopodiaceae representatives from Mongolia and Kazakhstan. *Botanicheskiy Zhurnal* 88: 113–115.
- Lomonosova, M.N., Sukhorukov, A.P. & Sinelnikova, N.V. (2012) Chromosome numbers of Chenopodiaceae/Amaranthaceae species from Spain, Egypt, Syria and Nepal. *Botanicheskiy Zhurnal* 97: 99–102.
- Long, D.G. (1984) Chenopodiaceae. In: Grierson, A.J.C. & Long, D.G. (Eds.) *Flora of Bhutan including a record of plants from Sikkim*. vol. 1(2). Royal Bot. Garden, Edinburgh, pp. 216–219.
- McNeill, J., Bassett, I.J., Crompton, C.W. & Taschereau, P.M. (1983) Taxonomic and nomenclatural notes on *Atriplex* L. (Chenopodiaceae). *Taxon* 32: 549–556.
<http://dx.doi.org/10.2307/1221723>
- McNeill, J., Barrie, F.R., Buck, W.R., Demoulin, V., Greuter, D.L., Hawksworth, D.L., Herendeen, P.S., Knapp, S., Marhold, K., Prado, J., Proud'Homme van Reine, W.F., Smith, J.F. & Wiersema, J.H. (Eds.) (2012) *International Code of Nomenclature for algae, fungi and plants (Melbourne Code): Adopted by the Eighteenth International Botanical Congress, Melbourne, Australia, July 2011*. Regnum Vegetabile 154. Ruggell.
- Meyer, C.A. (1829) *Chenopodieae*. In: Ledebour, C.F. (Ed.) *Flora Altaica*. vol. 1. Reimer, Berlin, pp. 370–418.
<http://dx.doi.org/10.5962/bhl.title.6618>
- Miller, P. (1754) The Gardeners disctionary. 4th edition, corrected and enlarged. London, without pagination.
- Moench, C. (1794) *Methodus plantas horti botanici et agri Marburgensis, a staminum situ describendi*. Officina nova libraria academiae, Marburgi, 368 pp.
<http://dx.doi.org/10.5962/bhl.title.304>
- Moquin-Tandon, A. (1834) Descriptiones de plusieur nouveaux genres de Chenopodees. [ser. 2] *Annales des Sciences Naturelles (Botanique)* 1: 289–294.
- Moquin-Tandon, A. (1840) *Chenopodearum monographica enumeratio*. Loss, Paris, 183 pp.
<http://dx.doi.org/10.5962/bhl.title.15484>
- Moquin-Tandon, A. (1849) Salsolaceae [Chenopodiaceae]. In: de Candolle, A. (Ed.) *Prodromus systematis naturalis regni vegetabilis*. vol. 13(2). Typ. Masson, Paris, pp. 43–219.
- Mosyakin, S.L. & Clemants, S.E. (2002) New nomenclatural combinations in *Dysphania* R. Br. (Chenopodiaceae): taxa occurring in North America. *Ukrainskiy Botanicheskiy Zhurnal* 59: 380–385.
- Mosyakin, S.L. & Clemants, S.E. (2008) Further transfers of glandular-pubescent species from *Chenopodium* subgen. *Ambrosia* to *Dysphania* (Chenopodiaceae). *Journal of the Botanical Research Institute of Texas* 2: 425–431.
- Murr, J. (1902) *Chenopodium*-Beiträge. *Magyar Botanikai Lapok* 1: 337–344; 359–369.
- Murr, J. (1923) Neue Übersicht über die Farn- und Blütenpflanzen von Vorarlberg und Liechtenstein mit Hervorhebung der geobotanischen Verhältnisse und mit Berücksichtigung der Nachbargebiete. vol. 1. Kommissionsverlag Unterberger, Bregenz, 507 pp.
- Pandeya, S.C. & Pandeya, A. (2003) Further contribution to biosystematics of *Chenopodium*, reporting three new species from North Indian plains. *Journal of the Bombay Natural History Museum* 100: 87–92.
- Paul, T.K. (2012) A synopsis of the family Chenopodiaceae in India. *Pleione* 6(2): 273–297.
- Pax, F. (1889) Caryophyllaceae (incl. Dysphanieae). In: Engler, A. & Prantl, K. (Eds.) *Die Natürlichen Pflanzensammlungen*. vol. 3(1b). Verlag Engelmann, Leipzig, pp. 61–94.
- Poiret, J.L.M. (1810) *Amaranthus persicarioides* [description]. In: Lamarck, J.B. & Poiret, J.L.M. (Eds.) *Encyclopedie Methodique*

- Botanique*. supplement vol. 1. Agasse, Paris, 311 pp.
<http://dx.doi.org/10.5962/bhl.title.49178>
- Press, J.R., Shrestha, K.K. & Sutton, D.A. (2000) *Annotated checklist of the flowering plants of Nepal*. The Natural History Museum, London, 430 pp.
- Probatova, N.S., Rudyka, E.G., Kozhevnikov, A.E. & Kozhevnikova, Z.V. (2004) Chromosome numbers of some representatives of the flora of the Primorsky Territory. *Botanicheskiy Zhurnal* 89(7): 1209–1217.
- Rafinesque, C.S. (1837) *Flora Telluriana*. vol. 3. Probasco, Philadelphia.
<http://dx.doi.org/10.5962/bhl.title.7751>
- Rahiminejad, M.R. (2006) Chenopodiaceae. In: Marhold, K. (ed.) *IAPT/IOPB chromosome data. Taxon* 55: 444–445.
<http://dx.doi.org/10.2307/25065590>
- Rechinger, K.-H. (Ed.) (1997) *Flora des Iranischen Hochlandes und der umrahmenden Gebirge*. vol. 172. Akad. Druck- u. Verlagsanstalt, Graz, 371 pp.
- Reynier, A. (1905) *Chenopodium album* L. subspecies *amaranticolor* Coste & Reynier [a new subspecies]. [ser. 2] *Bulletin de l'herbier Boissier* 5: 979–980.
- Reynier, A. (1907) Les *Chenopodium amaranticolor* Coste et Reynier et *Chenopodium pedunculare* Bertoloni, dans les Bouches-du-Rhône. *Bulletin de la Société botanique de France* 54: 178–183.
<http://dx.doi.org/10.1080/00378941.1907.10831249>
- Roth, A.W. (1821) *Nova plantarum species praesertim Indiae Orientalis*. Vogler, Halberstadt, 412 pp.
<http://dx.doi.org/10.5962/bhl.title.10723>
- Ruas, C.F., Ruas, P.M., Stutz, H.C. & Fairbanks, D.J. (2001) Cytogenetic studies in the genus *Atriplex* (Chenopodiaceae). *Caryologia* 54: 129–145.
<http://dx.doi.org/10.1080/00087114.2001.10589220>
- Runemark, H. (1996) Mediterranean chromosome number reports 6 (590–678). *Flora Mediterranea* 6: 223–243.
- Schrader, H.A. (1809) Über Pallas's Halophyta, mit besonderer Rücksicht auf die Gattungen *Salsola* und *Suaeda*. *Neues Journal für die Botanik* 3: 58–92.
- Schrader, H.A. (1822) *Acroglochin* [description]. In: Roemer, J.J. & Schultes, J.A. (Eds.) *Mantissa in volumen primum systematis vegetabilium Caroli a Linne*. vol. 1. Cottae, Stuttgart, 69 pp.
<http://dx.doi.org/10.5962/bhl.title.552>
- Schultes, J.A. (1820) *Systema vegetabilium: Secundum classes, ordines, genera, species. Cum characteribus differentiis et synonymis. Editio nova, speciebus inde ab editione XV. Detectis aucta et locupletata*. vol. 6. Cottae, Stuttgart, 857 pp.
<http://dx.doi.org/10.5962/bhl.title.825>
- Schweinfurth, G. (1867) *Beitrag zur Flora Aethiopiens*. Georg Reimer, Berlin, 311 pp.
- Scott, A.J. (1978a) A review of the classification of *Chenopodium* L. and related genera. *Botanische Jahrbücher für Systematik* 100: 205–220.
- Scott, A.J. (1978b) A revision of the Camphorosmoideae (Chenopodiaceae). *Feddes Repertorium* 89: 101–119.
<http://dx.doi.org/10.1002/fedr.19780890202>
- Smith, J.E. (1800) *Flora Britannica*. vol. 1. Typ. Davis, London, 458 pp.
<http://dx.doi.org/10.5962/bhl.title.6641>
- Steudel, E. (1821) *Nomenclator botanicus: enumerans ordine alphabetico nomina atque synonyma, tum generica tum specifica, et a Linnaeo et recentioribus de re botanica scriptoribus plantis phanerogamis imposita*. Cottae, Stuttgartiae & Tubingae, 900 pp.
<http://dx.doi.org/10.5962/bhl.title.544>
- Sukhorukov, A.P. (2005) Karpologische Untersuchungen der *Axyris*-Arten (Chenopodiaceae) im Zusammenhang mit ihrer Diagnostik und Taxonomie. *Feddes Repertorium* 116: 168–176.
<http://dx.doi.org/10.1002/fedr.200511070>
- Sukhorukov, A.P. (2011) *Axyris* (Chenopodiaceae s.str. or Amaranthaceae s.l.) in the Himalayas and Tibet. *Willdenowia* 41(1): 75–82.
<http://dx.doi.org/10.3372/wi.41.41108>
- Sukhorukov, A.P. (2012a) Taxonomic notes on *Dysphania* and *Atriplex* (Chenopodiaceae). *Willdenowia* 42: 169–180.
<http://dx.doi.org/10.3372/wi.42.42202>
- Sukhorukov, A.P. (2012b) The reproductive organ morphology of *Dysphania* R.Br. (Chenopodiaceae) in the context of geographic isolation. In: Mavrodiev, E.V. (ed.) *Analytical approaches in floristic studies and methods of biogeography*: Proc. 1st Int. Conference. Astrakhan State Univ., Astrakhan, pp. 64–67 (in Russian with English summary).
- Sukhorukov, A.P. (2014) *The carpology of the Chenopodiaceae with reference to the phylogeny, systematics and diagnostics of its representatives*. Grif & Co., Tula, 405 pp. (in Russian with English summary).
- Sukhorukov, A.P., Uotila, P., Zhang, M., Zhang, H.-X., Speranskaya, A.S. & Krinitsyna, A.A. (2013) New combinations in Asiatic *Oxybasis*

- (Amaranthaceae s.l.): evidence from morphological, carpological and molecular data. *Phytotaxa* 144: 1–12.
<http://dx.doi.org/10.11646/phytotaxa.144.1.1>
- Sukhorukov, A.P. & Zhang, M. (2013) Fruit and seed anatomy of *Chenopodium* and related genera (Chenopodioideae, Chenopodiaceae/ Amaranthaceae): Implications for evolution and phylogeny. *PLoS One* 8: 1–18 (e61906).
<http://dx.doi.org/10.1371/journal.pone.0061906>
- Sukhorukov, A.P., Zhang, M. & Nilova, M.V. (2014) The carpology and taxonomy of some Chinese *Corispermum*. *Phytotaxa* 172: 81–93.
<http://dx.doi.org/10.11646/phytotaxa.172.2.3>
- Thiers, B. (2008) *Index herbariorum: A global directory of public herbaria and associated staff*. New York Botanical Garden, Bronx, NY.
Available from: <http://sweetgum.nybg.org/ih/> (accessed 20 June 2013).
- Tzvelev, N.N. (1960) About some new and little known species from Pamir. *Botanicheskiye Materialy Gerbariya Botanicheskogo Instituta V.L. Komarova Akademii Nauk SSSR* 20: 413–439.
- Ulrich, E. (1934) Chenopodiaceae. In: Engler, A. & Prantl, K. (Eds.) *Die Natürlichen Pflanzenfamilien*. ed. 2, 16c. Duncker & Humblot, Leipzig, pp. 379–584.
- Uotila, P. (1993) Taxonomic and nomenclatural notes on *Chenopodium* in the Flora Iranica area. *Annales Botanici Fennici* 30: 189–194.
- Uotila, P. (1997) *Chenopodium* (s.l.). In: Rechinger, K.H. (Ed.) *Flora des Iranischen Hochlandes und der umrahmenden Gebirge*, vol. 172. Akademische Druck- und Verlagsanstalt, Graz, pp. 24–59.
- Uotila, P. (2001a) *Chenopodium*. In: Jonsell, B. (Ed.) *Flora Nordica*. vol. 2. Bergius Foundation, Stockholm, pp. 4–31.
- Uotila, P. (2001b) *Chenopodium*. In: Ali, S.I. & Qaiser, M. (Ed.) *Flora of Pakistan*. vol. 204. Univ. of Karachi & Missouri Bot. Garden, Karachi & Missouri, pp. 13–52.
- Uotila, P. (2013) *Dysphania* sect. *Botryodes* (Amaranthaceae s.lat.) in Asia. *Willdenowia* 43: 65–80.
<http://dx.doi.org/10.3372/wi.43.43107>
- Werker, E. (1997) *Seed anatomy*. – Berlin & Stuttgart: Borntraeger Verlag.
- Yonekura, K. (2008) Chenopodiaceae. In: Ohba, H., Iokawa, Y. & Sharma, L.R. (eds.) *Flora of Mustang, Nepal*. Kodansha Scientific Ltd., Tokyo, pp. 53–61.
- Zhu (Chu), G.-L., Mosyakin, S.L. & Clemants, S.E. (2003) Chenopodiaceae. In: Wu, Z. & Raven, P.H. (Eds.) *Flora of China*. vol 5. *Ulmaceae–Basellaceae*. St. Louis, pp. 351–414.
- Zhukova, P.G. & Petrovsky, V.V. (1980) Chromosome numbers and taxonomy of some species of the Anyui Mts. *Botanicheskiy Zhurnal* 65: 651–659.

Appendix. Specimens of *Dysphania kitiae* re-identified in the herbaria visited:

Gansu prov.: Lanzhou county, 27 September 1956, *Expedition team s.n.* (PE-00510830); Yongchang county, near Wu-wei, 1958, *anonym s.n.* (PE); Yongdeng, Liancheng, 2999 m, 14 September 1958, *Ganqing Team Zhong Buqiu 10147* (WUK). **Ningxia** prov.: Yanchi, Yangliubu, 1310 m, 18 September 1984, *Yang Jinxiang 5222* (WUK). **Inner Mongolia** prov.: Rehe Chifeng vicinity (Zhifang Village), 570 m, 21 September 1952, *Liou Tchen-ngo 5220* (WUK); Fengzhen, Xuegangshan Mountains, 1230 m, 28 September 1984, *Huangtu Team (Jin) 4197* (WUK). **Hebei** prov.: Lingshankou, Choluhsien, 7 September 1930, *Hsia 2448* (PE-00510747); Eastern Tombs, 16 September 1930, *Liou 1603* (PE-00510746); Neiqiu County, Xiaolingdi village, Bulaoqing, 16 September 1950, *Liu Ying 13801* (PE); [near Beijing] Baihuashan, Huang'antuo, 19 September 1950, *Wang Wen-cai 744* (PE); Huailai county, Zhaojiapeng, Tianqiaoshan, 19 September 1959, *anonym 172* (PE). **Shansi** prov.: Luyashan, Ning-wu, 5000 ft, 20 August 1929, *Hsia 1616* (PE-00510771); Chieh-hsiu [Jiexiu] distr., Sung-Lin-miao, in cultis, ca. 800 m, 3 October 1924, *Smith 7899* (PE-00510602); Sihsien, Shikowtze, 1500 m, 3 September 1935, *Wang 3449* (PE-00510781); Shangchuang, Sihsien, 1200 m, 12 September 1935, *Wang 3600* (PE-00510772). **Shaanxi** [Shensi] prov.: along Wei river, 10 September 1932, *Hao 4526* (PE-00510872); Ningshen-hsien, 900 m, 19 July 1933, *Kung 3105* (PE-00510797); Tsashui Hsien [Zhàshuǐ Xiān], 8 October 1933, *Wang 1991* (PE-00510790); Yulin, 10 August 1982, *Yang Jinxiang 3988* (PE); Wuqi County, 1200 m, 13 September 1984, *Yang Jinxiang 5115* (WUK); Zhouzhi, Taibeishan, 1500 m, 6 July 1999, *Zhu & Xu 1546* (PE-00235085). **Sichuan** prov.: Xiao, October 1914, *Handel-Mazzetti s.n.* (W); NW part, Rumichango, 2000 m, 29 October 1922, *Smith 4872* (PE-00510698); Mu-li, Ko-pa-ting, 2500 m, 17 July 1937, *Yu 7290* (PE-00510910); Sungpan [Songpan], Heiichiao, 1600 m, 23 October 1937, *Wang 7939* (PE-00510903). **Yunnan** prov.: Wei-se Hsien [Wēixīn Xiān], 2600 m, 19 September 1934, *Tsai 57965* (PE-00510952); Tehching, Tsawalungla, 2800 m, 18 September 1937, *Yu 10331* (PE-00510961).