



Notes on Early Land Plants Today. 54. A transfer in Lepidoziaceae (Marchantiophyta)

ENDYMION D. COOPER¹, LARS SÖDERSTRÖM², ANDERS HAGBORG³ & MATT VON KONRAT³

¹ CMNS-Cell Biology and Molecular Genetics, 2107 Bioscience Research Building, University of Maryland, College Park, MD 20742-4451, USA; endymion.dante.cooper@gmail.com.

² Department of Biology, Norwegian University of Science and Technology, N-7491 Trondheim, Norway; lars.soderstrom@ntnu.no

³ Department of Science and Education, The Field Museum, 1400 South Lake Shore Drive, Chicago, IL 60605–2496, USA; hagborg@pobox.com, mvonkonrat@fieldmuseum.org

When Cooper *et al.* (2013) reorganized species among genera in Lepidoziaceae Limpricht (1876: 310), one taxon, *Lepidozia leratii* Stephani (1922: 333), was mistakenly combined under *Neolepidozia* Fulford & Taylor (1959: 81). The transfer was not based on any morphological or molecular evidence placing it in *Neolepidozia*. On the contrary, molecular phylogenetic studies (Cooper *et al.*, 2011, 2012; Heslewood & Brown, 2007) all place *Lepidozia leratii* in close proximity of *Tricholepidozia pulcherrima* (Stephani 1909: 600) E.D.Cooper in Cooper *et al.* (2013: 60) the type of *Tricholepidozia* (Schuster 1963: 256) E.D.Cooper in Cooper *et al.* (2013: 58). This error is corrected here.

Tricholepidozia leratii (Steph.) E.D.Cooper, **comb nov.** Basionym:—*Lepidozia leratii* Steph., *Sp. Hepat.* (Stephani) 6: 333 (Stephani 1922). Type:—New Caledonia, summit of Mt Mou, July 1909, *Lerat* (PC-0102363, lectotype by Hürlimann 1985 [<http://coldb.mnhn.fr/catalognumber/mnhn/pc/pc0102363>]).

≡ *Neolepidozia leratii* (Steph.) E.D.Cooper, *Phytotaxa* 97: 56, 2013 (Cooper *et al.* 2013).

Acknowledgements

The Early Land Plants Today project (ELPT) has been generously supported in part by the Global Biological Information Facility (GBIF) Seed Money Award No.2007-41, activities facilitated in part by funding from the Biodiversity Synthesis Center of the Encyclopedia of Life (BioSynC), partial funding from the National Science Foundation (Award No's 0749762, 1115002), the Warwick Foundation, and the Negaunee Foundation.

References

- Cooper, E.D., Henwood, M.J. & Brown, E.A. (2012) A molecular phylogeny of the *Lepidozia* generic complex supports re-circumscription of the Lepidozioideae. *Molecular Phylogenetics and Evolution* 65: 10–22
<http://dx.doi.org/10.1016/j.ympev.2012.05.008>
- Cooper, E.D., Shaw, A.J., Shaw, B., Henwood, M.J., Heslewood, M.M. & Brown, E.A. (2011) A multi-locus molecular phylogeny of the Lepidoziaceae: Laying the foundations for a stable classification. *Molecular Phylogenetics and Evolution* 59: 489–509.
<http://dx.doi.org/10.1016/j.ympev.2011.02.006>
- Cooper, E.D., Söderström, L., Hagborg, A. & von Konrat, M.J. (2013) Notes on Early Land Plants Today. 38. New combinations and synonyms in Lepidoziaceae (Marchantiophyta). *Phytotaxa* 97: 52–62.
<http://dx.doi.org/10.11646/phytotaxa.97.2.5>
- Fulford, M.H. & Taylor, J. (1959) The segregate genera of the *Lepidozia* complex (Hepaticae). Part 1. {*Sprucella*} Steph. and *Neolepidozia* gen. nov. *Brittonia* 11: 77–85.
<http://dx.doi.org/10.2307/2805175>
- Heslewood, M.M. & Brown, E.A. (2007) A molecular phylogeny of the liverwort family Lepidoziaceae Limpr. in Australasia. *Plant Systematics and Evolution* 265: 193–219.
<http://dx.doi.org/10.1007/s00606-006-0512-z>
- Hürlimann, H. (1985) Hepaticae aus dem Gebiete des südlichen Pazifik. VIII. *Bauhinia* 8: 101–118.

- Limpricht, K.G. (1876) Lebermoose. *In*: Cohn, F. (Ed), *Kryptogamen-Flora von Schlesien, Vol. 1, heft 2*. J. U. Kern's Verlag: Breslau, pp. 225–471.
- Schuster, R.M. (1963) Studies on antipodal Hepaticae. I. Annotated keys to the genera of antipodal Hepaticae with special reference to New Zealand and Tasmania. *Journal of the Hattori Botanical Laboratory* 26: 185–309.
- Stephani, F. (1909) *Species Hepaticarum* 3. George & Cie, Genève & Bale, pp. 517–693.
- Stephani, F. (1922) *Species Hepaticarum* 6. George & Cie, Genève & Bale, pp. 241–368.