



http://dx.doi.org/10.11646/phytotaxa.170.2.8

A new combination in *Mcneillia* (Caryophyllaceae) for the Italian flora

FABRIZIO BARTOLUCCI^{1*}, FABIO CONTI¹, DUILIO IAMONICO² & E. DEL GUACCHIO³

¹Centro Ricerche Floristiche dell'Appennino (Università di Camerino-Parco Nazionale del Gran Sasso e Monti della Laga), 67021 Barisciano (AQ), Italy; e-mail fabrizio.bartolucci@gmail.com

²Laboratory of Phytogeography and Applied Geobotany, Section Environment and Landscape, Department PDTA, University of Rome Sapienza, 00196 Rome, Italy

³Bagnolifutura S.p.A., 80124 Naples, Italy

*Corresponding author

On the basis of DNA sequences of nrITS and plastid *matK* from 160 ingroup and 105 outgroup samples, Dillenberger & Kadereit (2014) reconstructed the phylogeny and the evolution of the relevant morphological characters of the genus *Minuartia* Linnaeus (1753: 89), to show that this genus is highly polyphyletic. A revised circumscription of *Minuartia* was proposed with the recognition of *Cherleria* Linnaeus (1753: 425), *Eremogone* Fenzl (1833: 13), *Facchinia* Reichenbach (1841: 63), *Mononeuria* Reichenbach (1841: 118), *Rhodalsine* Gay (1845: 25), *Sabulina* Reichenbach (1832: 24) and *Triplateia* Bartling (1830: 305), and recognition of new genera *Mcneillia* Dillenberger & Kadereit (2014: 83), *Minuartiella* Dillenberger & Kadereit (2014: 84) and *Pseudocherleria* Dillenberger & Kadereit (2014: 84).

Among the Italian endemic Caryophyllaceae (Iamonico 2013, Peruzzi *et al.* 2014), six entities are included in *Minuartia* as traditionally recognized: *M. glomerata* (Bieberstein 1808: 350) Degen in Fritsch (1910: 319) subsp. *trichocalycina* (Tenore & Gussone in Tenore 1835: 16) F. Conti (1997: 75), *M. graminifolia* (Arduino 1764: 25) Jávorka (1914: 22) subsp. *graminifolia*, *M. graminifolia* (Ard.) Jáv. subsp. *rosanoi* (Tenore 1811: 26) Mattfeld (1921: 57), *M. grignensis* (Reichenbach 1842: 88) Mattfeld (1922: 141), *M. laricifolia* Schinz & Thellung (1907: 403) subsp. *ophiolitica* Pignatti (1973: 207), and *M. moraldoi* F. Conti (2001: 193). *Minuartia glomerata* subsp. *trichocalycina* is the only taxon that remains in *Minuartia* [see Dillenberger & Kadereit (2014: Appendix S3)]. Concerning *M. graminifolia* s.s., and *M. moraldoi*, new combinations were proposed [respectively as *Mcneillia graminifolia* (Ard.) Dillenberger & Kadereit (2014: 83) and *Mcneillia moraldoi* (F. Conti) Dillenberger & Kadereit (2014: 84)], while the names *M. grignensis* and *M. laricifolia* subsp. *ophiolitica* are treated in an ongoing work by M. Dillenberger [*M. grignensis* as a member of *Facchinia*] and by A. Moore [*M. laricifolia* subsp. *ophiolitica* to be placed in *Cherleria*] (M. Dillenberger, and A. J. Moore pers. comm.). Concerning *M. graminifolia* subsp. *rosanoi* all of its diagnostic features match those of *Mcneillia* (see Dillenberger & Kadereit l.c.), specifically: bracts similar to stem leaves, sepals acuminate, 5-many-veined with marginal veins conspicuous beyond the middle of the sepal length, and petals obovate, narrowly cuneate at the base that are longer than the sepals. Considering that *Minuartia graminifolia* is the type *Mcneillia* (Dillenberger & Kadereit 2014), and in agreement with the taxonomic treatment of this species suggested by Conti (2003) in which the subsp. *rosanoi* is re-evaluated, we propose the following new combination:

***Mcneillia graminifolia* (Ard.) Dillenb. & Kadereit subsp. *rosanoi* (Ten.) F. Conti, Bartolucci, Iamonico & Del Guacchio, comb. nov.**

Basionym: *Arenaria rosanoi* Tenore (1811: XXVI).

Type (lectotype designated by Conti & Santangelo 2001: 195, Fig. 2):—ITALY, Basilicata, *sine die, sine collector s.n.* (NAP!, exsiccatum on the bottom-right).

≡ *Minuartia graminifolia* (Ard.) Jáv. subsp. *rosanoi* (Ten.) Mattf.

Note:—The spelling of the epithet “*rosanoi*” follows the Art. 60.7 of the ICN (McNeill *et al.* 2012).

Acknowledgements

Thanks are due to M. Dillenberger (Johannes Gutenberg-Universität, Mainz, Germany) and to A.J. Moore (Brown University, Rhode Islands, USA) for constructive comments about the name here studied and their ongoing researches.

References

- Arduino, P. (1764) *Animadversionum botanicarum specimen alterum*. Ex *Typographia Sansoniana*, Venetiis, 42 pp.
- Bartling, F.G. (1830) *Ordines naturales plantarum eorumque characteres et affinitates adjecta generum enumeratione*. Sumtibus Dieterichianis, Gottingae, 498 pp.
- Beiberstein, F.A.M. von (1808) *Flora Taurico-Caucasica*, vol. 1. Typis Academicis, Charkouiae, 478 pp.
<http://dx.doi.org/10.5962/bhl.title.10825>
- Conti, F. (1997) *Minuartia glomerata* subsp. *trichocalycina* comb. & stat. nov. (Caryophyllaceae), a central Apennine endemic. *Willdenowia* 27: 73–79.
- Conti, F. (2001) *Minuartia moraldoi* (Caryophyllaceae), a new species from Cilento (Campania, S. Italy). *Plant Biosystems* 135: 191–194.
- Conti, F. (2003) *Minuartia graminifolia* (Ard.) Jav. (Caryophyllaceae), a south-east European species. *Botanical Journal of the Linnean Society* 143: 419–432.
- Conti, F. & Santangelo, A. (2001) Lectotypification of the names *Arenaria graminifolia* Ard., *Arenaria rosani* Ten. and *Arenaria alpicola* Ten. (Caryophyllaceae). *Taxon* 50: 193–199.
<http://dx.doi.org/10.2307/1224523>
- Fenzl, E. (1833) *Versuch einer Darstellung der Geographischen Verbreitungs- und Vertheilungs-Verhaltnisse der Naturlichen Familie der Alsineen*. J.B. Wallishausser, Wien, 70 pp.
- Gay, J. (1845) Holostei, caryophyllearum alsinearum generis, monographia. *Annales des Sciences Naturelles, Botanique*, sér. 3, 4: 23–44.
- Fritsch, K. (1910) Neue Beiträge zur Flora der Balkan-halbinsel, insbesondere Serbiens, Bosniens und der Herzegowina. *Mitteilungen des Naturwissenschaftlichen Vereines für Steiermark* 46: 294–328.
- Dillenberger, M.S. & Kadereit, J.W. (2014) Maximum polyphyly: Multiple origins and delimitation with plesiomorphic characters require a new circumscription of *Minuartia* (Caryophyllaceae). *Taxon* 63: 64–88.
<http://dx.doi.org/10.12705/631.5>
- Iamónico, D. (2013) Italian endemic Caryophyllaceae: A review. *Lagascalia* 33: 275–298.
- Jávorka, S. (1914) 142. *Minuartia graminifolia* (Ard.) Jav. ssp. *M. hungarica* Jav. *Jegyzék Magyarország Növényeinek Gyűjteményéhez: Schedae ad Floram Hungaricam Exsiccatam Centuria II*: 22–23.
- Linnaeus, C. (1753) *Species Plantarum*, vol. 1. Salvius, Stockholm, 560 pp.
- Mattfeld, J. (1921) Beitrag zur Kenntnis der systematischen Gliederung und geographischen Verbreitung der Gattung *Minuartia*. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie*. 57 Beiblätter 126: 13–63.
- Mattfeld, J. (1922) Geographisch-genetische Untersuchungen über die Gattung *Minuartia* (L.) Hiern. *Repertorium Specierum Novarum Regni Vegetabilis Beihefte* 15: 1–228.
- McNeill, J., Barrie, F.R., Buck, W.R., Demoulin, V., Greuter, D.L., Hawksworth, D.L., Herendeen, P.S., Knapp, S., Marhold, K., Prado, J., Proud'Homme van Reine, W.F., Smith, J.F. & Wiersema, J.H. (eds.) (2012) International Code of Nomenclature for algae, fungi and plants (Melbourne Code): Adopted by the Eighteenth International Botanical Congress, Melbourne, Australia, July 2011. *Regnum Vegetabile* 154: 1–274.
- Peruzzi, L., Conti, F. & Bartolucci, F. (2014). An inventory of vascular plants endemic to Italy. *Phytotaxa* 168: 1–75.
- Pignatti, S. (1973) Note critiche sulla Flora d'Italia. 1. Appunti miscellanei. *Giornale Botanico Italiano* 107: 207–221.
<http://dx.doi.org/10.1080/11263507309426323>
- Reichenbach, H.G.L. (1832) *Flora germanica excursoria*, vol. 1(1). Carolum Cnoblock, Leipzig, 434 pp.
- Reichenbach, H.G.L. (1841) *Icones florae Germanicae et Helveticae: simul Pedemontanae, Tyrolensis, Istriacae, Dalmaticae, Austriacae, Hungaricae, Transylvanicae, Moravicae, Boressicae, Holsatica, Belgicae, Hollandicae, etgo Mediale Europae*, vol. 5. Ambrosi Abel, Lipsiae, 38 pp. + 103 plates.
<http://dx.doi.org/10.5962/bhl.title.6353>
- Schinz, H. & Thellung A. (1907) Begründung vorzunehmender Namensänderungen an der zweiten Auflage der «Flora der Schweiz» von Schinz und Keller. *Bulletin de l'Herbier Boissier*, sér. 2, 7: 387–406.
- Tenore, M. (1811) *Flora Napolitana*, vol. 1(2). Stamperia Reale, Napoli, 40 pp. (XII–XXXII + 21240), 5 pl.
- Tenore, M. (1835) *Ad florae Neapolitanae Syllogem, appendix quarta; loca natalia quaedam, observationes, plantasque nuperrime detectas vel antehac omissas complectens*. Tipografia del Fibreno, Napoli, 52 pp.