

<http://dx.doi.org/10.11646/phytotaxa.186.3.6>

***Radula multiflora* var. *reflexilobula* is a synonym of *R. decurrens* (Radulaceae, Marchantiophyta)**

CHATCHABA PROMMA & SAHUT CHANTANAORRAPINT*

Department of Biology, Faculty of Science, Prince of Songkla University, Hat Yai, Songkhla 90112, Thailand.

* Corresponding author; e-mail: sahut.c@psu.ac.th

Radula Dumortier (1822: 112) is one of the largest genera of the liverworts with about 200 currently accepted species (Frey & Stech 2009), but some of them are poorly known and their status is still unclear. *Radula javanica* Gottsche (in Gottsche *et al.* 1844: 257) is one of the oldest names of *Radula* which is widespread in the tropical and subtropical Asia, the Pacific Islands and Australasia (Yamada 1979, Renner 2014). So (2006) reduced a highly variable species *R. multiflora* Gottsche ex Schiffn. (in Engler 1889: 20) to *R. javanica* Gottsche. However, Renner (2014) suggested that *R. multiflora* might be different from *R. javanica* but similar to *R. reflexa* Nees & Mont. (in Montagne 1843: 255).

Yamada and Grolle described *R. multiflora* var. *reflexilobula* (Yamada 1982: 326) based on a collection made by B.O. van Zanten from New Guinea. This variety differs from the type variety by the large trigones of basal cells of leaf-lobes and the lobule strongly reflexed at the apex. Yamada (1982) also noted that *R. multiflora* var. *reflexilobula* is similar to *R. reflexa* and *R. decurrens* Mitt. (in Seemann 1873: 410) in appearance. However, *R. reflexa* differs from this variety by the medium-size trigones of the leaf-lobe and the arched interior margin of the leaf-lobule that fully covering the ventral stem surface, and *R. decurrens* is distinguished by its densely imbricate and strongly recurved leaf-lobules.

During our examination of the type specimens of *R. decurrens* and *R. multiflora* var. *reflexilobula* in the Herbarium Haussknecht (JE), however, we found that the shape of leaf-lobes of the fertile branch (Fig. 1) of *R. multiflora* var. *reflexilobula* is indistinguishable from *R. decurrens*. Thus, we here propose *R. multiflora* var. *reflexilobula* as new synonym of *R. decurrens*. Unfortunately, we have not seen the type specimens of *R. reflexa*, a putatively closely related species.

Formal treatment

The format of this note follows Söderström *et al.* (2012).

Radula decurrens Mitt. in Seemann, Fl. Vitiensis.: 410. 1873.

Type:—Samoa Island. s.d., T. Powell s.n. (holotype: NY; isotype: JE!, NY, YU)

= *Radula multiflora* Gottsche ex Schiffn. var. *reflexilobula* Grolle & Yamada in Yamada, J. Hattori Bot. Lab. 51: 326.

1982. Type:—New Guinea. Eastern Highlands. 1968, B.O. van Zanten 68-2953 (holotype NICH; isotypes L, JE!) *syn. nov.*

For further synonyms, see So (2006).

Acknowledgments

We would like to thank the curators and staffs of JE for the possibility to study type specimens. This work was supported by the Faculty of Science Research Fund, and the Graduate School, Prince of Songkla University. The first author would like to express her sincere appreciation to the Development and Promotion of Science and Technology Talent Project (DPST) for funding.




FIGURE 1. *Radula decurrens* Mitt.: A–B. Portions of sterile plants, ventral views; C. Portion of sterile plants, dorsal view; D. Portion of plant with andoecium; E. Portion of plant with gynoecium; F. Apical cells of leaf lobe; G. Median cells of leaf lobe; H. Basal cells of the leaf lobe. All from B.O. van Zanten 68-2953 (isotype of *Radula multiflora* Gottsche ex Schiffn. var. *reflexilobula* Grolle & Yamada, JE).

References

- Dumortier, B.C. (1822) Commentationes Botanicae. *Tournay, Ch. Casterman-Dien* 10: 112.
<http://dx.doi.org/10.5962/bhl.title.10534>
- Frey, W. & Stech, M. (2009) Marchantiophyta, Bryophyta, Anthocerotophyta. In: Frey, W. (Ed.) *Syllabus of Plant Families. A. Engler's Syllabus der Pflanzenfamilien, 13th ed., Part 3. Bryophytes and seedless Vascular Plants*, Berlin, Gebrüder Borntraeger, pp. 9–263.
- Gottsche, C.M. (1844) *Radula*. In: Gottsche, C.M., Lindenberg, J.B.G. & Nees ab Esenbeck, C.G. (Eds.) *Synopsis Hepaticarum*, Hamburg, Germany, 257 pp.
<http://dx.doi.org/10.5962/bhl.title.15221>
- Mitten, W. (1873) Ordo CIX. Jungermanniae. In: Seemann, B. (Ed.) *Flora Vitiensis*. London, pp. 410.
- Montagne, D.M. (1843) Quatrième centurie de Plantes cellulaires exotiques nouvelles. *Annales des Sciences Naturelles; Botanique*, ser. 2 19: 238–266.

- Renner, M.A.M. (2014) *Radula* subg. *Radula* in Australasia and the Pacific (Jungermanniopsida). *Telopea* 17: 107–167.
<http://dx.doi.org/10.7751/telopea20147553>
- Schiffner, V.F. (1889) Platiphyllae. In: Engler H.G.A. (Ed.) *Die Forschungsreise S.M.S. "Gazelle"*, IV. Teil. Botanik, 4, Berlin, Germany, 20 pp.
- So, M.L. (2006) *Radula* (Radulaceae, Marchantiophyta) in the South Pacific. *Journal of the Hattori Botanical Laboratory* 99: 207–232.
- Söderström, L., Hagborg, A. & von Konrat, M. (2012) Notes on Early Land Plants Today. *Phytotaxa* 65: 41–42.
- Yamada, K. (1979) A revision of the Asian taxa of *Radula* (Hepaticae). *Journal of the Hattori Botanical Laboratory* 45: 201–322.
- Yamada, K. (1982) Some new species of *Radula* (Hepaticae). *Journal of the Hattori Botanical Laboratory* 51: 323–328.