



Molecular phylogenetic analysis reveals two new species of *Discosia* from Italy

WEN-JING LI^{1,2,3,4}, JIAN-KUI LIU^{3,4}, D. JAYARAMA BHAT^{3,5}, ERIO CAMPORESI⁶, DONG-QING DAI^{3,4}, PETER E MORTIMER^{1,2}, JIAN-CHU XU^{1,2}, KEVIN D. HYDE^{1,2,3,4,7} & PUTARAK CHOMNUNTI^{3,4}

¹World Agroforestry Centre, East and Central Asia, 132 Lanhei Road, Kunming 650201, China

²Key Laboratory of Economic Plants and Biotechnology, Kunming Institute of Botany, Chinese Academy of Sciences, Lanhei Road No 132, Panlong District, Kunming, Yunnan Province, 650201, PR China

³Institute of Excellence in Fungal Research, Mae Fah Luang University, Chiang Rai 57100, Thailand

⁴School of Science, Mae Fah Luang University, Chiang Rai 57100, Thailand

⁵Formerly, Department of Botany, Goa University, Goa 403206, India

⁶A.M.B. Gruppo Micologico Forlivese “Antonio Cicognani”, Via Roma 18, Forlì, Italy

⁷Botany and Microbiology Department, College of Science, King Saud University, Riyadh, KSA 11442, Saudi Arabia

Corresponding author: Putarak Chomnunti, email address: putarak.cho@mfu.ac.th

Abstract

Two fresh collections of *Discosia* were made from dead leaves of *Fagus sylvatica* in Italy. As these collections could not be cultured, the fruiting bodies were directly used for sequencing using a Forensic DNA Extraction Kit. Based on analyses of the concatenated internal transcribed spacer regions of the nrDNA operon (ITS) and large subunit rDNA (LSU) gene sequences, as well as morphological characters, the fresh collections are introduced as two new species, namely *D. italica* and *D. fagi*. Phylogenetically, these two species are distinct from all other *Discosia* species. Morphologically, *D. italica* is somewhat similar with *D. fagi*, but can be distinguished using dimension of conidiomata and conidiogenous cells. Descriptions and illustrations of the new taxa are provided herein.

Key words: Asexual morphs, Amphisphaeriaceae, Phylogeny, Taxonomy

Introduction

Discosia (Amphisphaeriaceae) is a widely distributed genus of coelomycetes (Sutton 1980, Vanev 1992b, Nag Raj 1993, Wołczańska *et al.* 2004). Most species are endophytes or saprobes on various vascular plants in the tropical and temperate regions (Subramanian & Reddy 1974, Sutton 1980, Vanev 1992b, 1996, Nag Raj 1993, Okane *et al.* 1998). Some species in this genus are pathogens, such as *Adisciso yakushimense* Kaz. Tanaka, Okane & Hosoya on *Symplocos prunifolia* (Tanaka *et al.* 2011), as well as *A. kaki* Kaz. Tanaka, J. Yamam. & Toy. Sato on *Diospyros kaki* (Yamamoto *et al.* 2012), forming leaf spots on terrestrial plants from June to September.

The genus *Discosia* is characterized by uni- to multilocular conidiomata with multi-layered walls, occurring singly or in clusters. Conidiogenous cells are monoblastic, phialidic to annellidic. Conidia are almost hyaline to pale brown, and with polar or subpolar appendages inserted in the median part of the end cells (Sutton 1980, Nag Raj 1993).

Traditionally, morphological characters and host-association have been used to classify *Discosia* to species (Subramanian & Reddy 1974, Sutton 1980, Vanev 1991, 1992a, 1996, Nag Raj 1993). However, most species in this genus have overlapping characters, such as the location of the conidial septa and appendages, varying proportional lengths of conidial cells, and overall conidium size (Sutton 1977, 1980, Nag Raj 1993, Jeewon *et al.* 2002, Barber *et al.* 2011, Tanaka *et al.* 2011). The classification, validity, and delimitation of this genus have been problematic (Subramanian & Reddy 1974, Vanev 1991, 1992b, Sutton 1980, Nag Raj 1993, Jeewon *et al.* 2002). Subramanian & Reddy (1974) divided the taxa under *Discosia* into four sections based on the conidial morphology (size, septation and pigmentation). Vanev (1991) expanded this concept and grouped the species of *Discosia* into six sections, viz., *D. sect. Discosia*, *D. sect. Laurina* Vanev, *D. sect. Clypeata* Vanev, *D. sect. Libertia* Vanev, *D. sect. Strobilina* Vanev and *D. sect. Poikilo-*

D. aff. pleurochaeta (GenBank, AB593713, Identities = 806/808 (99%), no gaps) and *Discosia* sp. 1' (GenBank, AB593710, Identities = 806/808 (99%), no gaps). The closest hits using the ITS sequence had similarity to *Discosia* sp. (GenBank, AF405303, Identities = 549/560 (98%), Gaps = 2/560 (0%)), *D. aff. artocreas* (GenBank, AB594772, Identities = 525/536 (98%), Gaps = 3/536 (0%)) and *D. pseudoartocreas* (GenBank, KF777161, Identities = 557/569 (98%), Gaps = 4/569 (0%)). However, phylogenetic analyses derived from combined ITS and LSU sequences show that *D. fagi* is clearly distinct from all other *Discosia* species (Figure 1). Morphologically, *D. fagi* resembles *D. italica*, but can be distinguished using dimension of conidiomata, being slightly larger than *D. italica* (100–250 µm diam., and 20–45 µm high). Furthermore, the conidiogenous cells of *D. fagi* also longer than *D. italica* 2–5 (\bar{x} = 3) µm. Based on the substantive evidence provided above, *D. fagi* is introduced as novel species of *Discosia*.

In this paper, it has not been possible to apply LSU alone in resolving the species within *Discosia*, especially for species where very few morphological differences exist. Therefore, analyses of the combined sequences of ITS and LSU genes are used to delimit the species within *Discosia*. They also provide a significant evidence to justify that *D. italica* and *D. fagi* are distinct from each other. Nevertheless, the identification of interspecific relationship is problematic within the genus *Discosia*. *D. pini*, *D. aff. brasiliensis* and the type species *D. artocreas* clustered together with high bootstrap support, and this is in congruence with the observations of Tanaka *et al.* (2011). Thus, the interspecific relationship within *Discosia* should be studied further using multi-gene data. Moreover, on the bases of molecular data (ITS, LSU), the dimension of conidiomata and morphology of the conidiogenous cells and conidia, interspecies relationship can be distinguished. It would be premature to discuss the phylogenetic significance of various species, given that sequence data is available for only a small number of *Discosia* species. Future studies should focus on larger sampling of *Discosia* species and evaluation by molecular sequence data.

Acknowledgments

We would like to thank Humidtropics, a CGIAR Research Program that aims to develop new opportunities for improved livelihoods in a sustainable environment, for partially funding this work. KD Hyde thanks The Chinese Academy of Sciences, project number 2013T2S0030, for the award of Visiting Professorship for Senior International Scientists at Kunming Institute of Botany.

References

- Altschul, S.F., Gish, W., Miller, W., Myers, E.W. & Lipman, D.J. (1990) Basic local alignment search tool. *Journal of Molecular Biology* 215: 403–410.
[http://dx.doi.org/10.1016/S0022-2836\(05\)80360-2](http://dx.doi.org/10.1016/S0022-2836(05)80360-2)
- Barber, P.A., Crous, P.W., Groenewald, J.Z., Pascoe, I.G. & Keane, P. (2011) Reassessing *Vermisporium* (Amphisphaeriaceae), a genus of foliar pathogens of eucalypts. *Persoonia* 27: 90–118.
<http://dx.doi.org/10.3767/003158511X617381>
- Cai, L., Hyde, K.D., Taylor, P.W.J., Weir, B., Waller, J., Abang, M.M., Zhang, J.Z., Yang, Y.L., Phoulivong, S., Liu, Z.Y., Prihastuti, H., Shivas, R.G., Mckenzie, E.H.C. & Johnston, P.R. (2009) A polyphasic approach for studying *Colletotrichum*. *Fungal Diversity* 39: 183–204.
- Crous, P.W., Wingfield, M.J., Guarro, J., Cheewangkoon, R., van der Bank, M., Swart, W.J., Stchigel, A.M., Cano-Lira, J.F., Roux, J., Madrid, H., Damm, U., Wood, A.R., Shuttleworth, L.A., Hodges, C.S., Munster, M., de Jesús Yáñez-Morales, M., Zúñiga-Estrada, L., Cruywagen, E.M., de Hoog, G.S., Silvera, C., Najafzadeh, J., Davison, E.M., Davison, P.J.N., Barrett, M.D., Barrett, R.L., Manamgoda, D.S., Minnis, A.M., Kleczewski, N.M., Flory, S.L., Castlebury, L.A., Clay, K., Hyde, K.D., Maússe-Sitoe, S.N.D., Chen, Shuaifei, Lechat, C., Hairaud, M., Lesage-Meessen, L., Pawłowska, J., Wilk, M., Śliwińska-Wyrzychowska, A., Mętrak, M., Wrzosek, M., Pavlic-Zupanc, D., Maleme, H.M., Slippers, B., Mac Cormack, W.P., Archuby, D.I., Grünwald, N.J., Tellería, M.T., Dueñas, M., Martín, M.P., Marinowitz, S., de Beer, Z.W., Perez, C.A., Gené, J., Marin-Felix, Y. & Groenewald, J.Z. (2013) Fungal Planet description sheets: 154–213. *Persoonia* 31: 188–296.
<http://dx.doi.org/10.3767/003158513X675925>
- Farr, D.F. & Rossman, A.Y. (2011) Fungal databases, systematic mycology and microbiology laboratory, ARS, USDA. Retrieved February 15, 2011, from /fungal databases/.
- Jeewon, R., Liew, E.C.Y. & Hyde, K.D. (2002) Phylogenetic relationships of *Pestalotiopsis* and allied genera inferred from ribosomal DNA sequences and morphological characters. *Molecular Phylogenetics and Evolution* 25: 378–92.

[http://dx.doi.org/10.1016/S1055-7903\(02\)00422-0](http://dx.doi.org/10.1016/S1055-7903(02)00422-0)

- Hall, T.A. (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98.
- Hillis, D.M. & Bull, J.J. (1993) An empirical test of bootstrapping as a method for assessing confidence in phylogenetic analysis. *Systematic Biology* 42: 182–192.
<http://dx.doi.org/10.2307/2992540>
- Li, W.J., Liu, J.K., Bhat, D.J., Camporesi, E., Chukeatirote, E., Xu, J.C. & Hyde, K.D. (2014) Introducing the novel species, *Dothiorella symphoricarposicola*, from snowberry in Italy. *Cryptogamie, Mycologie* 35(3): 257–270.
<http://dx.doi.org/10.7872/crym.v35.iss3.2014.257>
- Liu, J.K., Phookamsak, R., Doilom, M., Wikee, S., Li, Y.M., Ariyawansa, H., Boonmee, S., Chomnunti, P., Dai, D.Q., Bhat, J.D., Romero, A.I., Zhuang, W.Y., Monka, J., Jones, E.B.G., Chukeatirote, E., Ko-Ko, T.W., Zhao, Y.C. & Wang, Y. (2012) Towards a natural classification of Botryosphaerales. *Fungal Diversity* 57: 149–210.
<http://dx.doi.org/10.1007/s13225-012-0207-4>
- Maharachchikumbura, S.S.N., Hyde, K.D., Groenewald, J.Z., Xu, J.C. & Crous, P.W. (2014) *Pestalotiopsis* revisited. *Studies in Mycology* 79: 121–186.
<http://dx.doi.org/10.1016/j.simyco.2014.09.005>
- Nag Raj, T.R. (1993) *Coelomycetous anamorphs with appendage-bearing conidia*. Mycologue Publications, Waterloo, Canada, 1101 pp.
- Nylander, J.A.A. (2004) MrModeltest 2.0. Program distributed by the author. Evolutionary Biology Centre, Uppsala University.
- Okane, I., Nakagiri, A. & Ito, T. (1998) Endophytic fungi in leaves of ericaceous plants. *Canadian Journal of Botany* 76: 657–663.
<http://dx.doi.org/10.1139/b98-032>
- Page, R.D.M. (1996) TreeView: an application to display phylogenetic trees on personal computers. *Computer Applications in the Biosciences* 12: 357–358.
<http://dx.doi.org/10.1093/bioinformatics/12.4.357>
- Phookamsak, R., Liu, J.K., Chukeatirote, E., McKenzie, E.H.C. & Hyde K.D. (2013) Phylogeny and morphology of *Leptosphaerulina saccharicola* sp. nov. and *Pleosphaerulina oryzae* and relationships with *Pithomyces*. *Cryptogamie, Mycologie* 34: 303–319.
<http://dx.doi.org/10.7872/crym.v34.iss4.2013.303>
- Ronquist, F. & Huelsenbeck, J.P. (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574.
<http://dx.doi.org/10.1093/bioinformatics/btg180>
- Silvestro, D. & Michalak, I. (2011) raxmlGUI: a graphical front-end for RAxML. *Organisms Diversity and Evolution* 12: 335–337.
<http://dx.doi.org/10.1007/s13127-011-0056-0>
- Subramanian, C.V. & Reddy, K.R.C. (1974) The genus *Discosia*. I, Taxonomy. *Kavaka* 2: 57–89.
- Sutton, B.C. (1977) Coelomycetes. VI. Nomenclature of generic names proposed for coelomycetes. *Mycological Papers* 141: 1–253.
- Sutton, B.C. (1980) *The Coelomycetes-Fungi imperfecti with pycnidia, acervuli and stromata*. Commonwealth Mycological Institute, Kew, UK, 496 pp.
- Swofford, D.L. (2002) PAUP: phylogenetic analysis using parsimony. version 4.0 b10. Sinauer Associates, Sunderland MA.
- Tanaka, K., Endo, M., Hirayama, K., Okane, I., Hosoya, T. & Sato, T. (2011) Phylogeny of *Discosia* and *Seimatosporium*, and introduction of *Adisciso* and *Immersidiscosia* genera nova. *Persoonia* 26: 85–98.
<http://dx.doi.org/10.3767/003158511X576666>
- Thompson, J.D., Gibson, T.J., Plewniak, F., Jeanmougin, F. & Higgins, D.G. (1997) The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research* 25: 4876–4882.
<http://dx.doi.org/10.1093/nar/25.24.4876>
- Vanev, S.G. (1991) Species conception and sections delimitation of genus *Discosia*. *Mycotaxon* 41: 387–396.
- Vanev, S.G. (1992a) Comparative morphological studies of *Discosia artocras* and *Discosia faginea*. *Mycotaxon* XLIV: 461–470.
- Vanev, S.G. (1992b) On the typification and nomenclature of some *Discosia* species. *International Journal of Mycology and Lichenology* 5: 293–300.
- Vanev, S.G. (1996) Fungi of the genus *Discosia* (Deuteromycotina) in the Mediterranean area. *Bocconea* 5: 351–357.
- Vilgalys, R. & Hester, M. (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172: 4238–4246.
- White, T., Bruns, T., Lee, S. & Taylor, J. (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *In: PCR protocols: a guide to methods and applications*. pp. 315–322.
- Wolczańska, A.G.W., Ozłowska, M.O.K., Iątek, M.A.P. & Ulenko, W.I.M. (2004) Survey of the genus *Discosia* (anamorphic fungi) in Poland. *Polish Botanical Journal* 49: 55–61.
- Yamamoto, J., Tanaka, K., Ohtaka, N. & Sato, T. (2012) Black leaf spot of Japanese persimmon (*Diospyros kaki*), a new disease caused by *Adisciso kaki* sp. nov. *Journal of General Plant Pathology* 78: 99–105.
<http://dx.doi.org/10.1007/s10327-012-0367-9>