



<http://dx.doi.org/10.11646/phytotaxa.195.3.5>

Type studies on *Archilejeunea* (Lejeuneaceae, Marchantiophyta): five new synonyms and a new combination

XUE-QIN SHI^{1,2}, S. ROBBERT GRADSTEIN³ & RUI-LIANG ZHU^{1,4}

¹ Bryology Laboratory, School of Life Sciences, East China Normal University, 3663 Zhong Shan North Road, Shanghai 200062, China

² School of Life Sciences, Anhui Normal University, 1 Beijing East Road, Wuhu, Anhui, 241000, China

³ Muséum National d'Histoire Naturelle, Dept. Systématique et Evolution, Case Postale 39, 57 rue Cuvier, 75231 Paris cedex 05, France

⁴ Tiantong National Station of Forest Ecosystem, Shanghai Key Lab for Urban Ecological Processes and Eco-Restoration, East China Normal University, 500 Dongchuan Road, Shanghai 200241, China

Author for correspondence: Rui-Liang Zhu, rlzhu@bio.ecnu.edu.cn

Archilejeunea (Spruce 1884: 88) Stephani (1888: 113) is a pantropical genus with about 28 species in two subgenera, subg. *Archilejeunea* with five species and the remaining ones in subg. *Dibrachiella* (Spruce 1884: 90) Schiffner (1893: 130) (Gradstein & Buskes 1985). The genus has been well-studied in the Neotropics and Australia (Gradstein & Buskes 1985; Thiers & Gradstein 1989; Gradstein 1994; Gradstein & Schäfer-Verwimp 2012) but the treatments of the African and Asiatic species are incomplete or outdated (e.g., Stephani 1911; Verdoorn 1934; Vanden Berghen 1951; Udar & Awasthi 1981a, 1982; Mizutani 1993; Gradstein *et al.* 2002). Several species have not been studied since their first publication. Moreover, recent molecular-phylogenetic studies indicate that *Archilejeunea* is not monophyletic (Wilson *et al.* 2007). Members of *A.* subg. *Archilejeunea* were resolved in a clade with *Caudalejeunea* (Stephani 1890: 18) Schiffner (1893: 129) and *Schiffnerolejeunea* Verdoorn (1933: 89), while those of *A.* subg. *Dibrachiella* resolved in a distant clade with species of *Spruceanthus* Verdoorn (1934: 151) and *Ptychanthus* Nees (1838: 211). As a result, the genus urgently needs revision.

In the framework of a molecular and morphological study of *Archilejeunea*, we have examined the type specimens of several *Archilejeunea* species. Based on the results of the type study five new synonyms and one new combination are proposed here. *Archilejeunea helenae* Pócs & Gyarmati (2006: 104), *A. linguifolia* Stephani (1911: 708), and *A. mauritiana* Stephani (1895: 60) are synonymous with *A. brevilibula* Stephani (1911: 706), *A. elobulata* Stephani (1911: 707), and *A. abbreviata* (Mitten 1863: 167) Vanden Berghen (1951: 117), respectively. *Archilejeunea apiculifolia* Stephani (1924: 558) and *A. falcifolia* Stephani (1910: 196) are synonyms of *A. planiuscula* (Mitten 1861: 111) Spruce ex Pearson (1892: 4). *Archilejeunea nobilis* (Gottsch 1882: 353) Spruce ex Pearson (Pearson 1892: 4) is transferred to *Marchesinia* Gray (1821: 689) and is an older name for *Marchesinia madagassa* Stephani (1912: 144).

Formal treatment

The format of this note follows Söderström *et al.* (2012).

Archilejeunea abbreviata (Mitt.) Vanden Berghen, *Rev. Bryol. Lichénol.* 20: 117. 1951 (Vanden Berghen 1951).

≡ *Lejeunea abbreviata* Mitt., *J. Proc. Linn. Soc., Bot.* 7: 167. 1864 [1863] (Mitten 1864 [1863]). Type:—Cameroun. Bagroo River, 1861, Mann s.n. (holotype NY!).

= *Archilejeunea mauritiana* Steph., *Hedwigia* 34: 60. 1895 (Stephani 1895). Type:—Mascarenes. “Insula Maurice”, J.B. Maugeot s.n., hb. Lindenberg 6904 (holotype G!), **syn. nov.**

For further synonyms, see Vanden Berghen (1951).

Archilejeunea brevilibula Steph., *Sp. Hepat.* 4: 706. 1911 (Stephani 1911). Type:—Madagascar. 1904, Voeltzkow s.n. (holotype G!).

= *Archilejeunea helenae* Pócs & Gyarmati, *Cryptog., Bryol.* 27: 104. 2006 (Pócs & Gyarmati 2006). Type:—Madagascar. Mananara Nord Biosphere Reserve, wet lowland rainforest, epiphyllous, 250–350 m, Aug. 1998, Pócs & Szabó 9878/FM (holotype EGR!), **syn. nov.**

Archilejeunea elobulata Steph., *Sp. Hepat.* 4: 707. 1911 (Stephani 1911). Type:—Central African Republic. 1902, *Chevalier s.n.*, hb.

Corbière 59 (holotype G!).

= *Archilejeunea linguifolia* Steph., *Sp. Hepat.* 4: 708. 1911 (Stephani 1911). Type:—Cameroon. Bipindi, Zenker 3539 (isotypes BR!, E!), *syn. nov.*

Archilejeunea planiuscula (Mitt.) Steph., *Sp. Hepat.* 4: 731. 1911 (Stephani 1911).

≡ *Lejeunea planiuscula* Mitt., *J. Proc. Linn. Soc., Bot.* 5: 111. 1861 (Mitten 1861) ≡ *Ptychocoleus planiusculus* (Mitt.) Verd., *Ann. Bryol.* Suppl. 4: 126. 1934 (Verdoorn 1934). Type:—Burma. Rangoon, *McClelland s.n.* (holotype NY!).

= *Archilejeunea apiculifolia* Steph., *Sp. Hepat.* 6: 558. 1924 (Stephani 1924). Type:—India. Karnataka, 1911, *Pfleiderer 20* (holotype G!).

= *Archilejeunea falcifolia* Steph., *Akad. Wiss. Wien, Math.-Naturwiss. Kl., Denkschr.* 85: 196. 1910 (Stephani 1910). Type:—Papua New Guinea. Rechinger 4614 (holotype G!), *syn. nov.*

For further synonyms, see Thiers & Gradstein (1989) and Gradstein *et al.* (2002).

Archilejeunea planiuscula is a polymorphic species, widely distributed in Indo-pacific regions. Verdoorn (1934) reduced *A. apiculifolia* to a synonym under his *A. mariana* (= *Archilejeunea planiuscula*) [non *A. mariana* (Gottscche 1845: 337) Stephani (1911: 729) (Gottscche *et al.* 1845; Stephani 1911)], but Udar & Awasthi (1981a, 1981b) treated *A. apiculifolia* as a separate taxon on the basis of mostly apiculate and only occasionally rounded leaf apices and a leaf-lobe vs. lobule ratio of 2:1. As both characters are present in the variability of *A. planiuscula*, *A. apiculifolia* becomes a synonym of *A. planiuscula*.

Marchesinia nobilis* (Gottscche) X.Q. Shi, R.L. Zhu & Gradst., *comb. nov.

≡ *Lejeunea nobilis* Gottscche *Abh. Naturwiss. Vereins Bremen* 7: 353. 1882 (Gottscche 1882) ≡ *Archilejeunea nobilis* (Gottscche) Spruce ex Pearson, *Forh. Vidensk.-Selsk. Kristiania* (8): 4. 1892 (Pearson 1892). Type:—Madagascar. *Rutenberg* 1570 (holotype G!).

= *Marchesinia madagassa* Steph., *Sp. Hepat.* 5: 144. 1912 (Stephani 1912). Type:—Madagascar. 1898, *Leloutres.n.* (holotype G, n.v.), *syn. nov.*

= *Marchesinia moelleriana* Pearson, *Ark. Bot.* 19(5): 10. 1924 (Pearson 1924). Type:—Kenya: Mt. Elgon, Kaimosi. 1920, *Lindblom* (holotype S, n.v.) (syn. fide Jones 1970).

This species is well-recognized by rounded-entire leaf apices, plane underleaf margins and entire female bracts and bracteoles (Wigginton 2004; Gradstein 2012).

Acknowledgements

This research was sponsored by the National Natural Science Foundation of China (nos. 31370238, 31170190), the Special Program for the National Basic Work of the Ministry of Science and Technology, China (no. 2012FY110600), and 211 Project for The East China Normal University. We also thank the curators of BR, E, EGR, G, MO, HSNU and NY for making type specimens of *Archilejeunea* and further collections available for our study through loans.

References

- Gottscche, C.M. (1882) Reliquiae Rutenbergianae. Lebermoose. *Abhandlungen herausgegeben vom Naturwissenschaftlichen Vereins zu Bremen* 7: 338–365.
- Gottscche, C.M., Lindenberg, J.B.G. & Nees von Esenbeck, C.G. (1845) *Synopsis Hepaticarum, fasc. 3.* Meissnerianis, Hamburg, pp. 305–464.
<http://dx.doi.org/10.5962/bhl.title.15221>
- Gradstein, S.R. (1994) Lejeuneaceae: Ptychantheae, Brachiolejeuneae. *Flora Neotropica, Monograph* 62: 1–216.
- Gradstein, S.R. (2012) An overview of the genus *Marchesinia* (Lejeuneaceae), with special attention to the species described from Asia and Oceania. *Polish Botanical Journal* 57: 69–79.
- Gradstein, S.R. & Buskes, G.M.C. (1985) A revision of Neotropical *Archilejeunea* (Spruce) Schiffn. *Beihefte zur Nova Hedwigia* 80: 89–112.

- Gradstein, S.R. & Schäfer-Verwimp, A. (2012) A new species of *Archilejeunea* (Spruce) Schiffn. (Lejeuneaceae) from Ecuador. *Cryptogamie, Bryologie* 32: 107–112.
<http://dx.doi.org/10.7872/cryb.v33.iss2.2012.107>
- Gradstein, S.R., He, X.L., Piippo, S. & Mizutani, M. (2002) Bryophyte flora of the Huon Peninsula, Papua New Guinea. LXVIII. Lejeuneaceae subfamily Ptychanthoideae (Hepaticae). *Acta Botanica Fennica* 174: 1–88.
- Gray, S.F. (1821) *A natural arrangement of British plants, according to their relation to each other*. Baldwin, Cradock, and Joy, London, 824 pp.
<http://dx.doi.org/10.5962/bhl.title.43804>
- Jones, E.W. (1970) African Hepaticas XXII. *Dicranolejeunea* and *Marchesinia*. *Transactions of the British Bryological Society* 6 (1): 72–81.
<http://dx.doi.org/10.1179/006813870804146545>
- Mitten, W. (1861) Hepaticae Indiae Orientalis: An enumeration of the Hepaticae of the East Indies. *Journal of the Proceedings of the Linnean Society, Botany* 5: 109–128.
<http://dx.doi.org/10.1111/j.1095-8312.1861.tb01343.x>
- Mitten, W. (1864) On the Musci and Hepaticae from the Cameroun mountains and from the River Niger. *Journal of the Proceedings of the Linnean Society, Botany* 7: 147–169.
<http://dx.doi.org/10.1111/j.1095-8312.1863.tb01066h.x>
- Mizutani, M. (1993) Notes on the Lejeuneaceae 18. Japanese species of the genus *Archilejeunea*. *Journal of the Hattori Botanical Laboratory* 73: 175–182.
- Nees von Esenbeck, C.G. (1838) *Naturgeschichte der Europäischen Lebermoose mit besonderer Beziehung auf Schlesien und die Oertlichkeiten des Riesengebirgs*, vol. 3. Grass, Barth & Co., Breslau, 593 pp.
- Pearson, W.H. (1892) *Lejeuneae Madagascariensis. Forhandlinger i Videnskabs-Selskabet i Kristiania* 1892 : 3–9.
- Pearson, W.H. (1924) A collection of Hepaticae from Mount Elgon. *Arkiv för Botanik* 19(5): 1–16.
- Pócs, T. & Sass-Gyarmati, A. (2006) New or little known epiphyllous liverworts, XII. *Archilejeunea helenae* Pócs & Gyarmati, sp. nova. *Cryptogamie, Bryologie* 27: 103–109.
- Schiffner, V. (1893) Hepaticae. In: Engler, H.G.A. & Prantl, K. (Eds.) *Die Natürlichen Pflanzenfamilien, Teil. I, Abt. 3*. Engelmann, Leipzig, pp. 1–144.
- Söderström, L., Hagborg, A. & von Konrat, M.J. (2012) Notes on early Land Plants Today. *Phytotaxa* 65: 41–42.
- Spruce, R. (1884) Hepaticae of the Amazon and the Andes of Peru and Ecuador. I. *Transactions and Proceedings of the Botanical Society of Edinburgh* 15: 1–308.
- Stephani, F. (1888) Hepaticae africanae. *Hedwigia* 27: 106–113.
- Stephani, F. (1890) Die Gattung *Lejeunea* im Herbarium Lindenberg. *Hedwigia* 29: 1–23.
- Stephani, F. (1895) Hepaticarum species novae VII. *Hedwigia* 34: 43–65.
- Stephani, F. (1910) Hepaticae. In: Rechinger, K.H. (Ed.) *Botanische und Zoologische Ergebnisse einer Wissenschaftlichen Forschungsreise nach den Samoa-Inseln, den Neu-Guinea-Archipel und den Salomoninseln von März bis Dezember 1905. III. Teil. vol. 85*. Akademie der Wissenschaften in Wien. Mathematisch-Naturwissenschaftliche Klasse, Denkschriften, pp. 193–201.
- Stephani, F. (1911) *Species Hepaticarum* 4. Genève & Bale, pp. 465–736.
- Stephani, F. (1912) *Species Hepaticarum* 5. Genève & Bale, pp. 1–176.
- Stephani, F. (1924) *Species Hepaticarum* 6. Genève & Bale, pp. 433–622.
- Thiers, B.M. & Gradstein, S.R. (1989) Lejeuneaceae (Hepaticae) of Australia. I. Subfamily Ptychanthoideae. *Memoirs of the New York Botanical Garden* 52: 1–79.
- Udar, R. & Awasthi, U.S. (1981a) The genus *Archilejeunea* (Spruce) Schiffn. in India. *Geophytology* 11: 72–79.
- Udar, R. & Awasthi, U.S. (1981b) Observations on *Archilejeunea apiculifolia* St. from India. *Journal of Bryology* 11: 709–714.
<http://dx.doi.org/10.1179/jbr.1981.11.4.709>
- Udar, R. & Awasthi, U.S. (1982) The status of *Spruceanthus marianus* (Gott.) Mizut. in India. *Journal of Bryology* 12: 33–36.
<http://dx.doi.org/10.1179/jbr.1982.12.1.33>
- Vanden Berghen, C. (1951) Contribution a l'étude des espèces africaines du genre *Archilejeunea* (Spr.) Schiffn. *Revue Bryologique et Lichénologique* 20: 112–121.
- Verdoorn, F. (1933) Hepaticae selectae et criticae, series V et VI. *Annales Bryologici* 6: 95–104.
- Verdoorn, F. (1934) Studien über Asiatische Jubuleae. *Annales Bryologici*, suppl. 4: 1–231.
- Wigginton, A. (2004) *E.W. Jones's Liverwort and Hornwort Flora of West Africa*. National Botanical Garden (Belgium), Meise, 443 pp.
- Wilson, R., Gradstein, S.R., Schneider, H. & Heinrichs, J. (2007) Unravelling the phylogeny of Lejeuneaceae (Jungermanniopsida): evidence for four main lineages. *Molecular Phylogenetics and Evolution* 43: 270–282.
<http://dx.doi.org/10.1016/j.ympev.2006.10.017>