377)

George Perry's molluscan taxa and notes on the editions of his *Conchology* of 1811

RICHARD E. PETIT



RICHARD E. PETIT

George Perry's molluscan taxa and notes on the editions of his *Conchology* of 1811 (*Zootaxa* 377)

72 pp.; 30 cm.

8 December 2003

ISBN 1-877354-22-8 (Paperback)

ISBN 1-877354-23-6 (Online edition)

PUBLISHED BY

Magnolia Press

P.O. Box 41383 St. Lukes

Auckland 1030

New Zealand

e-mail: zootaxa@mapress.com

http://www.mapress.com/zootaxa/

© 2003 Magnolia Press

All rights reserved.

No part of this publication may be reproduced, stored, transmitted or disseminated, in any form, or by any means, without prior written permission from the publisher, to whom all requests to reproduce copyright material should be directed in writing.

This authorization does not extend to any other kind of copying, by any means, in any form, and for any purpose other than private research use.

ISSN 1175-5326 (Print edition)

ISSN 1175-5334 (Online edition)

George Perry's molluscan taxa and notes on the editions of his *Conchology* of 1811

RICHARD E. PETIT

806 St. Charles Road, North Myrtle Beach, SC 29582-2846, USA; email: r.e.petit@att.net

Table of contents

Abstract	. 3
George Perry	
The Conchology	. 5
Plates-Artists-Engraver	. 5
Printings-Editions-Watermarks	. 7
Index	. 9
Treatment	. 9
Sources of material	16
Sources of names	
Perry's taxa	
Genus-group names	
Species-group names	
Acknowledgements	57
Literature cited	58
Plate	70

Abstract

The molluscan taxa named by George Perry in his two publications, the *Arcana*; or the museum of natural history:, etc., published in 1810–1811, and the *Conchology* of 1811 are listed with notations on their current status, some figures being identified to the species level for the first time, some being placed in the synonymy of earlier names and others declared nomina dubia. Five species that are identified as previously unused senior subjective synonyms of names in current usage and one unused senior primary homonym are declared nomina oblita. Those declared herein to be nomina oblita are: *Triplex cornuta* Perry, 1811; *Buccinella quadrata* Perry, 1811; *Cassis rotundata* Perry, 1811; *Aranea tentacula* Perry, 1811; *Buccinella tuberculata* Perry, 1811; and *Solen viride* [sic; correctly viridis]. The names conserved are, respectively: *Murex brevifrons* Lamarck, 1822; *Cancellaria spengleriana* Deshayes, 1830; *Cassis madagascariensis* Lamarck, 1822; *Murex cabritii* Bernardi, 1859; *Turbinella* [now *Vasum*] globulus Lamarck, 1816; and *Solen viridis* Say, 1822.



Three Perry species are shown to be senior synonyms of names in current use but do not qualify for nomen oblitum status under the Code. They are: Tellina aurea Perry, 1811; Triplex rosaria Perry, 1811; and Voluta pattersonia Perry, 1811. Junior subjective synonyms affected by these three are, respectively: Tellina vulsella Hanley, 1845; Chicoreus palmarosae (Lamarck, 1822); and Lyria nucleus (Lamarck, 1811). Type species designations for Perry's genera are listed and type species are designated for Columna and Strigula. The treatment of Perry's Conchology by other authors is discussed. Three distinct editions of the Conchology are identified. Perry's Arcana is the subject of another paper now in preparation but its mollusks are treated herein.

Key words. Mollusca, Perry, *Arcana*, *Conchology*, validity of taxa, type species designations, *nomina oblita*

George Perry

Almost nothing is known about George Perry except that he published two natural history works. The first of these (Perry, 1810–1811) is the *Arcana; or the museum of natural history:*, etc., a monthly publication issued from January 1810 through September 1811 that included birds, mammals, fish, reptiles, insects, and other animals. This rare serial is the subject of a paper now in preparation but its mollusks are treated herein. His other, better known, work (Perry, 1811) is the *Conchology, or the natural history of shells: containing a new arrangement of the genera and species, illustrated by coloured engravings executed from the natural specimens, and including the latest discoveries.* It is this latter work that is the primary subject of this paper. All of Perry's molluscan taxa are listed showing their present status. Many of his taxa are placed in synonymy of other taxa for the first time and others are declared to be either *nomina dubia* or *nomina oblita*.

Unfortunately, not even birth and death dates are known for Perry although he must have lived in, or close to, London. The only record found of any George Perry who might be the same is in the Membership Lists of the Palaeontographical Society, founded in 1847 in London. These lists show that George Perry, Esq., was an early member. He was not in the first list for 1848 but was in the second list published that year with his address given as 39 Spencer-street, Northampton-square, a listing repeated in the 1851 list. In the 1853 list his address was changed to Croydon, and in 1856 it was emended to Croydon and Charterhouse. The 1859 list is the last in which he appears (the next list was not published until 1865). In 1848 James de C. Sowerby and G. B. Sowerby, Jr., were both members, the former remaining a member until his death in 1871. The last list on which G. B. Sowerby, Jr., appears is for 1853. G. B. Sowerby, Sr. was never a member. Whether or not this is the George Perry who wrote the works in question has not been determined. The improbability of Perry and G. B. Sowerby, Sr. being members of the same scientific society will become obvious in the following discussion.

The Conchology



Perry's Conchology, or the natural history of shells: containing a new arrangement of the genera and species, illustrated by coloured engravings executed from the natural specimens, and including the latest discoveries [hereinafter referred to as Conchology] was published in 1811. The introduction is dated January 1, 1811, but Sherborn (1922: ci) dates it as April, 1811. Unfortunately, no reason is given for that date, but we accept it herein as have all other authors. However, as early as April 1, 1810, Perry, in his Arcana, referred to a species in "Mr. Perry's Work on Conchology" and in September, 1810, he referred to genera "recently described and established by the Editor of this work, in a large work on the History of Shells, shortly intended to be published...." It appears that the Conchology was in production for a long period. Nevertheless it remained in print for many years after publication as will be discussed below.

The *Conchology* is folio in size with 61 hand-colored plates. Referred to for many years as the only work on shells printed by the expensive hand-colored aquatint process, another such work has now been so identified (Dance & Heppell, 1991: 46). The text, other than the four-page introduction and an index at the end, consists only of unnumbered pages facing each plate describing or commenting on the shells figured, one page facing each plate.

Plates-Artists-Engraver

Most of the figures in the Conchology, despite past criticisms which will be discussed below, are of good quality and are identifiable. Others show shells that are garishly colored and poorly drawn. The name of the artist for the Conchology is not stated but a reading of the introduction leads to the conclusion that they are by Perry, who stated that "the plates are engraved, and coloured after the original drawings, by Mr. John Clarke." There is so much difference in the quality of the plates that it is impossible to attribute them all to Perry. Perry was a very good artist as attested to by the many plates in his Arcana drawn by him. Perry drew many of the plates for his Arcana, all of which were engraved by T. L. Busby. Most of the Arcana plates drawn by Perry are realistic renditions. Although the two works were being produced together for some time there was no crossover between the engravers. Some plates in the Arcana were not drawn by Perry, but all are attributed. Several dealers in prints advertising on the Internet incorrectly list Conchology plates as "original drawings by John Clarke, engravings by William Miller." These errors are due to a misreading of the introduction and to the publisher's name being on the plates. All plates are imprinted at the bottom "London, Pub. by W. Miller, 1810" with minor variations in placement and typography.

The figures in the Arcana, and some in the Conchology, are excellent and are equal to most of those produced by the Sowerbys, the best known natural history artists of the time, and far superior to many other works. It is here speculated that many of the Perry figures were drawn by lesser artists who were given the task of either drawing an available shell or copying an existing figure. Some of the odder figures in the Conchology are copied with varying degrees of accuracy from older works. Some authors have maligned a Perry figure as unidentifiable, but at the same time expressed no problem with a similar Martini or Chemnitz figure. A discussion appears later herein. There are three basic problems with some of the figures in the Conchology: the anterior portion of the aperture is sometimes distorted as if drawn by someone only vaguely familiar with gastropods trying to draw one from memory; excessive and incorrect color; and the "free form" shape of many of the bivalves. Many of the bivalve figures are among the worst, but a few are good. Surely the Arca figures were not from the pen of the same person who drew the Venus figures. It is remarkable that Perry's work has been vilified because some figures are not identifiable and others are stylized. Such comments are never heard about the works on which the names of Linnaeus, Gmelin, Röding, and others are based. Many of their nomina are based on figures as outrageous as many in Perry. Although Montfort's wood-block figures are poorly executed, distorted and stylized, there is no criticism of them to be found. Rafinesque, considered by many to be a malacological giant, drew poor illustrations of imaginary creatures which he then named (see Moore, 1981; Petit, 1985; Markle, 1997). Those criticizing Perry never mention others whose works are even worse. Also, the fact that painted shells were sometimes passed off as genuine cannot be ignored. For a discussion of painted and altered shells, see Dance (1966: 80–86; 1986: 54–57).

Several authors have indicated that some of the figures in the *Conchology* were copied from Martini and Chemnitz, but Perry also copied from other sources. Within the *Conchology* only three prior illustrations are mentioned. Perry states that his *Cassidea cornuta* (pl. 34, fig. 3) "has been figured by Regenfus [sic; = Regenfuss, 1758] in his large Work of Conchology." Comparison shows that Perry's figure is only a slightly changed version of Regenfuss' figure (pl. 5, lower figure 49). Regenfuss' figure is of *Melongena melongena* (Linnaeus, 1758) and was added to the citations for that species by Linnaeus (1767: 1220).

Perry also mentioned that his *Melania acuta* (pl. 29, fig. 2) had been previously illustrated by Born. Martens (1872: 5) identified Perry's figure as *Lymnaea stagnalis* (Linnaeus, 1758), which seems unlikely until Born's figure (1780: pl. 16, fig. 16) is consulted and then it is easy to see how it could have been transformed into the stylized Perry figure. The only other illustration referenced is under *Buccinella coerulea* where Perry states that it has been "described, but not well figured, by the German Conchologist Martini." Although stated to be from a specimen in the British Musem, it is quite likely that Perry's figure is an "improved" copy of Martini (1777, pl. XCV, fig. 916) which it matches in color.

6

No measurements are given for any of the figures and they are drawn to different scales, often complicating identification. There are many highly stylized figures that are impossible to identify and are herein considered to be *nomina dubia*. Three newly identified figures that have junior synonyms in common use are herein declared to be *nomina oblita* as is a previously unnoticed senior primary homonym. Two previously noted synonyms, declared to be *nomena oblita* in ways not meeting the current Code, are also formally declared to be such. Other names are discussed that are senior synonyms but do not meet the criteria of Article 23.9.1 of the Code (International Commission on Zoological Nomenclature, 1999b).

There was evidently a mix-up in the plate numbering at the beginning of the work as some plates are misnumbered. Plates 5, 6, and 23 are engraved with the numbers 3, 4, and 5 respectively. A second plate 3 appears as plate 54 and is listed on the index as a "supplementary plate" although it is obvious that it should have followed plate 2 illustrating similar shells.

Printings-Editions-Watermarks

As will be shown below, there are three easily identifiable editions of the Conchology. That there are at least two editions of the *Conchology* was shown by Johnson (1970: 287) who examined a number of copies, including one in the British Museum (Natural History) [now The Natural History Museum] that had been included in a paper on printings by Hopwood (1946: 152-153). Although Hopwood went into more detail about the watermarks in the various printings, and listed eight printings, he did not note that the type had been reset, stating that the type was "kept standing." Unfortunately, Hopwood's listing of printings is not tenable; the watermarks in various editions are not uniform as watermarked paper seems to have been used randomly. Johnson (1970) correctly stated that the type had been reset and identified two editions. It cannot be determined exactly when the "second edition" came into being but the first so identified are those with the text watermarked 1818. A copy of that printing is on hand with most of the text pages watermarked "H Smith 1818" and the plates not watermarked, seemingly identical to Hopwood's example 4. This "second edition", said to have "minor differences in the text and letterpress" by Johnson (1970: 287), is reset in a slightly larger type in most places, causing a noticeable difference in format, but no words have been found to be altered. It is here reported for the first time that there is a "third edition" in which the type has again been reset in a still larger font. This edition was referred to by Johnson as possibly a "second printing of the second edition", as he only saw one copy. These editions can easily be determined by the text to Plate I. In the 1st edition the block of text under "Remarks" consists of 15 lines, in the 2nd there are 16 lines with the single word "criterion" being the final line, and in the 3rd there are 16 lines with the final line consisting of the two words "indubitable criterion."

377

Also, there is no date on the title page of the 3rd edition. The earliest watermark found for this 3rd edition is 1820, although it could not have been printed until 1821 or later. The 2nd edition was printed as late as 1821, as a copy exists with J. Whatman 1820 and 1821 watermarks (personal communication, S. Jordan, August 26, 2003). The book continued to be printed as late as 1836; plates with that watermark date were cited by Hopwood and are present in the copy in the library of the Academy of Natural Sciences (personal communication, G. Rosenberg, November 11, 1998). One plate in a copy of the 3rd printing which includes 1827 watermarked plates and 1823 watermarked text has one plate with a "J Whatman / Turkey Mill / 1826" watermark (personal communication, R. Cowie, October 21, 2003). This is the only report of the well-known "Turkey Mill" watermark. It is obviously impossible to date an edition from the watermarks, as older paper was often used. Judging from the variously dated watermarks (or lack thereof), each of the three editions seems to have been in print for rather long stretches at a time. Therefore, the number of printings can never be known.

Hopwood, for the original issue, listed the plates as being watermarked "J Whatman" and dated 1808, 1809, and 1810. Watermarks on text pages were not mentioned. A copy at hand has plates watermarked "J Whatman 1806," "J Whatman 1809," J Whatman 1810" and some with the three-line watermark "1808 / J Whatman / W Balston." This 1806 watermark is the earliest reported for this work. The text in this copy is on paper watermarked "W Turner & Son." The "W. Miller, London 1810" watermark reported by Johnson has not been seen by this author nor was it listed by Hopwood.

The title pages of the first and second editions show the place of printing as London, followed by "Printed for William Miller, Albemarle-Street / by W. Bulmer and Co. Cleveland-Row, St. James's / 1811." The title page of the third edition is imprinted "Published by William Miller, Albemarle-Street / "Howlett and Brimmer, printers, Frith Street, Soho" and is not dated.

Early issues of the *Conchology* have the index page at the end printed by "W. Bulmer and Co., Cleveland-row, St. James." The 2nd edition with 1818 watermarked paper has the index page printed by "J. M'Creery, Black-Horse-Court, Fleet-Street." The index page in the 3rd edition is printed by "Howlett and Brimmer, 10, Frith-Street, Soho." Also, in the first printing there is a two-page (one leaf) advertisement headed: "A catalogue of the principal books published by William Miller, Albemarle-Street, up to the spring of 1811." There is no such advertisement in later printings.

Hopwood (1946: 152) reports that the book was first issued at 16 guineas (=£16 16 shillings). That price must have remained unchanged for many years as the University of Michigan has a 3rd edition in original boards with a printed label on the front cover showing that price. Hopwood gives antiquarian book dealers' prices for various years from 1912 (£1 8 s.) to 1944 (£15 10s.). In the early 1960s it was routinely listed at US\$100 but increased steadily. A decade or so ago the price began escalating as antiquarian print dealers discovered these large colorful plates. In June 2003 a search of web sites selling such



prints located two dealers listing Perry plates, all of which were priced from \$275 to \$325 each with the average being over \$300. With \$300 per plate as an average price, the 61 plates in one book would bring in \$18,300! It is no wonder that the *Conchology* is now being listed by dealers for over \$8,500 (up from \$5,500 two years ago).

Index

Immediately following the final plate, there is a full page "Index to the plates and genera" on which all plates are listed and with three different symbols used to indicate the genera: (1) "adopted from Linnaeus", (2) "from Bruguière and Lamarck," and (3) those "now first invented and adopted by the Author of this work." Two genera indicated as "new" are Linnaean names (*Argonauta* and *Patella*) and another is of Müller (*Pecten*). Another Linnaean genus, *Ostrea*, is denoted as being both (1) and (3)! There is much that is inexplicable in Perry. In the text he decries Linnaeus' failure to separate *Pecten* from *Ostrea* and states that "as most of the succeeding Conchologists have chosen to separate it, I have adopted the same mode and have taken the definition from Mr. Montague" Also, the genus *Cassidea* is marked as new and in the text is defined without mention of Bruguière. One of Perry's new names, *Monoplex*, is indicated as being in category (2). Others of Perry's new names are homonyms of earlier names. See additional comments under "Sources" below.

Treatment

Perry's work was ignored in the molluscan literature until 1827, when the first mention of it appears. The treatment of Perry's works in the 19th Century is well-documented as will be shown below, but there has been no speculation as to why it was ignored before 1827. In his major work of 1817, Dillwyn did not include Perry in the extensive list of books to which he had referred. That a book the size of the Conchology, published in London, could escape the notice of everyone writing about mollusks is implausible. Here a guess will be made as to the reason for this early rejection (a second reason for later rejection will be given below). In the late 18th and early 19th Centuries the work of Linnaeus was considered by almost all English writers to be inviolate and any "improvement" of the Linnaean systems (i.e., Jussieu's system in botany and Lamarck's expansion of zoological genera) was the subject of scorn. The prime mover of the pro-Linnaean movement was James Edward Smith (1759–1828), who founded the Linnean Society of London after purchasing the Linnaean collections. His contempt for the French systems was such that authors feared to adopt Jussieuan or Lamarckian taxa as it could endanger their membership in the Linnean Society, the foremost scientific society in England. In 1821 James Edward Gray wrote the systematic part of A natural arrangement of British plants, a work utilizing the genera of

Jussieu, issued under the authorship of his father, Samuel Frederick Gray. It was so wellknown that he was the author of the systematic part of that work that when proposed by friends for membership in the Linnean Society in 1822 he was "black-balled" and denied membership as being opposed to the Linnaean System (Gunther, 1974: 64, 73). It was only after Smith's death that members of the Society could feel free to embrace a newer concept (also, see Dance, 1966: 116–117; 1986: 81–83 for a discussion of this problem). As Perry, in both the Arcana and the Conchology, spoke highly of Lamarck and utilized genera proposed by Bruguière and Lamarck, he would have been anathema to Smith and therefore to the writers of that time (Wood, Dillwyn, Brown, et al.). This is reinforced by the fact that, although Perry must have been on good terms with William Bullock as he utilized so much of his material, he is not mentioned in Bullock's 1812 work on his museum. Neither did Bullock use any of Perry's plates from the Arcana, although many of the plates were the same size and format and based on the same animals. Bullock was not only a member of the Linnean Society but he lists (Bullock, 1812: 125) among his shells "several articles which were once the property of the celebrated Sir Charles Linnaeus; ... presented by Dr. Smith, President of the Linnaean [sic] Society." [The inconsistent spellings "Linnaean" and "Linnean" arise from the selective use of his given name, "Linnaeus," which was changed when he was ennobled to "von Linné". Bullock did not correctly spell the name of the Society on the title page of his book. Also, see Dance, 1966: 116; 1986: 82.] G. B. Sowerby I was a member of the Linnean Society from March 5, 1811 (Cleevely, 1974: 510) and thereafter his name was always followed by "F.L.S."

John Edward Gray published a monograph on *Cypraea* in the *Zoological Journal* in parts (Gray, 1824–28a). Dillwyn (1827: 315) wrote a short note concerning Gray's treatment of some of the *Cypraea*, and mentioned *Cypraea dama* Perry. On the same page as Dillwyn's article was a footnote by G. B. Sowerby I in which he stated: "As to Perry's work, the worst of all bad books, it ought never to be cited." Dillwyn's use is the earliest citation of a Perry name that has been located except by Perry himself in the *Arcana*.

Gray (1828b) published, in the same journal, a paper giving "additions and corrections." In that paper he acknowledged and accepted Perry's *Conchology* with the comment: "I have ventured to refer to this work, as I consider that it is just that every author should be quoted; and this author has anticipated Lamarck, Swainson, and Sowerby in several species." Gray had no hesitation in suppressing his own *Cypraea princeps* in favor of the earlier *C. valentia* Perry. However, when Sowerby realized how acceptance of Perry would affect names he had proposed, he, in today's vernacular, "went ballistic." Gray knew, of course, of Sowerby's rejection of Dillwyn's use of a Perry name (Sowerby, 1827) and was probably aware that Sowerby would take exception to his recognition of Perry. His use of "ventured" seems appropriate.

Sowerby immediately wrote a paper (1828) in the form of a letter to W. J. Broderip, one of the editors of the *Zoological Journal* and, incidentally, describer of *Cypraea nivosa* which Gray suppressed in favor of the prior *C. dama* Perry. It is doubtful that there is a

more vituperative epistle in all malacological literature. Reading it over a century and a half after publication, one is still embarrassed for Perry. Sowerby's tirade was much too long, so much so that it is easy to imagine some previous unpleasantness between Sowerby and Perry of a more serious nature.

To add further insult, there is appended to Sowerby's five-page article a "Note by the Editor" in which it is stated that "it is our unanimous opinion that Perry's *Conchology* is not a work worthy of being cited as authority." The Editor at that time was N. A. Vigors, to whom Dance (1966: 121; 1986: 88) attributed the note. It is impossible to determine who actually wrote the note, but Broderip, G. B. Sowerby I and J. de C. Sowerby were all "coeditors".

His acceptance of Perry's work probably cost Gray the authorship of *The Conchological Illustrations* which was originally planned to be written by Gray and illustrated by G. B. Sowerby II. For details on this aborted authorship, refer to Sherborn (1909: 331–332). G. B. Sowerby I was, of course, in charge of the project even though his son was the illustrator. When Gray delivered the first 24 pages of text, unfortunately on *Cypraea*, Sowerby wrote him that he did not like the printing job. Evidently Sowerby and Gray failed to agree on the matter as within a few weeks of Sowerby's letter the first issues were published by Sowerby with no mention of Gray. There was more to this decision than bad printing; the proofs still exist and the few typographical errors could have been corrected by the printer in a matter of minutes. The problem, in all probability, was Gray's insistence on using Perry's names.

I would like to interject here a note about the authorship of the *Cypraea* monograph in *The Conchological Illustrations*. Almost all modern authors show G. B. Sowerby II as author, which indicates that they did not read the introductory text of either the *Cypraea* monograph or of the work as a whole. G. B. Sowerby II was responsible for only four of the monographs and G. B. Sowerby I wrote all the others, including the one on *Cypraea*.

After Gray, the next person to recognize Perry's work was G. P. Deshayes (1838–1845) in the second edition of Lamarck's *Histoire Naturelle des Animaux sans Vertèbres*. Deshayes possibly did Perry more harm than good. Although he recognized many of Perry's taxa as being ones later named by Lamarck, Deshayes would not surrender a Lamarckian name. References to Perry's figures were simply placed in synonymy with the only reference being plate and figure numbers, without date and usually without name. As a result later workers assumed that these figures in synonymy post-dated Lamarck and were ignored. Deshayes' treatment of Lamarckian names has been mentioned in the literature before (Petit, 1984).

Apparently Deshayes did not acquire a copy of the *Conchology* until after some of the mollusk section of his edition of Lamarck had been written. His first mention of a Perry name is in volume 8 at page 305 and the second did not appear until volume 9 at page 284. Deshayes recognized 124 Perry species. He credited Perry with authorship of only three [Fusus forceps, Cypraea valentia and Cypraea camelopardalis], and used Perry's names



but with himself as author on two [Fusus variegatus and Ranella perca]. Only two other Perry names were listed by name in the synonymies, all of the other listings being plate and figure reference only. In keeping with the custom of the time, Deshayes usually assumed authorship of any name he placed in a different genus from that in which it was originally described. However, some of his actions are inexplicable. He changed Perry's Murex forceps to Fusus forceps and attributed authorship to Perry, but when he changed Murex variegatus Perry to Fusus variegatus, he assumed authorship for himself, putting his own F. laticostatus in synonymy. He also assumed authorship of Biplex perca when he placed it in Ranella.

Deshayes had two comments about Perry. One is quoted by Hopwood (1946: 152) but the original source has not been determined. Fide Hopwood, Deshayes wrote: "Cet ouvrage, d'un grand luxe, offre des figures gravées et coloriés faites pour plaire aux yeux des personnes peu versées dans science, et non à ceux des vrais naturalistes, car la plupart n'ont rien d'exact dans la forme et la couleur; quant au texte, on y remarque des erreurs nombreuses ..." ["This publication, of a great luxury, offers engraved and colored figures, made to be pleasant to the eyes of people with little scientific knowledge, but not to those (eyes) of true naturalists, for most of them have nothing correct in the shape or color, and concerning the text, one remarks numerous errors therein ..." (translation by Stéphane Pras)].

Deshayes also stated (1843: 466), under *Fusus forceps* Perry: "Nous rendons à cette espèce son premier nom que M. Kiener n'aurait pas dû lui ôter. Quoique l'ouvrage de Perry soit peu estimé, il contient cependant quelques bonnes figures, et celle qu'il a donnée du *Fusus forceps* fait reconnaître l'espèce avec la plus grande facilité." [We give back to that species its first name, that Mr. Kiener should not have removed. Although Perry's publication is held in low esteem, it nevertheless contains a few good figures, and that one he gave for *Fusus forceps* allows recognition of the species with the greatest ease." (translation by Stéphane Pras)].

Reeve, who was publishing books illustrated by the Sowerbys, took exception to Deshayes and under his listing of *Ranella pulchra* (Reeve, 1844, *Ranella* sp. 47) stated "I cannot agree with M. Deshayes in giving priority to the specific name assigned to 'The finned Frog' by Mr. Perry. That author has long forfeited the notice of scientific men by his absurd names and pantomimic display of figures." That is an amazingly disingenuous statement as Perry named the shell *Biplex perca* and it is only by placing the species in *Ranella* that "frog" comes into the picture. In his effort to humiliate Perry, Reeve neglected to mention that in his description Perry stated that "... the fins or spines certainly resemble those of the perch, from which circumstance we have given it the name of *Biplex perca*."

Reeve was unrelenting and the following year (Reeve, 1845, Cypraea Sp.17, *Cypraea melanostoma* Sowerby) he reached new depths of disparagement when he wrote that "Since Mr. Sowerby described this species in the Tankerville Catalogue, under Mr. Leathes' manuscript name of *Cypraea melanostoma*, it has been published by one or two

authors under that of *Cypraea camelopardalis*, from its having thus been originally named by one Mr. Perry; the writings of this person have, however, been censured a mass of buffoonery wholly unworthy of credit. So much has already been said on this subject by Mr. Sowerby, (Zoological Journal vol. iv, p. 218) that it only remains to state how fully I concur in the opinion there recorded. If every charlatan who sets himself up for a naturalist, with brush in one hand, and writing-tool in the other, is to be regarded as an authority on scientific matters; all the zoological sign-boards in the kingdom might be quoted as figures of reference. For my own part, I consider that the 'Camelopard of Perry' should no more enter the nomenclature of zoology, than the 'Red Lion of Brentford'."

Still not satisfied, he then (Reeve, 1847, *Fusus laticostatus* Deshayes, Fusus Sp. 33) stated that "M. Deshayes has adopted the title of *Fusus variegatus* for this species in his edition of the Anim. sans vert., owing to its having been so named under the head of *Murex* by Mr. Perry; the labours of that author are, however, unworthy of credit and have long ceased to be acknowledged in this country."

Gray (1847) treated some but not all of Perry's genera. The Adams brothers, in their important work (H. & A. Adams, 1853–1858) mentioned only about half of Perry's new genera, placing all but one in the synonymy of older names. Only a few Perry species are mentioned in the entire work. The only Perry genus accepted by them was *Columna* and even there neither of Perry's included species was listed. Later workers treated Perry's names in a variety of ways, mostly "picking and choosing" which Perry names they would treat. Monographers consistently treated some of Perry's names in the group being monographed and ignored others (e.g., Rehder on *Harpa*; Radwin & D'Attilio on *Murex*, Burgess on *Cypraea*, Powell on *Patella*, etc.). Many of these omissions are inexplicable. As one example from many, in recent years workers on *Distorsio* Röding, 1798 have shown *Distorta* Schumacher, 1817, as a synonym while ignoring *Distorta* Perry, 1811, but treating Perry's *Distorta* species in the same work. Cossmann (1895–1925) in his *Essais de Paléoconchologie Comparée* treated only some of the genera proposed by Perry. Why authors ignored some Perry names but utilized others is not known.

There is little, if any, further mention of Perry's taxa until Martens wrote a paper on the Mollusca collected during Cook's travels, or shells that probably came from his travels, that were described by various authors. He wrote (1872: 4–5): "Perry's Conchology, 1811, fol., is of little value for our purpose; although there are various indications of collecting localities that point towards Cook's travels, such as New Zealand or [Tonga], the reliability that the figured shells stem from the indicated localities is considerably lower with Perry than with Martyn and others." Martens continued with an example of a Florida shell stated by Perry to be "obtained via our latest discoveries in the southern seas" and two other examples of incorrect localities. [translations by R. Bieler]. However, he did identify some of Perry's figures. Perry was certainly not the only author of that time to have incorrect localities and many noted authors described new taxa without stated localities.

377

Jousseaume (1880) recognized two Perry genera in his division of the "purpuridés" and Melvill (1888) published a list of *Cypraea* in which he recognized Perry species. Melvill (1888: 198–199) took Perry to task for not consulting previous authors, etc. He then stated that Perry "has fallen under the lash of subsequent authors, especially Messers Sowerby and Reeve, who ignore all his descriptions, but in justice to him and to the law of priority, where his species can be deciphered, many of the paintings being very fanciful, it is only right to attach his names...." In a footnote Melvill notes that Perry was not careful but neither were authors subsequent to 1811 who either ignored or were unaware of his work.

The Conchology, after being consigned to oblivion by Sowerby et al., and effectively hidden by Deshayes, was neglected as a source of valid molluscan names, except for a few scattered usages, until 1902 when Hedley and Gatliff independently published papers on Perry's Australian species. Hedley (1902: 25) remarked about the *Conchology*: "The work is post-Linnean, the species are named binomially, and are systematically figured and described. In short, the nomenclature fulfils the requirements of modern zoological legislation, and must take precedence of younger names. The illustrations vary in merit; some are excellent, others are wretched; but most, if not all, are recognisable." He continued with: "Sowerby and Reeve were not disposed to acknowledge any work which upset their species. They ungenerously denied Perry's right to be cited, and assailed his nomenclature with hatred and abuse. So powerful and evil was their influence that Perry's 'Conchology' has been practically suppressed by London writers. Even the brothers Adams, who offended against modern usage by adopting names from polynomial and pre-Linnean writers, yet refused to admit Perry's names." Hedley's reference to the influence of Sowerby and Reeve being "powerful and evil" is not an overstatement.

Gatliff (1902: 75), in his paper identifying Australian species for which Perry's names are earlier than those then in use, discussed the *Conchology* and its figures and stated: "It therefore follows that when the figures are past recognition the work is useless so far as it relates to them; but other figures are excellent."

Another half-century passed with little notice to the *Conchology* until there was a flurry of notes in America. Clench & Turner (1952: 320), in comments on *Epitonium greenlandicum* (Perry) wrote: "Perry evidently attempted to copy the figure in Chemnitz and embellished his drawing with a few characters certainly not found in any specimen of this species that we have seen. All of Perry's figures on plate 28 are overdrawn and he also gave free rein to his imagination when coloring them." Perry's figure is indeed almost a duplicate of the one in Chemnitz but with the addition of improbable red dots on the posterior ends of the axial ribs! John Clarke exercised a lot of "artistic license" when coloring the plates.

John Q. Burch, a California shell and book dealer who edited the *Minutes of the Conchological Club of Southern California*, wrote several notes on Perry's names. In 1955 (Burch, 1955) a short note objected to the introduction of old names, with reference to a



15

Conus described by Born and to Hexaplex Perry, followed by another note about Hexaplex in 1956 (Burch, 1956a). These two papers are notable for perpetuating a modern myth that "the Sowerbys ... were very thorough scholars" and wrongly stating that Sowerby, "as early as 1825 [sic]" expressed the opinion that "Perry's names were too vaguely defined for serious consideration." Sowerby (1827, 1828) objected to Perry's names being used on several grounds, but "vaguely defined genera" was not mentioned as a reason. Burch's first note about Perry was probably due to advance notice of Hertlein & Strong's work printed later the same month as his second (1956a) was definitely the result of their (1955:255) use of *Hexaplex*. This was a rather mild paper as Hertlein and Strong were west coast scientists of repute whom Burch did not wish to offend with harsh criticism. In a letter Butot (1956: 1–2) took exception to Burch's 1955 comments and placed blame on Sowerby and Reeve for not "giving due credit to Perry." Strangely, in his prefatory remarks preceding Butot's letter, Burch (1956b: 1) stated that "it was not the intention to challenge the great contribution to conchology made by Perry, but only to question certain generic terms as being too vaguely defined." Obviously Burch's opinion was affected by more than genera as the next year when Hollister (1957) reintroduced Fasciolaria hunteria (Perry) it produced an instant response from Burch (1957a: 9). Burch's criticism will not be quoted at length as it suffices to say that it is sophomoric and unnecessarily derisive. Burch, in company with many others, was either not aware, or did not care, that Deshayes (1845: 433) had identified as F. distans Lamarck the readily recognizable Perry figure about which Burch states: "No living shell from anywhere could be classified from the figure or the description." A mild rebuttal from Dr. H. B. Baker (1957: 7-8) appeared a few months later. Baker pointed out that the figure in Lister that is basis for Lamarck's name is not as recognizable as the one in Perry. Later in the same year Burch (1957b) again took exception to the use of *Hexaplex* and referred to Perry's figures as "modern art."

Burch (1958) took another shot at Perry when he reviewed Wilkens (1957) who commented both on Perry and on Burch's articles. Wilkins (1957b) wrote an excellent paper on the Cracherode collection in which he addressed the Conchology and its treatment. He gave a good account of the denigration of Perry by Sowerby and Reeve and quoted some of their statements. He also commented (Wilkins 1957b: 135): "While it must be admitted that much of the letterpress in both the Arcana and the later Conchology is not particularly good, the figures (although at times fantastic and of almost dream-like quality) are sometimes excellent and clearly recognizable. The descriptions meagre or prolix according to the fancy of the author, are sufficiently clear for there to have been no real reason to reject Perry's valid but unwelcome names, even if it meant upsetting those established by the various British authors who resolutely refused to study their predecessor's book." He also (Wilkins 1957b: 136) commented: "Both Sowerby and Reeve realized only too well that much of their careful work would be upset by the general adoption of Perry's nomenclature, and their opinions were therefore not without bias. By the force of their displeasure they succeeded in keeping this awkward volume in the background for the rest of their lives, an obscurity from which it did not emerge until referred to by Melvill in 1888 and by

(377)

both Hedley and Gatliff in 1902." Wilkins then addressed Burch's comments with: "John Q. Birch [sic; =Burch] has published a note on the resurrection of early names ... in which he makes a belated attempt to uphold the dictum of G. B. Sowerby, W. J. Broderip and his followers in suppressing Perry's names. He apparently ignores the happenings of the last fifty years...."

Emerson & Old (1963: 3) in describing Cymatium perryi commented: "Nevertheless, Perry's illustrations, poorly executed as they may be are certainly no worse than those in some recently issued works that are offered as scientific endeavors."

Even now authors speak ill of Perry's work. Dance (1966: 120-121; 1986: 88-89) wrote: "The reasons for the vilification which was later heaped upon the Conchology are not only attributable to its figures, however, but to its text, where there is an assortment of odd-sounding new generic names: Monoplex, Biplex, Hexaplex, Polyplex, etc., besides a number of new specific names." He adds that "the acceptance of Perry's names would have upset much of the work of both Sowerby and Reeve so that they had some reason to denigrate him." That statement should have been continued to point out that the Conchology was published before G. B. Sowery I ever described a new species and before Reeve was born. If Reeve had wished to simply ignore Perry's work, he could have done so without comment but he was probably catering to Sowerby's hatred as the Sowerbys were providing the plates for Reeve's Conchologia Iconica and other works. Dance also quoted some of the derogatory comments made by Sowerby and Reeve and continued to deride Perry unnecessarily when he stated (p. 103) that "Perry also based some of his nightmarish illustrations in the Conchology on specimens in the Jennings cabinet...." In another work Dance (1978: 90-93) reproduces a small, uncolored version of one of the worst Perry plates with the comment that "the shells are so badly drawn and so garishly coloured that they have become caricatures", as well as other demeaning remarks. It is unfortunate that he did not have before him at that writing a copy of his later book on classic shell prints (Dance & Heppell, 1991), in which plates by other authors that are much more garish and over-colored are presented. In that work one of the better Perry plates (pl. 13) is reproduced but the authors evidently could not resist including a plate of Oliva (pl. 41) that has, in addition to a fine central figure of Oliva porphyria (Linnaeus), four smaller figures, one of which is justifiably called "odd-looking." They also state that Perry's plates are "like no others in the literature of natural history, the bold lines and garish colouring often giving many of the shells a surreal quality," a statement belied by some of the other plates reproduced.

Sources of material

Perry seems to have known a large number of collectors who allowed him access to their collections. In the Conchology he acknowledges some 29 collectors and museums in addition to his own, which is mentioned 24 times. The British Museum is listed as the source

PETIT

of 12 specimens, followed by Bullock's Museum with 11 citations. The individuals with the most mentions are Dr. Lettsom (7), Mr. Latham and Lord Valentia (6 each), Mr. Spence (5), and Mr. Jennings and Dr. Combe (4 each). The remaining individuals are mentioned only once or twice each and include the Museum of Mr. Humphrey and Mr. Bligh's collection.

None of Perry's types are known to be extant. It is possible that the type of *Conus fasciatus* may be a specimen in the Sloan Collection (BMNH 2788) but Kohn (1986: 7) was not certain. Wilkins (1957b) found no Perry types in the Cracherode collection. However, some Perry types may still be identifiable in the British Museum (now The Natural History Museum) or elsewhere.

The localities cited by Perry are often incorrect, as noted by Martens and others cited above. This is not unique with Perry. Some species in the Conchology are incorrectly stated to be from South Africa. Dr. R. N. Kilburn of the Natal Museum has advised (personal communication October 4, 1982) that he has worked through the Conchology and concluded that "few, if any, actually came from South Africa. In essence, by 1811 there were no settlements here except in Table Bay, although nearby bays ... were sometimes used in bad weather. In other words, by this date practically all shells originating from here came from the cold temperate Western Cape Province. Warmer water Eastern Cape Province shells began to come to light after 1820, when the first settlement was established in Algoa Bay, Tropical Natal shells were unknown until the 1840's. This means that Indo-Pacific species such as [Cypraea species; Cypraeacassis rufa, etc.] must have come from elsewhere. Possible Cape species are Argonauta grandiformis [=argo], which does wash up in False Bay, (futhermore Bullock's Museum did have some South African material). ... The only other two 'Cape' species that I noted were Volutella nigricans and Natica quadrata; whatever their identity there is nothing in South Africa that agrees with them. It is curious that Perry had no Cape limpets, which were conspicuous, accessible and available." However, Kilburn did not notice the misnumbering on Plate 42 (see Argonauta rotunda in taxa list) and expressed doubt about Carinaria. However, it was not the Carinaria that was stated to be from South Africa but Argonauta rotunda (= A. nodosum [Lightfoot, 1786]) which is sometimes found there. Also, *Haliotis sinuata* Perry, 1811, described without any locality data, was tentatively placed in the synonymy of the South African Haliotis spadicea Donovan, 1808 by Geiger (1998: 100, 108).

Sources of names

Dance's objection to Perry's names on the basis that they are "odd-sounding" is remarkable. There are few genus-group names in zoology that are as descriptive and memorable as some of the ones proposed by Perry. Who can look at a muricid with three varices and not think of *Triplex*? Montfort's name for this genus-group, now in use, is *Chicoreus*, a

name that does not commend itself to either the ear or to ease of spelling. Perry's *Distorta* was displaced by the earlier *Distorsio* Röding, 1798. Certainly *Proscenula* is as descriptive as the earlier *Crepidula*, and it cannot be disputed that *Aculea* is more descriptive than the equivalent *Haustator*. A long list could be given of names proposed in recent years that have no such redeeming features. In Dance and Heppell (1991: 42) Perry's choice of names is again criticized in the statement that "Perry's names were imaginative and idiosyncratic."

Wilkins (1957b) discussed both the *Arcana* and the *Conchology* as Perry mentioned the Cracherode collection in both works and figured some Cracherode shells. Wilkins shows samples of labels from the Calonne collection prepared by Humphrey, and which became part of the Cracherode collection. One such label is for a species of *Murex* labeled "Triplex pinnatus." It was from this museum label to which he had access, or directly from Humphrey's *Museum Calonnianum*, that Perry picked up the genus name *Triplex*. Another genus name from the same source is *Haustellum* in addition to several species names. There are also names that evidently came from Müller (1773–1774) but may have been picked up indirectly. Perry took some of his names from early nonbinominal works, as did innumerable other authors.

Wilkins' 1957 paper is recommended for many reasons, one being his depiction of the state of the mollusk collection at the British Museum and its curation at the dawn of the 19th Century.

It should be noted that Schumacher (1817) used, as new, some of the same genus and species names used by Perry.

Perry's taxa

Perry did not attribute the names in his works and did not indicate, except for the genera (listed in the Index), names that he was newly introducing. There are numerous instances where Perry used an older name but illustrated a different shell. As most such cases have been treated by later workers as introductions of new names and not as misidentifications they are so treated herein [e.g., the shell figured as *Cassidea cornuta* is not *C. cornuta* Bruguière, 1892 but is *Melongena corona* (Gmelin, 1791) and is treated as *C. cornuta* Perry, 1811]. In the species listings below, when the first entry is an earlier name of the same spelling, the name is attributable to the author listed. An authority is cited after such names only when the genus has changed or its status is arguable. For names attributable to Perry where there is little or no question about the identity of the species, only one or two references are given to their placement.

A few authors designated some Perry species as *nomina oblita* (singular, *nomen oblitum*) but such unilateral action taken between January 1, 1973 and January 1, 2000 has no effect as it was not allowed by the Code then in effect. In fact, the term *nomen oblitum* was

19

not even in the Code during that period. Some of Perry's species have been referred to by various authors as "nude names" or "nomina nuda", usually in check lists. This is unfortunate and misleading. A nude name is one that is not accompanied by a description or a figure. A name accompanied by a description, a figure, or a reference to a figure, cannot be a nomen nudum. The figure does not have to be recognizable. It is more appropriate to refer to such an unrecognizable name as a nomen dubium. This is an important distinction as a nude name is not an available name and does not enter into priority.

In the *Arcana* the ligatures 'ae' and 'oe' are used rather indiscriminantly. There is a species of bird, *Certhia coerulea*, with the species name as "*coerulea*" on the plate and in the text heading, but as "*caerulea*" in the text. There is no such confusion in the *Conchology*. Perry used *coerulea* as a specific name in five different genera, but they are almost always rendered as "*caerulea*" when mentioned in the literature. Although the two spellings are equivalent, both in Latin and under the Code, the original spelling cannot be changed. The ligature used for the dipthong in "*coerulea*" is definitely 'oe' as is obvious when compared with the 'ae' ligature in *Strombus laevis* in the *Conchology*. In the *Arcana* the ligature in *Cerithium laevis* is 'oe' but in view of the ambiguity within the *Arcana*, coupled with the fact that "loevis" is not a Latin word, *Cerithium laevis* is considered to be the intended spelling. It was so cited by Mathews & Iredale (1912: 11). Also, see comments under *Monoplex oboesus* in the species list below.

Of the 24 genus-group names in Mollusca introduced by Perry, or attributed to him, 8 are in current use; two are considered to be emendations of Linnaean names; five are junior homonyms; 7 are junior synonyms, one has been suppressed by the I.C.Z.N.; and one has been shown to be a species-group name.

In his two works Perry treated 352 nominal species-group taxa of Mollusca, of which 35 are attributable to earlier authors; 166 are subjective synonyms of earlier names (including some that are both synonyms and homonyms); six are homonyms of earlier names; ten are *nomina oblita* (six so designated herein); and 98 are considered to be unidentifiable and are listed as *nomina dubia*. The remaining 37 taxa are considered valid, all but three of which are in general use. These three are *Voluta pattersonia*, *Triplex rosarius*, and *Tellina rosea*. They are discussed below. Also listed are five taxa in the *Conchology* named in molluscan genera but are shown to be non-molluscan.

Conventions used in the following lists:

Arcana. Neither the text pages nor plates are numbered. The pages are therefore paginated by signature letter and a number indicating position in signature. Plate numbers of the *Arcana* are in Roman numerals following Mathews & Iredale (1912). It was found that parts 1 and 21 were incorrectly arranged by Mathews & Iredale and the resulting rearrangement affects species on plates III and LXXXII, formerly plates II and LXXXIV. Names often appear on the plates in the *Arcana*.



The plates are dated and all but three were issued on the engraved date. Publication date is given for each name.

Conchology. Names do not appear on the plates but are given, together with a short description or comments, on a facing page of unnumbered letterpress. Nomina are therefore referred to by plate and figure number. In some cases where there is more than one genus on a plate each genus has its own numbering so that the same number can appear two or three times on a plate. However, the way the text is laid out opposite the plate there can be no confusion. All names in the *Conchology* date from April 1, 1811.

Names attributable to Perry and in current usage are in **bold type**.

Genus-group names

Aculea—Arcana, signature H₅ (Pl. XV), April 1, 1810.

Suppressed by ICZN Opinion 1677 (1992); = *Haustator* Montfort, 1810.

Aquaria—Conchology, Pl. 52.

Type species, by subsequent designation (L.A. Smith, 1962), *Aquaria radiata* Perry, 1811. Subjective synonym of *Penicillus* Bruguière, 1789.

Aranea—Arcana, signature Aa₅ (Pl. XLVII), December 1, 1810.

Type species, by monotypy, *Aranea gracilis* Perry, 1810 (= *Murex pecten* [Lightfoot, 1786]). A junior homonym of *Aranea* Linnaeus, 1758 (Arachn.) and therefore unavailable. A synonym of *Murex s.s.* fide Vokes (1964: 4).

Biplex—*Arcana*, signature M₇ [text to Pl. XXIII; genus without species], June, 1810; Conchology, Pls. 4, 5.

Type species, by subsequent designation (Gray, 1847: 133), *Biplex perca* Perry, 1811. Used as a subgenus of *Gyrineum* Link, 1807 by Beu (1971: 101).

Buccinella—Conchology, Pl. 27.

Type species, by subsequent designation (Abbott, 1950: 203), *Buccinella coerulea* Perry, 1811 [= *Turbinella pyrum* (Linnaeus, 1767)]. For discussion of this type designation and prior treatment see Petit & Harasewych (1990: 4).

Cassidea—Conchology, Pl. 34.

Type species never designated. A junior homonym of *Cassidea* Bruguière, 1789 and therefore unavailable. Includes species now placed in *Phalium* and *Melongena*.

Columna—Conchology, Pl. 51.

Type species, here designated, *Columna marmorea* Perry, 1811, considered to be a junior synonym of *Buccinum columna* Müller, 1774. Treated as genus by Thiele (1931: 560), type not mentioned. Zilch (1959: 372) shows as type, by monotypy, *C. columna* (O.F. Müller) but that is not correct as Perry had two species in his new genus, neither identified as *C. columna*. Gray (1847:177) also cited *Helix columna* as

type. Perry figured and described two species, *C. grisea* and *C. marmorea*. Details on these two species will be found under the species listings. In the evident absence of a valid type species designation one is made here.

Cymbium—Conchology, Pl. 37.

Type species never designated. A junior homonym of *Cymbium* Röding, 1798 and therefore unavailable. Of the three species figured under this genus, two are species of *Melo* and the other is a very stylized, unidentifiable shell, possibly intended to be an *Oliva*.

Dentalia—Conchology, Pl. 52.

Type species, by subsequent designation (Emerson, 1952: 298), *Dentalia viridis* Perry, 1811 (= *Dentalium elephantinum* Linnaeus, 1758). Although treated as a valid name by Emerson it is an emendation of *Dentalium* as Perry indicated on his "Index to the plates and genera" that it was one of the genera adopted from Linnaeus.

Distorta—Conchology, Pl. 10.

Type species, by subsequent designation (Emerson & Puffer, 1953: 96), *Distorta rotunda* Perry, 1811 [= *Distorsio anus* (Linnaeus, 1758)]. Subjective synonym of *Distorsio* Röding, 1798.

Haustrum—Conchology, Pl. 44.

Type species, by subsequent designation (Iredale, 1915: 474), *Haustrum zealandicum* Perry, 1811; as *Buccinum haustrum* Martyn, 1784 (non-binominal), = *Buccinum haustrum* Gmelin, 1791.

Hexaplex—Arcana, signature M₇ [text to Pl. XXIII; genus without species], June, 1810; Conchology, Pl. 8.

Type species, by subsequent designation (Iredale, 1915: 469), *Hexaplex foliacea* Perry, 1811 (= *H. cichoreus* (Gmelin, 1791)); = *Murex cichoreum* Gmelin, 1791. Type species designation often attributed to Jousseaume (1880: 335), but he designated *Murex cichoreum* Gmelin, 1791, not an originally included species.

Monoplex—*Arcana*, signature M₇ [text to Pl. XXIII; genus without species], June, 1810; *Conchology*, Pl. 9.

Type species, by subsequent designation (Dall, 1904: 138), *Monoplex australasiae* Perry, 1811 (= Cymatium parthenopeum (Salis Marschlins, 1793)).

Nerites—Conchology, Pl. 34.

= *Nerita* Linnaeus, 1758. Listed by Sherborn (1928: 4297) and Neave (1940, III: 319), but not by Thiele (1929–1935). Two species were described by Perry in this genus, neither identifiable as to species but the figures are obviously intended to be of *Nerita*. On his "Index to the plates and genera" Perry indicates that this is a Linnaean genus and *Nerites* is therefore considered to be an unjustified emendation.

Pinus—Arcana Pl. LXXXII, September 1, 1811

Listed as a genus by Sherborn (1928: 4985) and by Neave (1940, III: 765). Shown to be a species-group name by Petit & Le Renard, 1990: 32, q.v.

Planorbis—Conchology, Pl. 51.

Type species never designated. A junior homonym of *Planorbis* Müller, 1774 and therefore unavailable. Perry indicated that *Planorbis* was a new name, but that was

377

possibly an error on the index page. As the three shells figured under this genus are unidentifiable it cannot be determined if the two names are synonyms.

Polyplex—Arcana, signature M₇ [text to plate XXIII; genus without species], June 1, 1810; *Conchology*, Pl. 9.

Type species, by subsequent designation (I.C.Z.N. 1970, Opinion 911), Polyplex bulbosa Perry, 1811 [= Trophon geversianus (Pallas, 1774)]; = Trophon Montfort, 1810. In a petition to the I.C.Z.N. Keen (1964b) addressed the then recent designation of a type species for *Polyplex Perry*. The species selected, *P. purpurascens*, had been stated to be the mollusk from which dye was made thus fixing it as a synonym of Murex trunculus Linnaeus, 1758. Keen did not approve of the designation for three reasons. Her first objection was that the species was not the only one from which dye was made and her second was that the designation would displace *Trunculariopsis* Cossmann, 1921, then in use sparingly as a subgenus. Her third objection was that, of the species first placed in *Polyplex* by Perry which had previously appeared as a "genus without species," P. purpurascens was not from the locality given when the genus name was first mentioned. Keen, after discussing other alternatives, then declared that "there is another of the five species figured by Perry that seems to be identifiable, although I do not find that anyone has made this suggestion: The figure of P. bulbosa, as well as the description, can be matched very satisfactorily by specimens of Trophon geversianus (Pallas, 1774), the type of the genus *Trophon*." She obviously overlooked Deshayes (1843: 590) and Martens (1872: 8) who had identified Perry's very good figure. Keen asked that the earlier type designation be voided and that *Polyplex bulbosa* Perry be declared the type species of *Polyplex*. Her tenuous reasons were not questioned by the Commission and her petition was approved. For additional discussion regarding history of this name see Vokes (1964: 12).

Pomacea—Arcana, signature G₅ (text to Pl. XII), March 1, 1810.

Type species, by monotypy, *Pomacea maculata* Perry, 1810. For discussion, see Pain (1956: 79), Cowie (1999) and I.C.Z.N. (1999a, Opinion 1913).

Proscenula—*Arcana*, signature B₅ [genus without species], January 1, 1810; *Conchology*, Pl. 53.

Genus first described in *Arcana*, B₅, Jan. 1, 1810, but without species. Férussac, 1820, cited as *Proxenula* in error fide Neave (1940, III: 924, 949; Férussac not seen). Thiele (1931: 738) lists as: "*Proscenula* (Perry) oder *Proxenula* Férussac 1820 ist synonym mit *Crepidula*" [*Proscenula* (Perry) or *Proxenula* Férussac 1820 is synonymous with *Crepidula*]. None of the three species described in this genus by Perry are identifiable to species, but are obviously species of *Crepidula* Lamarck, 1799. No type designation has been made for this genus as no included species are identifiable.

Septa—Arcana, signature B₂ (text to Plate III), January, 1810.

Type species, by monotypy, *Septa scarlatina* Perry, 1810 [= *Cymatium (Septa) rubecula* (Linnaeus, 1758)].

Strigula—Conchology, Pl. 15.

22

Type species, here designated, *Strigula maculata* Perry, 1811 [= *Pythia scarabaeus* (Linnaeus, 1758)]. Listed as a synonym of *Pythia* Röding, 1798, by Thiele (1931: 468)

and Zilch (1959: 72). No prior type designations have been located.

Trigona—Conchology, Pl. 51

Type species, by monotypy, *Trigona pellucida* Perry, 1811 [= *Trigonostoma scalare* (Gmelin, 1791)]. A junior homonym of *Trigona* Jurine, 1807 (Hymenoptera fide Neave, 1940 [Jurine not seen]) and therefore unavailable. A synonym of *Trigonostoma* Blainville, 1827.

Triplex—Arcana, signature M₇ (text to Pl. XXIII), June, 1810.

Type species, by monotypy, *Triplex foliatus* Perry, 1810. Used as a subgenus of Chicoreus by Houart, 1992. In a petition having to do with the identity of the type species of the genus Chicoreus Montfort, 1810 (May 28), which is based on a woodcut that is barely identifiable as a mollusk, Keen (1964b) properly showed that *Chicoreus* "has clear priority" over Triplex Perry, 1810 (June 1). She then stated that "the type species of Triplex is T. foliatus, better known under the latter [sic] name Murex palmarosae Lamarck, 1822. Preservation of the latter and suppression of T. foliatus as a nomen oblitum would be desirable." Why it would be desirable is not stated as Perry's name not only has twelve years priority, but it is based on an excellent illustration that was probably never seen by Keen. Not mentioned by Keen is the fact that this synonymy had been noted by Deshayes (1843: 572) and Mathews & Iredale (1912: 11). Also, two months before Keen's petition was published, Vokes (1964: 8) discussed this synonymy at length and ended her discussion with the statement that "the name M. palmarosae must be replaced by T. foliatus Perry." Keen's petition had so many items that caused prolonged debate, as shown in the published Opinion 911 (I.C.Z.N., 1970: 20–26), that the request to sink Triplex foliatus was uncontested although other items were debated. It is unfortunate that Keen did not know that eventually Triplex would be used as a subgenus of *Chicoreus*. Article 81.2.1 (I.C.Z.N., 1999b) allows a nomen oblitum to be the type species of a genus, but it is certainly not a desirable situation. See Plate 1 herein.

Volutella—*Arcana*, signature B₂ (text to Pl. III), January 1, 1810.

Type species, by monotypy, *Volutella divergens* Perry, 1810 (= *Voluta muricata* Born, 1778); = *Vasum* Röding, 1798.

Species-group names

abbatis (Mitra)—Conchology, Pl. 39, figs. 2, 3. = Mitra stictica (Link, 1807) fide Cerhohorsky (1967b: 148). abortiva (Triplex)—Conchology, Pl. 6, fig. 5.

Nomen dubium fide Houart, 1992: 13. Cernohorsky (1967a: 119) placed Perry's name in the synonymy of *Chicoreus carneolus* (Röding, 1798) together with *C. torrefactus* (Sowerby, 1841). Vokes (1970: 185), while admitting that Perry's figure might possibly be identified as *C. torrefactus*, objected to the introduction of either *C. carneolus* or *C. abortiva* as she considered them to be *nomina oblita*. Houart (1992:13) treated both of these names as *nomina dubia*, an assessment here considered to be correct.

377

achatina (Cypraea)—Conchology, Pl. 20, fig. 6.

= Cypraea ventriculus Lamarck, 1810, fide Martens (1872: 11).

aculeata (Aranea)—Conchology, Pl. 46, fig. 2.

Nomen dubium. This name has never been used as a senior synonym. The figure and a prior mention of the name are discussed by Vokes (1970:183). She there considered the figure to probably represent *Murex carbonnieri* (Jousseaume, 1881). Later Ponder & Vokes (1988:23) declared that "Perry's figure is so poor that accurate identification is impossible and this name is best regarded as a *nomen nudum*." As discussed herein, the name cannot be a *nomen nudum*. Perry's figure is here considered to be unidentifiable and must remain a *nomen dubium*.

aculeatus (Strombus)—Conchology, Pl. 13, fig. 2.

= Lambis crocata (Link, 1807) fide Abbott (1961: 158, 159) who adds that it is "perhaps a nomen dubium". Perry's figure is quite adequate for identification and the comment by Abbott was because it could possibly represent the subspecies L. crocata pilsbryi Abbott being described at the time [as stated by Abbott (1961: 159)].

acuminatus (Trochus)—Conchology, Pl. 47, fig. 1.

= Cantharidus opalus (Martyn, 1784). Placed in synonymy of Trochus iris Gmelin, 1791 (= C. opalus (Martyn)) by Martens (1872: 18). Iredale (1915: 437) stated: "This synonym of Cantharidus opalus Martyn, 1784, is not included by Suter. In Perry's Conchology ... an easily recognizable figure is given." Placed in synonymy of C. opalus (Martyn, 1784) by Marshall (1998: 112) and a neotype designated making it a junior objective synonym of C. opalus (Martyn, 1784).

acuta (Buccinella)—Conchology, Pl. 27, fig. 6.

Nomen dubium. This name has not appeared in subsequent literature and the figure is unidentifiable. Based on the enlarged posterior columellar fold and "knobby" spire it may be a very poor rendering of *Cancellaria cassidiformis* Sowerby, 1832 but it certainly cannot be so identified.

acuta (Distorta)—Conchology, Pl. 10, Distorta fig. 1.

= Distorsio reticularis (Linnaeus, 1758) fide Beu (1998: 195).

acuta (Melania)—Conchology, Pl. 29, fig. 2.

=Lymnaea stagnalis (Linnaeus, 1758) fide Martens (1872: 5). Perry stated that his species had been illustrated by Born and comparison shows that Perry's figure could be based on Born's figure (1780: pl. 16, fig. 16).

acuta (Pleurotoma)—Conchology, Pl. 54, fig. 5.

= Lophiotoma acuta (Perry, 1811) fide Powell (1964: 303).

acutus (Strombus)—Conchology, Pl. 12, fig. 2.

=Strombus vomer (Röding, 1798) fide Abbott (1960: 131). Very much like the Chemnitz figure with which Deshayes (1843: 716) and Martens (1872: 20) equated it.

adunca (Patella)—Conchology, Pl. 43, fig. 5.

Nomen dubium. Pilsbry (1891: 118) placed with a query under Nacella aenea var. deaurata Gmelin. In family Fissurellidae fide Powell (1973: 84). The figure is here considered to be unidentifiable.

aladraconis (Rostellaria)—Conchology, Pl. 11, fig. 6, (as ala-draconis).

- = *Aporrhais pespelecani* (Linnaeus, 1758). This synonymy has not previously been noted. The unusual blue coloration is possibly the result of burial in mud.
- alba (Ancilla)—Conchology, Pl. 31, fig. 1.
 - = Bullia vittata (Linnaeus, 1767) fide Cernohorsky (1984: 30).
- alba (Cassis)—Conchology, Pl. 33, fig. 2.
 - = Cassis flammea (Linnaeus, 1758) fide Deshayes (1844: 23).
- alba (Ovula)—Conchology, Pl. 53, Ovula fig. 1.
 - = Ovula ovum (Linnaeus, 1758) fide Deshayes (1844: 467).
- albida (Pleurotoma)—Conchology, Pl. 32, fig. 4.
 - = *Polystira albida* (Perry, 1811) fide Abbott & Dance (1982: 239). Placed in the synonymy of the later *Pleurotoma virgo* Lamarck by Deshayes (1843: 350) who would not, as mentioned elsewhere, surrender a Lamarck name. Type species, by original designation, of *Polystira* Woodring, 1928.
- alga (Cypraea)—Conchology, Pl. 23, fig. 1.
 - =*Cypraea mappa alga* Perry, 1811, fide Schilder & Schilder (1971: 33) and Cernohorsky (1978: 55). Some later authors (e.g., Abbott & Dance, 1982: 96) consider this to be only a synonym of *C. mappa* Linnaeus, 1758.
- altus (Trochus)—Conchology, Pl. 47, fig. 3.
 - = *Tectus conus* (Gmelin, 1791), fide OBIS Indo-Pacific Molluscan Database (OBIS, 2003). A senior primary homonym of *Trochus altus* Philippi, 1851 which is also a synonym of *Tectus conus* (Gmelin) fide OBIS (2003).
- anatomica (Hexaplex)—Conchology, Pl. 8, fig. 2.
 - = Homalocantha anatomica (Perry, 1811) fide Abbott & Dance (1982: 139).
- angulata (Aculea)—Arcana, signature H₅, Pl. XV, fig. 2, April 1, 1810.
 - = Turritella (Haustator) imbricataria Lamarck, 1804, fide Petit & Le Renard, 1990: 28.
- angulatus (Conus)—Arcana, signature H₅, Pl. XV, fig. 1, April 1, 1810.
 - = Conus deperditus Bruguière, 1792 fide Kohn (1986: 4).
- angulatum (Solen)—Conchology, Pl. 57, Solen fig. 2.
 - *Nomen dubium.* This name has not appeared in subsequent literature and the figure is unidentifiable.
- annularis (Cypraea)—Conchology, Pl. 22, fig. 6.
 - = *Cypraea annulus* Linnaeus, 1758 fide Deshayes (1844: 539).
- annularis (Pomacea)—Conchology, Pl. 38, fig. 1.
 - Nomen dubium. Not a Pomacea but is a helicid fide Pilsbry (1927: 63).
- apiaria (Trochus)—Arcana, signature B₃, Pl. III, fig. 4, January 1, 1810; Conchology, Pl. 47, fig. 5.
 - = *Tectus fenestratus* (Gmelin, 1791). This synonymy has not previously been noted but the short green spiral segments arranged in axial rows is distinctive. Both Perry figures match color photographs of *T. fenestratus*. Incorrectly listed as a synonym of *Trochus conus* (Gmelin, 1791) on OBIS Indo-Pacific Molluscan Database (OBIS, 2003).
- arabica (Cypraea)—Conchology, Pl. 21, fig. 1.



377

= Cypraea arabica Linnaeus, 1758.

argillaceous (Conus)—Conchology, Pl. 24, fig. 6.

Nomen dubium fide Kohn (1986: 6; as argillaceus [sic]).

argus (Cypraea)—Conchology, Pl. 20, fig. 7.

= Cypraea argus Linnaeus, 1758.

asellus (Cypraea)—Conchology, Pl. 19, fig. 3.

= Cypraea asellus Linnaeus, 1758.

aspera (Ovula)—Conchology, Pl. 53, Ovula fig. 3.

= Volva volva (Linnaeus, 1758) fide Deshayes (1844: 475).

aurantia (Melania)—Conchology, Pl. 29, fig. 1.

= *Plekocheilus aurissileni* (Born, 1778) fide Martens (1872: 26) [as *Bulimus auris Sileni*]. One of Born's figures (1780, pl. 9, fig. 4), slightly reoriented, may have served as the model for Perry's figure. Born's two figures were copied by Chemnitz (1795: pl. 176, figs. 1701, 1702). However, Perry states that his figure is "From a delineation of a specimen in Mr. Jennings's Museum."

aurantia (Murex)—Conchology, Pl. 1, fig. 1.

= Latrius gibbulus (Gmelin, 1791) fide Adam & Leloup (1938: 186).

aurantia (Nerites)—Conchology, Pl. 34, Nerites fig. 1.

Nomen dubium. Listed by Martens (1889: 135) in an index as "*Neritina pulligera*" but the figure is not close to that species fide Eichhorst (2003: pers. comm.). Aside from the listing by Martens this name has not appeared in the literature.

aurantia (Turbo)—Conchology, Pl. 49, fig. 4.

= Turbo petholatus Linnaeus, 1758. Perry's figure is listed with a query in synonymy of Turbo personatus Linnaeus by Mörch (1852: 162). It is hard to follow Hanley (1855: 328) but he appears to consider T. personatus the same as T. variabilis Reeve, 1842. The latter name is preoccupied and equals T. reevei [sic] Philippi, 1846 fide Tryon (1888: 194) as a variety of T. petholatus. Turbo reevei [sic] Philippi was listed as a full species by Abbott & Dance (1982:46). The name T. personatus seems to have disappeared from the literature, Philippi (1846: 48) stating Linnaeus' species to be entirely unknown to him (although it is based on a very good figure in Rumphius). The specific name of Philippi's taxon T. reevii is almost universally misspelled. Perry's species is here considered to be a synonym of Turbo petholatus Linnaeus, 1758 because of its shape and markings, the perceived differences between T. petholatus and T. reevii being too subtle to be recognized in Perry's figure.

aurantia (Voluta)—Conchology, Pl. 18, no. 2.

= *Harpulina arausiaca* [Lightfoot, 1786] fide Deshayes (1844: 402; as *Voluta vexillum* Chemnitz). The only other citation located is that of Wagner & Abbott (1978: 20-703) who cited as a nude name, an inappropriate term. Not *Voluta aurantia* Gmelin, 1791 which is in Mitridae.

aurea (Tellina)—Conchology, Pl. 55, Tellina fig. 2.

= Tellina (Pharaonella) aurea Perry, 1811. Hedley (1902: 28) stated: "...evidently T. vulsella Chemnitz, = T. rostrata of other authors than Linnaeus, not T. (Phylloda) aurea Schumacher (1817). Since by Hanley's [1855] showing ... the Linnaean name

has been generally misapplied, and that of Chemnitz was not binomial, Perry's name should be utilised." Hedley was unaware that this species had already been named *Tellina vulsella* Hanley, 1846 but his admonition that Perry's name should be utilized remains true. As Perry's name was treated as valid by Hedley after 1899, and has been used in more recent years (e.g. Kira, 1964: 158; Matsukuma et al, 1991:187) it cannot be dismissed as a *nomen oblitum* even if Hanley's name can be demonstrated to have been used in 25 works in the last 50 years which is very doubtful. Hedley's mention of *Tellina aurea* Schumacher is in error as that name was originally proposed in the genus *Phylloda*, is a junior objective synonym of *Phylloda foliacea* (Linnaeus, 1758), and has no place in this discussion.

australasia (Biplex)—Conchology, Pl. 4, figs. 2, 4.

= Ranella australasia (Perry, 1811) fide Henning & Hemmen (1993: 34). Type species of Mayena Iredale, 1917 by subsequent designation of Powell, 1937.

australasia (Pyrula)—Conchology, Pl. 54, fig. 4.

= *Pleuroploca australasia* (Perry, 1811) fide Wilson & Gillett (1971: 98). *australasiae* (*Monoplex*)—*Conchology*, Pl. 3, fig. 3.

= Cymatium (Monoplex) parthenopeum (Salis Marschlins, 1793) fide Beu (1970b: 229). Type species of Monoplex Perry, 1811 by subsequent designation of Dall, 1904. autumnalis (Cypraea)—Conchology, Pl. 21, fig. 2.

= Cypraea mus Linnaeus, 1758, fide Deshayes (1844: 500).

babylonica (Pleurotoma)—Conchology, Pl. 32, no. 5.

= *Turris babylonia* (Linnaeus, 1758), fide Deshayes (1843: 351). Perry stated that this species had "formerly been placed by Linnaeus under the genus *Murex*" so it is obviously a spelling error. This is the only species figured on two different plates in the *Conchology* (under the same species name) demonstrating that the plates were not all prepared at one time. Perry had already figured this species as *Murex babylonicus* on Pl. 2. Not listed by Sherborn so he evidently also considered this to be an error for *babylonia* Linnaeus.

babylonicus (Murex)—Conchology, Pl. 2, fig. 2.

=Turris babylonia (Linnaeus, 1758), fide Deshayes (1843: 351). See under *Pleurotoma* babylonica above.

bandarius (Murex)—Conchology, Pl. 1, fig. 5

= Pugilina morio (Linnaeus, 1758), fide Deshayes (1843: 451).

bandata (Cassidea)—Conchology, Pl. 34, Cassidea fig. 2.

= Phalium bandatum (Perry, 1811) fide Abbott (1968a: 83).

bandata (Cypraea)—Conchology, Pl. 20, fig. 2.

= Cypraea caputserpentis Linnaeus, 1758, fide Adam & Leloup (1938: 134).

bandata (Dentalia)—Conchology, Pl. 52, Dentalia fig. 4.

Nomen dubium. This name has not appeared in subsequent literature. It is a stylized scaphopod unidentifiable to species.

bandatum (Cerithium)—Conchology, Pl. 36, fig. 4.

= Rhinoclavis fasciata (Bruguière, 1792) fide Houbrick (1978: 41–47).

bandatum (Cymbium)—Conchology, Pl. 37, Cymbium fig. 2 (No. 4 on plate explanation).



Nomen dubium. This name has not appeared in subsequent literature. It is a stylized *Melo* that cannot be specifically identified.

bandatus (Conus)—Conchology, Pl. 25, fig. 4; Arcana, signature Pp₇, Pl. LXXX, August, 1811

Nomen dubium fide Kohn (1986: 7). The figures in the two works are different but neither is identifiable.

bandatus (Murex)—Conchology, Pl. 1, fig. 4.

= Pugilina morio (Linnaeus, 1758) fide Deshayes (1843: 451).

bicolor (Trochus)—Conchology, Pl. 47, fig. 4.

= *Trochus radiatus* Gmelin, 1791. This synonymy has not previously been noted but the axial bands of color and the shells' overall shape agree perfectly with Gmelin's species. Perry's name has not appeared in subsequent literature.

brachiatus (Turbo)—Conchology, Pl. 49, fig. 2.

= *Turbo cornutus* [Lightfoot, 1786]. This synonymy has not previously been noted although the spiral sculpture and prominant spines make Perry's figure easily identifiable. Perry's name has not appeared in subsequent literature.

brunella (Scalaria)—Conchology, Pl. 28, fig. 2.

Nomen dubium. This name has not appeared in subsequent literature. It is an unidentifiable epitoniid.

bulbosa (Polyplex)—Conchology, Pl. 9, fig. 5.

= Trophon geversianus (Pallas, 1774) fide Deshayes (1843: 590; as Murex magellanicus Lamarck [sic; = Trophon magellanicus (Gmelin, 1791)]. Type species of Polyplex Perry, 1811. See under Polyplex above and also see Polyplex gracilis Perry below.

caerulea—see under coerulea and also comments under "Perry's Taxa" above.

camelopardalis (Cypraea)—Conchology, Pl. 19, fig. 5.

- = *Cypraea* (*Lyncina*) *camelopardalis* Perry, 1811 fide Schilder & Schilder (1971: 39). *canulata* (*Buccinella*)—*Conchology*, Pl. 27, fig. 1.
 - = Cancellaria reticulata (Linnaeus, 1767) fide Deshayes (1843: 402).

capitata (Voluta)—Conchology, Pl. 17, fig. 4.

= Fulgoraria rupestris (Gmelin, 1791) fide Weaver & du Pont (1970: 35).

capitatus (Monoplex)—Conchology, Pl. 3, fig. 4.

= *Tudicla spirillus* (Linnaeus, 1767) fide Deshayes (1843: 513; as *Pyrula spirillus* Lamarck).

caputserpentis (Cypraea)—Conchology, Pl. 21, fig. 4 (as caput-serpentis).

= Cypraea caputserpentis Linnaeus, 1758.

carinata (Buccinella)—Conchology, Pl. 27, fig. 5.

Nomen dubium. This name has not been used in subsequent literature but it is probably Latirolagena smaragdula (Linnaeus, 1758). Although there is extra emphasis on the columellar folds the closely spaced brown spirals on a white shell make this identification reasonable. As its identification is not positive it must be considered a nomen dubium.

carinatum (Cerithium)—Conchology, Pl. 35, fig. 3.

= Terebralia palustris (Linnaeus, 1767) fide Deshayes (1843: 284) and Houbrick

(1991: 305; as carnaticum [sic]).

ZOOTAXA 377

- carinatus (Bulimus)—Conchology, Pl. 30, fig. 1.
 - = Phasianotrochus eximius (Perry, 1811). For discussions see Pilsbry (1901: 8; 1902: 72), Gatliff (1902: 76), Pritchard & Gatliff (1906: 65), and Hedley (1908: 465). Not Bulimus carinatus Bruguière, 1789.
- carnatis (Melania)—Conchology, Pl. 29, fig. 3.
 - = *Megalobulimus oblongus* (Müller, 1774). Although placed in synonymy of *Stro-phocheilus oblongus* (Müller, 1774) with a query by Bequaert (1948: 67), there is little question about this synonymy.
- carnea (Terebra)—Conchology, Pl. 16, Terebra fig. 1.
 - = Terebra dimidiata (Linnaeus, 1758) fide Deshayes (1844: 241).
- cedonulli (Conus)—Conchology, Pl. 24, fig. 1 (as cedo nulli).
 - = Conus cedonulli Linnaeus, 1767.
- chalcedonia (Cypraea)—Conchology, Pl. 19, fig. 6.
 - = *Cypraea helvola* Linnaeus, 1758, fide Deshayes (1844: 533).
- cicercula (Cypraea)—Conchology, Pl. 23, fig. 7.
 - = Cypraea cicercula Linnaeus, 1758.
- cincta (Helix)—Conchology, Pl. 15, Helix fig. 3.

Nomen dubium. Figure not identifiable but possibly an embellished rendering of a form of *Polymita picta* (Born, 1778). Not *Helix cincta* Müller, 1774.

- cinctoria (Bulla)—Conchology, Pl. 40, Bulla fig. 1.
- = Hydatina zonata [Lightfoot, 1786] fide Cernohorsky (1972: 206) who, under H. albocincta (Hoeven, 1839), stated: "Iredale and McMichael (1962) consider H. cinctoria (Perry, 1811) to be an earlier name for albocincta, but I share Pilsbry's view that Perry's figure probably represents the Indian Ocean species zonata Lightfoot, 1786...." cinerea (Aranea)—Conchology, Pl. 46, fig. 4.
 - = *Bolinus brandaris* (Linnaeus, 1758) fide Adam & Leloup (1938: 157). Listed without comment by Vokes (1971: 32) but misspelled as *cinera*.
- circularis (Chama)—Conchology, Pl. 59, Chama fig. 2.

Nomen dubium. This figure is not identifiable and the name has not appeared in subsequent literature.

- circularis (Natica)—Conchology, Pl. 48, Natica fig. 2.
 - *Nomen dubium.* This figure is not identifiable and the name has not appeared in subsequent literature.
- clathrus (Scalaria)—Conchology, Pl. 28, fig. 3.
 - = Epitonium clathrus (Linnaeus, 1758).
- coerulea (Ancilla)—Conchology, Pl. 31, fig. 6.

Nomen dubium. This name has appeared in subsequent literature only when it was stated to be a *Nassarius* by Burch & Burch (1960: 8; as *caerulea*) in their catalogue of Olividae. Not listed by Cernohorsky (1984) in his monograph on Nassariidae. This figure is not identifiable.

- coerulea (Buccinella)—Conchology, Pl. 27, fig. 4.
 - = Turbinella pyrum (Linnaeus, 1767) fide Abbott (1950: 203; as type species of Buc-

377)

cinella Perry, 1811; as caerulea). This is a slightly altered copy of Martini's figure (1777, pl. XCV, fig. 916) that is also blue. Perry states: "Shell of a sky blue color ... described, but not well figured, by the German Conchologist Martini. The original is in the noble collection of the British Museum." Abbott's reference to Winckworth (1945: 145) for the identification of this species is enigmatic as no Perry species are mentioned in the reference cited.

coerulea (Cypraea)—Conchology, Pl. 22, fig. 8.

= *Cypraea annulus* Linnaeus, 1758, fide Deshayes (1844: 539).

coerulea (Patella)—Conchology, Pl. 43, fig. 3.

= *Patella caerulea* Linnaeus, 1758. This is the only *Patella* for which there is no description given and is probably intended for Linnaeus's species.

coerulea (Pyrula)—Conchology, Pl. 50, fig. 2.

Nomen dubium. Stated to be *Fasciolaria tulipa* (Linnaeus, 1758) by Deshayes (1843: 432) but it could just as easily be *F. hunteria* (Perry, 1811) which has a bluish cast more often than does *F. tulipa*.

collapsus (Planorbis)—Conchology, Pl. 51, fig. 5.

Nomen dubium. This figure is not identifiable and the name has not appeared in subsequent literature.

colubrina (Helix)—Conchology, Pl. 15, Helix fig. 4.

Nomen dubium. This figure is not identifiable and the name has not appeared in subsequent literature.

colus (Murex)—Conchology, Pl. 2, fig. 5.

= Fusinus colus (Linnaeus, 1758).

comma (Cypraea)—Conchology, Pl. 21, fig. 5.

= Cribrarula cribraria comma (Perry, 1811) fide Lorenz & Hubert (2000: 177).

concavum (Pecten)—Conchology, Pl. 55, Pecten fig. 1.

Nomen dubium. This figure is not identifiable and the name has not appeared in subsequent literature.

conspicua (Aranea)—Arcana, Pl. LIV (Feb. 1, 1811); Conchology, Pl. 46, fig. 3.

= Bolinus cornutus (Linnaeus, 1758) fide Vokes (1971: 35).

contorta (Pleurotoma)—Conchology, Pl. 32, fig. 1.

= Turricula javana (Linnaeus, 1758) fide Deshayes (1843: 354; as *Pleurotoma nodifera* Lamarck) and Powell (1969: 236).

contrastriata (Cypraea)—Conchology, Pl. 20, fig. 3.

= Lyncina argus contrastriata (Perry, 1811) fide Lorenz & Hubert (2000: 77).

coralloides (Conus)—Conchology, Pl. 25, fig. 6.

Nomen dubium fide Kohn (1986: 7).

cornucervi (Triplex)—Conchology, Pl. 7, fig. 4 (as cornu-cervi).

= Chicoreus banksii (Sowerby, 1841) fide Houart (1992: 69). Earlier considered to be

C. damicornis Hedley, 1903 by Vokes (1971: 36). Not *C. cornucervi* (Röding, 1798). *cornuta* (*Cassidea*)—*Conchology*, Pl. 34, *Cassidea* fig. 3.

= Melongena melongena (Linnaeus, 1758). Figure copied from Regenfuss (1758, pl.

5, fig. 49). Not a synonym of M. corona (Gmelin, 1791) as stated by Abbott (1968a:

30

48). A junior primary homonym of *Cassidea cornuta* Bruguière, 1792. *cornuta* (*Triplex*)—*Conchology*, Pl. 7, fig. 3.

Nomen oblitum. Vokes (1971: 36) listed as "? = Chicoreus brevifrons (Lamarck, 1822)" and Roland Houart (personal communication, July, 23, 2003) agrees that it may be that species. Triplex cornuta Perry, 1811 is therefore a senior subjective synonym of Murex brevifrons Lamarck, 1822 now placed in the genus Chicoreus. To insure that Perry's name cannot replace the later well known synonym, action is here taken in accordance with Article 23.9.2. An exhaustive search of the literature reveals no usages of Triplex cornuta Perry, 1811 as a senior synonym. Listed here are 25 usages of Chicoreus brevifrons (Lamarck) during the past 50 years: Abbott, 1974: 174; Abbott & Dance, 1982: 135; Abbott & Morris, 1995: 209; Altena, 1969: 20, 43; Altena, 1975: 42; Bayer, 1971: 157; Bullis, 1964: 105; Dance, 1974: 123; Eisenberg, 1981: 88; Fair, 1976: 27; Habe & Okutani, 1985: 139; Kaicher, 1973: 131; Macsotay & Campos Villarroel, 2001: 74; Usticke, 1959: 62; Radwin & D'Attilio, 1976: 34; Rios, 1970: 78; Rios, 1975: 85; Rios, 1985: 82; Rios, 1994: 108; Vokes, 1989: 33; Vokes, 1990b: 34; Vokes, 1995: 119; Vokes, 1996: 67; Wagner & Abbott, 1978: 13-802; Warmke & Abbott, 1961: 105. Triplex cornuta Perry, 1811 is declared to be a nomen oblitum and Murex brevifrons Lamarck, 1822 becomes a nomen protectum.

cornutus (Monoplex)—Conchology, Pl. 3, fig. 1.

Nomen dubium. Hedley (1902: 26) considered this to be a senior synonym of *Triton exaratus* Reeve, 1844, but Pritchard & Gatliff (1906: 42) disagreed, citing Dall's (1904: 123) statement that Perry's species is "unidentifiable, certainly not *exaratum* Reeve, to which it has been referred." Kesteven (1902: 460) treated a species, for which he figured only the protoconch, as *Lotorium cornutum* (Perry). This Kesteven reference is included by Kilias (1973: 96) in his synonymy of *Cymatium exaratum* (Reeve) but he does not mention Perry's species elsewhere in his monograph of Cymatiidae. Not listed in the synonymy of *C. (M.) exaratum* (Reeve, 1844) by Beu (1998: 89–90). Perry's figure is here considered to be unidentifiable.

cornutus (Strombus)—Conchology, Pl. 12, fig. 4.

= Strombus pugilis Linnaeus, 1758 fide Deshayes (1843: 696).

corrugata (Biplex)—Conchology, Pl. 5, fig. 1.

= Bursa corrugata (Perry, 1811) fide Abbott (1974: 167).

costata (Chama)—Conchology, Pl. 59, Chama fig. 3.

Nomen dubium. This figure is not identifiable and the name has not appeared in subsequent literature.

costatum (Cardium)—Conchology, Pl. 57, Cardium fig. 2.

= *Cardium costatum* Linnaeus, 1758. Figure possibly copied from Chemnitz, 1782, pl. 15, fig. 151.

crenata (Donax)—Conchology, Pl. 58, Donax fig. 2.

Nomen dubium. This figure is not identifiable and the name has not appeared in subsequent literature.

crenatus (Polyplex)—Conchology, Pl. 9, fig. 3.

Nomen dubium. This figure is not identifiable but is probably a species in Trophoni-



377

nae. This name has not appeared in subsequent literature.

cruentum (Cardium)—Conchology, Pl. 57, Cardium fig. 1.

= *Corculum cardissa* (Linnaeus, 1758). Perry's figure may be a poor copy of Chemnitz, 1782, pl. 14, fig. 1. Combined with his description ("heart-shaped and pointed," etc.) there is little doubt of Perry's intention.

curvirostra (Cerithium)—Conchology, Pl. 35, fig. 2.

= Cerithium nodulosum Bruguière, 1792 fide Deshayes (1843: 287).

cypridium (Patella)—Conchology, Pl. 43, fig. 6.

= Helcion pellucidus (Linnaeus, 1758) fide Powell (1973: 144).

dama (Cypraea)—Conchology, Pl. 23, fig. 3.

= *Cypraea vitellus* Linnaeus, 1758, fide Schilder & Schilder (1971: 39).

decarinatus (Trochus)—Conchology, Pl. 47, fig. 2.

=Nomen dubium. Martens (1872: 16) thought it possible that this figure was intended for *Trochus cunninghami* Gray, 1834 [=Maurea selecta (Dillwyn, 1817)] and later Martens (1873: 31) stated "appears to be the same" and that the name is intended to signify "that the whorls are separated by a keel." Placed in synonymy of *Calliostoma canaliculatum* [Lightfoot, 1786] with a query by Pilsbry (1889: 361). Perry's figure is here considered to be unidentifiable.

decussata (Orthoceras)—Conchology, Pl. 52, Orthoceras fig. 2.

=Nautilus fascia Linnaeus, 1758 which is not a Nautilus and is not a mollusk. Perry's figure is a slightly stylized copy of Martini, 1769, Vignette 1, figure D, which in turn is a copy of Gualtieri, 1742, pl. 19, figure O. Linnaeus based his N. fascia on the Gualtieri figure which had been placed in a group called Orthoceras by that non-binominal author. Orthoceras Bruguière, 1789 is a genus of Paleozoic cephalopods of considerable size, all being several inches or more in length. Both Gualtieri and Martini include a life-size drawing indicating that the specimen is only 4 or 5 mm in length. This and most other Linnaean Nautilus stated to be Foraminifera by Hanley (1855: 154).

deflecta (Pinna)—Conchology, Pl. 61, Pinna fig. 2.

Nomen dubium fide Rosewater (1961: 185).

delicata (Harpa)—Conchology, Pl. 40, Harpa fig. 2.

Placed on Official Index of Rejected and Invalid Specific Names in Zoology (I.C.Z.N., 1989, Opinion 1518). Listed as a senior synonym of *Harpa articularis* Lamarck, 1822 by Rehder (1973: 250) who further stated it to be a *nomen oblitum*. At that time the rules concerning *nomina oblita* were very confusing. In the 1985 Code (I.C.Z.N., 1985) the rules were again changed. As a result, under Article 79c(iii) of the 1985 Code his action was invalid. A petition was submitted (Rehder & Petit, 1987) to ratify his action, resulting in Opinion 1518 (I.C.Z.N., 1989).

denacta (Dentalia)—Conchology, Pl. 52, Dentalia fig. 5.

Nomen dubium. This name has not appeared in subsequent literature. It is a stylized scaphopod unidentifiable to species.

denactus (Mytilus)—Conchology, Pl. 61, Mytilus fig. 3.

Nomen dubium. This figure is not identifiable and the name has not appeared in subse-

quent literature.

ZOOTAXA 377

- dentata (Ostrea)—Conchology, Pl. 60, Ostrea fig. 1.
 - = Ostrea denticulata Born, 1778. Perry's figure is obviously a stylized copy of Born's figure (1780, pl. 6, fig. 9). The figures are almost identical in size, the smaller upper valve sits well into the lower valve, and the digitations extending from the margins of the lower valve are alike in number and spacing. This synonymy has not previously been noted.
- dentex (Haustrum)—Conchology, Pl. 44, fig. 3.

Placed on Official Index of Rejected and Invalid Specific Names in Zoology (I.C.Z.N., 1969, Opinion 886). Recognized as a synonym of *Purpura columellaris* (Lamarck, 1822), by Deshayes (1844: 62) (see discussion herein concerning Deshayes' treatment of Perry's taxa) and by Martens (1872: 43). Keen (1964a) petitioned the I.C.Z.N. for clarification of the status of the genera *Purpura*, *Ocenebra*, and *Muricanthus*. Although having nothing to do with the status of these genera, she included a request, without comment or reason, to declare as *nomina oblita* the species *Haustrum pictum* Perry, 1811 and *H. dentex* Perry, 1811, stated to be senior synonyms of *Purpura planospira* Lamarck, 1822 and *P. columellaris* Lamarck, 1822 respectively. It was not mentioned that over 100 years earlier Perry's names had been shown by Deshayes and others to be senior synonyms of the Lamarck names. The resulting Opinion was published (I.C.Z.N., 1969: 128–132) showing no dissent on the suppression of the two Perry names.

- dentula (Rostellaria)—Conchology, Pl. 10, Rostellaria fig. 3.
 - = *Tibia insulaechorab* (Röding, 1798) fide Deshayes (1843: 654; as *R. curvirostris* Lam.).
- denudata (Aranea)—Conchology, Pl. 45, fig. 1.
 - = Haustellum haustellum (Linnaeus, 1758) fide Deshayes (1843: 568).
- denudata (Triplex)—Conchology, Pl. 7, fig. 2.
 - = *Chicoreus denudatus* (Perry, 1811) fide Radwin & D'Attilio (1976: 36). Hedley (1902: 26) recognized *T. denudata* and *T. frondosa* Perry to be conspecific and recommended the use of *T. denudata*. Type species of *Torvamurex* Iredale, 1936 by original designation.
- despectum (Cerithium)—Conchology, Pl. 35, fig. 1.
 - = *Rhinoclavis vertagus* (Linnaeus, 1767) fide Houbrick (1978: 33).
- detusa (Pinna)—Conchology, Pl. 61, Pinna fig. 1.

Nomen dubium. This figure is not identifiable and the name has not appeared in subsequent literature.

- digitatum (Pecten)—Conchology, Pl. 55, Pecten fig. 2.
 - = *Manupecten pesfelis* (Linnaeus, 1758). This is an obvious copy of Born's 1780 (pl. 6, fig. 2) figure of *Ostrea elongata* Born, 1778, considered to be a form of *Manupecten pesfelis* (L.) by Rombouts (1991: 46). Mentioned only as preoccupying *Pecten digitatus* Hinds, 1845 by Hertlein (1935: 306) and by Grau (1959: 122, 123). This synonymy has not been previously noted.
- digitatus (Spondylus)—Conchology, Pl. 59, Spondylus fig. 3.

377

= *Spondylus foliaceus* Schreibers, 1793, fide Lamprell & Willan (2000: 4). It was earlier placed in the synonymy of *S. sinensis* Schreibers, 1793 by Abbott & Dance (1982: 316) and Lamprell (1986: 52). Although Schreibers work is non-binominal his names are in common usage.

digitatus (Strombus)—Conchology, Pl. 13, fig. 1.

= Lambis digitata (Perry, 1811) fide Abbott (1961: 163).

dilatum (Buccinum)—Arcana Pl. LVIII (March, 1811)

= Buccinum orbiculare Perry, 1811, q.v.

dilatus (Planorbis)—Conchology, Pl. 51, fig. 4.

Nomen dubium. This figure is not identifiable and the name has not appeared in subsequent literature.

discoloratum (Pecten)—Conchology, Pl. 55, Pecten fig. 3.

Nomen dubium. This figure is not identifiable and the name has not appeared in subsequent literature.

disjecta (Venus)—Conchology, Pl. 58, Venus fig. 3.

= Bassina disjecta (Perry, 1811) fide Hedley (1902: 28).

disjuncta (Scalaria)—Arcana, signature P₁, Pl. XXVIII (unnumbered fig. labeled Scalaria), July 1, 1810; Conchology, Pl. 28, figs. 4, 5.

= *Epitonium scalare* (Linnaeus, 1758) fide Mathews & Iredale (1912: 11). The figure in the *Arcana* is much superior to the one in the *Conchology*.

distentum (Buccinum)—Arcana, signature B₃, Pl. LXVI, May 1, 1811 (plate incorrectly dated May 1, 1810).

Nomen dubium. Possibly intended for a specimen of *Turbo petholatus* Linnaeus, 1758 fide Mathews & Iredale (1912: 13).

divaricatus (Planorbis)—Conchology, Pl. 51, fig. 3.

Nomen dubium. This figure is not identifiable and the name has not appeared in subsequent literature.

divergens (Strombus)—Arcana, signature Nn₅, Pl. LXXV, July, 1811.

= Lambis chiragra arthritica Röding, 1798 fide Abbott (1961: 173; Pl. 74 cited in error). This was named *S. divergens* on the plate and *S. nigricans* in the text.

divergens (Volutella)—Arcana, Pl. III, fig. 1, Jan. 1, 1810; Conchology, Pl. 26, fig. 3.

= *Vasum muricatum* Born, 1778, fide Abbott (1959: 15). Type species of *Volutella* Perry, 1810 by monotypy.

elegans (Arca)—Conchology, Pl. 60, Arca fig. 1.

Nomen dubium. This figure is not identifiable and the name has not appeared in subsequent literature.

elegans (Biplex)—Conchology, Pl. 5, fig. 3.

= Gyrineum natator (Röding, 1798) fide Beu (1998: 57).

elongata (Cypraea)—Conchology, Pl. 22, fig. 5.

= Erronea caurica elongata (Perry, 1811) fide Lorenz & Hubert (2000: 138).

episcopalis (Mitra)—Conchology, Pl. 39, no. 4.

= Mitra episcopalis (Linnaeus, 1758); = M. mitra (Linnaeus, 1758).

exanthemata (Cypraea)—Conchology, Pl. 22, fig. 7.

= Cypraea zebra Linnaeus, 1758, fide Schilder & Schilder (1971: 33). Possibly intended as an emendation of *C. exanthema* Linnaeus, 1767 which is now considered a synonym of *C. zebra* L.

eximius (Bulimus)—Conchology, Pl. 30, fig. 2.

= *Phasianotrochus eximius* (Perry, 1811) fide Pilsbry (1901: 8, as *Cantharidus eximus* [sic]); Wilson & Gillett (1971: 26).

fasciata (Ancilla)—Conchology, Pl. 31, Ancilla fig. 2.

Nomen dubium. Probably a buccinid but the figure is not identifiable. This name has not appeared in subsequent literature except for its citation as a homonym. Kilburn (1993: 370–371) rejected *Ancillaria fasciata* Reeve, 1864 as a secondary junior homonym of *Ancilla fasciata* Perry, 1811 as Reeve's species was being placed in the genus *Ancilla*. Kilburn's action was improper under Article 59(c) of the Code then in force (I.C.Z.N. 1985) as the two taxa in question are not considered congeneric. The name *Ancilla (Sparella) fasciata* (Reeve, 1864) must take precedence over *A. (S.) ordinaria* E.A. Smith, 1906 contrary to Kilburn's usage.

fasciata (Aranea)—Conchology, Pl. 46, fig. 1.

Nomen dubium. This name has appeared in the subsequent literature only when Vokes (1971: 47) listed it as "? = *Murex messorius* (G. B. Sowerby, II, 1841)." Perry's figure is here considered to be unidentifiable, a position endorsed by Roland Houart (personal communication, July 23, 2003).

fasciatus (Conus)—Conchology, Pl. 24, fig. 3.

= Conus genuanus Linnaeus, 1758 fide Kohn (1986: 7). A junior primary homonym of *C. fasciatus* Schröter, 1803.

fasciata (Cypraea)—Conchology, Pl. 22, fig. 9 (misprinted as No. 2 on plate explanation).

= *Cypraea erosa* Linnaeus, 1758 fide Adam & Leloup (1938: 133). A junior primary homonym of *C. fasciata* Gmelin, 1791.

ferruginosa (Bulla)—Conchology, Pl. 40, Bulla fig. 2.

= *Hydatina albocincta* Hoeven, 1839 fide Pilsbry (1893: 389) who stated that the use of Perry's name is barred as it is a junior primary homonym of *Bulla ferruginosa* Gmelin, 1791.

ferruginosum (Cerithium)—Conchology, Pl. 36, fig. 1.

= *Pyrazus ebeninus* (Bruguière, 1792) fide Hedley (1906: 529–531; as *P. herculeus* Martyn, 1784 [non-binominal]).

fissurella (Rostellaria)—Conchology, Pl. 11, fig. 4.

=Rimella fissurella (Linnaeus, 1767). That this is a Linnaean species has not previously been noted, probably as it is a Tertiary fossil. Perry described fossils from Grignon (Petit & Le Renard, 1990) and species from there are of this same uniform brown color. The name is not attributed by Perry, nor is any location given.

flammeum (Solen)—Conchology, Pl. 57, Solen fig 3.

Nomen dubium. This figure is not identifiable and the name has not appeared in subsequent literature.

flavescens (Pecten)—Conchology, Pl. 55, Pecten fig. 4.

Nomen dubium. Not a synonym of Flexopecten flavescens (Gmelin, 1791), considered



377

by Rombouts (1991: 41) to be a form of F. glaber Linnaeus. Perry's figure is not identifiable and the name has not appeared in subsequent literature.

flavicunda (Triplex)—Arcana, signature S₇, Pl. XXXV, Sept. 1, 1810; Conchology, Pl. 6,

= Chicoreus brunneus (Link, 1807) fide Deshayes (1843: 574; as Murex adustus Lamarck, 1822).

flexuosa (Triplex)—Conchology, Pl. 7, fig. 1.

= Naquetia triqueter (Born, 1778) fide Vokes (1964: 16).

foliacea (Hexaplex)—Conchology, Pl. 8, fig. 4.

= Hexaplex cichoreum (Gmelin, 1791) fide Cernohorsky (1967a: 128). Type species of Hexaplex Perry, 1811, q.v.

foliatus (Triplex)—Arcana, signature M₇, Pl. XXIII, June 1, 1810. [Plate 1, figure C, hereinl

Placed on Official Index of Rejected and Invalid Specific Names in Zoology (I.C.Z.N., 1970, Opinion 911). Type species of *Triplex Perry*, 1810. See discussion under genus Triplex above and under Triplex rosaria below.

forceps (Murex)—Conchology, Pl. 2, fig. 4.

= Fusinus forceps (Perry, 1811) fide Deshayes (1843: 466). This was one of the first of Perry's names to be treated as a senior synonym by Deshayes who placed Fusus turricula Kiener, 1840 in synonymy.

formosus (Monoplex)—Conchology, Pl. 3, fig. 5.

= Cymatium gutturnium (Röding, 1798) fide Cernohorsky (1972: 115).

fragum (Cardium)—Conchology, Pl. 57, Cardium fig. 3.

= Fragum fragum (Linnaeus, 1758). Perry's figure is based on Chemnitz, 1784, pl. 16, fig. 166.

frondosa (Triplex)—Conchology, Pl. 6, fig. 1.

= Chicoreus denudata (Perry, 1811) fide Vokes (1971: 51). See comments under T. denudata.

fulgens (Ancilla)—Conchology, Pl. 31, fig. 4.

= Ancilla glabrata Linnaeus, 1758 fide Deshayes (1843: 233; as Eburna).

fuliginosa (Cypraea)—Conchology, Pl. 22, fig. 1.

= Cypraea mauritiana Linnaeus, 1758 fide Schilder & Schilder (1971: 35). A junior primary homonym of C. fuliginosa Röding, 1798.

furbellata (Venus)—Conchology, Pl. 58, Venus fig. 1.

Nomen dubium. This figure is not identifiable and the name has not appeared in subsequent literature.

fusca (Hexaplex)—Conchology, Pl. 8, fig. 3.

= Homalocantha scorpio Linnaeus, 1758 fide Deshayes (1843: 586; as Murex).

fusca (Oliva)—Conchology, Pl. 41, fig. 1.

= Oliva vidua Röding, 1798 fide Deshayes (1844: 609; as Oliva maura Lamarck, 1810). Incorrectly stated by Burch & Burch (1960: 16) and Wagner & Abbott (1978: 18-806) to be a nomen nudum. Listed as a nude name by Tursch & Greifeneder (2001: 445) on the authority of the listing by Wagner & Abbott. A junior primary homonym of both O. fusca Link, 1807 O. fusca Fischer von Waldheim, 1807.

fusca (Terebra)—Conchology, Pl. 16, Terebra, fig. 3.

Nomen dubium. Placed in synonymy of *Terebra subulata* (Linnaeus, 1767) by Cernohorsky (1967b: 205). Listed under *T. taurina* [Lightfoot, 1786] by Rosenberg (2003) with statement that it is "generally considered to be a synonym of the Indo-Pacific *Terebra subulata*; but description and illustration no less consistent with *T. taurina* [Lightfoot, 1786]."

fuscum (Cerithium)—Conchology, Pl. 36, fig. 3.

Nomen dubium. This name has not appeared in subsequent literature. Possibly a stylized *Terebralia palustris* Linnaeus, 1767).

fusiformis (Strigula)—Conchology, Pl. 15, Strigula fig. 2.

Nomen dubium. This name has not appeared in subsequent literature. Probably a species of *Pythia*.

gemmosa (Cypraea)—Conchology, Pl. 23, fig. 5 (figure 6 on plate caption in error).

= Cypraea nucleus Linnaeus, 1758 fide Schilder & Schilder (1971: 66).

gigantea (Chama)—Conchology, Introduction, p. 2, not figured.

= *Tridacna gigas* (Linnaeus, 1758) fide Rosewater (1965: 373).

gloriamaris (Conus)—Arcana, signature I₇, Pl. XVI, April 1, 1810; Conchology, Pl. 25, fig. 1 (as gloria maris).

= *Conus textile* Linnaeus, 1758, fide Deshayes (1845: 124). Not *C. gloriamaris* Chemnitz, 1777. This was not intended as the introduction of a new name, but is an obvious misidentification by Perry as he referred to "that superb shell which was once in the collection of the Duchess of Portland and later sold to Lord Tankerville."

gracilis (Aranea)—Arcana, signature Aa₅, Pl. XLVII, Dec. 1, 1810.

= Murex pecten [Lightfoot, 1786], fide Ponder & Vokes (1988: 70).

gracilis (Polyplex)—Conchology, Pl. 9, fig. 4.

Nomen dubium. Placed in synonymy of *Trophon multicostatus* (Eschscholtz, 1829) by Gabb (1869: 70) although Perry's name has priority. That name is considered by some (e.g., Abbott, 1979: 189) to be a synonym of *Boreotrophon clathratus* (Linnaeus, 1767). Deshayes (1843: 590) considered both *P. gracilis* Perry and *P. bulbosa* Perry to be synonyms of *Murex magellanicus* Lamarck [sic; = Gmelin], now *Trophon geversianus* (Pallas, 1774). Martens (1872: 8) separated the two Perry figures and names and placed *P. gracilis* in the synonymy of "*Trophon laciniatus* Martyn." Cernohorsky (1977: 117) placed Perry's species in the synonymy of *Trophon plicatus* [Lightfoot, 1786]. Houart (personal communication, July 23, 2003) thinks Perry's figure is probably *Trophon multicostatus* (Eschscholtz). In view of the various placements of Perry's species, and the complexity of the Trophoninae, Perry's name should be considered a *nomen dubium*.

grandiformis (Argonauta)—Conchology, Pl. 42, fig. 4.

= Argonauta argo Linnaeus, 1758, fide Deshayes (1845: 356).

grandiformis (Harpa)—Conchology, Pl. 40, Harpa fig. 1.

= Harpa major Röding, 1798, fide Rehder (1973: 247).

greenlandica (Scalaria)—Conchology, Pl. 28, fig. 8.

- **377**
- = *Epitonium (Boreoscala) greenlandicum* (Perry, 1811), fide Clench & Turner (1952: 320). Figure copied from Chemnitz (1795, pl. 195a, fig. 1879). As shown by Clench & Turner (1952: 320) the species name sometimes is misspelled as *groenlandica* as Chemnitz' nonbinominal name for it was "Turbo clathrus groenlandicus."
- grisea (Columna)—Conchology, Pl. 51, fig. 6 (Columna No. 1 on plate explanation).
 - = Columna columna (Müller, 1774) fide Deshayes (1838: 305; as Achatina columellaris Lamarck). Figure probably copied from Chemnitz, 1786, pl. 112, fig. 955. Also see Columna marmorea herein.
- grisea (Helix)—Conchology, Pl. 15, Helix fig. 2.

Nomen dubium. This figure is not identifiable and the name has not appeared in subsequent literature.

grisea (Proscenula)—Conchology, Pl. 53, Proscenula fig. 3.

Nomen dubium. This name has not appeared in subsequent literature. It is an unidentifiable species of *Crepidula*.

hunteria (Pyrula)—Conchology, Pl. 50, fig. 4.

= Fasciolaria lilium hunteria (Perry, 1811) fide Abbott (1974: 228). Type species of Cinctura Hollister, 1957 by original designation.

imbricata (Aquaria)—Conchology, Pl. 52, Aquaria fig. 4.

Nomen dubium. Listed by L.A. Smith (1962: 173) as *P[enicillus]* (?*F[oegia]*) *imbricatus*. Improperly listed as a *nomen oblitum* by B.J. Smith (1976: 200) who stated that it cannot be referred to a particular species with certainty.

imbricatus (Spondylus)—Conchology, Pl. 59, Spondylus fig. 1.

Nomen dubium. This figure is not identifiable and the name has not appeared in subsequent literature.

indentata (Pyrula)—Conchology, Pl. 50, fig. 3.

Nomen dubium. This name has not appeared in subsequent literature. The figure is extremely stylized and cannot even be placed to family.

ionica (Rostellaria)—Conchology, Pl. 11, fig. 5.

= *Tibia fusus* (Linnaeus, 1758) fide Deshayes (1843: 655; as *R. rectirostris* Lamarck, 1822).

isabella (Cypraea)—Conchology, Pl. 19, fig. 7.

= Cypraea isabella Linnaeus, 1758.

jenningsia (Cypraea)—Conchology, Pl. 19, fig. 4.

= Cypraea limacina Lamarck, 1810 fide Schilder & Schilder (1971: 65).

labiata (Cassidea)—Conchology, Pl. 34, Cassidea fig. 1.

= Phalium labiatum (Perry, 1811) fide Abbott & Dance (1982: 114).

labiata (Cassis)—Conchology, Pl. 33, fig. 4.

= Cypraecassis rufa (Linnaeus, 1758) fide Adam & Leloup (1938: 144).

labiatus (Strombus)—Conchology, Pl. 12, fig. 3.

= *Strombus gibberulus* Linnaeus, 1758 fide Deshayes (1843: 698). A junior primary homonym of S. *labiatus* Röding, 1798.

laevis (Cerithium)—Arcana, signature H₅, Pl. XV, fig. 3, April 1, 1810.

= Rhinoclavis (Pseudovertagus) striatus (Bruguière, 1792) fide Petit & Le Renard

200TAXA

(1990:30).

laevis (Strombus)—Conchology, Pl. 13, fig. 4.

= Strombus bulla (Röding, 1798) fide Abbott (1960: 128).

lazarus (Chama)—Conchology, Pl. 59, Chama fig. 1.

= Chama lazarus (Linnaeus, 1758).

legumen (Solen)—Conchology, Pl. 57, Solen fig. 4.

= Pharus legumen (Linnaeus, 1758).

lentiginosus (Strombus)—Conchology, Pl. 12, fig. 5.

= Strombus lentiginosus Linnaeus, 1758.

leveriana (Oliva)—Conchology, Pl. 41, fig. 3.

= Oliva porphyria (Linnaeus, 1758) fide Deshayes (1844: 605).

lignarius (Murex)—Conchology, Pl. 1, fig. 2.

= Latirus polygonus (Gmelin, 1791) fide Cernohorsky (1972: 154, 155). A homonym but not a synonym of *Murex lignarius* Linnaeus, 1767 (= *M. ligniarius* Linnaeus, 1758).

linearis (Pomacea)—Conchology, Pl. 38, fig. 2.

Nomen dubium. Since its publication this name has been mentioned only twice in the literature, once by Say (1829: 260; fide Binney, 1858: 148) who listed under *Ampullaria flagellata* with the comment that he is "unacquainted with the *A. linearis* Perry, which he supposes to be a native of the coast of North America" and by Pilsbry (1927: 63) with the comment that it is not a *Pomacea*. The figure is not identifiable.

lineata (Aculea)—Conchology, Pl. 16, Aculea fig. 1.

Nomen dubium. This name has not appeared in subsequent literature. It is an unidentifiable species of *Turritella*.

lineata (Ancilla)—Conchology, Pl. 31, fig. 7.

= Nassarius glans (Linnaeus, 1758) fide Deshayes (1844: 159; as Nassa).

lineata (Murex)—Conchology, Pl. 54, fig. 3.

= Fasciolaria trapezium (Linnaeus, 1758) fide Deshayes (1843: 433).

lineata (Pleurotoma)—Conchology, Pl. 32, fig. 2.

Nomen dubium. This name has not appeared in subsequent literature and the figure is not identifiable. This name does not preoccupy the West African *Clavatula lineata* Lamarck, 1816, now placed in the genus *Perrona*.

lineata (Volutella)—Conchology, Pl. 26, fig. 5.

Nomen dubium. This name has not appeared in subsequent literature. It is an unidentifiable species of Vasidae.

lineatus (Bulimus)—Conchology, Pl. 30, fig. 5.

Nomen dubium. This name has not appeared in subsequent literature. May be a stylized *Limicolaria flammea* (Müller, 1774).

lineatum (Terebellum)—Conchology, Pl. 37, Terebellum fig. 1.

= Terebellum terebellum (Linnaeus, 1758) fide Jung & Abbott (1967: 449–453).

lycophagus (Mytilus)—Conchology, Pl. 61, Mytilus fig. 2.

Nomen dubium. This figure is not identifiable and the name has not appeared in subsequent literature.

- 377
- lynx (Cypraea)—Conchology, Pl. 22, fig. 3.
 - = Cypraea lynx Linnaeus, 1758.
- maculata (Ancilla)—Conchology, Pl. 31, fig. 5.
 - = Babylonia areolata (Link, 1807) fide Altena & Gittenberger (1981: 16).
- maculata (Cypraea)—Conchology, Pl. 20, fig. 5.
 - = *Cypraea angustata* Gmelin, 1791, fide Schilder & Schilder (1971: 47).
- maculata (Natica)—Conchology, Pl. 48, Natica fig. 3.
 - = *Natica tigrina* (Röding, 1798) fide Cernohorsky (1971: 203). Not *Natica maculata* (Salis Marschlins, 1793).
- *maculata* (*Pomacea*)—*Arcana*, signature G₅, Pl. XII, Mar. 1, 1810; *Conchology*, Pl. 38, fig. 3.
 - = *Pomacea maculata* Perry, 1810 fide Pain (1956: 79). Type species of *Pomacea* Perry, 1810 by monotypy.
- maculata (Strigula)—Conchology, Pl. 15, Strigula fig. 3.
 - = *Pythia scarabaeus* (Linnaeus, 1758). Although stylized, Perry's figure is identifiable as this species because of its coloration and the odd structure of the aperture. This synonymy has not previously been noted. This species name has not appeared in subsequent literature. Designated herein as type species of *Strigula* Perry, 1811.
- maculata (Terebra)—Conchology, Pl. 16, Terebra fig. 2.
 - = *Terebra crenulata* (Linnaeus, 1758) fide Deshayes (1844: 241). A junior primary homonym but not a synonym of *Terebra maculata* (Linnaeus, 1758).
- maculatus (Conus)—Conchology, Pl. 24, fig. 4.
 - = Conus marmoreus Linnaeus, 1758 fide Kohn (1986: 8). A junior primary homonym of Conus maculatus Bosc, 1801.
- magnifica (Aculea)—Conchology, Pl. 16, Aculea fig. 2.
 - = *Turritella terebra* (Linnaeus, 1758). Although the attenuated shell with sculpture of spiral cords makes it immediately identifiable as *Turritella terebra*, this synonymy has not previously been noted and the name has not appeared in subsequent literature.
- magnifica (Voluta)—Conchology, Pl. 18, no. 1.
 - = Cymbiola magnifica (Gebauer, 1802) fide Deshayes (1844: 397; as V. magnifica Chemnitz).
- marmoratus (Pholas)—Conchology, Pl. 56, Pholas figs. 1–3.
 - = *Pholas dactylus* Linnaeus, 1758. Although the figure is stylized, with an evenly rounded instead of slightly sinuate anterior margin, Perry's locality information and his long discussion of the shell's boring habits makes this identification probable. The name has not appeared in subsequent literature.
- marmorea (Columna)—Conchology, Pl. 51, fig. 7 (Columna No. 2 on plate explanation).
 - = *Columna columna* (Müller, 1774). Designated herein as type species of *Columna* Perry, 1811. See notes under *Columna* in the genus list herein and also *C. greisea* in the species list. This figure is a slightly stylized copy of Chemnitz, 1795, pl. 213, figure 3021.
- marmorea (Voluta)—Conchology, Pl. 18, fig. 3.
 - Nomen dubium. Martens (1872: 31) stated that this figure much resembles Voluta lap-

200TAXA 377

ponica Linnaeus, 1767, a synonymy not noted elsewhere. Not mentioned by Weaver & duPont, 1970.

marmoreus (Conus)—Conchology, Pl. 25, fig. 3.

Nomen dubium fide Kohn (1986: 8). Not Conus marmoreus Linnaeus, 1758.

maroccana (Scalaria)—Conchology, Pl. 28, fig. 1.

Nomen dubium. This name has not appeared in subsequent literature. Figure is not identifiable and probably does not represent any species of Epitoniidae.

mauritania (Cypraea)—Conchology, Pl. 21, figs. 6, 7.

= Cypraea mauritiana Linnaeus, 1758 fide Schilder & Schilder (1971: 131).

melissus (Trochus)—Conchology, Pl. 47, fig. 6.

= *Calliostoma conulum* (Linnaeus, 1758). The attenuated shape, color, and white band bordering the suture in Perry's figure make this synoymy likely. This probable synonymy has not previously been noted.

minimus (Bulimus)—Conchology, plate 30 explanation page, as subspecies of B. phasianus; not figured.

Nomen dubium. Not mentioned by Robertson (1958) in his treatment of *B. phasianus*. This name has not appeared in subsequent literature.

misella (Cypraea)—Conchology, Pl. 23, fig. 6 (figure 5 on plate caption in error).

= Palmadusta ziczac misella Perry, 1811 fide Lorenz (2002: 291).

moneta (Cypraea)—Conchology, Pl. 22, fig. 4.

= Cypraea moneta Linnaeus, 1758.

nigra (Volutella)—Conchology, Pl. 26, fig. 1.

= Vasum turbinellus (Linnaeus, 1758) fide Adam & Leloup (1938: 192).

nigricans (Strombus)—Arcana, signature Nn₅, Pl. LXXV, July, 1811.

= Lambis chiragra arthritica (Röding, 1798); = S. divergens Perry, 1811 (S. divergens on plate; S. nigricans in text).

nigricans (Volutella)—Conchology, Pl. 26, fig. 6.

Nomen dubium. May be *Vasum turbinellus* Linnaeus, 1758 but definitive identification is not possible. This name has not appeared in subsequent literature.

nitida (Tellina)—Conchology, Pl. 55, Tellina fig. 1.

= *Tellina astula* Hedley, 1917 fide Hedley (1917: 690). A junior primary homonym of *Tellina nitida* Poli, 1791.

nonpareil (Melania)—Conchology, Pl. 29, fig. 4.

= Chilonopsis aurisvulpina (Holten, 1802). Perry's figure is a copy of Chemnitz, 1795, pl. 210, fig. 2087. Two specimens were listed and figured as Chilinopsis [sic] nonpareil (Perry, 1811) by Woodward, 1991: 1, 4.

nucleata (Proscenula)—Conchology, Pl. 53, Proscenula fig. 4.

Nomen dubium. This name has not appeared in subsequent literature. It is an unidentifiable species of *Crepidula*.

oblonga (Patella)—Conchology, Pl. 43, fig. 4.

Nomen dubium. Pilsbry (1891: 156) listed as "unknown and unknowable." Not mentioned by Powell (1973) in his monograph of *Patella*.

oboesus (Monoplex)—Conchology, Pl. 3, fig. 2.



= Cymatium (Ranularia) oboesum (Perry, 1811) fide Beu (1985: 59) and Henning & Hemmen (1993: 80), both as a senior synonym of *Triton retusum* Lamarck, 1822. Mörch (1852: 110) listed as *Ranularia obesus* [sic] Perry, 1811, with *T. retusum* Lamarck, 1822, in synonymy. Deshayes (1843: 635) listed *T. retusum* and *M. oboesus* as synonyms, but gave preference to Lamarck's name as he always did. In his monograph of Cymatiidae, Kilias (1973) did not mention *M. oboesus* Perry, but he did treat *C. retusum* (Lamarck, 1822) in detail. Kilias' omission of this Perry species is inexplicable as he treated the other two species of Cymatiidae on *Conchology* plate 3.

In naming *Monoplex oboesus*, Perry used a ligature for "oe" just as in his usages of the specific name *coerulea*. In cataloguing this species Sherborn (1929) listed "oboesus" with a reference to "obesus" where *M. oboesus* is listed but with its spelling unaltered. There are no other listings for usages of *oboesus* by any author. As pointed out by Harry G. Lee (personal communication: July, 2003) the epithet *oboesus* does not appear in any classical language or biological glossary resource books and no etymon approaching "oboe" existed in Latin or Greek. The possibility of the name being based on the musical instrument oboe, itself a corrupt word, has been dismissed for several reasons. Mörch (1852: 110) cited the name as "*obesus*" but even if that is taken as an emendation instead of an error, it would be an unjustified emendation. Unfortunately the Code addresses ligatures only in company with diacritic marks, hyphens, and apostrophes with the statement that they are to be corrected. There is no provision under which either of the connected vowels can be dropped. The spelling must be left as "oboesus" and be declined as an adjective.

obtusa (Cypraea)—Conchology, Pl. 19, fig. 1.

= Cypraea pantherina Solander, 1786 fide Schilder & Schilder (1971: 37).

orbata (Pomacea)—Conchology, Pl. 38, fig. 5.

= *Pila orbata* (Perry, 1811), fide Mörch (1852: 50) and Pilsbry (1927: 63) as *Ampullaria orbata* (Perry).

orbiculare (Buccinum)—Arcana, signature Gg5, Pl. LVIII, March, 1811.

= Tonna dolium (Linnaeus, 1758). Named Buccinum dilatum on the plate but the text is headed B. orbiculare. The latter name was selected by Mathews & Iredale (1912:13) who stated that it is "a good figure of the shell later named Dolium maculatum by Lamarck [= T. dolium (L.)]." No other usage of Perry's names has been located. Winckworth & Tomlin (1933) did not include either in their list of Recent species of Tonna, nor did Kilias (1962) in his monograph of Tonnidae. The figure is of a slightly broken shell.

ornata (Strigula)—Conchology, Pl. 15, Strigula fig. 1.

Nomen dubium. This name has not appeared in subsequent literature. Possibly a specimen of the variable *Pythia scarabaeus* (Linnaeus, 1758).

ovata (Anomia)—Conchology, Pl. 60, Anomia fig. 1.

Nomen dubium. This name has not appeared in subsequent literature. It is an extremely stylized brachiopod.

ovata (Cypraea)—Conchology, Pl. 21, fig. 3.

= Cypraea turdus Lamarck, 1810 fide Deshayes (1844: 520). A junior primary hom-

onym of *C. ovata* Gmelin, 1791.

ovata (Hexaplex)—Conchology, Pl. 8, fig. 6.

Nomen dubium. Listed as a Hexaplex by Vokes (1971: 77) but not listed by Radwin & D'Attilio (1976). Wagner & Abbott (1978: 13-806) listed as "an earlier nomen oblitum for stainforthi Reeve, 1843." Their statement that it is a nomen oblitum is incorrect as the Code in force at that time had no provision for such declaration. Houart (personal communication, July 23, 2003) does not consider it a synonym of Murex stainforthi Reeve, 1843 nor does he consider Perry's figure to be identifiable. That assessment is here endorsed.

pacifica (Voluta)—Arcana, signature Q₅, Pl. XXX, August, 1810.

= *Alcithoe arabica* (Gmelin, 1791) fide Weaver & duPont (1970: 109). The name *Voluta pacifica* was taken from either the Portland Catalogue [Lightfoot, 1786: 190] or the Museum Calonnianum [Humphrey, 1797: 18], which refers to the Portland Catalogue, in both of which it is a *nomen nudum*.

pallas (Scalaria)—Conchology, Pl. 28, fig. 7.

Nomen dubium. This name has not been used in subsequent literature. It does not preoccupy *Scalaria pallasi* Kiener, 1838 as "*pallas*" is a noun in apposition and "*pallasi*" is more properly in the genitive singular.

pallida (Aculea)—Conchology, Pl. 16, Aculea fig. 3.

Nomen dubium. This name has not appeared in subsequent literature. It is possibly intended to be *Turritella terebra* (Linnaeus, 1758).

pallida (Ancilla)—Conchology, Pl. 31, fig. 3.

= Babylonia spirata (Linnaeus, 1758) fide Altena & Gittenberger (1981: 34).

pallida (Aranea)—Conchology, Pl. 46, fig. 5.

Nomen dubium. Vokes (1971: 78) listed with a query as *Murex brevispina* Lamarck, 1822. Not listed in discussion or synonymy of *Murex brevispina* Lamarck by Ponder & Vokes (1988: 60) in their treatment of the species. Other than Vokes' 1971 listing this name has not appeared in the later literature. Perry's figure is considered to be unidentifiable.

pallidus (Turbo)—Conchology, Pl. 49, fig. 5.

= *Turbo argyrostomus* Linnaeus, 1758. The spiral sculpture ornamented with spines is identifiable with a specimen figured by Abbott & Dance (1982: 46) as a "long-spined variation" of *T. argyrostomus* L. This synonymy has not previously been noted and Perry's name has not appeared in subsequent literature.

papalis (Mitra)—Conchology, Pl. 39, fig. 1.

= Mitra papalis (Linnaeus, 1758).

parkinsonia (Septa)—Conchology, Pl. 14, fig. 1.

= Austrotriton (Austrosassia) parkinsonia (Perry, 1811) fide Powell (1979: 165). Type species of Austrosassia Finlay, 1931 by original designation.

particolor (Conus)—Arcana, signature U₅, Pl. XXXIX, Oct. 1, 1810; (particolar on plate).

= *Conus aulicus* Linnaeus, 1758 fide Mathews & Iredale (1912: 12, listing plate XXIX in error) and Kohn (1986: 6).

377

particolor (Mya)—Conchology, Pl. 56, Mya figs. 1, 2.

= Panopea glicimeris (Born, 1778). Sometimes shown as an example of Perry's imagination, these two figures are simply highly stylized copies of the two figures in Born (1780, pl. 1, fig. 8 [2 figs.]) and are easily matched. In Born's figure of the hinge there is a strange protruding spoon-like structure (that does not exist in nature) with a depression or hole in its center that is duplicated by Perry. Other features of the hinge also agree. Although Perry's figure shows a symmetrical rounded shell instead of an asymmetrical oblong shell, the synonymy is obvious when the two works are compared. Perry's figures 1 and 2 are reversed as figure 1 is of a hinge and the plate caption has the hinge as figure 2. This synonymy has not previously been noted and Perry's name has not appeared in subsequent literature.

pattersonia (Voluta)—Conchology, Pl. 17, fig. 1.

= Lyria pattersonia (Perry, 1811). Hedley (1902) recognized this as being Voluta nucleus Lamarck, 1811b but stated that since they had the same date [sic] "there is no need to disturb the accepted nomenclature." Weaver & du Pont (1970: 23) list V. pattersonia as a junior synonym of Lyria nucleus (Lamarck, 1811b) but do not show exact dates of publications in the synonymy nor do they give any reason for using the later name. However, in their bibliography they show the date of Lamarck as July, 1811 and of Perry as April, 1811. It is difficult to accept that use of the earlier name would cause much confusion as this species is seldom cited, nor do we understand their apparent attempt to conceal this priority in what was supposed to be a definitive monograph. Brazier (1888: 996) had gone even further when he placed Perry's species in synonymy and stated that "if Perry had been a good authority on the subject of Conchology, his name might have taken precedence of Lamarck and other authors...." Wilson & Gillett (1971: 126), under L. nucleus, state that "the Australian form (as illustrated) is sometimes known as L. pattersonia Perry, 1811 but that name is generally regarded as a synonym." As Perry's name is in limited use it cannot be declared a nomen oblitum and no declaration is being made as to its status. Such action will be left to workers on the Australasian fauna and/or Volutidae who are more familiar with any ramifications that might arise from reinstatement of the older name and who can decide whether or not to petition the Commission.

pellucida (Trigona)—Conchology, Pl. 51, Trigona figs. 1, 2.

= *Trigonostoma scalare* (Gmelin, 1791) fide Petit (1984: 58). This species appeared in the literature for many years under the synonym *T. trigonostoma* (Lamarck, 1822). *perca* (*Biplex*)—*Conchology*, Pl. 4, fig. 5.

= *Biplex perca* (Perry, 1811) fide Beu (1998: 28). Type species of *Biplex* Perry, 1811 by subsequent designation of Gray, 1847..

perforatus (Turbo)—Conchology, Pl. 49, fig. 3.

= *Ninella torquata* (Gmelin, 1791). This is a very nice figure representing a shell well within the range of variation in *Ninella torquata* (Gmelin). The size, shape, sculpture and details of the umbilical area make this identification possible. This synonymy has not previously been noted and the name has not appeared in subsequent literature.

pespelicani (Rostellaria)—Conchology, Pl. 10, Rostellaria fig. 2 (as pes-pelicani).

- = Aporrhais pespelicani (Linnaeus, 1758).
- pharetra (Ovula)—Conchology, Pl. 53, Ovula fig. 2.
 - = Cyphoma gibbosa (Linnaeus, 1758) fide Deshayes (1844: 472).
- phasianus (Bulimus)—Arcana, signature Y₇, Pl. XLIII, November, 1810; Conchology, Pl. 30, fig. 4.
 - = Phasianella australis (Gmelin, 1791) fide Robertson (1958: 256).
- picta (Patella)—Conchology, Pl. 43, fig. 7.

Nomen dubium. Placed in synonymy of *P. granatina* (Linnaeus, 1758) with a query by Pilsbry (1891: 106); not mentioned by Powell (1973) in his treatment of that species or elsewhere in his monograph.

pictoria (Helix)—Conchology, Pl. 15, Helix fig. 1.

Nomen dubium. This name has not appeared in subsequent literature. Possibly an embellished rendering of a form of *Polymita picta* (Born, 1778).

pictum (Haustrum)—Conchology, Pl. 44, fig. 2.

Placed on Official Index of Rejected and Invalid Specific Names in Zoology (I.C.Z.N., 1969, Opinion 886). Recognized as a synonym of *Purpura planospira* (Lamarck, 1822) by Deshayes (1844: 71). See discussion under *H. dentex* above. This is another Perry species rejected at the request of Dr. Keen.

pictus (Mytilus)—Conchology, Pl. 61, Mytilus fig. 4.

= *Perna picta* (Born, 1778). Perry's figure is an almost exact copy of the figure of *Mytilus pictus* Born, 1778 in Born (1780, plate 7, fig. 7). It has not previously been noted that this is a Born species.

pinnata (Triplex)—Conchology, Pl. 7, fig. 5.

= Ceratostoma foliatum (Gmelin, 1791) fide Martens (1872: 42) and Vokes (1970: 186).

Not *Murex pinnatus* Swainson, 1822 which is a synonym of *Pterynotus alatus* (Röding, 1798).

- pinus (Rostellaria)—Arcana, signature Rr₃₋₄, Pl. LXXXII, September 1, 1811.
 - = Clavilithes pinus (Perry, 1811) fide Petit & Le Renard (1990: 32). This is a senior subjective synonym of *C. macrospira* Cossmann, 1889.
- polygonus (Murex)—Conchology, Pl. 2, fig. 1.
 - = Latirus infundibulum (Gmelin, 1791) fide Deshayes (1843: 386; as Turbinella infundibulum Lamarck). A junior homonym, but not a synonym, of Murex polygonus Gmelin, 1791.
- porcellana (Voluta)—Conchology, Pl. 17, fig. 2.
- = Marginella ventricosa Fischer von Waldheim, 1807 fide Wagner & Abbott (1978: 22-015). A junior homonym, but not a synonym, of *Voluta porcellana* Gmelin, 1791. porphyracea (Oliva)—Conchology, Pl. 41, fig. 2.
 - = Oliva miniacea (Röding, 1798) fide Burch & Burch (1960: 34). Not Oliva porphyria (Linnaeus, 1758) as stated by Wagner & Abbott (1978: 18-810) and Petuch & Sargent (1986: 113).
- punctuata (Hexaplex)—Conchology, Pl. 8, fig. 5.

Nomen dubium. Vokes (1971: 87) listed with a query as Hexaplex fulvescens (Sow-

377

erby, 1834) but there is little resemblance. Not listed by Radwin & D'Attilio (1976). Here considered to be unidentifiable.

purpurascens (Polyplex)—Conchology, Pl. 9, fig. 1.

= *Hexaplex trunculus* (Linnaeus, 1758) fide Vokes (1971: 88). See under *Polyplex* above.

purpurea (Strigula)—Conchology, Pl. 15, Strigula fig. 4.

Nomen dubium. This name has not appeared in subsequent literature. Probably a species of *Pythia* but the figure is not identifiable to species.

purpureum (Cymbium)—Conchology, Pl. 37, Cymbium fig. 3 (No. 5 on plate explanation). Nomen dubium. This name has not appeared in subsequent literature. This figure is highly stylized and may have been intended for an *Oliva* but it is not identifiable. pyriformis (Voluta)—Conchology, Pl. 17, fig. 5.

= *Turbinella pyrum* (Linnaeus, 1758) fide Deshayes (1843: 378; cited as of Lamarck). *quadrata* (*Buccinella*)—*Conchology*, Pl. 27, fig. 3.

Nomen oblitum. Only when it is realized that Perry's figure is a stylized copy of Chemnitz, 1795, pl. 179, fig. 1728, can it be identified. It is Cancellaria spengleriana Deshayes, 1830a, now placed in the genus Sydaphera. Perry's name was placed in the synonymy of S. spengleriana with a query by Verhecken (1986: 57) and listed as a nomen inquirendum by Petit & Harasewych (1990: 36). In view of this synonymy, to insure that Perry's name cannot replace the later well known synonym, action is here taken in accordance with Article 23.9.2. An exhaustive search of the literature reveals no usages of Buccinella quadrata Perry, 1811 as a senior synonym. Listed here are 25 usages of Sydaphera spengleriana (Deshayes, 1830a) during the past 50 years: Abbott & Dance, 1982: 226; Dance, 1974: 192; Habe, 1961: 434; Habe & Kosuge, 1967: 90; Habe & Okutani, 1985: 230; Hasegawa, 2000: 581; Hayasaka, 1962: 212; Higo & Goto, 1993: 276; Higo, Callomon & Goto, 1999: 294; Ikeda & Tada, 1963: 25; Kaicher, 1978: 1957; Kaseno & Matsuura, 1965: 51; Kira, 1962: 91; Kira, 1964: 82; Kuroda, Habe & Oyama, 1971: 202; Lindner, 1977: 214; Marcy & Bot, 1969: 220; Ogasawara, 1977: 139; Oliver, 1975: 262; Oyama & Takemura, 1963: Cancellaria plate; Shimaka & Horikoshi, 1963: 90, 110; Shuto, 1962: 72; M. Smith, 1961: 160; Verhecken, 1986: 57; Yoo, 1976: 149. Buccinella quadrata Perry, 1811 is declared to be a nomen oblitum and Cancellaria spengleriana Deshayes, 1830 becomes a nomen protectum.

quadrata (Natica)—Conchology, Pl. 48, Natica fig. 1.

Nomen dubium. This name has not appeared in subsequent literature. Perry's figure is a slightly stylized copy of Chemnitz, 1795, pl. 197, fig. 1902 that was placed in the synonymy of *Natica rufa* (Born, 1778) by Deshayes (1838: 639–640).

quadratus (Conus)—Conchology, Pl. 24, fig. 5.

= Conus ebraeus Linnaeus, 1758 fide Kohn (1986: 8). Not Conus quadratus (Röding, 1798).

quadratus (Strombus)—Conchology, Pl. 12, fig. 1.

= Strombus raninus Gmelin, 1791 fide Deshayes (1843: 691; as S. bituberculatus Lamarck, 1822).

radiata (Aquaria)—Conchology, Pl. 52, Aquaria fig. 3.

= *Penicillus penis* (Linnaues, 1758), fide L.A. Smith (1962: 172). Placed in same synonymy, although in genus *Brechites*, by B.J. Smith (1976: 201) as *B. radiatus* [sic]. Type species of *Aquaria* Perry, 1811, q.v.

radiata (Patella)—Conchology, Pl. 43, fig. 1.

= *Patella vulgata* Linnaeus, 1758, fide Powell (1973: 95). A junior primary homonym, but not a synonym, of *P. radiata* Born, 1778.

radiata (Venus)—Conchology, Pl. 58, Venus fig. 2.

=Pitar dione (Linnaeus, 1758) fide Martens (1872: 26).

rana (Biplex)—Conchology, Pl. 5, fig. 4.

= Bufonaria rana (Linnaeus, 1758).

reteaureum (Conus)—Conchology, Pl. 25, fig. 5 (as rete aureum).

= Conus textile Linnaeus, 1758 fide Kohn (1986: 9).

reticulatum (Buccinum)—Conchology, Pl. 48, Buccinum fig. 1.

= Buccinum undatum Linnaeus, 1758 fide Deshayes (1844: 154).

reticulatum (Cerithium)—Conchology, Pl. 35, fig. 4.

Nomen dubium. This name has not appeared in subsequent literature. It may be *Cerithium litteratum* (Born, 1778).

reticulatus (Conus)—Conchology, Pl. 24, fig. 2.

= Conus araneosus [Lightfoot, 1786] fide Kohn (1986: 9). Not C. reticulatus Born, 1778.

rosa (Biplex)—Conchology, Pl. 4, fig. 1.

= Bursa rosa (Perry, 1811) fide Beu (1998: 166). Misspelled as rosea by Martens (1872: 38) who placed it in the synonymy of "Ranella bufonia Gmelin."

rosaria (Triplex)—Conchology, Pl. 6, fig. 3. [Plate 1, figure D, herein]

= *Chicoreus rosarius* (Perry, 1811) fide Springsteen & Leobrera (1986: 134); also a synonym of *T. foliatus* Perry, q.v. The figures of *T. rosaria* and *T. foliatus* are probably of the same specimen fide Roland Houart (personal communication, July 23, 2003), although they were considered to be "the same figure" by Mathews & Iredale (1912: 11). Although *Triplex rosaria* Perry was shown to be a synonym of the later *Murex palmarosae* Lamarck, 1822 (e.g., by Deshayes, 1843: 572) many authors used the later name as they considered *Triplex rosaria* to have been suppressed in the same Opinion as *T. foliatus* (e.g., Houart, 1992: 52). However, it was not mentioned therein and continues to be an available name. A number of modern authors also showed this synonymy but used Perry's name as the senior synonym (e.g., Cernohorsky, 1978: 65).

Vokes (1978: 386) treated *Chicoreus palmarosae* (Lamarck) and wrote that "the Commission has finally reached a decision on the complicated issue of secondary homonymy, declaring that any name rejected as a secondary homonym before 1961 could not be restored (Art. 59bi, Bull. Zool. Nomen., v. 31, 1974). As *Triplex rosaria* had long been considered a secondary homonym of *Purpura rosarium* Röding, 1798, the problem has at last been satisfactorily settled and the long known name given by Lamarck continues to apply to this lovely species." Unfortunately Vokes gave no references for her statement and an extensive search has failed to find any rejection of

377

Perry's name due to this alleged secondary homonomy. Vokes certainly never mentioned any possibility of homonymy in her 1964 paper (see under *Triplex* herein) nor in her catalogue of species (Vokes, 1971). Inasmuch as both names are in use it appears that *Triplex rosaria* Perry must replace *Murex palmarosae* Lamarck, 1822. The latter name cannot be retained without action of the Commission.

rotunda (Argonauta)—Conchology, Pl. 42, fig. 1; not figure 2 as printed.

= Argonauta nodosus [Lightfoot, 1786], fide Cernohorsky (1972: 243; as nodosa but Argonauta is masculine). Cernohorsky did not notice the error in the numbering on this plate, citing A. vitrea Perry as being the synonym. See comments under A. vitrea below.

rotunda (Distorta)—Conchology, Pl. 10, Distorta fig. 2.

= *Distorsio anus* (Linnaeus, 1758) fide Emerson & Puffer (1953: 96, 103). Type species of *Distorta* Perry, 1811, q.v.

rotundata (Buccinella)—Conchology, Pl. 27, fig. 2.

= Cancellaria cancellata (Linnaeus, 1767) fide Deshayes (1845: 380).

rotundata (Cassis)—Conchology, Pl. 33, fig. 1.

Nomen oblitum. Placed in the synonymy of Cassis tuberosa (Linnaeus, 1758) by Deshayes (1844: 22) and Clench (1944: 11). However, it is not that species but is a rather good representation of C. madagascariensis spinella Clench, 1944 a taxon considered to be only a form of C. madagascariensis Lamarck, 1822 by Abbott (1968a: 52). In his discussion of Lamarck's species Abbott declared C. rotundata Perry, 1811 to be a nomen oblitum. However, his action fails to meet the requirements of Articles 23.12 of the current Code as he did not explicitly apply Article 23b of the Code then in force. To validate Abbott's action and insure that Perry's name cannot replace the later well known synonym, action is here taken in accordance with Article 23.9.2. An exhaustive search of the literature reveals no uses of C. rotundata Perry, 1811 as a senior synonym. Listed here are 25 usages of C. madagascariensis Lamarck, 1822 and/or C. m. spinella Clench, 1944 during the past 50 years: Abbott, 1968a: 52; Abbott, 1974: 161; Abbott & Dance, 1982: 110; Dance, 1974: 107; Eisenberg, 1981: 79; Emerson & Jacobson, 1976: 118; Habe & Okutani, 1985: 114; Kaicher, 1983: 3678; Lozet & Pétron, 1977: 58; Lyons, 1998: 21; Marcy & Bot, 1969: 170; Moore, 1956: 1; Oliver, 1975: 128; Porter, 1974: 196; Porter, 1975: 39; Porter & Houser, 1999: 98; Redfern, 2001: 58; Ruppert & Fox, 1988: 388; Shikama & Horikoshi, 1963: 61; M. Smith, 1961: 86; Sutty, 1986: 50; Turgeon et al., 1988: 82; Wagner & Abbott, 1964: 82; Wagner & Abbott, 1978: 12-401; Warmke & Abbott, 1961: 98. Cassis rotundata Perry, 1811 is declared to be a nomen oblitum and Cassis madagascariensis Lamarck, 1822 becomes a nomen protectum.

rotundata (Haliotis)—Conchology, Pl. 52, Haliotis fig. 1.

Nomen dubium fide Geiger, 1998: 96, 100.

rubecula (Septa)—Arcana, Signature H₈, April, 1810 (nomen nudum); Conchology, Pl. 14, fig. 5.

= Cymatium rubeculum (Linnaeus, 1758). Although not attributed directly to Linnaeus, Perry states that it "has been mistakenly called a Murex by some authors who

48



have delineated and described it. No reference to this figure has been located in synonymies. Even Deshayes (1843: 640) listed Perry's Pl. 14, fig. 2 (*Septa scarlatina*) in synonymy but did not list figure 5.

- rubescens (Volutella)—Conchology, Pl. 26, fig. 4.
 - = Vasum capitellum (Linnaeus, 1758) fide Deshayes (1843: 382; as Turbinella capitellum Lamarck).
- rubicola (Biplex)—Conchology, Pl. 5, fig. 5.
 - = *Bursa granularis* (Röding, 1798) fide Beu (1998: 150).
- rubicunda (Murex)—Conchology, Pl. 54, fig. 2.
 - = Leucozonia nassa (Gmelin, 1791). Vokes (1971: 92) listed as a Leucozonia and as not = Triplex rubicunda Perry, Pl. 6, fig. 4. From the shape and distinctive color of Perry's figure it is obvious that Vokes considered the figure to be L. nassa (Gmelin), commonly known as the Chestnut Latirus.
- rubicunda (Nerites)—Conchology, Pl. 34, Nerites fig. 2.

Nomen dubium. Listed in an index of Neritidae by Martens (1889: 139) as unrecognizable. The name has not appeared elsewhere.

rubicunda (Rostellaria)—Arcana, signature B₃, Pl. III, fig. 3, January, 1810; Conchology, Pl. 11, fig. 1.

Nomen dubium. Mathews & Iredale (1912: 9, 10; R. rubicauda [sic] on p. 9) considered this to be the same shell figured by Chemnitz (1795, figs. 1874–1875) later named Strombus erythrinus Dillwyn, 1817. Abbott (1960: 80) did not consider the figures to be equivalent and stated that Perry's "extremely poor figure probably represents some sort of buccinid." Although Perry's figures may be stylized copies of the Chemnitz figure, they are certainly not identifiable.

- rubicunda (Septa)—Conchology, Pl. 14, fig. 4.
 - = Charonia lampas rubicunda (Perry, 1811) fide Beu (1970a: 215).
- *rubicunda (Triplex)*—*Arcana*, signature S₇, Pl. XXXV, September, 1810; Conchology, Pl. 6, fig. 4.
 - = Chicoreus brunneus (Link, 1807) fide Cernohorsky (1967a: 117).
- rubra (Arca)—Conchology, Pl. 60, Arca fig. 2.

Nomen dubium. This name has not appeared in subsequent literature. The figure is not identifiable.

- rugosus (Polyplex)—Conchology, Pl. 9, fig. 2.
 - = *Nucella lamellosa* (Gmelin, 1791) fide Abbott (1974: 182). Recognized by Martens (1872: 43) as the Chemnitz figure (Chemnitz, 1788: 176, vignette 21, fig. A) on which *N. lamellosa* is based. Wilkins (1957b: 136) recognized Perry's figure as a *Nucella*, but did not realize that the figure was a copy. He stated that he could not locate the original although it was supposedly "drawn from the collection of Mr. Cracherode, now in the British Museum."
- rustica (Patella)—Conchology, Pl. 43, fig. 2.
 - = *Patella rustica* Linnaeus, 1758. This reference does not seem to have been used in any synonymy of *Patella rustica*. Perry's location is correct and the figure is appropriately colored.

377

sanguineum (Pecten)—Arcana, signature Ll₅, Pl. LXXI, June 1, 1811.

Nomen dubium. This name has not appeared in subsequent literature. Probably not intended for *Ostrea sanguinea* Linnaeus, 1758 (Pectinidae). Perry's figure is not identifiable.

scalaria (Orthoceras)—Conchology, Pl. 52, Orthoceras fig. 1

=Nautilus raphanus Linnaeus, 1758 which is not a Nautilus and is not a mollusk. Perry's figure is a stylized copy of Martini, 1769, Vignette 1, figure A, which in turn is a copy of Gualtieri, 1742, pl. 19, figure L. Linnaeus based his *N. raphanus* on the Gualtieri figure. Actual size as shown by Gualtieri and Martini is only 4 to 6 mm in length. See discussion under *Orthoceras decussata* above. Stated to be a species of Foraminifera by Hanley (1855: 154).

scarlatina (Septa)—Arcana, signature B₂, Pl. III, fig. 2, January 1, 1810; Conchology, Pl. 14, fig. 2.

= Cymatium rubeculum (Linnaeus, 1758) fide Deshayes (1843: 640; as *Triton rubecula* Lamarck). Type species of *Septa* Perry, 1810 by monotypy.

serrata (Rostellaria)—Conchology, Pl. 11, fig. 2.

Nomen dubium. Perry's figure is copied from Chemnitz (1795: 141, pl. 195A, fig. 1869) who copied it from Favanne (1780, pl. 79, fig. Y). Three additional authors later named the Chemnitz figure: Strombus fissus Dillwyn (1817: 656); Rostellaria serrata Swainson, 1840: 310; and *R. favannii* Pfeiffer, 1840: 106. Romagna-Manoja (1977: 3) figured a specimen from the Persian Gulf that he considered being this species. That specimen was designated as a neotype of R. serrata Perry (K. N[icolay] in Romagna-Manoja, 1977: 4) but that designation is not valid as the conditions of Article 75(b) then in effect were not met. Dekker (2002:34) stated that Romagna-Manoja's specimen is conspecific with Tibia curta (Sowerby, 1842) and contended that the neotype designation is not valid as "names based on teratological specimens are invalid (ICZN, Article 1.3.2). Otherwise the well known name T. curta should have been replaced by T. serrata." This is incorrect as the Code does not exclude names based on abnormal (teratological) specimens if they were not known to be such at the time they were proposed. What the Code does prohibit are names for abnormal specimens as such (i.e., an abnormal specimen cannot have a valid name of its own but must take the same name as normal specimens of the same species; if the earliest name is based on a specimen not known to be abnormal when proposed, that name must be used) (see Article 1b and Eschmeyer, 1998: 2848). Wagner & Abbott (1978: 87-001) list as "a malformed [Tibia] insulaechorab [Röding]," a conclusion accepted by all recent authors except Romagna-Manoja and Dekker. It is here considered that Rostellaria serrata Perry cannot be identified with any described species and must remain a nomen dubium.

sinensis (Rostellaria)—Conchology, Pl. 10, Rostellaria fig. 1.

= *Tibia fusus* (Linnaeus, 1758) juvenile fide Wagner & Abbott (1978: 09-657). *sinuata* (*Haliotis*)—*Conchology*, Pl. 52, *Haliotis* fig. 2

?= *Haliotis spadicea* Donovan, 1808 fide Geiger (1998: 100). *sinuata* (*Rostellaria*)—*Conchology*, Pl. 11, fig. 3.

- = *Strombus vittatus* Linnaeus, 1758. This synonymy has not been previously noted but there is little question of it. This is not *Strombus sinuatus* Perry (see below).
- sinuatus (Strombus)—Conchology, Pl. 13, fig. 3.
 - = Lambis scorpius indomaris Abbott, 1961. Abbott (1961: 165–166) proposed this as a replacement name because of the prior Strombus sinuatus [Lightfoot, 1786]. As L. s. indomaris is a replacement name, Perry's figured specimen must remain the holotype. Abbott's statement about lectotypes and paratypes is incorrect. Perry gave no indication that additional specimens were included.
- solitaris (Strombus)—Arcana, signature Dd₁, Pl. LII, January 1, 1811.
 - = Strombus gallus Linnaeus, 1758 fide Mathews & Iredale (1912: 12). A very good figure but not mentioned by Clench & Abbott (1941: 4) in their treatment of *S. gallus* L.
- spengleri (Septa)—Conchology, Pl. 14, fig. 3.
 - = Cabestana spengleri (Perry, 1811) fide Powell (1979: 164). Type species of Cymatilesta Iredale, 1936 by original designation.
- spicatum (Cerithium)—Conchology, Pl. 36, fig. 2.
 - = Tympanotonus fuscatus (Linnaeus, 1758) fide Deshayes (1843: 293; as Cerithium muricatum Bruguière, 1792).
- spinosa (Biplex)—Conchology, Pl. 5, fig. 6.
 - = *Bufonaria echinata* (Link, 1807) fide Deshayes (1843: 546; as *Ranella spinosa* Lamarck, 1816). This species was given the name *spinosa* by three authors: *Biplex spinosa* Perry, 1811; *Ranella spinosa* Lamarck, 1816; and *Bufonaria spinosa* Schumacher, 1817; all preceded by Link's name.
- spinosum (Cymbium)—Conchology, Pl. 37, Cymbium fig. 1 (No. 3 on plate explanation).
 - = *Melo amphora* [Lightfoot, 1786]. The prominent shoulder spines together with the color pattern of interrupted spiral bands make this synonymy possible. This synonymy has not previously been noted and the name has not appeared in subsequent literature.
- spinosus (Spondylus)—Conchology, Pl. 59, Spondylus fig. 2
 - *Nomen dubium.* This figure is not identifiable. This is a homonym but not a synonym of *Spondylus spinosus* Schreibers, 1793 which was treated as a valid name by Lamprell (1986: 44). See comment under *S. digitatus* Perry.
- spirale (Terebellum)—Conchology, Pl. 37, Terebellum fig. 2.
 - = Terebellum terebellum (Linnaeus, 1758) fide Jung & Abbott (1967: 449.
- stellata (Cypraea)—Conchology, Pl. 22, fig. 2.
 - = Cypraea nebrites Melvill, 1888 fide Schilder & Schilder (1971: 64). Not Cypraea stellata Gmelin, 1791.
- stolida (Cypraea)—Conchology, Pl. 23, fig. 4.
 - = Cypraea stolida Linnaeus, 1758.
- striata (Argonauta)—Conchology, Pl. 42, fig. 3.
 - *Nomen dubium.* This name has not appeared in subsequent literature. The figure is not identifiable to species.
- striata (Melania)—Conchology, Pl. 29, fig. 5.
 - Nomen dubium. This name has not appeared in subsequent literature. It is probably a



stylized representation of *Ellobium aurisjudae* (Linnaeus, 1758) or some other ellobiid.

striatum (Haustrum)—Conchology, Pl. 44, fig. 1.

Nomen dubium. This name has not appeared in subsequent literature. The figure obviously represents a *Thais* but is not indentifiable to species.

striatus (Mytilus)—Conchology, Pl. 61, Mytilus fig. 1.

Nomen dubium. This name has not appeared in subsequent literature. The figure is not identifiable to species.

subflavus (Turbo)—Conchology, Pl. 49, fig. 1

= *Turbo chrysostomus* Linnaeus, 1758. The overall shape, sculpture and coloration of Perry's figure make it easily identifiable. This synonymy has not previously been noted and the name has not appeared in subsequent literature.

subveridis (Helix)—Conchology, Pl. 15, Helix fig. 5.

Nomen dubium. Is a species of Turbo fide Martens (1872: 26).

subveridis (Oliva)—Conchology, Pl. 41, fig. 5.

Nomen dubium fide Tursch & Greifeneder (2001: 464). Listed as a color form of *Oliva tricolor* Lamarck, 1811a [Jan.—Mar.] by Petuch & Sargent (1986: 168). Incorrectly listed as a *nomen nudum* by Burch & Burch (1960: 40).

sulcata (Donax)—Conchology, Pl. 58, Donax fig. 3.

Nomen dubium. This name has not appeared in subsequent literature. The figure is not identifiable to species.

sulcata (Volutella)—Conchology, Pl. 26, fig. 2.

Nomen dubium. This name has not appeared in subsequent literature. It is an unidentifiable *Vasum*.

surinamensis (Cypraea)—Conchology, Pl. 20, fig. 4.

= *Proadusta surinamensis* (Perry, 1811) fide Lorenz (2002: 287). Type species of *Propustularia* Schilder, 1927.

tentacula (Aranea)—Conchology, Pl. 45, fig. 4.

Nomen oblitum. Sherborn (1932: 6414) mispelled as tentaculata. Vokes (1971: 106) listed simply as "Murex s.s." Roland Houart (personal communication, July 23, 2003) agrees with this writer that Perry's figure is identifiable as Vokesimurex cabritii (Bernardi, 1859). In view of this synonymy, to insure that Perry's name cannot replace the later well known synonym, action is here taken in accordance with Article 23.9.2. An exhaustive search of the literature reveals no usages of Aranea tentacula Perry, 1811 as a senior synonym. Listed here are 25 usages of Vokesimurex cabritii (Bernardi, 1859) during the past 50 years: Abbott, 1974: 171; Abbott & Dance, 1982: 131; Abbott & Morris, 1995: 208; Bayer, 1971: 154; Eisenberg, 1981: 88; Fair, 1976: 28; Habe & Okutani, 1985: 135; Kaicher, 1973: 116; Lipe & Abbott, 1991: 28; Lyons, 1998: 23; Macsotay & Campos Villarroel, 2001: 72; Marcy & Bot, 1969: 194; Merrill & Petit, 1965: 63; Perry & Schwengel, 1955: 152; Ponder & Vokes, 1988: 136; Porter, 1974: 204; Radwin & D'Attilio, 1976: 28; Rice & Korniker, 1965: 123; M. Smith, 1961: 99; Sunderland, 1990: 15; Vokes, 1963: 96; Vokes, 1988: 20; Vokes, 1990a: 26; Vokes, 1990b: 23; Wagner & Abbott, 1978: 13-803. Aranea tentacula Perry, 1811 is

200TAXA 377

declared to be a *nomen oblitum* and *Murex cabritii* Bernardi, 1859 becomes a *nomen protectum*.

- tenuis (Hexaplex)—Conchology, Pl. 8, fig. 1.
- = Homalocantha scorpio Linnaeus, 1758 fide Deshayes (1843: 586; as Murex). testudinosa (Cypraea)—Conchology, Pl. 20, fig. 1.
- = *Cypraea testudinaria* Linnaeus, 1758 fide Schilder & Schilder (1971: 40). *trapezium (Murex)—Conchology*, Pl. 1, fig. 3.
 - = *Pleuroploca filamentosa* (Röding, 1798) fide Vokes (1971: 108). Not *Murex trape-zium* Linnaeus, 1758 which is a different species of *Pleuroploca*.
- triangularis (Septa)—Conchology, Pl. 14, fig. 6.

Nomen dubium. Listed as a synonym of Cymatium (Lotoria) perryi Emerson & Old, 1963 by Henning & Hemmen (1993: 55). Emerson & Old (1963: 3) considered Perry's species "to be a species inquirendae [sic]." See their discussion and description of Cymatium perryi Emerson & Old. If specialists determine that Perry's species is the same as that named C. perryi, Perry's name should take precedence.

- tribulus (Aranea)—Conchology, Pl. 45, fig. 2.
 - = *Murex spicatus* Ponder & Vokes, 1988 fide Ponder & Vokes (1988: 78). A secondary homonym, but not a synonym, of *Murex tribulus* Linnaeus, 1758.
- triremis (Aranea)—Conchology, Pl. 45, fig. 3.
 - = Murex pecten [Lightfoot, 1786] fide Deshayes (1843: 566; as M. tenuispina Lamarck, 1822).
- tritonia (Septa)—Arcana, signature D₅, Pl. VI, February, 1810.
- = *Charonia tritonis* (Linnaeus, 1758) fide Mathews & Iredale (1912: 10). *tuberculata (Buccinella)—Conchology*, Pl. 27, fig. 7.

Nomen oblitum. Recognized as a synonym of Vasum globulum [sic] (Lamarck, 1816) by Abbott (1950: 216) and by Vokes (1966:19). The specific name globulus is a Latin noun in apposition not subject to change in termination. Lamarck's introduction is usually cited incorrectly or incompletely. The name appears on page 7 of the 1816 Liste with a reference to Plate 431bis*, figure 2. There is no name on the plate and no locality is given in the text or on the plate. Watters et al. (1999: 84) incorrectly stated that the original type locality given by Lamarck was "West Africa." Lamarck's species, the nominal subspecies Vasum globulus nuttingi Henderson, 1919 and V. g. antiguensis Usticke, 1969 are now considered to be the same. Buccinella tuberculata was declared a nomen oblitum by Vokes (1966:20) but her action fails to meet the requirements of Article 23.12 of the current Code (I.C.Z.N., 1999) as she did not explicitly apply Article 23b of the Code then in force. To validate Voke's action and insure that Perry's name cannot replace the later well known synonym, action is here taken in accordance with Article 23.9.2. An exhaustive search of the literature reveals no uses of Buccinella tuberculata Perry, 1811 as a senior synonym. Listed here are 25 usages of Vasum globulus and its synonyms listed above during the past 50 years: Abbott, 1959: 15; Abbott, 1962: 90; Abbott, 1974: 242; Abbott & Dance, 1982: 209; Boyko & Cordeiro, 2001: 128; Eisenberg, 1981: 237; Faber, 1988: 71; Habe & Okutani, 1985: 213; Hutsell et al., 1997: 68; Huttsell et al., 1999: 96; Hutsell et al., 2001:

377

116; Kaicher, 1989: 5678; Lozet & Pétron, 1977: 86; Shuster & Bode, 1961: 1; Sutty, 1986: 70; Usticke, 1969: 19; Usticke, 1971: 15; Vermeij & Rosenberg, 1993: 186; Vokes, 1966: 19; Wagner & Abbott, 1964: 80; Wagner & Abbott, 1967: 114; Wagner & Abbott, 1978: 20-403; Wagner & Abbott, 1990: 65; Watters et al., 1999: 84; Wolfe, 1976: 4. *Buccinella tuberculata* Perry, 1811 is declared to be a *nomen oblitum* and *Turbinella globulus* Lamarck, 1816 becomes a *nomen protectum*.

tuberculatum (Haustrum)—Conchology, Pl. 44, fig. 4.

= *Purpura patula* (Linnaeus, 1758) fide Deshayes (1844: 61). *tuberculus (Biplex)—Conchology*, Pl. 4, fig. 3.

= *Tutufa rubeta* (Linnaeus, 1758) fide Beu (1998: 178).

tulipa (Pyrula)—Conchology, Pl. 50, fig. 1.

= Fasciolaria tulipa (Linnaeus, 1758) fide Deshayes (1843: 432).

turriformis (Cerithium)—Conchology, Pl. 36, fig. 5.

= *Pseudovertagus aluco* (Linnaeus, 1758) fide Deshayes (1843: 291; as *Cerithium*). Not mentioned by Houbrick (1978) in his monograph on *Pseudovertagus*.

turritum (Buccinum)—Conchology, Pl. 48, Buccinum fig. 2.

Nomen dubium. ?= Bullia (Buccinanops) cochlidia (Dillwyn, 1817). Perry's figure appears to be a stylized copy of the Chemnitz figure named by Dillwyn. This was recognized by Deshayes (1844: 186-188) when he renamed Buccinum cochlidium "Chemnitz" Kiener as B. gradatum and also listed B. cochlidium Chemn[itz] where he placed Perry's figure with a query as well as Dillwyn's reference. The most recent author to treat Buccinum cochlidium, now placed in the subgenus Buccinanops in the genus Bullia, in any detail was Allmon (1990). He discussed the history of these and other names associated with this variable species, placing all in the synonymy of Bullia cochlidia (Dillwyn) with the exception of Perry's name which was not mentioned. There is obvious uncertainty about species of Buccinanops as most of the recent references listed by Allmon were cited either as B. gradatum (Deshayes) or B. lamarckii (Kiener) and it does not appear to be firmly established whether all of the available names apply to one or more species. Making the problem more difficult is the evident lack of type material for any of these names as no types are listed by Allmon. Seemingly the first available name after Perry's unused name is Bullia cochlidia (Dillwyn, 1817). In Allmon's extensive chresonymy there are only three listings, in addition to his, of Dillwyn's name being used as a senior synonym in the past 50 years and less than 25 since its introduction. Perry's name cannot be declared a nomen oblitum as the conditions of Article 23.9.1.2 cannot be met. At the same time it does not seem wise to introduce ito this complex another named based solely on a figure. Therefore, the only action that can be taken at this time is to state that Buccinum turritum Perry, 1811 is a nomen dubium and leave it to specialists to take appropriate action. It is to be noted that Buccinum turritum Perry, 1811 is shown above as ?= Bullia cochlidia (Dillwyn, 1817) and not as a senior synonym.

undulata (Anomia)—Conchology, Pl. 60, Anomia fig. 2

Nomen dubium. This name has not appeared in subsequent literature. It is an extremely stylized brachiopod.

```
undulata (Pyrula)—Conchology, Pl. 54, fig. 1.
```

= Fusinus undulatus (Perry, 1811) fide Abbott & Dance (1982: 187).

undulata (Voluta)—Conchology, Pl. 17, fig. 3.

= Voluta undulata Lamarck, 1804 fide Deshayes (1844: 401).

urniformis (Harpa)—Conchology, Pl. 40, Harpa fig. 3.

Nomen dubium. Deshayes (1844: 131) placed this in the synonymy of *Harpa ventricosa* Lamarck, 1816 and for that reason it was originally included in an application to the I.C.Z.N. (Rehder & Petit, 1987) for the conservation of *Harpa* names in long usage. However, during the application process it was learned that *Harpa ventricosa* Lamarck, 1801 is quite possibly a different species, not treated by Rehder (1973) in his monograph. The application was rewritten by the I.C.Z.N. without approval or proofs being seen by the authors (Rehder & Petit). As a result of that I.C.Z.N. action the application incorrectly states that *H. urniformis* threatens *H. articularis* Lamarck, 1822. See Opinion 1518 (I.C.Z.N., 1989:53).

ursellus (Cypraea)—Conchology, Pl. 19, fig. 2.

= Cypraea ursellus Gmelin, 1791.

valentia (Cypraea)—Conchology, Pl. 23, fig. 2.

= Leporicypraea valentia (Perry, 1811), fide Lorenz (2002: 290).

valentiana (Eburna) Perry—Mörch, 1852:76

Not a Perry species; error for Eburna valentiana Swainson, 1822.

variegata (Biplex)—Conchology, Pl. 5, fig. 2.

= Gyrineum gyrinum (Linnaeus, 1758) fide Beu (1998: 44).

variegata (Cassis)—Conchology, Pl. 33, fig. 3.

= Phalium flammiferum (Röding, 1798) fide Abbott & Dance (1982: 111).

variegata (Donax)—Conchology, Pl. 58, Donax fig. 1.

= *Meretrix meretrix* (Linnaeus, 1758). Wilkins (1957b: 136) stated that it "can safely be assigned to *Egeria radiata* (Lamarck) taken from a specimen from which the thick brown periostracum had been removed by polishing" However, Perry's figure is obviously a stylized copy of *Venus paradoxa* Born, 1778 as figured by Born (1780, pl. 4, fig. 12). Perry's figure has the same area of solid color at the umbo, the same small undulation of the lip, and almost identical striping including similarly placed double and triple stripes. Born's species is a junior synonym of *M. meretrix* (L.). The synonymy of Perry's name with *M. meretrix* (L.) has not previously been noted.

variegata (Pomacea)—Conchology, Pl. 38, fig. 4.

Nomen dubium. Pilsbry (1927: 63) stated that it "is a helicid, probably not determinable."

variegata (Scalaria)—Conchology, Pl. 28, fig. 6.

Nomen dubium. This name has not appeared in subsequent literature. A stylized epitoniid, not identifiable to species.

variegata (Serpula)—Conchology, Pl. 53, Serpula fig. 1.

Nomen dubium. This name has not appeared in subsequent literature. This highly stylized figure may represent an annelid, but it cannot definitely be identified with any known animal. It is not a mollusk.

377

variegatus (Murex)—Conchology, Pl. 2, fig. 3.

= Fusinus nicobaricus (Gmelin, 1791) fide Deshayes (1843: 468; as Fusus variegatus Desh[ayes], with Fusus laticostatus Deshayes, [1830b] in synonymy). Deshayes used Perry's name as it was earlier than his, but he assumed authorship as he placed it is a different genus, a common practice of the time.

verrucosa (Cassis)—Arcana, signature H₅, Pl. XV, fig. 4, April 1, 1810.

= *Cassis (Morionella) verrucosa* Perry, 1810, fide Petit & Le Renard (1990: 30). Is *C. cancellata* Lamarck, 1803, not *C. cancellata* Röding, 1798.

vespertina (Tellina)—Conchology, Pl. 55, Tellina fig. 3.

= *Siliqua radiata* (Linnaeus, 1758). Perry's figure is identical in shape, coloration, and marking to *S. radita* (Linnaeus) as figured by Abbott & Dance (1982: 339). This name has not appeared in subsequent literature. This synonymy has not previously been noted.

viride (Solen)—Conchology, Pl. 57, Solen fig. 1.

Nomen oblitum. This figure is not recognizable but it is certainly not referable to Solenidae. The epithet "viride" is a third declension adjective declined consistent with Perry's treatment of the genus Solen as a neuter noun (e.g., Solen angulatum and S. flammeum). However, Solen is a masculine Greek noun introduced by Linnaeus (1758) who treated it as such and Perry's taxon must become Solen viridis Perry. This name has not appeared in subsequent literature but is a senior primary homonym of Solen viridis Say, 1822, a well-known species from eastern North America. In order to enable continued usage of Say's taxon, action is here taken in accordance with Article 23.9.2. An exhaustive search of the literature reveals no usages of *Solen viridis* Perry, 1811 (as either S. viride or S. viridis). Listed here are 25 usages of Solen viridis Say, 1822 during the past 50 years: Abbott, 1968b: 252; Abbott, 1974: 495; Abbott & Dance, 1982: 338; Abbott & Morris, 1995: 98; Altena, 1971: 78; Andrews, 1977: 223; Campbell et al., 1975: 90; Coomans, 1962: 93; Dance, 1974: 254; Eisenberg, 1981: 173; Emerson & Jacobson, 1976: 400; Fox & Ruppert, 1985: 248; Habe & Okutani, 1985: 342; Lyons, 1998: 12; Maurer et al., 1974: 39; Mikkelsen et al., 1995: 124; Porter, 1971: 33; Porter, 1974: 104; Porter, 1975: 44; Porter & Houser, 1999: 58; Porter & Tyler, 1971: 10; Ruppert & Fox, 1988: 152; Shoemaker et al., 1978: 132; Turgeon et al., 1988: 40; Weisbord, 1964: 379. Solen viridis Perry, 1811 is declared to be a nomen oblitum and Solen viridis Say, 1822 becomes a nomen protectum.

viridis (Dentalia)—Conchology, Pl. 52, Dentalia fig. 3.

= Dentalium elephantinum (Linnaeus, 1758) fide Emerson (1952: 298).

viridis (Proscenula)—Conchology, Pl. 53, Proscenula figs. 1, 2.

Nomen dubium. This name has not appeared in subsequent literature. It is an unidentifiable species of *Crepidula*.

vitifera (Conus)—Conchology, Pl. 25, fig. 2.

Nomen dubium fide Kohn (1986: 9).

56

vitrea (Argonauta)—Conchology, Pl. 42, fig. 2; not figure 1 as printed.

= Argonauta vitreus Gmelin, 1791 = Carinaria cristata (Linnaeus, 1767). The transposition of the descriptions of figures 1 and 2 on Plate 42 is so obvious that little dis-

200TAXA 377

cussion is necessary. It was noticed by Lamarck who did not even comment on the transposition. Perry describes, with an adequate figure, a specimen of *Carinaria cristata* (Linnaeus, 1767) using the Gmelin name *Argonauta vitreus* (improperly emended to *vitrea*; *Argonauta* is masculine). Perry discusses the rarity of the species, mentioning the Lyanet [sic; = Lyonet] specimen, one belonging to Mr. Jennings, and "another in the British Musuem from which this was delineated." He mentions Humphrey's placement which must have been a physical placement within the British Museum collection as it does not match that of the Museum Calonnianum where the species is described in great detail. The specimen listed therein by Humphrey (1797: 6) was sold to Lyonet fide Dance (1966; 1986) who mentioned this species several times. When Dance discussed the specimen in the Jennings collection he took the opportunity to state that "Perry also based some of the nightmarish illustrations in the Conchology on specimens from the Jennings cabinet and also named *Cypraea jenningsia* after him." Why Dance did not comment on Perry's figure of this rare species is not known, nor is the significance of the Jennings shells as only four figures in Perry are so attributed.

- zealandicum (Haustrum)—Conchology, Pl. 44, fig. 5.
 - = Haustrum haustorium (Gmelin, 1791) fide Powell (1979: 180). Type species of Haustrum Perry, 1811, q.v.
- zebra (Bulimus)—Arcana, signature L₅, Pl. XIX, May 1, 1810; Conchology, Pl. 30, fig. 3.
 - = Achatina zebra (Gmelin, 1791) fide Martens (1872: 5) and Mathews & Iredale (1912: 11).
- zebra (Pleurotoma)—Conchology, Pl. 32, fig. 3.
 - *Nomen dubium.* This name has not appeared in subsequent literature. It is a stylized turrid, not identifiable.
- *zebra* (*Trochus*)—*Arcana*, signature Ii₅, Pl. LXII, April 1, 1811 (plate dated March 1, 1811 in error).
 - = *Trochus niloticus* Linnaeus, 1767 fide Mathews & Iredale (1912: 13).
- zigzag (Oliva)—Conchology, Pl. 41, fig. 4.
 - = Oliva reticulata (Röding, 1798) fide Tursch & Greifeneder (2001: 460). Listed as a color form of Oliva elegans Lamarck, 1811a [Jan.–Mar.] by Petuch & Sargent (1986: 159).

Acknowledgements

A number of persons assisted in the preparation of this paper. For translations I am indebted to Dr. Rüdiger Bieler and M. Stéphane Pras. For correspondence, literature and/or discussions about some of Perry's taxa thanks are due to Dr. E. V. Coan, Dr. Rob Cowie, Mr. Tom Eichhorst, Mr. Marien Faber, Mr. Lindsey Groves, Dr. M. G. Harasewych, Mr. Roland Houart, Dr. R. N. Kilburn, and Mr. Gijs Kronenberg. Dr. Rob Cowie of the University of Hawaii, Dr. Paula Mikkelsen of the American Museum of Natural History, Dr. Gary Rosenberg of the Academy of Natural Sciences, and Dr. Angel Valdés of the Museum of Natural History of Los Angeles County provided information on the copies of the *Con*-



chology in the libraries of their institutions. Mr. John Callomon, Mr. Paul Callomon, Dr. Harry G. Lee and Mr. Scott Jordan provided details about their copies of the Conchology. Mr. Paul Ensom, Secretary of The Palaeontographical Society, provided details of their membership lists. Dr. Harry G. Lee and Dr. Gary Rosenberg both read and commented on early drafts of this paper and their discussions on some of the problems involved, as well as their corrections and suggestions, contributed materially to the final product as did reviews by Dr. Rüdiger Bieler and Dr. Daniel Geiger.

Literature cited

Abbott, R.T. (1950) The genera Xancus and Vasum in the western Atlantic. Johnsonia, 2(28), 201-219.

Abbott, R.T. (1959) The family Vasidae in the Indo-Pacific. Indo-Pacific Mollusca, 1(1), 15-32, pls. 1-10.

Abbott, R.T. (1960) The genus Strombus in the Indo-Pacific. Indo-Pacific Mollusca, 1(2), 33-146, pls. 11-117.

Abbott, R.T. (1961) The genus Lambis in the Indo-Pacific. Indo-Pacific Mollusca, 1(3):147-174, pl. 118-134.

Abbott, R.T. (1962) Sea shells of the world. Golden Press, New York. 160 pp.

Abbott, R.T. (1968a) The helmet shells of the world (Cassidae). Part 1. *Indo-Pacific Mollusca*, 2(9), 7–202, pls. 1–186.

Abbott, R.T. (1968b). Seashells of North America. Golden Press, New York. 280 pp.

Abbott, R.T. (1974) *American seashells. Second edition*. Van Nostrand Reinhold Company, New York. 663 pp., 24 pls.

Abbott, R.T. & Dance, S.P. (1982) Compendium of Seashells. ix + 411 p. E.P. Dutton, New York.

Abbott, R.T. & Morris, P.A. (1995) A field guide to shells of the Atlantic and gulf coasts and the West Indies. Houghton Mifflin, Boston. xxxiii + 350 pp. + 74 pls.

Adam, W. & Leloup, E. (1938) Résultats scientifiques du voyage aux Indes Orientales Néerlandaises de LL. AA. RR. le Prince et la Princesse Léopold de Belgique. Prosobranchia et Opistobranchia. *Musée Royal d'Histoire Naturelle de Belgique, Mémoires*, 2(19), 1–209, pls. 1–8.

Adams, H. & Adams, A. (1853–58) *The genera of Recent Mollusca; arranged according to their organization*. John van Voorst, London. 3 vols. 1, i–x1 + 1–484 pp.; 2, 1–660 pp.; 3, 137 pls.

Allmon, W.D. (1990) Review of the *Bullia* Group (Gastropoda: Nassariidae) with comments on its Evolution, Biogeography, and Phylogeny. *Bulletins of American Paleontology*, 99(335), 1–179.

Altena, C.O.v.R. (1969) The marine Mollusca of Suriname (Dutch Guiana) Holocene and Recent. Part 1. General Introduction. *Zoologische Verhandelingen*, 101, 1–49, pls. 1–4.

Altena, C.O.v.R. (1971) On six species of marine Mollusca from Suriname, four of which are new. Zoologische Mededelingen, 45(5), 75–86.

Altena, C.O.v.R. (1975) The marine Mollusca of Suriname (Dutch Guiana) Holocene and Recent. Part III. Gastropoda and Cephalopoda. *Zoologische Verhandelingen*, 139, 1–104, pls. 1–11.

Altena, C.O.v.R. & Gittenberger, E. (1981) The genus *Babylonia* (Prosobranchia, Buccinidae). *Zoologische Verhandelingen* 188, 1–57, pls. 1–11.

Andrews, J. (1977) Shells and shores of Texas. University of Texas Press, Austin. xx + 365 pp.

Baker, H.B. (1957) Fasciolaria distans Lamarck [a letter]. Minutes of the Conchological Club of Southern California, 170, 7–8.

Bayer, F.M. (1971) New and unusual mollusks collected by R/V John Elliott Pillsbury and R/V Gerda in the tropical western Atlantic. *Bulletin of Marine Science*, 21, 11–236.

Bequaert, J.C. (1948) Monograph of the Strophocheilidae, a neotropical family of terrestrial mollusks. *Bulletin of the Museum of Comparative Zoology*, 100(1), 1–210, pls. 1–32.

Bernardi, A.C. (1859) Description d'espèces nouvelles. Journal de Conchyliologie, 7, 301–302, pl. 10.

Beu, A.G. (1970a) The Mollusca of the genus *Charonia* (family Cymatiidae). *Transactions of the Royal Society of New Zealand* 11(16), 205–223, 5 pls.

Beu, A.G. (1970b) The Mollusca of the subgenus Monoplex (family Cymatiidae). Transactions of the Royal

- Society of New Zealand 11(17), 225–237, 5 pls.
- Beu, A.G. (1971) Some synonymies in the molluscan family Cymatiidae. *Journal of the Malacological Society of Australia*, 2(2), 101–111.
- Beu, A.G. (1985) Check list of living world Ranellidae and Bursidae. *Conchologists of America Bulletin* 13(4), 56–66.
- Beu, A.G. (1998) Indo-West Pacific Ranellidae, Bursidae and Personidae (Mollusca: Gastropoda). *Mémoires du Muséum National de Histoire Naturelle*, 178, 1–255.
- Binney, W.G. (1858) *The complete writings of Thomas Say on the Conchology of the United States.* H. Balliere, New York. vi + 252 pp., 38 pls.
- Blainville, H.M.D. de (1825–1827) *Manuel de Malacologie et de Conchyliologie*. Paris. 1–647, 1825; 649–664, 109 pls., 1827.
- Born, I. (1778) *Index rerum naturalium Musei Caesarei Vindobonensis. Pars I. Testacea*. Vienna. [xlii] + 458 + [82] pp., 1 pl.
- Born, I. (1780) Testacea musei caesarei vindobonensis. Vienna. xxxvi + 442 + [17] p., 18 pls.
- Bosc, L.A.G. [1801] *Histoire naturelle des coquilles, Contenant leur description, les moeurs des animaux qui les habitent et leurs usages.* Vol. 5. Crapelet, Paris. 256 pp. [This work is dated *An* X which is September 23, 1801 through September 22, 1802. However, Dodge (1950, *The Nautilus* 63, 138–139; 64, 29–33) has shown that it was published in 1801.]
- Boyko, C.B. & Cordeiro, J.R. (2001) Catalog of Recent type specimens in the Division of Invertebrate Zoology, American Museum of Natural History. V. Mollusca, Part 2 (Class Gastropoda [exclusive of Opisthobranchia and Pulmonata], with supplements to Gastropoda [Opisthobranchia], and Bivalvia). *Bulletin of the America Museum of Natural History*, 262, 170 pp.
- Brazier, J. (1888) III. Mollusca. Proceedings of the Linnean Society of New South Wales (2)2, 993-1001.
- Bruguière, J.G. (1789–92) *Histoire Naturelle des Vers*. Encyclopedié Méthodique, 1(1), i–xviii, 1–344, 1789; 1(2), 345–757, 1792. [For complete collation of this work see Evenhuis & Petit, 2003]
- Bullis, H.R., Jr. (1964) Muricidae (Gastropoda) from the Northeast coast of South America, with descriptions of four new species. *Tulane Studies in Zoology*, 11(4), 99–107.
- Bullock, W. (1812) A companion to Mr. Bullock's London Museum and Pantherion; etc., 12th Edition. Privately printed, London. xii + 136 pp., 31 pls.
- Buonanni, F. (1681) Recreatione dell'occhio e della mente nell' osseruation' delle Chiocciole, proposta a' Curiosi delle Opere della Natura. Varese, Roma. [16] + 384 + [16] pp., 106 pls. (319 figures).
- Butot, L.J.M. (1956) [letter]. Minutes of the Conchological Club of Southern California, 155, 1-2.
- Burch, J.Q. (1955) Resurection [sic] of ancient names. *Minutes of the Conchological Club of Southern California*, 152, 2–3.
- Burch, J.Q. (1956a) Publications. Minutes of the Conchological Club of Southern California 154, 3.
- Burch, J.Q. (1956b) [no title]. Minutes of the Conchological Club of Southern California 155, 1-2.
- Burch, J.Q. (1957a) A dissenting view on Fasciolaria distans Lamarck. Minutes of the Conchological Club of Southern California, 166, 9.
- Burch, J.Q. (1957b) Biplex, Triplex, Polyplex, and other Plexes. Minutes of the Conchological Club of Southern California, 171, 11.
- Burch, J.Q. (1958) Comments on publications. *Minutes of the Conchological Club of Southern California* 178, 1–5.
- Burch, J.Q. & Burch, R.L. (1960) Catalogue of Recent and fossil olives. *Minutes of the Conchological Club of Southern California*, 196, 1–46.
- Burgess, C.M. (1970) The Living Cowries. A.S. Barnes, New York. 389 pp.
- Campbell, L., Campbell S., Colquhoun D., Ernisse J. & Abbott, W. (1975) Plio-Pleistocene faunas of the central Carolina Coastal Plain. *Geologic Notes*, 19(3), 51–124.
- Cernohorsky, W.O. (1967a) The Muricidae of Fiji. Pt. 1. Subfamilies Muricinae and Tritonaliinae. *The Veliger*, 10(2), 111–132.
- Cernohorsky, W.O. (1967b) Marine shells of the Pacific. Pacific Publications, Sydney. 248 pp., 60 pls.
- Cernohorsky, W.O. (1971) The family Naticidae (Mollusca: gastropoda) in the Fiji Islands. *Records of the Auckland Institute and Museum*, 8, 169–208.
- Cernohorsky, W.O. (1972) Marine shells of the Pacific, Vol. II. Pacific Publications, Sydney. 411 pp., 68 pls.



- Cernohorsky, W.O. (1977) The taxonomy of some Southern Ocean Mollusca (Gastropoda) mainly Antarctic and Subantarctic. *Records of the Auckland Institute and Museum*, 14, 105–119.
- Cernohorsky, W.O. (1978) Tropical Pacific Marine Shells. Pacific Publications, Sydney. 352 pp., 68 pls.
- Cernohorsky, W.O. (1984) Systematics of the family Nassariidae (Mollusca: Gastropoda). *Auckland Institute and Museum, Bulletin*, 14, i–iv, 1–356, 51 pls.
- Chemnitz, J.H. (1777) Von einer ausserordentlich seltenen Art walzenförmiger Tuten oder Kegelschnekken, welche den Namen Gloria maris führt. Beschäftigungen der Berlinischen Gesellschaft Naturforschender Freunde, 3, 321–331.
- Chemnitz, J.H. (1780–1795) *Neues systematisches Conchylien Cabinet*. G.N. Raspe, Nürnberg. Vols. 4–11 [4, [24] + 344 pp., pls. CXXII–CLIX, 1780; 5, [20] + 324 pp., pls. CLX–CLXXXXIII, 1781; 6, [12] + 375 pp., pls. 1–36, 1782; 7, [12] + 356 pp., pls. 37–69, 1784; 8, [16] + 372 pp., pls. 70–102, 1785; 9(1), [12] + 151 pp., pls. 103–116, 1786; 9(2), xxvi + 194 pp., pls. 117–136, 1786; 10, [20] + 376 pp., pls. 137–173, 1788; 11, [20] + 310 pp., pls. 174–213, 1795]. Also, see F.H.W. Martini.
- Cleevely, R.J. (1974) A provisional bibliography of natural history works by the Sowerby family. *Journal of the Society for the Bibliography of Natural History*, 6(6), 482–559.
- Clench, W.J. (1944) The genera *Casmaria*, *Galeodea*, *Phalium* and *Cassis* in the western Atlantic. *Johnsonia*, 1(16), 1–16.
- Clench, W.J. & Abbott, R.T. (1941) The genus Strombus in the western Atlantic. Johnsonia, 1(1), 1-15.
- Clench, W.J. & Turner, R.D. (1952) The genera *Epitonium* (Part II), *Depressiscala, Cylindriscala, Nystiella* and *Solutiscala* in the western Atlantic. *Johnsonia*, 2(31), 289–356.
- Coomans, H.E. (1962) The marine mollusk fauna of the Virginian area as a basis for defining zoogeographical provinces. *Beaufortia*, 9(98), 83–104.
- Cossmann, M. (1889) Catalogue illustré des coquilles fossiles de l'Eocene des environs de Paris, 4. *Annales de la Société royale Malacologique de Belgique*, 24, 7–385, 11 pls.
- Cossmann, M. (1895–1925) Essais de Paléoconchologie Comparée. Privately printed, Paris. 13 vols. [1, 160 pp., 7 pls., 1895; 2, 180 pp., 8 pls., 1896; 3, 201 pp., 8 pls., 1899; 4, 294 pp., 10 pls., 1901; 5, 216 pp., 9 pls., 1903; 6, 152 pp., 9 pls., 1904; 7, 262 pp., 14 pls., 1906; 8, 248 pp., 4 ps., 1909; 9, 216 pp., 10 pls., 1912; 10, 292 pp., 12 pls., 1916; 11, 388 pp., 11 pls., 1918; 12, 348 pp., 6 pls., 1921; 13, 345 pp., 11 pls., 1925].
- Cossmann, M. (1921) Rectifications de nomenclature. Revue Critique de Paléozoologie et de Paléophytologie, 25(2), 79–80.
- Cowie, R.H. (1999) *Pila* Röding, 1798 and *Pomacea* Perry, 1810 (Mollusca, Gastropoda): proposed placement on the Official List, and Ampullariidae Gray, 1824: proposed confirmation as the nomenclaturally valid synonym of Pilidae Preston, 1915. *Bulletin of Zoological Nomenclature*, 54(2), 83–88.
- Dall, W.H. (1904) An historical and systematic review of the frog-shells and tritons. Smithsonian Miscellaneous Collections, 47, 114–144.
- Dance, S.P. (1966) Shell Collecting: An Illustrated History. Faber and Faber, London. 344 pp., 35 pls.
- Dance, S.P. (1974) The collector's encyclopedia of shells. McGraw-Hill, New York. 288 pp.
- Dance, S.P. (1978) The art of natural history. The Overlook Press, New York. 224 pp.
- Dance, S.P. (1986) A history of shell collecting. E.J. Brill, Leiden. xv, 265 pp., 32 pls.
- Dance, S.P. & Heppell, D. (1991) Classic natural history prints, shells. Studio Editions Ltd., London. 128 pp.
- Dekker, H. (2002) A note on the distribution of two species of *Tibia* in Yemen (Gastropoda, Rostellariidae). *Vita Malacologica*, 1, 33–37.
- Deshayes, G.P. (1830) *Histoire Naturelle des Vers*. Encyclopédie Méthodique, 2(1), i–vii, 1–256, 1830a; 2(2), 1–144, 1830b. [For complete collation of this work see Evenhuis & Petit, 2003]
- Deshayes, G.P. (1838) In G.P. Deshayes & Milne Edwards, H. Histoire naturelle des animaux sans vertèbres. Deuxieme Édition, Tome 8. J.B. Ballière, Paris. 660 pp.
- Deshayes, G.P. (1843) In G. P. Deshayes & Milne Edwards, H. Histoire naturelle des animaux sans vertèbres. Deuxieme Édition, Tome 9. J.B. Ballière, Paris. 728 pp.
- Deshayes, G.P. (1844) In G. P. Deshayes & Milne Edwards, H. Histoire naturelle des animaux sans vertèbres. Deuxieme Édition, Tome 10. J.B. Ballière, Paris. 638 pp.
- Deshayes, G.P. (1845) In G. P. Deshayes & Milne Edwards, H. Histoire naturelle des animaux sans vertèbres. Deuxieme Édition, Tome 11. J.B. Ballière, Paris. 665 pp.

- Dillwyn, L.W. (1817) A descriptive catalogue of Recent shells, arranged according to the Linnaean method; with particular attention to the synonymy. John and Arthur Arch, London. i–xii + 1–1092 + [1–29 (index)] pp.
- Dillwyn, L.W. (1827) Remarks on Cypraeae described by Mr. Gray. The Zoological Journal, 3, 315.
- Donovan, E. (1808) In (1802–1820) The new Cyclopaedia or Universal Dictionary of Arts and Sciences. [not seen].
- Eisenberg, J.M. (1981) A collector's guide to seashells of the world. McGraw-Hill, New York. 239 pp.
- Emerson, W.K. (1952) Generic and subgeneric names in the molluscan class Scaphopoda. *Journal of the Washington Academy of Sciences* 42(9), 296–303.
- Emerson, W.K. & Jacobson, M.K. (1976) *The American Museum of Natural History guide to shells—Nova Scotia to Florida*. Alfred A. Knopf, Inc., New York. 482 + xviii pp., 47 pls.
- Emerson, W.K. & Old, W.E., Jr. (1963) A new subgenus and species of *Cymatium* (Mollusca, Gastropoda). *American Museum Novitates*, 2137, 1–13.
- Emerson, W.K. & Puffer, E.L. (1953) A catalogue of the molluscan genus *Distorsio* (Gastropoda: Cymatiidae). *Proceedings of the Biological Society of Washington*, 66, 93–108.
- Eschmeyer, W.M. (1998) Appendix A. Species and Genera of Fishes and the International Code of Zoological Nomenclature, pp. 2847–2881 *In* Eschmeyer, W.M., editor, Catalog of Fishes. *Center for Biodiversity Research and Information, California Academy of Sciences, Special Publication No.1*, 3 volumes. 2,905 pp.
- Eschscholtz, F. (1829) Zoologischer Atlas, enthaltend Abbildungen und Beschreibungen neuer Thierarten, während des Flottcapitains von Kotzebue zweiter Reise um die Welt, auf der Russisch-Kaiserlichen Kriegsschlupp Predpriaetië in den Jahren 1823-1826. Zweites Heft. G. Reimer, Berlin. 18 pp., pls. 6–15. [not seen].
- Evenhuis, N.L. & Petit, R.E. (2003) Corrections and additions to the dating of the "Histoire Naturelle des Vers" and the Tableau Encyclopédie (Vers, coquilles, mollusques et polypiers) portions of the Encyclopédie Méthodique. Zootaxa, 207, 1–4.
- Faber, M.J. (1988) Studies on West Indian marine mollusks 13. The malacological taxa of Gordon W. Nowell-Usticke. *De Kreukel*, 24(4–5), 67–102
- Fair, R.H. (1976) The Murex Book: an illustrated catalogue of Recent Muricidae (Muricinae, Muricopsinae, Ocenebrinae). Privately printed. 138 pp.
- Favanne, J. de & Favanne, J.G. de (1780) La conchyliologie, histoire naturelle des coquillages de mer, d'eau douce, terrestres et fossiles ...par M. Desallier d'Argenville...troisième édition. De Bure, Paris. 2 volumes + Atlas, 80 pls..
- Finlay, H.J. (1931) On *Austrosassia, Austroharpa*, and *Austrolithes*, new genera; with some remarks on the gastropod protoconch. *Transactions of the New Zealand Institute*, 62, 7–19.
- Fischer von Waldheim, G. (1807) Muséum Démidoff, ou catalogue systématique er raisonné des curiosités de la nature et de l'art.Tome 3. Moscou. [Mollusques, pp. 102–280; other parts not seen].
- Fox, R.S. & Ruppert, E.E. (1985) Shallow-water marine benthic macroinvertebrates of South Carolina. *The Belle W. Baruch Library in Marine Science*, Number 14, 1–330.
- Gabb, W.M. (1869) Palaeontology of California. Vol. II. Section I. Tertiary invertebrate fossils. Part III. Synopsis of the Tertiary invertebrate fossils of California. Geological Survey of California, Palaeontology, 2, 65–124.
- Gatliff, J.H. (1902) Notes on Perry's "Conchology." Victorian Naturalist, 19(5), 75–76.
- Gebauer, J.J. (1802) Systematisches Verzeichniß der Seesterne, Seeigel, Conchylien und Pflanzenthiere, nach Linné Systema Naturae... Halle. xii + 150 pp. [not seen].
- Geiger, D.L. (1998) Recent genera and species of the family Haliotidae Rafinesque, 1815 (Gastropoda: Vertigastropoda). *The Nautilus*, 111(3), 85–116.
- Gmelin, J.F. (1791) Caroli a Linné Systema Naturae per regna tria naturae. Edition decima tertia. Lipsiae. 1(6)(Vermes), 3021–3910.
- Grau, G. (1959) Pectinidae of the eastern Pacific. Allan Hancock Pacific Expeditions, 23, i-viii, 1-308.
- Gray, J.E. (1824–28a) Monograph on the Cypraeideae, a family of testaceous Mollusca. *The Zoological Journal*, 1, 71–80, 137–152, 367–391, 489–518, pls. 7, 12 (1825); 3, 363–371 (1827); 3, 567–576 (1828a).
- Gray, J.E. (1828b) Additions and corrections to a Monograph on Cypraea, a genus of

- testaceous Mollusca. The Zoological Journal, 4, 66-88.
- Gray, J.E. (1847) List of the genera of Recent Mollusca, their synonyma and types. *Proceedings of the Zoological Society of London*, 1847, 129–219.
- Gualtieri, N. (1742) *Index testarum conchyliorum*. Typographia Caietani Albizzini, [Florentiae]. xxiii pp. + 110 pls. with descriptive letterpress.
- Gunther, A.E. (1974) A note on the autobiographical manuscripts of John Edward Gray (1800–1875). *Journal of the Society for the Bibliography of Natural History*, 7(1), 35–76.
- Habe, T. (1961) Description of four new cancellariid species, with a list of the Japanese species of the family Cancellariidae. *Venus*, 21(4), 431–441, pls. 23–24.
- Habe, T. & Kosuge, S. (1967) 'The standard book of Japanese shells' [in Japanese]. Hoikusha, Osaka. xviii + 223 pp. + 64 pls.
- Habe, T. & Okutani, T. (1985) 'Compendium of seashells' [in Japanese]. 444 pp. [see Higo, Callomon & Goto, 1999: 576].
- Hanley, S. (1846) A monograph of the genus Tellina. Thesaurus Conchyliorum, 1(6-7), 221-236, pls. 56-66.
- Hanley, S. (1855) *Ipsa Linnaei conchylia. The shells of Linnaeus, determined from his manuscripts and collection.* William and Norgate, London. 556 pp., 5 pls.
- Hasegawa, K. (2000) Cancellarioidea, pp. 581–585, pls. 289–291 In Okutani, T. (ed.), Marine Mollusks in Japan. Tokai University Press, Tokyo. 1173 pp.
- Hayasaka, S. (1962) Summary of the geology and paleontology of the Atsumi Peninsula, Aichi Prefecture, Japan. Science Reports of the Tohoku University, Sendai, Japan, Second Series (Geology), Special Volume, No. 5, 195–217.
- Hedley, C. (1902) Studies on Australian Mollusca. Part VI. Proceedings of the Linnean Society of New South Wales, 17(1), 7–29, pls. 1–3.
- Hedley, C. (1903) Scientific results of the trawling expedition of H.M.C.S. "Thetis" off the coast of New South Wales, in February and March, 1898. Mollusca. Part II. Scaphopoda and Gastropoda. *Australian Museum, Memoir*, 4, 327–402, pls. 36–38.
- Hedley, C. (1906) Studies on Australian Mollusca. Part IX. Proceedings of the Linnean Society of New South Wales, 20(4), 520–546, pls. 31–33.
- Hedley, C. (1908) Studies on Australian Mollusca. Part X. Proceedings of the Linnean Society of New South Wales, 23(3), 456–489, pls. 7–10.
- Hedley, C. (1917) Studies on Australian Mollusca. Part XIII. *Proceedings of the Linnean Society of New South Wales*, 41(4), 679–719, pls. 46–52.
- Henning, T. & Hemmen, J. (1993) *Ranellidae & Personidae of the world*. Verlag Christa Hemmen, Wiesbaden. 263 pp.
- Hertlein, L.G. (1935) The Templeton Crocker Expedition of the California Academy of Sciences, 1932. *Proceedings of the California Academy of Sciences*, 4th Series, 21(25), 301–328, pls. 18–19.
- Hertlein, L.G. & Strong, A.M. (1955) Marine mollusks collected during the "Askoy" expedition to Panama, Colombia, and Ecuador in 1941. *Bulletin of the American Museum of Natural History*, 107(2), 159–318, pls. 1–3. (November 28)
- Higo, S. & Goto, Y. (1993) A systematic list of molluscan shells from the Japanese Is. and the adjacent area. Elle Corp., Osaka. 3 + 22 + 693 + 13 + 149 pp.
- Higo, S., Callomon, P. & Goto, Y. (1999) Catalogue and bibliography of the marine shell-bearing Mollusca of *Japan*. Elle Scientific Publications, Osaka. 749 pp.
- Hinds, R.B. (1844–1845) *The zoology of H.M.S. Sulphur, under the command of Captain Sir Edward Belcher, R.N., C.B., F.R.G.S., etc., during the years 1836–42. Vol. II. Mollusca.* Smith, Elder and Co., London. v + 72 pp., 21 pls. [Issued in parts: Pt. 1, 1–24, pls. 1–7, July 1844; pt. 2, 25–48, pls. 8–14, October 1844; pt. 3, i–v, 49–72, pls. 15–21, January, 1845].
- Hoeven, J.v.d. (1839) Tijdschrift voor Natuurlijke Geschiedenis en Physiologie. Vol. 6. [not seen].
- Hollister, S.C. (1957) On the status of Fasciolaria distans Lamarck. The Nautilus, 70(3), 73-84, pl. 6.
- Holten, H.S. (1802) Enumeratio systematica conchyliorum beat. J.H. Chemnitzii ... K.H. Scidelini, Havniae. [vi] + 88 pp.
- Hopwood, A.T. (1946) Miscellaneous notes. Proceedings of the Malacological Society of London, 26, 152–153.

- Houart, R. (1992) The genus *Chicoreus* and related genera (Gastropoda: Muricidae) in the Indo-West Pacific. *Memoires du Muséum National d'Histoire Naturelle, Zoologie* (A), 154, 1–188.
- Houbrick, R.S. (1978) The family Cerithiidae in the Indo-Pacific. Part 1: The genera *Rhinoclavis, Pseudovertagus* and *Clavocerithium. Monographs of Marine Mollusca*, 1, 1–130, pl. 1–97.
- Houbrick, R.S. (1991) Systematic review and functional morphology of the mangrove snails *Terebralia* and *Telescopium* (Potamididae; Prosobranchia). *Malacologia*, 33(1–2), 289–338.
- Hutsell, K.C., Hutsell, L.L. & Pisor, D.L. (1997) *Registry of world record size shells*. Snail's Pace Productions, San Diego. [ii] + 101 + [4] pp.
- Hutsell, K.C., Hutsell, L.L. & Pisor, D.L. (1999) *Registry of world record size shells*. 2nd *Edition*. Snail's Pace Productions, San Diego. [ii] + vii +131 + [3] pp.
- Hutsell, K.C., Hutsell, L.L. & Pisor, D.L. (2001) *Registry of world record size shells*. 3rd Edition. Snail's Pace Productions, San Diego. [ii] + vii + 158 + [2] pp.
- [Humphrey, G.] (1797) Museum Calonnianum. London. viii + 84 pp.
- Ikeda, Y. & Tada, T. (1963) 'Checklist of the Mollusca of northern Yamaguchi Prefecture' [in Japanese]. Publication of the Hagi City Museum, Yamaguchi Prefercture, Japan. 70 pp.
- International Commission on Zoological Nomenclature (1969) Opinion 886. *Purpura* Bruguière and *Muricanthus* Swainson (Gastropoda): Designations of type-species under the Plenary Powers with grant of precedence to Thaididae over Purpuridae. *Bulletin of Zoological Nomenclature*, 26(3/4), 128–132.
- International Commission on Zoological Nomenclature (1970) Opinion 911. Six misidentified type-species in the superfamily Muricacea (Gastropoda). *Bulletin of Zoological Nomenclature*, 27(1), 20–26.
- International Commission on Zoological Nomenclature (1985) *International Code of Zoological Nomenclature. Third edition.* I.T.Z.N., London. xx + 338 pp.
- International Commission on Zoological Nomenclature (1989) Opinion 1518. *Harpa articularis* Lamarck, 1822 (Mollusca, Gastropoda): specific name conserved. *Bulletin of Zoological Nomenclature*, 46(1), 53.
- International Commission on Zoological Nomenclature (1992) Opinion 1677. *Haustator* Montfort, 1810 (Mollusca, Gastropoda): conserved. *Bulletin of Zoological Nomenclature*, 49(2), 159.
- International Commission on Zoological Nomenclature (1999a) Opinion 1913. *Pila* Röding, 1798 and *Pomacea*, Perry, 1810 (Mollusca, Gastropoda) placed on the Official List, and Ampullariidae Gray, 1824: confirmed as the nomenclaturally valid synonym of Pilidae Preston, 1915. *Bulletin of Zoological Nomenclature*, 56(1), 74–76.
- International Commission on Zoological Nomenclature (1999b) *International Code of Zoological Nomenclature. Fourth edition.* I.T.Z.N., London. xxix, 306 pp.
- Iredale, T. (1915) A commentary on Suter's "Manual of the New Zealand Mollusca." *Transactions of the New Zealand Institute*, 47, 417–497.
- Iredale, T. (1936) Australian Molluscan Notes No. 2. *Records of the Australian Museum*, 29(5), 267–340, pls. 20–24.
- Iredale, T. & McMichael, D.F. (1962) A reference list of the marine Mollusca of New South Wales. *The Australian Museum*, Memoir 11, 1–109.
- Johnson, R.I. (1970) Perry's Conchology (1811) more than one edition. *Journal of the Society for the Bibliog-raphy of Natural History*, 5(4), 287.
- Jousseaume, F. (1880) Division méthodique de la famille des purpuridés. *Le Naturaliste*, Année 2(42), 335–336.
- Jousseaume, F. (1881) Diagnoses de mollusques nouveaux. Le Naturaliste, Année 3(44), 349-350.
- Jung, P. & Abbott, R.T. (1967) The genus *Terebellum* (Gastropoda: Strombidae). *Indo-Pacific Mollusca*, 1(7), 445–454, pls. 318–327.
- Kaicher, S.D. (1973) Pack #2—Muricidae Part I. Card catalogue of world-wide shells. Cards 98–195.
- Kaicher, S.D. (1978) Pack #19—Cancellariidae. Card catalogue of world-wide shells. Cards 1859-1964.
- Kaicher, S.D. (1983) Pack #36—Cassidae & Oocorythidae. Card catalogue of world-wide shells. Cards 3665–3770
- Kaicher, S.D. (1989) Pack #55—Turbinellidae Part I. Card catalogue of world-wide shells. Cards 5578–5685.
- Kaseno, Y. & Matsuura, N. (1965) Pliocene shells from the Omma Formation around Kanazawa City, Japan. *Science Reports of Kanazawa University*, 10(1), 27–62, pls. 1–20.



- Keen, A.M. (1964a) *Purpura, Ocenebra*, and *Muricanthus* (Gastropoda): Request for clarification of status. Z.N.(S.)1621. *Bulletin of Zoological Nomenclature*, 21(3), 235–239.
- Keen, A.M. (1964b) Six misidentified type-species in the superfamily Muricacea (Gastropoda). Z.N.(S.)1623. Bulletin of Zoological Nomenclature, 21(6), 422–428.
- Kesteven, H.C. (1902) Notes on Prosobranchia. No. I.- Lotorium. Proceedings of the Linnean Society of New South Wales, (2)27(3), 443–482, pl. 17.
- Kiener, L.C. (1838) Species général et iconographie des coquilles vivantes. Genre Scalaire. J.B. Baillière et fils, Paris. 22 pp., 7 pls.
- Kilburn, R.N. (1993) Notes on some South African Ancillinae with descriptions of five new species of *Amalda* (Mollusca: Gastropoda: Olividae). *Annals of the Natal Museum*, 34(2), 369–389.
- Kilias, R. (1962) Gastropoda/Prosobranchia: Tonnidae. Das Tierreich 77, 1-64.
- Kilias, R. (1973) Gastropoda/Prosobranchia: Cymatiidae. Das Tierreich 92, i–viii, 1–235.
- Kira, T. (1962) Shells of the western Pacific in color. Hoikusha, Osaka. [vii] + 224 pp., 72 pls.
- Kira, T. (1964) Coloured illustrations of the shells of Japan. Enlarged and Revised Edition. (8th printing). Hoikusha, Osaka. [6] + vii + [1] + 240 pp., [1] +71 pls.
- Kohn, A.J. (1986) Type specimens and identity of the described species of *Conus* VII. The species described 1810–1820. *Zoological Journal of the Linnean Society*, 86, 1–41.
- Kuroda, T., Habe, T. & Oyama, K. (1971) *The Sea Shells of Sagami Bay*. Maruzen Co., Tokyo. Pp. xix, 1–714 (Japanese), 121 pls., 1–489 (English), 1–53 (index).
- Lamarck, J.B.P.A. (1799) Prodrome d'une nouvelle classification des coquilles, comprenant une rédaction appropriée des caractères génériques, et l'établissement d'un grand nombre de genres nouveaux. *Mémoires de la Société d'Histoire Naturelle de Paris*, 1, 63–91.
- Lamarck, J.B.P.A. (1803) Mémoires sur les fossiles des environs des Paris. Suite. Annales du Muséum d'Histoire naturelle, 2, 163–169.
- Lamarck, J.B.P.A. (1804) Mémoires sur les fossiles des environs des Paris. Suite. Annales du Muséum d'Histoire naturelle, 4, 212–222.
- Lamarck, J.B.P.A. (1811a) Suite de la détermination des espéces de mollusques testaces: continuation du genré ovule, tarrière, ancillaire et olive. *Annales du Muséum d'Histoire naturelle*, 16, 89–114, 300–328..
- Lamarck, J.B.P.A. (1811b) Suite de la détermination des espéces de mollusques testaces, *Voluta. Annales du Muséum d'Histoire naturelle*, 17, 54–80.
- Lamarck, J.B.P.A. (1816) *Liste des objets représentés dans les planches de cette livraison*. Tableau Encyclopédique et Méthodique des Trois Règnes de la Nature. Part 23, 1–16, pls. 391–488. [For complete collation of this work see Evenhuis & Petit, 2003]
- Lamarck, J.B.P.A. (1822) Histoire naturelle des animaux sans vertèbres. Vol. 7. Paris. 711 pp.
- Lamprell, K. (1986) Spondylus, *spiny oyster shells of the world*. Robert Brown & Associates (Aust.) Pty. Ltd., Bathurst, Australia. 84 pp.
- Lamprell, K.L. & Willan, R.C. (2000) Rectification of nomenclature for three species of *Spondylus* Linnaeus (Bivalvia: Pectinoidea: Spondylidae) from the Indo-Pacific based on re-examination of type specimens. *Vita Marina*, 47(1), 1–8.
- [Lightfoot, J.] (1786) A catalogue of the Portland Museum, lately the property of the Duchess Dowager of Portland, deceased, which will be sold at auction, by Mr. Skinner and Co., ... [London]. viii + 194 pp.
- Lindner, G. (1977) Field guide to seashells of the world. Van Nostrand Reinhold, New York. 271 pp.
- Link, H.F. (1807) Beschreibung der Naturalien-Sammlung der Universität zu Rostock. Rostock. 2, 82–100; 3, 101–160, 4, 5–23.
- Linnaeus, C. (1758) Systema naturae per regna tria naturae. Editio decima, reformata. Vol. 1, Regnum animale. Stockholm. 824 pp.
- Linnaeus, C. (1767) Systema naturae per regna tria naturae. Editio duodecima, reformata. Stockholm. 1(2), 533–1327.
- Lipe, R.E. & Abbott, R.T. (1991) Living shells of the Caribbean and Florida Keys. American Malacologists, Melbourne, Florida. 80 pp.
- Lister, M. (1770) *Historiae sive synopsis methodicae conchyliorum et tabularum anatomicarum. Editio altera*. (W. Huddesford, ed.). Claredoniano, Oxonii. iv + 6 + 7 + [2] + 12 + 77 pp. + 1,059 pls. + 22 pls. [plates 1–

- 1059 are from the same engravings as the 3rd edition; see Wilkins, 1957a]
- Lorenz, F. (2002) New Worldwide Cowries. Schriften zur Malakozoologie (Cismar) 20, [i-iv], 1-292.
- Lorenz, F. & Hubert, A. (2000) A guide to worldwide cowries. Second enlarged and completely revised edition. ConchBooks, Hackenheim. 584 pp., 127 pls.
- Lozet, J.B. & Pétron, C. (1977) Shells of the Caribbean. Les éditions du pacifique, Tahiti. 138 pp.
- Lyons, W.G. (1998) Checklist of shallow-water marine Mollusca of Florida. Florida Department of Environmental Protection Technical Report, TR-3, 5–78.
- Macsotay, O. & Campos Villarroel, R. (2001) *Moluscos representativos de la plataforma de Margarita, Venezuela*. Editora Rivolta, Valencia, Venezuela. iii + 280 pp.
- Marcy, J. & Bot, J. (1969) Les coquillages. Les gastéropodes marins. La Vie Rustique, Bruxelles. 281 pp.
- Markle, D.F. (1997) Audubon's hoax: Ohio River fishes described by Rafinesque. *Archives of Natural History*, 24(3), 439–447.
- Marshall, B.A. (1998) The New Zealand Recent species of *Cantharidus* Montfort, 1810 and *Micrelenchus* Finlay, 1926 (Mollusca: Gastropoda: Trochidae). *Molluscan Research*, 19(1), 107–156.
- Martens, E. von (1872) Conchylien von Cook's Reisen. Malakozoologische Blätter, 19, 1-48.
- Martens, E. von (1873) Critical list of the Mollusca of New Zealand contained in European collections, with references to descriptions and synonyms. Colonial Museum and Geological Survey Department, Wellington. 51 + viii pp.
- Martens, E. von (1889) Die Gattungen Nerita und Neritopsis. Systematisches Conchylien-Cabinet von Martini und Chemnitz, [Kuster Edition], 2(11), 105–147, pls. 14–15, A [Theil 369].
- Martini, F.H.W. (1769–77). *Neues systematisches Conchylien Cabinet*. G.N. Raspe, Nürnberg. Vols. 1–3 [1, [18] + xxviii + 408 pp., pls. I–XXXI, 1769; 2, [6] + xvi + 362 pp., pls. XXXII–LXV; 3, [6] + vi + 434 pp., pls. LXV–CXXI, 1777; continued by J.H. Chemnitz, q.v.].
- Martyn, T. (1784–89) The universal conchologist exhibiting the figure of every known shell accurately drawn and painted after nature with a new systematic arrangement. London. Frontispiece + [3] + 39 + [2] pp. + 80 pls. [see Opinion 456, I.C.Z.N., 1957, Opinions and Declarations, 15: 393–418 and Opinion 479, I.C.Z.N., 1957, Opinions and Declarations, 16: 365–416].
- Mathews, G.M. & Iredale, T. (1912) "Perry's Arcana" An overlooked work. Victorian Naturalist, 29, 7-13.
- Matsukuma, A., Okutani, T. & Habe, T. (1991) World seashells of rarity and beauty. National Science Museum, Tokyo. viii, 206 pp.
- Maurer, D., Watling, L. & Aprill, G. (1974) The distribution and ecology of common marine and estuarine pelecypods in the Delaware Bay area. *The Nautilus*, 88(2), 38–45.
- Melvill, J.C. (1888) A survey of the genus *Cypraea* (Linné), its nomenclature, geographical distribution, and distinctive affinities, with descriptions of two new species, and several varieties. *Memoirs and Proceedings of the Manchester Literary and Philosophical Society*, (4)1, 184–252.
- Merrill, A.S. & Petit, R.E. (1965) Mollusks new to South Carolina. The Nautilus, 79(2), 58-66.
- Mikkelsen, P.M., Mikkelsen, P.S. & Karlen, D.J. (1995). Molluscan biodiversity in the Indian River lagoon, Florida. *Bulletin of Marine Science*, 57(1), 94–127.
- Montfort, D. de (1810) Conchyliologie systématique, et classification méthodique des coquilles. Tome 2. F. Schoell, Paris. 676 p. (pre-May 28, 1810; see Iredale, 1915: 457).
- Moore, D.R. (1956) Observations of predation on echinoderms by three species of Cassididae. *The Nautilus*, 69(3), 73–76.
- Moore, D.R. (1981) A long lost (extinct?) mollusc of the Ohio River. *The Bulletin of the American Malacological Union, Inc.*, 1980, 70.
- Mörch, O.A.L. (1852) Catalogus conchyliorum quae reliquit D. Alphonso D'Aguirra & Gadea Comes de Yoldi,... Fasciculus primus. Ludovici Kleini, Hafniae. 170 pp.
- Müller, O.F. (1773–74) *Vermium terrestrium et fluviatilium*. Havniae & Lipsiae. xxx + [1] + 214 + [2] + [8 (index)] pp.
- Neave, S.A. (1939–40) *Nomenclator Zoologicus*. Zoological Society of London, London. 4 volumes. xiv + 957 + 1,025 + 1,065 + 758 pp.
- OBIS (2003) Indo-Pacific Molluscan Database at the Academy of Natural Sciences. http://data.acnatsci.org/obis/search2.php/6685 (accessed November 2003).
- Ogasawara, K. (1977) Paleontological analysis of Omma Fauna from Toyama-Ishikawa Area, Horiku Prov-





- ince, Japan. Science Reports of the Tohoku University, Sendai, Second Series (Geology), 47(2), 43-156.
- Oliver, A.P.H. (1975) The Henry Holt Guide to shells of the world. Henry Holt, New York. 320 pp.
- Oyama, K. & Takemura, Y. (1963) *The Molluscan Shells. Vol. 6.* Resources Exploitation Institute, Tokyo. (unnumbered plates; see Goto, Callomon & Gotô, 1999: 609).
- Pain, T. (1956) Notes on the generic names Pomacea and Ampullarius. Journal of Conchology 24(3), 79.
- Pallas, P.S. (1774) Spicilegia Zoologica, quibus novae imprimis et obscurae animalium species iconibus, descriptionibus atque commentariis illustromtur. Berolini. Fasc.10, 1–41, pls. 1–4. [not seen].
- Perry, G. (1810–11) Arcana; or the Museum of Natural History: containing the most recent discovered objects. James Stratford, London. 84 pls. with unnumbered text [issued in parts, pls. 1–48 in 1810, pls. 49–84 in 1811].
- Perry, G. (1811) Conchology, or the natural history of shells; containing a new arrangement of the genera and species. W. Miller, London. 4 + [61] + [1] pp., 61 pls.
- Perry, L.M. & Schwengel, J.S. (1955) *Marine shells of the western coast of Florida*. Paleontological Research Institution, Ithaca. 318 pp.
- Petit, R.E. (1984) Some early names in Cancellariidae. American Malacological Bulletin, 2, 57-61.
- Petit, R.E. (1985) The trivalved mollusk. Shells and Sea Life 17(40), 132-133.
- Petit, R.E. & Harasewych, M.G. (1990) Catalogue of the superfamily Cancellarioidea Forbes and Hanley, 1851 (Gastropoda: Prosobranchia). *The Nautilus*, Supplement 1, 1–69.
- Petit, R.E. & Le Renard, J. (1990) George Perry's fossil molluscan taxa, published in the 'Arcana' (1810–1811). *Contributions to Tertiary and Quaternary Geology*, 27(1), 27–35, pls. 1–3.
- Petit, R.E. & Le Renard, J. (1991) *Haustator* Montfort, 1810 (Mollusca, Gastropoda): proposed conservation. *Bulletin of Zoological Nomenclature*, 48(1), 25–26.
- Petuch, E.J. & Sargent, D.M. (1986) *Atlas of the living olive shells of the world*. Coastal Education & Research Foundation, Charlottesville, VA. xiii + 253 pp., 39 pls.
- Pfeiffer, L. (1840) Kritisches Register zu Martini und Chemnitz's systematischem Konchylien-Kabinet. Th. Fischer, Kassel. viii + 112 pp.
- Philippi, R.A. (1846) Die Kreiselschnecken oder Trochoideen (Gattungen *Turbo, Trochus, Solarium, Rotella, Delphinula, Phasianella*). Systematisches Conchylien-Cabinet von Martini und Chemnitz, [Küster edition], von Bauer & Raspe, Nuremberg. 2(3), 98 pp., 19 pls.
- Pilsbry, H.A. (1889) Manual of Conchology. Vol. XI. Trochidae, Stomatiidae, Pleurotomariidae, Haliotidae. Academy of Natural Sciences, Philadelphia. 519 pp., 67 pls.
- Pilsbry, H.A. (1891) Manual of Conchology. Vol. XIII. Acmaeidae, Lepetidae, Patellidae, Titiscaniidae. Academy of Natural Sciences, Philadelphia. 195 pp., 74 pls.
- Pilsbry, H.A. (1893) Manual of Conchology. Vol. XV. Polyplacophora, Acanthochitidae, Cryptoplacidae and appendix. Tectibranchiata. Academy of Natural Sciences, Philadelphia. 436 pp., 61 pls.
- Pilsbry, H.A. (1901) Note on two species of Cantharidus. The Nautilus, 15(1), 8.
- Pilsbry, H.A. (1902) Cantharidus peronii Phil. The Nautilus, 16(6), 72.
- Pilsbry, H.A. (1927) On *Pomacea Perry* (Ampullariidae). *The Nautilus*, 41(2), 63–64.
- Poli, G.S. (1791–95) Testacea utriusque Siciliae eorumque historia et anatome. Bodoni Ed., Parmae. 2 volumes [1, 257 pp., 18 pls., 1791; 2, 234 pp., 20 pls., 1795].
- Ponder, W.F. & Vokes, E.H. (1988) A revision of the Indo-West Pacific fossil and Recent species of *Murex s.s.* and *Haustellum* (Mollusca: Gastropoda: Muricidae). *Records of the Australian Museum*, Supplement 8, 1–160.
- Porter, H.J. (1971) Records sizes of North Carolina mollusks. List No. 2. *North Carolina Shell Club Bulletin*, 6, 29–34.
- Porter, H.J. (1974) *The North Carolina marine and estuarine Mollusca. An atlas of occurrence*. Institute of Marine Sciences, Morehead City. vi + 351 pp.
- Porter, H.J. (1975) Records sizes of North Carolina mollusks. List No. 4. North Carolina Shell Club Bulletin, 8, 38–44.
- Porter, H.J. & Houser, L. [1999] *Seashells of North Carolina*. North Carolina Sea Grant College Program, Raleigh. iv + 132 pp.
- Porter, H.J. & Tyler, J. (1971) *Sea Shells Common to North Carolina*. North Carolina Department of Natural and Economic Resources. 36 pp.



- Powell, A.W.B. (1964) The family Turridae in the Indo-Pacific. Part 1. The subfamily Turrinae. *Indo-Pacific Mollusca*, 1(5), 227–346, pls. 172–262.
- Powell, A.W.B. (1969) The family Turridae in the Indo-Pacific. Part 2. The subfamily Turriculinae. *Indo-Pacific Mollusca*, 2(10), 215–416, pls. 188–324.
- Powell, A.W.B. (1973) The patellid limpets of the world (Patellidae). *Indo-Pacific Mollusca*, 3(15), 75–206, pls. 63–182.
- Powell, A.W.B. (1979) New Zealand Mollusca. William Collins Publishers, Auckland. xiv + 500 pp.
- Pritchard, G.B. & Gatliff, J.H. (1906) Catalogue of the marine shells of Victoria. Pt. IX. *Proceedings of the Royal Society of Victoria*, 18(N.S.)2, 39–92.
- Radwin, G.E. & D'Attilio, A. (1976) Murex *shells of the world*. Stanford University Press, Stanford. 284 pp., 32 pls.
- Redfern, C. (2001) Bahamian Seashells. A thousand species from Abaco, Bahamas. Bahamianseashells.com, Inc., Boca Raton. ix + 280 pp. + 124 pls.
- Reeve, L.A. (1841–42) Conchologia Systematica, or complete system of conchology: in which the lepades and conchiferous Mollusca are described and classified according to their natural organization and habits. Longman, Brown, Green and Longmans, London. 2 vols. [1, 1–195, pls. 1–114, 114*, 115–129, 1841; 2, 1–337, pls. 130–300, 1842].
- Reeve, L.A. (1843) On a new species of the genus *Murex. Proceedings of the Zoological Society of London*, 10(114), 104.
- Reeve, L.A. (1844) Monograph of the genus *Ranella*. *Conchologia Iconica*, 2, [unpaginated text], pls. 1–8. Reeve, L.A. (1845–46) Monograph of the genus *Cypraea*. *Conchologia Iconica*, 3, [unpaginated text], pls. 1–27
- Reeve, L.A. (1847–48) Monograph of the genus *Fusus*. *Conchologia Iconica*, 4, [unpaginated text], pls. 1–21. Reeve, L.A. (1864) Monograph of the genus *Ancillaria*. *Conchologia Iconica* 15, [unpaginated text], pls. 1–12. Regenfuss, F.M. (1758) *Auserlesne Schnecken*, *Muscheln und andre Schaalthiere*. A.H. Godiche, Copenhagen. [xiv] + xiv + 22 + [lxxxvii] pp., 12 pls.
- Rehder, H.A. (1973) The family Harpidae of the world. *Indo-Pacific Mollusca*, 3(16), 207–274, pls. 183–247. Rehder, H.A. & Petit, R.E. (1987) *Harpa articularis* Lamarck, 1822 (Mollusca, Gastropoda): proposed conservation of the specific name. *Bulletin of Zoological Nomenclature*, 44(1), 19–20.
- Rice, W.H. & Kornicker, L.S. (1965). Mollusks from the deeper waters of the Northwestern Campeche Bank, Mexico. *Publications of Institute of Marine Science, Texas* 10, 108–172.
- Rios, E.C. (1970) Coastal Brazilian Seashells. Museu Oceanográfico de Rio Grande. 255 pp., 60 pls.
- Rios, E.C. (1975) Brazilian marine mollusks iconography. Museu Oceanográfico da FURG. 331 pp., 91 pls.
- Rios, E.C. (1985) Seashells of Brazil. Museu Oceanográfico da FURG. 328 pp., 102 pls.
- Rios, E.C. (1994) Seashells of Brazil. 2nd Edition. Museu Oceanográfico da FURG. 368 pp., 113 pls.
- Robertson, R. (1958) The family Phasianellidae in the western Atlantic. *Johnsonia*, 3(37), 245–283.
- Röding, P.F. (1798) Museum Boltenianum sive Catalogus cimeliorum e tribus regnis naturae quae olim collegerat Joa. Fried. Bolten, M.D.p.d. Johan. Christi. Trappii, Hamburgi. 199 pp.
- Rombouts, A. (1991) Guidebook to Pecten shells: Recent Pectinidae and Propeamussiidae of the world. Universal Book Services, Oegstgeest. xiv + 157 p.
- Romagna-Manoja, E. (1977) Tibia serrata is no longer a "Ghost species." La Conchiglia, 102-103, 3-4.
- Rosenberg, G. (2003) [*Terebra taurina*] in Western Atlantic Gastropod Database, Malacog Version 3.2.3. http://erato.acnatsci.org/wasp/search.php/7853 (access November 2003).
- Rosewater, J. (1961) The family Pinnidae in the Indo-Pacific. *Indo-Pacific Mollusca*, 1(4), 175–226, pls. 135–171.
- Rosewater, J. (1965) The family Tridacnidae in the Indo-Pacific. *Indo-Pacific Mollusca*, 1(6), 347–396, pls. 263–293.
- Ruppert, E.E. & Fox, R.S. (1988) Seashore Animals of the Southeast. University of South Carolina Press, Columbia. 429 pp.
- Salis Marschlins, H.U. von (1793) Reisen in verschiedenen Provinzen des Königsreichs Neapel. Ziegler, Zürich & Leipzig [not seen].
- Salis Marschlins, H.U. von (1795) *Travels through various provinces of the Kingdom of Naples in 1789*. Translated from the German by A. Aufrere. London. i–iv, i–viii, 527 p. [English translation of Salis Mar-

- schlins (1793)].
- Say, T. (1822) An account of some of the marine shells of the United States (cont'd). *Journal of the Academy of Natural Sciences of Philadelphia*, 2, 302–325.
- Schilder, F.A. (1927) Revision der Cypraeacea (Mollusca, Gastropoda). *Archiv für Naturgeschichte* 91A(10), 1–171.
- Schilder, M. & Schilder, F.A. (1971) A catalogue of living and fossil cowries. *Institut Royal des Sciences Naturelles de Belgique, Mémoires*, 85, 1–246.
- Schreibers, K. (1793) *Versuch einer vollständigen Conchylienkenntniss nach Linnes System*. Wien. 2 volumes. 1, [10] + 446 + [1] pp.; 2, [8] + 416 pp.
- Schröter, J.S. (1801–06) Neue Conchylienarten und Abänderungen, Anmerkungen, und Berichtigungen nach dem linnéischen Systerm der XII. Ausgabe. *Archiv für Zoologie und Zootomie*, 2(1), 82–110, 1801; 2(2), 107–148, 1802; 3(1), 125–166, 1802; 3(2), 33–78, 1803; 4(1), 7–44, 1804; 4(2), 75–137, 1805; 5(1), 129–181, 1806.
- Schumacher, C.F. (1817) Essai d'un nouveau système des habitations des vers testacés. Schultz, Copenhague. 287 pp., 22 pls.
- Sherborn, C.D. (1909) On "The Conchological Ilustrations," by George Brettingham Sowerby, Jun., London, 1832–41, and the "Descriptive Catalogue of Shells," by John Edward Gray, 1832. *Proceedings of the Malacological Society of London*, 8(6), 331–332.
- Sherborn, C.D. (1922–32) *Index Animalium sive index nominum quae ab A.D. MDCCLVIII generibus et specibus animalium imposita sunt. Sectio Secunda. A kalendis Ianuariis, MDCCCI usque ad finem Decembris, MDCCCL*. British Museum (Natural History), London. cxlvii + 7,056 + 114 pp.
- Shimaka, T. & Horikoshi, M. (1963). Selected shells of the world illustrated in colours. Hokuryukan, Tokyo. 8 + 102 pls. (with overleafs) + 154 pp.
- Shoemaker, A.H., Porter, H.J., Boothe, B., Petit, R.E. & Eyster, L.S. (1978) Marine Mollusks, pp. 123–135 *in* Zingmark, R.G. (ed.), *An Annotated Checklist of the Biota of the Coastal Zone of South Carolina*. University of South Carolina Press, Columbia. xii + 364 pp.
- Shuster, C.N., Jr. (1961) Observations on *Vasum globulus nuttingi*, with comments on other Caribbean vase shells. *The Nautilus*, 75(1), 1–7.
- Shuto, T. (1962) Buccinacean and Volutacen gastropods from the Miyazaki Group. *Memoirs of the Faculty of Science, Kyushu University, Series D, Geology*, 12(1), 27–85, pls. 6–13.
- Smith, B.J. (1976) Revision of the Recent species of the family Clavagellidae (Mollusca, Bivalvia). *Journal of the Malacological Society of Australia*, 3(3–4), 187–209.
- Smith, E.A. (1906) On South African marine Mollusca. *Annals of the Natal Government Museum*, 1(1), 19–71, pls. 7, 8.
- Smith, L.A. (1962) Revision of the Clavagellacea. The Veliger, 4(4), 167-174.
- Smith, M. (1961) *Universal Shells*. Alpine Press, Asheville. xiii + 254 pp.
- Sowerby, G.B., I (1827) [footnote to Dillwyn's article]. The Zoological Journal, 3, 315.
- Sowerby, G.B., I (1828) Notes on Mr. Gray's Additions and Corrections to a Monograph on *Cypraea*, (Zoological Journal, IV. p. 66,) in a Letter addressed to W. J. Broderip, Esq. *The Zoological Journal*, 4, 217–221.
- Sowerby, G.B., I (1832–33) *Cancellaria. The Conchological Illustrations*, Parts 9–13. London. 5 pls. with explanations [pages and plates unnumbered] + catalogue, 10 p. [Parts 9–12, figs. 1–35, 1832; pt. 13, figs. 36–44 and catalogue, 1833].
- Sowerby, G.B., II (1834) *Murex. The Conchological Illustrations*, Parts 58–67. London. 10 pls. with explanations [pages and plates unnumbered].
- Sowerby, G.B., II (1841) Descriptions of some new species of *Murex*, principally from the collection of H. Cuming, Esq. *Proceedings of the Zoological Society of London*, 8(94), 137–147.
- Sowerby, G.B. II (1842) Monograph of the genus Rostellaria. Thesaurus Conchyliorum, 1, 21-22, pl. 5.
- Springsteen, F.J. & Leobrera, F.M. (1986). *Shells of the Philippines*. Carfel Seashell Museum, Manila. 377 pp., 100 pls.
- Sunderland, K. (1990) Caribbean Muricidae—*Siratus* and *Haustellum*. *American Conchologist*, 18(1), 14–15. Sutty, L. (1986) *Seashell Treasures of the Caribbean*. E.P. Dutton, New York. 128 pp.
- Swainson, W. (1822) A catalogue of the ... shells which formed the collection of Mrs. Bligh, with an appendix containing ... descriptions of many new species. London. 58 pp., 2 pls. [not seen].


- Swainson, W. (1840) A treatise on malacology, or shells and shell-fish. Longman, Orme, Brown, Green, & Longmans, London. viii + 419 pp.
- Thiele, J. (1929–35) Handbuch der systematischen Weichtierkunde. Gustav Fischer, Jena. 1(1), 1–376, 1929; 1(2), i–vi, 377–778, 1931; 2(3), 779–1022, 1934; 2(4), i–vi, 1023–1154, 1935.
- Tryon, G.W., Jr. (1888) *Manual of Conchology. Vol. X, Part 2. Monographs of the Turbinidae and Trochidae*. Philadelphia. Pp. 161–290, pls. 39a–69.
- Turgeon, D.D., Bogan, A.E., Coan, E.V., Emerson, W.K., Lyons, W.G., Pratt, W.L., Roper, C.F.E., Scheltema, A., Thompson, F.G. & Williams, J.D. (1988) Common and scientific names of aquatic invertebrates from the United States and Canada: Mollusks. *American Fisheries Society Special Publication*, vii + 277 pp.
- Tursch, B. & Greifeneder, D. (2001) *Oliva Shells. The genus Oliva and the species problem.* L'Informatore Piceno, Ancona. x + 570 pp.
- Usticke, G.W.N. (1959) A check list of the marine shells of St. Croix, U. S. Virgin Islands with random annotations. Lane Press, Burlington. vi + 90 pp., 4 pls.
- Usticke, G.W.N. (1969) A supplementary listing of new shells (illustrated), to be added to the check list of the marine shells of St. Croix. Privately printed. 32 pp., 6 pls. [imprinted 1968 but published in 1969]
- Usticke, G.W.N. (1969) A supplementary listing of new shells (illustrated), revised edition, to be added to the check list of the marine shells of St. Croix. Privately printed. 32 pp., 6 pls.
- Verhecken, A. (1986) The Recent Cancellariidae of Indonesia (Neogastropoda, Cancellariacea). Gloria Maris, 25(2), 29–66.
- Vermeij, G.J. & Rosenberg, G. (1993) Giving and receiving: the tropical Atlantic as donor and recipient region for invading species. American Malacological Bulletin, 10(2), 181–194.
- Vokes, E.H. (1963) Cenozoic Muricidae of the western Atlantic region. Part I—Murex sensu stricto. Tulane Studies in Geology, 1(3), 95–123.
- Vokes, E.H. (1964) Supraspecific groups in the subfamilies Muricinae and Tritonaliinae (Gastropoda: Muricidae). *Malacologia*, 2(1), 1–41.
- Vokes, E.H. (1966) The genus *Vasum* (Mollusca: Gastropoda) in the New World. *Tulane Studies in Geology*, 5(1), 1–36.
- Vokes, E.H. (1970) Some comments on Cernohorsky's "Muricidae of Fiji" (The Veliger, 1967). *The Veliger*, 13(2), 182–187.
- Vokes, E.H. (1971) Catalogue of the genus *Murex* Linné (Mollusca: Gastropoda; Muricinae, Ocenebrinae. *Bulletins of American Paleontology*, 61(268), 1–141.
- Vokes, E.H. (1978) Muricidae (Mollusca: Gastropoda) from the eastern coast of Africa. *Annals of the Natal Museum*, 23(2), 375–418.
- Vokes, E.H. (1988) Muricidae (Mollusca: Gastropoda) of the Esmeraldas Beds, northwestern Ecuador. *Tulane Studies in Geology and Paleontology*, 21(1), 1–50.
- Vokes, E.H. (1989) Neogene paleontology in the northern Dominican Republic. 8. The family Muricidae (Mollusca: Gastropoda). *Bulletins of American Paleontology*, 97(332), 1–94, pls. 1–12.
- Vokes, E.H. (1990a). What ever happened to dear old Murex? American Conchologist, 18(1), 3-7.
- Vokes, E.H. (1990b) Cenozoic Muricidae of the western Atlantic region. Part VIII—Murex s.s., Haustellum, Chicoreus, and Hexaplex; additions and corrections. Tulane Studies in Geology and Paleontology, 23(1–3), 1–96.
- Vokes, E.H. (1995) Two additions to the Cenozoic Muricidae (Gastropoda: Muricidae) of the western Atlantic region. *Tulane Studies in Geology and Paleontology*, 28(4), 119–126.
- Vokes, E.H. (1996) A new species of *Poirieria (Paziella)* (Gastropoda: Muricidae) from the Punta Gavilan Formation, northern Venezuela. *Tulane Studies in Geology and Paleontology*, 29(3), 67–68.
- Wagner, R.J.L. & Abbott, R.T. (1964) Van Nostrand's Standard catalog of shells. First Edition. D. Van Nostrand Company Inc., Princeton. vii + 195 pp.
- Wagner, R.J.L. & Abbott, R.T. (1967) *Van Nostrand's Standard catalog of shells. Second Edition.* D. Van Nostrand Company Inc., Princeton. xi + 303 pp.
- Wagner, R.J.L. & Abbott, R.T. (1978) *Standard catalog of shells. Third edition.* American Malacologists, Greenville, Delaware. Looseleaf; irregular pagination.
- Wagner, R.J.L. & Abbott, R.T. (1990) World size records 1990. Standard Catalog of Shells. Supplement 4. American Malacologists, Melbourne, Florida. ii + 80 pp.



- Warmke, G.L. & Abbott, R.T. (1961) *Caribbean seashells*. Livingston Publishing Company, Narberth. x + 346 pp.
- Watters, D.R., Petuch, E.J. & Serrand, N. (1999) Anomalous distributions of *Vasum globulus*: Discordant records of an endemic gastropod in archaeological and modern contexts. *Caribbean Journal of Science*, 35(1–2), 83–92.
- Weaver, C.S. & du Pont, J.E. (1970) Living Volutes. *Delaware Museum of Natural History, Monograph* 1, 1–375, pls. 1–79.
- Wilkins, G.L. (1957a) Notes on the *Historia Conchyliorum* of Martin Lister (1638–1712). *The Journal of the Society for the Bibliography of Natural History*, 3(4), 196–205.
- Wilkins, G.L. (1957b) The Cracherode shell collection. *Bulletin of the British Museum (Natural History), Historical Series*, 1(4), 121–184, pls. 20–25.
- Weisbord, N.E. (1964) Late Cenozoic pelecypods from northern Venezuela. *Bulletins of American Paleontology*, 45(204), 564 pp.
- Wilson, B.R. & Gillett, K. (1971) Australian shells. Charles E. Tuttle Co., Rutland, Vermont. 168 pp., 106 pls. Winckworth, R. (1945) The types of the Boltenian genera. Proceedings of the Malacological Society of London, 26(4–5), 136–148.
- Winckworth, R. & Tomlin, J.R. le B. (1933) Recent species of the genus *Tonna* (=Dolium). Proceedings of the Malacological Society of London, 20(4), 206–213.
- Wolfe, C.S. (1976) Notes on the December identification page. Hawaiian Shell News, 24(12), 4, 7.
- Woodward, F.R. (1991) Dr. William B. Lorrain and St. Helena's endemic land shells. *Hawaiian Shell News*, 39(11), 1, 4.
- Yoo, J.S. (1976) 'Korean shells in colour' [in Korean]. Ijisa Co., Seoul. 196 pp., 36 pls.
- Zilch, A. (1959) Euthyneura. In: Handbuch der Paläozoologie. Gebrüder Borntraeger, Berlin. Lief. 1, 1–200, Lief 2, 201–400.
- **PLATE 1.** Reproduced here are the two figures upon which *Murex palmarosae* Lamarck, 1822 is based. Also reproduced are figures of two species from Perry's works considered to be synonyms of *M. palmarosae* Lamarck. All four figures are reproduced here near their original size (A. 9.6 cm; B. 9.8 cm; C. 11.5 cm; D. 11.6 cm).
- A. Figure (41)946 from Lister, 1770 [1688]. One of the two illustrations referred to by Lamarck in the description of *M. palmarosae* Lamarck, 1822.
- B. Figure 276 from Buonanni, 1681. One of the two illustrations referred to by Lamarck in the description of *M. palmarosae* Lamarck, 1822. Some, but not all, of the figures in Buonanni are reversed.
- C. *Triplex foliatus* Perry, 1810, *Arcana* Plate XXIII. See in text under genus *Triplex* and also under species *T. foliatus* and *T. rosaria*.
- D. *Triplex rosaria* Perry, 1811, *Conchology* Plate 6, figure 3. See in text under genus *Triplex* and also under species *T. foliatus* and *T. rosaria*.

The fact that the Lister figure is an almost exact copy of the Buonanni figure is the subject of a paper now in progress.





About the author



Richard E. Petit has been an active amateur malacologist and paleontologist for many years. His particular interests are in the gastropod superfamily Cancellarioidea and the molluscan literature. In connection with this hobby he has visited most major American museums, The Natural History Museum (London), the Royal Scottish Museum (Edinburgh), Muséum National d'Histoire Naturelle (Paris), Naturhistorisches Museum (Basel), National Science Museum (Tokyo & Ibaraki), and others. He is a Past-President of the American Malacological Society, Inc., and was made an Honorary Life Member in 1997. He served as a Trustee of the Paleontological Research Institution of which he is a Life Member. His publications include over 75 papers in various journals.