

Systematics of *Schinia chrysellus* (Grote) complex: Revised status of *Schinia alencis* (Harvey) with a description of two new species (Lepidoptera: Noctuidae: Heliothinae)

MICHAEL G. POGUE¹ AND CHARLES E. HARP^{2,3}

¹Systematic Entomology Laboratory, PSI, Agricultural Research Service, U. S. Department of Agriculture, c/o Smithsonian Institution, P.O. Box 37012, NMNH, MRC-168, Washington, DC, 20013-7012, USA
mpogue@sel.barc.usda.gov

²Research Associate: C. P. Gillette Museum of Arthropod Diversity, Colorado State University, Ft. Collins, CO.

³8834 West Quarto Ave., Littleton, CO, 80128, USA cehmoth@aol.com

Table of contents

Abstract	1
Introduction	2
Materials & methods	2
Key to species of the <i>Schinia chrysellus</i> complex	3
<i>Schinia chrysellus</i> (Grote)	4
<i>Schinia chryselloides</i> Pogue and Harp new species	10
<i>Schinia alencis</i> (Harvy) Revised Status	13
<i>Schinia ciliata</i> Smith	16
<i>Schinia rufocostulata</i> Pogue and Harp, new species	21
Acknowledgements	22
Literature Cited	22

Abstract

Based on morphological characters and host plant use, species related to *Schinia chrysellus* (Grote) are revised. *Schinia alencis* (Harvey) is considered a valid species, **revised status**. *Schinia chryselloides*, **new species** is closely allied with *S. chrysellus*, and *Schinia rufocostulata*, **new species** is related to *S. ciliata* Smith. Localities of collected adult moths and their asteraceous host plants are compared and illustrated with distribution maps. Adults and male and female genitalia are illustrated for each species.

Key words: systematics, genitalia, distribution, host plant, Asteraceae