

ZOOTAXA

1207

**Review of the Neotropical species of the genus *Stereodermus* Lacordaire,
1866 (Coleoptera: Brentidae, Stereodermini)**

ANTOINE MANTILLERI & ALESSANDRA SFORZI



Magnolia Press
Auckland, New Zealand

ANTOINE MANTILLERI & ALESSANDRA SFORZI
Review of the Neotropical species of the genus *Stereodermus* Lacordaire, 1866 (Coleoptera: Bren-tidae, Stereodermini)
(*Zootaxa* 1207)
64 pp.; 30 cm.
22 May 2006
ISBN 978-1-86977-008-2 (paperback)
ISBN 978-1-86977-009-9 (Online edition)

FIRST PUBLISHED IN 2006 BY

Magnolia Press
P.O. Box 41383
Auckland 1030
New Zealand
e-mail: zootaxa@mapress.com
<http://www.mapress.com/zootaxa/>

© 2006 Magnolia Press

All rights reserved.

No part of this publication may be reproduced, stored, transmitted or disseminated, in any form, or by any means, without prior written permission from the publisher, to whom all requests to reproduce copyright material should be directed in writing.

This authorization does not extend to any other kind of copying, by any means, in any form, and for any purpose other than private research use.

ISSN 1175-5326 (Print edition)
ISSN 1175-5334 (Online edition)

Review of the Neotropical species of the genus *Stereodermus* Lacordaire, 1866 (Coleoptera: Brentidae, Stereodermini)

ANTOINE MANTILLERI¹ & ALESSANDRA SFORZI²

¹Muséum national d'Histoire naturelle, Département Systématique et Evolution, USM 601, Entomologie, C.P. 50, 45 rue Buffon, F-75231 Paris cedex 05, France

²Museo di Storia Naturale, Sezione di Zoologia "La Specola", Università degli Studi di Firenze, via Romana 17, I-50125 Florence, Italy

Table of contents

Abstract	4
Résumé	4
Introduction	5
Material and methods	5
<i>Stereodermus</i> Lacordaire, 1866	6
1. <i>Stereodermus pygmaeus</i> (Gyllenhal, 1833)	9
2. <i>Stereodermus fessus</i> Kleine, 1927	13
3. <i>Stereodermus breviceps</i> Sharp, 1895	15
4. <i>Stereodermus chontalensis</i> Sharp, 1895	17
5. <i>Stereodermus similis</i> n. sp.	19
6. <i>Stereodermus jonathani</i> Mantilleri, 2004	21
7. <i>Stereodermus latirostris</i> Sharp, 1895	23
8. <i>Stereodermus eliseannaee</i> Mantilleri, 2004	26
9. <i>Stereodermus exilis</i> Suffrian, 1870	27
10. <i>Stereodermus maelae</i> Mantilleri, 2004	29
11. <i>Stereodermus paulettae</i> n. sp.	31
12. <i>Stereodermus nigriceps</i> Sharp, 1895	32
13. <i>Stereodermus denisi</i> n. sp.	33
14. <i>Stereodermus filum</i> Sharp, 1895	35
15. <i>Stereodermus attenuatus</i> n. sp.	36
16. <i>Stereodermus dentipennis</i> Sharp, 1895	38
17. <i>Stereodermus barclayi</i> Mantilleri, 2004	40
18. <i>Stereodermus barbirostris</i> Sharp, 1895	42
19. <i>Stereodermus nathaliae</i> Mantilleri, 2004	44
20. <i>Stereodermus imparicostatus</i> n. sp.	45
21. <i>Stereodermus dentipes</i> Sharp, 1895	46
22. <i>Stereodermus olivieri</i> Mantilleri, 2004	48

23. <i>Stereodermus calvus</i> Sharp, 1895	50
24. <i>Stereodermus explanatus</i> n. sp.....	52
25. <i>Stereodermus zunilensis</i> Sharp, 1895	54
26. <i>Stereodermus puncticollis</i> Sharp, 1895	56
27. <i>Stereodermus carinatus</i> Sharp, 1895	57
28. <i>Stereodermus godmani</i> Sharp, 1895	58
Discussion	60
Acknowledgments	64
References	64

Abstract

The genus *Stereodermus* is represented both in the Neotropical and Oriental regions. The 32 described neotropical species of the genus *Stereodermus* Lacordaire, 1866 are reviewed and an identification key and distributional maps are provided. Two new synonymies are proposed: *Stereodermus longiceps* Sharp, 1895 = *S. pygmaeus* (Gyllenhal, 1833), **n. syn.** and *S. mitratus* Sharp, 1895 = *S. dentipennis* Sharp, 1895, **n. syn.** All known species are redescribed and illustrated; six are new to science: *S. attenuatus* **n. sp.** (type locality: Peru, Iquitos), *S. denisi* **n. sp.** (type locality: Costa Rica, Provincia Puntarenas, Acosa, Estación Sirena), *S. explanatus* **n. sp.** (type locality: Costa Rica, Provincia Guanacaste, Volcan Cacao, Estacion Cacao), *S. imparicostatus* **n. sp.** (type locality: French Guyana, Roura Mountain, PK 41.1), *S. paulettae* **n. sp.** (type locality: Costa Rica, Provincia Puntarenas, Estación Sirena) and *S. similis* **n. sp.** (type locality: Brazil, Nova-Friburgo).

Key words: Brentidae, *Stereodermus*, neotropical, review, synonymies, new species, identification key

Résumé

Révision des espèces néotropicales du genre *Stereodermus* Lacordaire, 1866 (Coleoptera, Brentidae, Stereodermini). Le genre *Stereodermus* est représenté à la fois dans les régions Néotropicales et Orientales. Les 32 espèces néotropicales décrites du genre *Stereodermus* Lacordaire, 1866, sont révisées; une clef d'identification et des cartes de répartition sont proposées. Deux nouvelles synonymies sont établies: *Stereodermus longiceps* Sharp, 1895 = *S. pygmaeus* (Gyllenhal, 1833), **n. syn.** and *S. mitratus* Sharp, 1895 = *S. dentipennis* Sharp, 1895, **n. syn.** Toutes les espèces connues sont redécrisées et illustrées; six sont nouvelles pour la science: *S. attenuatus* **n. sp.** (localité type: Perou, Iquitos), *S. denisi* **n. sp.** (localité type: Costa Rica, Provincia Puntarenas, Acosa, Estación Sirena), *S. explanatus* **n. sp.** (localité type: Costa Rica, Provincia Guanacaste, Volcan Cacao, Estacion Cacao), *S. imparicostatus* **n. sp.** (localité type: Guyane Française, Montagne de Roura, PK 41.1), *S. paulettae* **n. sp.** (localité type: Costa Rica, Provincia Puntarenas, Estación Sirena), et *S. similis* **n. sp.** (localité type: Brésil, Nova-Friburgo).