

Dactylogyrids (Monogenoidea, Polyonchoinea) parasitizing the gills of snappers (Perciformes, Lutjanidae): Proposal of *Haliotrematoides* n. gen. and descriptions of new and previously described species from marine fishes of the Red Sea, the eastern and Indo-west Pacific Ocean, Gulf of Mexico and Caribbean Sea

DELANE C. KRITSKY¹, YANG TINGBAO² & SUN YUAN³

¹Department of Health and Nutrition Sciences, Campus Box 8090, Idaho State University, Pocatello, Idaho 83209, U.S.A.

E-mail: kritdela@isu.edu

²Research Center for Parasitic Organisms and State Key Laboratory of Biocontrol, School of Life Sciences, Sun Yat-sen University, Guangzhou, 510275, Peoples Republic of China. E-mail: lsyytb@mail.sysu.edu.cn

³School of Life Sciences, South China Agricultural University, Guangzhou 510642, Guangdong Province, Peoples Republic of China

Table of contents

Abstract	2
Introduction	3
Materials and methods	4
Results	6
Taxonomic revision	7
Class Monogenoidea Bychowsky, 1937	7
Subclass Polyonchoinea Bychowsky, 1937	7
Order Dactylogyridea Bychowsky, 1937	7
Dactylogyridae Bychowsky, 1933	7
<i>Haliotrematoides</i> n. gen.	7
<i>Haliotrematoides hatzenbuehlerae</i> n. sp.	9
<i>Haliotrematoides tainophallus</i> Kritsky & Justine n. sp.	12
<i>Haliotrematoides noncalcaroides</i> n. sp.	13
<i>Haliotrematoides brachyflagellocirrus</i> (Wang, Liu & Zhou, 2003) n. comb.	16
<i>Haliotrematoides noncalcaris</i> (Bychowsky & Nagibina, 1971) n. comb.	19
<i>Haliotrematoides potens</i> Kritsky & Justine n. sp.	20
<i>Haliotrematoides novaecaledoniae</i> Kritsky & Justine n. sp.	20
<i>Haliotrematoides patellacirrus</i> (Bychowsky & Nagibina, 1971) n. comb.	22
<i>Haliotrematoides nagibinae</i> n. sp.	25
<i>Haliotrematoides lanx</i> Kritsky & Justine n. sp.	25
<i>Haliotrematoides longitubocirrus</i> (Bychowsky & Nagibina, 1971) n. comb.	27
<i>Haliotrematoides calcaris</i> (Bychowsky & Nagibina, 1971) n. comb.	30
<i>Haliotrematoides shenzhenensis</i> (Wang, Liu & Zhou, 2003) n. comb.	30
<i>Haliotrematoides spinatus</i> Kritsky & Mendoza-Franco n. sp.	31
<i>Haliotrematoides gracilithamus</i> (Zhukov, 1976) n. comb.	32
<i>Haliotrematoides overstreeti</i> Kritsky & Bullard n. sp.	33
<i>Haliotrematoides zhukovi</i> Kritsky & Bullard n. sp.	35
<i>Haliotrematoides plectridium</i> Kritsky & Mendoza-Franco n. sp.	37
<i>Haliotrematoides guttati</i> (García-Vargas, Fajer-Ávila & Lamothe-Argumedo, 2008) n. comb.	37
<i>Haliotrematoides mastigion</i> Kritsky & Mendoza-Franco n. sp.	39
<i>Haliotrematoides magnigastrohamus</i> (Zhukov, 1976) n. comb.	40

<i>Haliotrematooides cornigerum</i> (Zhukov, 1976) n. comb.	42
<i>Haliotrematooides heteracantha</i> (Zhukov, 1976) n. comb.	42
<i>Haliotrematooides longihamus</i> (Zhukov, 1976) n. comb.	43
<i>Haliotrematooides</i> species from non-Lutjanid hosts	43
<i>Haliotrematooides caesionis</i> (Yamaguti, 1953) n. comb.	43
<i>Haliotrematooides isolens</i> n. sp.	45
<i>Haliotrematooides mediohamides</i> Kritsky & Mendoza-Franco n. sp.	45
<i>Haliotrematooides prolixohamus</i> Kritsky & Mendoza-Franco n. sp.	46
<i>Haliotrematooides mediohamus</i> (Zhukov, 1983) n. comb.	48
<i>Haliotrematooides parvicirrus</i> (Zhukov, 1983) n. comb.	48
Discussion	48
Acknowledgments	49
References	50

Abstract

Twenty six of 28 species of snappers (Lutjanidae), examined for dactylogyrids (Monogenoidea) from the Red Sea, the Indo-west and eastern Pacific Ocean, the Gulf of Mexico and the Caribbean Sea, were infected: 18 species with *Euryhaliotrema* spp., 17 with species of *Haliotrematooides* n. gen. and eight with species of undetermined dactylogyrid genera. Two species of fusiliers (Caesionidae) and one species of porgy (Sparidae) were also examined and found positive for species of *Haliotrematooides*. *Haliotrematooides* n. gen., supported by both 18S rDNA and 28S rDNA analyses, was proposed for 15 new and 15 previously described species of *Haliotrema* (s. l.) from lutjanid, caesionid and sparid hosts: *Haliotrematooides hatzenbuehlerae* n. sp., *Haliotrematooides brachyflagellocirrus* n. comb., *Haliotrematooides caesionis* n. comb., *Haliotrematooides calcaris* n. comb., *Haliotrematooides cornigerum* n. comb., *Haliotrematooides gracilhamus* n. comb., *Haliotrematooides guttati* n. comb., *Haliotrematooides heteracantha* n. comb., *Haliotrematooides isolens* n. sp., *Haliotrematooides lanx* n. sp., *Haliotrematooides longihamus* n. comb., *Haliotrematooides longitubocirrus* n. comb., *Haliotrematooides magnigastrohamus* n. comb., *Haliotrematooides mastigion* n. sp., *Haliotrematooides mediohamides* n. sp., *Haliotrematooides mediohamus* n. comb., *Haliotrematooides nagibinae* n. sp., *Haliotrematooides noncalcaris* n. comb., *Haliotrematooides noncalcaroides* n. sp., *Haliotrematooides novacaledoniae* n. sp., *Haliotrematooides overstreeti* n. sp., *Haliotrematooides parvicirrus* n. comb., *Haliotrematooides patellacirrus* n. comb., *Haliotrematooides plectridium* n. sp., *Haliotrematooides potens* n. sp., *Haliotrematooides prolixohamus* n. sp., *Haliotrematooides shenzhenensis* n. comb., *Haliotrematooides spinatus* n. sp., *Haliotrematooides tainophallus* n. sp., *Haliotrematooides zhukovi* n. sp. The new genus was characterized by dactylogyrids having 1) an unique hook distribution with pairs 1, 2, 3, 7 on the peduncle or trunk of the body, pair 5 associated with the point and distal shaft of the ventral anchors, and pairs 4 and 6 located at the level of the anchor bases or shifted anteriorly to the union of the haptor and peduncle; 2) dorsal anchors lacking well-developed deep roots and having perforated bases, elongate relatively straight shafts, and superficial grooves on the distal portions of the shafts and/or points; 3) ventral anchors with developed basal roots, an elongate shaft, and superficial grooves along the distal portions of the shafts and/or points; 4) a ventral bar usually with two submedial pockets along the anterior margin; and 5) a copulatory complex lacking an accessory piece. New host and locality records for some previously described species were reported.

Key words: Monogenoidea, Dactylogyridae, *Haliotrematooides* n.gen., Lutjanidae, *Lutjanus* spp., *Caesio cuning*, *Pterocaesio marri*, *Calamus brachysomus*, *Haliotrematooides hatzenbuehlerae* n. sp., *Haliotrematooides brachyflagellocirrus* n. comb., *Haliotrematooides caesionis* n. comb., *Haliotrematooides calcaris* n. comb., *Haliotrematooides cornigerum* n. comb., *Haliotrematooides gracilhamus* n. comb., *Haliotrematooides guttati* n. comb., *Haliotrematooides heteracantha* n. comb., *Haliotrematooides isolens* n. sp., *Haliotrematooides lanx* n. sp., *Haliotrematooides longihamus* n. comb., *Haliotrematooides longitubocirrus* n. comb., *Haliotrematooides magnigastrohamus* n. comb., *Haliotrematooides mastigion* n. sp., *Haliotrematooides mediohamides* n. sp., *Haliotrematooides mediohamus* n. comb., *Haliotrematooides nagibinae* n. sp., *Haliotrematooides noncalcaris* n. comb., *Haliotrematooides noncalcaroides* n. sp., *Haliotrematooides novacaledoniae* n. sp., *Haliotrematooides overstreeti* n. sp., *Haliotrematooides parvicirrus* n. comb., *Haliotrematooides patellacirrus* n. comb., *Haliotrematooides plectridium* n. sp., *Haliotrematooides potens* n. sp., *Haliotrematooides prolixohamus* n. sp., *Haliotrematooides shenzhenensis* n. comb., *Haliotrematooides spinatus* n. sp., *Haliotrematooides tainophallus* n. sp., *Haliotrematooides zhukovi* n. sp.