

## On Recent species of *Tonnacypris* Diebel & Pietrzeniuk, 1975 (Crustacea, Ostracoda), with new species descriptions from Mongolia

THIJS VAN DER MEEREN<sup>1,2</sup>, YONDON KHAND<sup>3</sup> & KOEN MARTENS<sup>1,2</sup>

<sup>1</sup>Ghent University, Department of Biology, Limnology unit, K. L. Ledeganckstraat 35, B-9000 Gent, Belgium

<sup>2</sup>Royal Belgian Institute for Natural Sciences, Freshwater Biology, Vautierstraat 29, B-1000 Brussels, Belgium

<sup>3</sup>Mongolian Academy of Sciences, Paleontological Centre, Enkhtaivan Avenue 63, Ulaanbaatar 210351, P.O.B. 260, Mongolia

### Table of contents

Abstract .....	1
Introduction .....	1
Material and methods .....	2
Taxonomy .....	4
Genus <i>Tonnacypris</i> Diebel & Pietrzeniuk, 1975 .....	4
<i>Tonnacypris estonica</i> (Järvekülg, 1960) .....	5
<i>Tonnacypris tonnensis</i> (Diebel & Pietrzeniuk, 1975) .....	12
<i>Tonnacypris edlundi</i> n. sp. .....	16
<i>Tonnacypris mazepovae</i> n. sp. ....	22
<i>Tonnacypris glacialis</i> (Sars, 1890) .....	28
<i>Tonnacypris lutaria</i> (Koch, 1838) .....	30
Discussion .....	33
Acknowledgements .....	39
References .....	40

### Abstract

The ostracod species *Tonnacypris estonica* (Järvekülg, 1960), *T. tonnensis* (Diebel & Pietrzeniuk, 1975), *T. edlundi* n. sp. and *T. mazepovae* n. sp. are here reported from Mongolia. Redescriptions of females of *T. glacialis* (Sars, 1890) and males of *T. lutaria* (Koch, 1838) are included. The revised generic diagnosis maintains the main character of the anteroventral peg on the calcified part of the inner lamella of the left valve, but this character has a significant level of morphological variability between species of this genus. This is also true for several characters of soft parts, which results in a broader concept for the genus. Different ecological strategies and reproductive modes are present. New records also change the evolutionary and zoogeographical scenario of the genus.

**Key words:** taxonomy, Ostracoda, morphology, zoogeography

### Introduction

There is an increased interest in Asian ostracod faunas, as some ostracod lineages that became extinct in Europe during the Pleistocene glaciations might have survived in the East-Palaearctic. Certain so-called ‘Lazarus-species’ were recently discovered alive in Asia, for example *Cyclocypris diebeli* Absolon, 1973 (Matzke-Karasz *et al.* 2004), *Cryptocandona brehmi* Klie, 1934 (Namiotko & Danielopol 2002), *Ilyocypris salebrosa* Stepanaitys, 1960 (Mischke & Schudack 2001), *Tonnacypris tonnensis* Diebel & Pietrzeniuk, 1975 and *T. convexa* Diebel & Pietrzeniuk, 1975 (Schornikov 2007). Moreover, in the last couple of years the