



Synopsis of the genus *Sangaris* Dalman in South America (Coleoptera, Cerambycidae), with description of two new species and new combinations

MIGUEL A. MONNÉ^{1,2} & MARCELA L. MONNÉ^{1,2}

¹ Museu Nacional, Universidade Federal do Rio de Janeiro. Quinta da Boa Vista, São Cristóvão, 20940-040 Rio de Janeiro, Rio de Janeiro, Brazil. E-mail: monne@uol.com.br

² CNPq fellow.

Abstract

A synopsis of the South American species of *Sangaris* Dalman, 1823 is presented. *Sangaris luteonotata* Monné & Monné **sp. nov.** and *S. ordinale* Monné & Monné **sp. nov.** are described from Ecuador. *Sangaris luctuosa* (Pascoe, 1859) **comb. nov.** and *Sangaris obtusicarinata* (Zajciw, 1962) **comb. nov.** are transferred from *Colobothea* Lepeletier & Audinet-Serville, 1825. New distributional data are given for *S. trifasciata* Melzer, 1928, *S. zikani* Melzer, 1931, and *S. luctuosa* (Pascoe, 1859). A key to the South American species is provided.

Key words: Colobotheini, Lamiinae, Neotropical, taxonomy, new species, new combination

Resumo

Sinopse das espécies sul-americanas de *Sangaris* Dalman, 1823 é apresentada. *Sangaris luteonotata* Monné & Monné **sp. nov.** e *S. ordinale* Monné & Monné **sp. nov.** são descritas do Equador. *Sangaris luctuosa* (Pascoe, 1859) **comb. nov.** e *Sangaris obtusicarinata* (Zajciw, 1962) **comb. nov.** são transferidas de *Colobothea* Lepeletier & Audinet-Serville, 1825. Novos dados de distribuição são fornecidos para *S. trifasciata* Melzer, 1928, *S. zikani* Melzer, 1931 e *S. luctuosa* (Pascoe, 1859). Chave para espécies sul-americanas é adicionada.

Palavras-chave. Colobotheini, Lamiinae, neotropical, taxonomia, espécies novas, novas combinações

Introduction

The Neotropical tribe of longhorn beetles Colobotheini Thomson, 1860 is represented by 13 genera: *Apechthes* Thomson, 1861 (2 species from Mexico), *Calliphenges* Waterhouse, 1880 (1 species from Ecuador and Peru), *Carneades* Bates, 1869 (13 species distributed from Mexico to the Amazon), *Carterica* Pascoe, 1858 (8 species distributed from Central America to Bolivia), *Cathexis* Thomson, 1860 (2 species from Brazil), *Colobothea* Lepeletier & Audinet-Serville, 1825 (117 species, widely distributed in the neotropics), *Colobothina* Hovore, 1989 (1 species from Central America), *Hilobothea* Monné & Martins, 1979 (2 species from Brazil), *Nodubothea* Monné & Monné, 2008 (2 species from Mexico and northern South America), *Priscilla* Thomson, 1864 [1 species, with a wide distribution from Honduras and Nicaragua to Brazil (Amazonas, Mato Grosso), Peru and Bolivia], *Sangaris* Dalman, 1823 (21 species distributed from Central America to Argentina, plus 2 described herein and another 2 transferred from *Colobothea*), *Sparna* Thomson, 1864 (7 species, mostly with Andean distributions), and *Sympleurotis* Bates, 1881 (4 species from Mexico and Central America).

The classification of tribes within the subfamilies of Cerambycidae has not been evaluated