

The slim mute crickets, Euscyrtinae (Orthoptera: Grylloidea; Gryllidae; Euscyrtinae) of Taiwan with three new species and lectotype designations of three of Shiraki's species

JENG-TZE YANG¹ & CHING-SHAN YANG

Department of Entomology, National Chung Hsing University, Taichung, 40227, Taiwan

¹Corresponding author. E-mail:jtyang@dragon.nchu.edu.tw

Table of contents

Abstract	1
Introduction	2
Systematics	2
Subfamily Euscyrtinae Gorochov, 1988	2
Key to genera of the Euscyrtinae from Taiwan	2
Genus <i>Euscyrtus</i> Guerin 1884	3
Key to species of <i>Euscyrtus</i> from Taiwan	3
<i>Euscyrtus hemelytrus</i> (Haan, 1842)	3
<i>Euscyrtus japonicus</i> Shiraki, 1930	8
<i>Euscyrtus ogatai</i> Shiraki, 1930	12
Genus <i>Patiscus</i> Stål, 1877	12
Key to species of <i>Patiscus</i> from Taiwan	13
<i>Patiscus malayanus</i> Chopard, 1969	13
<i>Patiscus cephalotes</i> (Saussure, 1877)	17
Genus <i>Beybienkoana</i> Gorochov, 1988	22
Key to species of <i>Beybienkoana</i> from Taiwan	22
<i>Beybienkoana formosana</i> (Shiraki, 1930)	22
<i>Beybienkoana Karnyi</i> (Shiraki, 1930)	27
<i>Beybienkoana splendida</i> Yang and Yang, n. sp.	31
<i>Beybienkoana luteola</i> n. sp.	36
<i>Beybienkoana gregaria</i> n. sp.	40
References	45

Abstract

Three genera and five species of Taiwanese Euscyrtinae possess developed tegmina but produce no sound at all. In this paper, we report on 10 species in 3 genera including 3 new species, i.e., *Beybienkoana splendida* n. sp., *B. luteola* n. sp. and *B. gregaria* n. sp. Lectotypes of three Shiraki's species, i.e., *Euscyrtus japonicus* Shiraki 1930, *Beybienkoana formosana* (Shiraki, 1930) and *Beybienkoana karnyi* (Shiraki, 1930) are designated. The types are deposited in the collection of Department of Entomology, National Taiwan University. In addition, *Euscyrtus japonicus* Shiraki, 1930 and *Patiscus malayanus* Chopard, 1969, are recorded from Taiwan for the first time. The morphology, including the ultra fine structure of metanotal gland, and ecology, including habitats and host plants, are applied in the taxonomy of this subfamily. The structure of the metanotal gland is described in detail using the Scanning Electronic Microscope. In this paper some taxonomic characters are used for the first time in this group. The definition of Euscyrtinae emended with the structure of metanotal glands presented as new character for subfamily definition. A key to genera and to species of Euscyrtinae from Taiwan provided in this paper as well.

Key words: Euscyrtinae, new species, lectotype, emendation, dorsal gland, Taiwan