

<http://dx.doi.org/10.11646/zootaxa.3637.1.4>
<http://zoobank.org/urn:lsid:zoobank.org:pub:9852C0E8-283F-4F8A-8A94-81911F86A698>

A new species of *Balera* Young and redescription of *Habralebra gillettei* Young (Hemiptera, Cicadellidae, Typhlocybinae) from Viçosa, Minas Gerais State, Brazil

LUCI BOA NOVA COELHO^{1,3}, JORGE LUIZ NESSIMIAN¹ & ELIDIOMAR RIBEIRO DA-SILVA²

¹Laboratório de Entomologia, Departamento de Zoologia, Instituto de Biologia, Universidade Federal do Rio de Janeiro. Caixa Postal 68044, CEP 21944-970, Rio de Janeiro, RJ, Brazil

²Laboratório de Insetos Aquáticos, Departamento de Zoologia, Instituto de Biociências, UNIRIO. CEP 22290-240, Rio de Janeiro, RJ, Brazil

³Corresponding author. E-mail: lucibncoelho@ufrj.br

Abstract

Balera fiuzai sp. nov. is described based on male specimens obtained from light trap collections at Mata do Paraíso, Municipality of Viçosa, Minas Gerais State, Brazil. An identification key to the species of *Balera* is given. Additionally, *Habralebra gillettei* Young is redescribed from the same locality. These specimens of *H. gillettei* differ somewhat from previously described specimens, but are considered intra-specific variations. Both genera, *Balera* Young and *Habralebra* Young, are recorded from Minas Gerais State for the first time.

Key words: Alebrini, Atlantic Forest, diversity, identification key, leafhoppers, taxonomy

Resumo

Balera fiuzai sp. nov. é descrita com base em machos obtidos a partir de coletas com armadilha luminosa, realizadas na Mata do Paraíso, município de Viçosa, Minas Gerais, Brasil. Uma chave de identificação para as espécies de *Balera* é apresentada. Adicionalmente, *Habralebra gillettei* Young é redescrita da mesma localidade. Esses exemplares de *H. gillettei* apresentam diferenças morfológicas em relação às descrições anteriores, sendo consideradas aqui como variações intra-específicas. Os gêneros *Balera* Young e *Habralebra* Young são, pela primeira vez, registrados para o Estado de Minas Gerais.

Palavras-chave: Alebrini, Mata Atlântica, diversidade, chave de identificação, cigarrinhas, taxonomia

Introduction

The Neotropical genus *Balera* Young was described by Young (1952) based on type species *Dikraneura pellucida* Osborn. Young (1957a) transferred *Empoasca emarginata* Osborn to *Balera*. He later (Young 1957b) presented a revised diagnosis for the genus, described two new species, *B. caraguatae* and *B. pusilla*, and provided a key to species. Ruppel (1959) added *B. bracata*. Freytag (1992) described *B. myersi*, *B. obtusa*, *B. ecuadora*, *B. napoensis*, and *B. plagata*, redescribed *B. caraguatae*, and included them in a new identification key. Dworakowska (1994) described an additional species, *B. signata*. Thus, prior to the present study, *Balera* included eleven species, recorded from Trinidad and Tobago, Panama, Ecuador, Bolivia, Colombia, Venezuela, and Brazil. Only *B. caraguatae* Young is recorded from Brazil (Mato Grosso and Rondônia States) (Young 1957b; Zanol & Menezes 1982; Freytag 1992). In this work a new species of *Balera* is described from the Atlantic Forest of Southeastern Brazil (Minas Gerais State) and a key for identification of the twelve described species is included.

Habralebra gillettei Young was described by Young (1957b) to accommodate two paratypes of *Alebra*