

<http://dx.doi.org/10.11646/zootaxa.3737.4.8>  
<http://zoobank.org/urn:lsid:zoobank.org:pub:B12F5D90-E13A-4ACA-8583-817DB0F6FD18>

## Redescription of *Alatina alata* (Reynaud, 1830) (Cnidaria: Cubozoa) from Bonaire, Dutch Caribbean

CHERYL LEWIS<sup>1,2,8</sup>, BASTIAN BENTLAGE<sup>2,3</sup>, ANGEL YANAGIHARA<sup>4</sup>, WILLIAM GILLAN<sup>5</sup>, JOHAN VAN BLERK<sup>6</sup>, DANIEL P. KEIL<sup>1</sup>, ALEXANDRA E. BELY<sup>7</sup> & ALLEN G. COLLINS<sup>2</sup>

<sup>1</sup>*Biological Sciences Graduate Program, University of Maryland, College Park, MD 20742, USA; dkeil@umd.edu; clames1@umd.edu*

<sup>2</sup>*National Systematics Laboratory, National Museum of Natural History, MRC-153, Smithsonian Institution, P.O. Box 37012, Washington, DC 20013-7012, USA. E-mail: collinsa@si.edu*

<sup>3</sup>*Department of Cell Biology and Molecular Genetics, University of Maryland, College Park, MD 20742, USA.  
E-mail: bastian.bentlage@gmail.com*

<sup>4</sup>*Department of Tropical Medicine, Medical Microbiology and Pharmacology John A. Burns School of Medicine, University of Hawai'i at Manoa, Honolulu, Hawai'i 96822, USA. E-mail: ayanagih@hawaii.edu*

<sup>5</sup>*Palm Beach County (FL) Schools, Boynton Beach Community High School, 4975 Park Ridge Boulevard, Boynton Beach, FL, 33426, USA. E-mail: william.gillan@palmbeachschools.org*

<sup>6</sup>*Kralendijk, Bonaire, The Netherlands. E-mail: blerkbonaire@hotmail.com*

<sup>7</sup>*Biology Department, University of Maryland, College Park, MD 20742, USA. E-mail: abely@umd.edu*

<sup>8</sup>*Corresponding author. E-mail: clames1@umd.edu*

### Abstract

Here we establish a neotype for *Alatina alata* (Reynaud, 1830) from the Dutch Caribbean island of Bonaire. The species was originally described one hundred and eighty three years ago as *Carybdea alata* in *La Centurie Zoologique*—a monograph published by René Primevère Lesson during the age of worldwide scientific exploration. While monitoring monthly reproductive swarms of *A. alata* medusae in Bonaire, we documented the ecology and sexual reproduction of this cubozoan species. Examination of forty six *A. alata* specimens and additional archived multimedia material in the collections of the National Museum of Natural History, Washington, DC revealed that *A. alata* is found at depths ranging from surface waters to 675 m. Additional studies have reported it at depths of up to 1607 m in the tropical and subtropical Atlantic Ocean. Herein, we resolve the taxonomic confusion long associated with *A. alata* due to a lack of detail in the original description and conflicting statements in the scientific literature. A new cubozoan character, the velarial lappet, is described for this taxon. The complete description provided here serves to stabilize the taxonomy of the second oldest box jellyfish species, and provide a thorough redescription of the species.

**Key words:** Neotype, *Carybdea alata*, Carybdeida, cubomedusae, box jellyfish, Atlantic Ocean, aggregation, sexual reproduction, deep-sea, taxonomy

### Introduction

The winged box jellyfish *Alatina alata* (Cnidaria: Cubozoa: Carybdeida: Alatinidae) has appeared in the scientific literature (as *Carybdea alata*) many times (see Gershwin 2005 for list of citations) since its original description (by Reynaud) and illustration (by Prêtre; reproduced herein as Fig 1a) appeared in the extensive monograph by French naturalist René Primevère Lesson in 1830. Reynaud's brief description gave no details about the collection events or the whereabouts of the specimen (or possibly multiple specimens), stating only that this box jellyfish “lives in the Atlantic Ocean”. Almost a century later, Bigelow (1918, 1938) redescribed *A. alata* (as *C. alata*) (original drawings reproduced herein as Fig 1b–g) from Jamaica and Cuba, effectively clarifying its identity in the Atlantic Ocean and the associated Gulf of Mexico and Caribbean Sea, where the name has since been in common usage (see additional reports by Arneson (1976) from Puerto Rico, Morandini (2003) from Brazil, Graham (1998, as *C. alata* var. *grandis*) and Larson *et al.* (1991) from the Gulf of Mexico). In the last decade, nine nominal species from

restaurant is located during the annual Jellyfish Jamboree on Bonaire, The Netherlands. Jonathan Lawley assisted with photographing preserved specimens. We are also thankful to the reviewers of an earlier draft of this manuscript who provided excellent advice for improvement, and to the Zootaxa editors.

## References

- Agassiz, L. (1862) *Contributions to the natural history of the United States of America. Vol. IV*. Little Brown, Boston, 380 pp., 19 pls. Available from: <http://www.biodiversitylibrary.org/item/54510#page/9/mode/1up> (accessed 2 Oct 2013)
- Arneson, A.C. (1976) Life history of *Carybdea alata* Reynaud, 1830 (Cubomedusae). *M.S. Thesis*, University of Puerto Rico, Mayaguez, Commonwealth of Puerto Rico, USA, 43 pp.
- Arneson, A.C. & Cutress, C.E. (1976) Life history of *Carybdea alata* Reynaud, 1830 (Cubomedusae). In: Mackie, G.O. (Eds.), *Coelenterate Ecology and Behavior*. Plenum Press, New York, NY, pp. 227–236.
- Bauchot, M.-L., Daget, J. & Bauchot, R. (1990) L'ichthyologie en France au début du XIX<sup>e</sup> siècle. L'Histoire naturelle des Poissons de Cuvier et Valenciennes. *Bulletin du Muséum national d'Histoire naturelle, Paris, section A*, supplément 12, 3–142.  
<http://dx.doi.org/10.3366/anh.1993.20.1.139a>
- Bentlage, B. (2010) *Carybdea alata* auct. (Cubozoa): rediscovery of the *Alatina grandis* type. *Zootaxa*, 2713, 52–54.
- Bentlage, B., Cartwright, P., Yanagihara, A.A., Lewis, C., Richards, G.S. & Collins, A.G. (2010) Evolution of box jellyfish (Cnidaria: Cubozoa), a group of highly toxic invertebrates. *Proceedings of the Royal Society B*, 277, 493–501.  
<http://dx.doi.org/10.1098/rspb.2009.1707>
- Bentlage, B. & Lewis, C. (2012) An illustrated key and synopsis of the families and genera of carybdeid box jellyfishes (Cnidaria: Cubozoa: Carybdeida), with emphasis on the “Irukandji family (Carukiidae). *Journal of Natural History*, 41–42, 2595–2620.  
<http://dx.doi.org/10.1080/00222933.2012.717645>
- Bigelow, H.B. (1918) Some Medusae and Siphonophora from the western Atlantic. *Bulletin of the Museum of Comparative Zoology of Harvard College*, 62, 363–442, pls 1–8. Available from: <http://www.biodiversitylibrary.org/item/21150#page/463/mode/1up> (accessed 14 Nov. 2013)
- Bigelow, H.B. (1938) Plankton of the Bermuda oceanographic expeditions. VII. Medusae taken during the years 1929 and 1930. *Zoologica. Scientific Contributions of the New York Zoological Society*, 23, 99–189.
- Collins, A.G., Bentlage, B., Gillan, W., Lynn, T.H., Morandini, A.C. & Marques, A.C. (2011) Naming the Bonaire banded box jelly, *Tamoya ohboya*, n. sp. (Cnidaria: Cubozoa: Carybdeida: Tamoyidae). *Zootaxa*, 2753, 53–68.
- De Meyer, K. (1997) Bonaire, Netherlands Antilles. Coastal Region and Small Island (CSI). *Environment and development in coastal regions and in small islands. Paper 3*, Available from <http://www.unesco.org/csi/pub/papers/demayer.htm> (accessed 2 Oct 2013)
- Gershwin, L. (2005) *Carybdea alata* auct. and *Manokia stiasnyi*, reclassification to a new family with description of a new genus and two new species. *Memoirs of the Queensland Museum*, 52, 501–523.
- Graham, W.M. (1998) First report of *Carybdea alata* var. *grandis* (Reynaud 1830) (Cnidaria: Cubozoa) from the Gulf of Mexico. *Gulf of Mexico Science*, 16, 1, 28–30.
- Haeckel, E. (1880) *System der Acraspeden – Zweite Hälfte des Systems der Medusen. Denkschriften der Medizinisch-Naturwissenschaftlichen Gesellschaft zu Jena*, Germany, 82 pp.
- Humann, P. & Deloach N. (2002) *Reef creature Identification: Florida, Caribbean, Bahamas* 2nd Ed. New World Publications Inc., 420 pp.
- ICZN (1999) *International Code of Zoological Nomenclature, 4th edition*. London: The International Commission on Zoological Nomenclature. Available from: <http://www.iczn.org/code> (accessed 23 September 2013)
- Kayal, E., Bentlage, B., Collins, A.G., Kayal, M., Pirro, S. & Lavrov, D.V. (2012) Evolution of linear mitochondrial genomes in Cnidaria. *Genome Biology and Evolution*, *Genome Biology and Evolution*, 4, 1, 1–12.  
<http://dx.doi.org/10.1093/gbe/evr123>
- Kramp, P.L. (1961) Synopsis of the Medusae of the World. *Journal of the Marine Biological Association of the United Kingdom*, 40, 1–469.  
<http://dx.doi.org/10.1017/s0025315400007347>
- Larson, R.J. (1976) Cubomedusa: feeding – functional morphology, behavior and phylogenetic Position. In: Mackie, G.O. (Eds.), *Coelenterate ecology and behavior*. Plenum Press, New York, pp. 237–245.
- Larson, R.J., Mills, C.E. & Harbison, G.R. (1991) Western Atlantic midwater hydrozoan and scyphozoan medusae: *in situ* studies using manned submersibles. *Hydrobiologia*, 216–217 (1),
- Lesson, R.P. (Ed.) (1830) *Centurie Zoologique*. Levrault, Paris, France, 244 pp.
- Lesson, R.P. (1837). *Prodrome d'une monographie des Méduses. Extraite d'une Histoire manuscrite des méduses, en 3 volumes 4<sup>e</sup> avec 200 planches coloriées, ouvrage entièrement terminé*, Rochefort, 63 pp., 200 pls.
- Lesson, R.P. (1843) *Historie naturelle des zoophytes. Acalèphes*. Encyclopédique de Roret, Paris, 663 pp.

- Linnaeus, C. (1758) *Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis*. Editio decima, reformata. Tomus I. Holmiae (Laurentii Salvii), i–iv, 824 pp.
- Mariscal, R.N. (1974) Nematocysts. In: Muscatine, L. & Lenhoff, H.M. (Eds.), *Coelenterate biology—reviews and new perspectives*. Academic Press, New York, pp. 129–178.
- Mayer, A.G. (1910) Medusae of the World. Vol. III. The Scyphomedusae. *Carnegie Institution of Washington Publication*, 109 (3), pp. 499–735
- Morandini, A.C. (2003) Deep-Sea medusae (Cnidaria: Cubozoa, Hydrozoa and Scyphozoa) from the coast of Bahia (Western south Atlantic, Brazil). *Mitteilungen aus dem hamburgischen zoologischen Museum und Institut*, 100, 13–25.
- Östman, C. (2000) A guideline to nematocyst nomenclature and classification, and some notes on the systematic value of nematocysts. In: Mills, C.E., Boero, F., Migotto A. & Gili, J.M. (Eds.), Trends in hydrozoan biology—IV. *Scientia Marina*, 64 (Supl. 1), 31–46  
<http://dx.doi.org/10.3989/scimar.2000.64s131>
- Reynaud, M. (1830) *Carybdea alata* n. sp. In: Lesson, R.P. (Ed), *Centurie Zoologique*. Levrault, Paris (France), pp. 95, pl. 33.
- Smith, D.R., Kayal, E., Yanagihara, A.A., Collins, A.G., Pirro, S. & Keeling, P.J. (2011) First complete mitochondrial genome sequence from a box jellyfish reveals a highly fragmented linear architecture and insights into telomere evolution. *Genome Biology and Evolution*, 4 (1), 52–58.  
<http://dx.doi.org/10.1093/gbe/evr127>
- Yanagihara, A.A., Kuroiwa, J.M.Y., Oliver, L.M., Chung, J.J., Kunkel, D.D. (2002) Ultrastructure of a novel eurytele nematocyst of *Carybdea alata* Reynaud (Cubozoa, Cnidaria). *Cell and Tissue Research*, 308, 307–318.  
<http://dx.doi.org/10.1007/s00441-002-0545-8>
- Yoshimoto, C.M. & Yanagihara, A.A. (2002) Cnidarian (coelenterate) envenomations in Hawai'i improve following heat application. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 96, 300–303.  
[http://dx.doi.org/10.1016/s0035-9203\(02\)90105-7](http://dx.doi.org/10.1016/s0035-9203(02)90105-7)