

Tersilochinae of Western Amazonia (Hymenoptera: Ichneumonidae). Genus *Stethantyx* Townes, part 1

ANDREY I. KHALAIM¹, ILARI E. SÄÄKSJÄRVI² & SANTIAGO BORDERA³

¹Zoological Institute, Russian Academy of Sciences, Universitetskaya nab. 1, St.Petersburg 199034, Russia. Facultad de Ingeniería y Ciencias, Universidad Autónoma de Tamaulipas, Cd. Victoria 87149, México. E-mail: ptera@mail.ru

²Zoological Museum, Section of Biodiversity and Environmental Sciences, Department of Biology, University of Turku, 20014 Turku, Finland. E-mail: ileesa@utu.fi

³Instituto de Investigación de Biodiversidad CIBIO, Universidad de Alicante, Ap. Corr. 99, 03080 Alicante, Spain.
E-mail: s.bordera@ua.es

Abstract

Sixteen species of the genus *Stethantyx* Townes having first and second sections of radius meeting at obtuse angle are recorded from Peru and Ecuador in Western Amazonia. Ten species, *S. amazonica* Khalaim & Sääksjärvi, sp. nov., *S. aurantia* Khalaim & Sääksjärvi, sp. nov., *S. juninensis* Khalaim & Bordera, sp. nov., *S. loretosa* Khalaim & Sääksjärvi, sp. nov., *S. neopropodeator* Khalaim & Sääksjärvi, sp. nov., *S. orellana* Khalaim & Sääksjärvi, sp. nov., *S. sinuata* Khalaim & Sääksjärvi, sp. nov., *S. tiputinia* Khalaim & Sääksjärvi, sp. nov., *S. trepida* Khalaim & Sääksjärvi, sp. nov. and *S. voluptaria* Khalaim & Sääksjärvi, sp. nov., are described as new. Six species, previously known only from Costa Rica, *S. alajuela* Khalaim & Broad, *S. heredia* Khalaim & Broad, *S. orosia* Khalaim & Broad, *S. propodeator* Khalaim & Broad, *S. puntarenasa* Khalaim & Broad and *S. sanjosea* Khalaim & Broad, are recorded from South America for the first time. A partial key to Western Amazonian species of *Stethantyx* is given.

Key words: Peru, Ecuador, Neotropics, new species, new records, taxonomy, key, fauna, parasitoids

Resumen

Se aborda el estudio de dieciséis especies del género *Stethantyx* Townes colectadas en la Amazonía peruana y ecuatoriana, caracterizadas por presentar la primera y segunda sección de la vena radial unidas en ángulo obtuso. Se describen diez nuevas especies: *S. amazonica* Khalaim & Sääksjärvi, sp. nov., *S. aurantia* Khalaim & Sääksjärvi, sp. nov., *S. juninensis* Khalaim & Bordera, sp. nov., *S. loretosa* Khalaim & Sääksjärvi, sp. nov., *S. neopropodeator* Khalaim & Sääksjärvi, sp. nov., *S. orellana* Khalaim & Sääksjärvi, sp. nov., *S. sinuata* Khalaim & Sääksjärvi, sp. nov., *S. tiputinia* Khalaim & Sääksjärvi, sp. nov., *S. trepida* Khalaim & Sääksjärvi, sp. nov. y *S. voluptaria* Khalaim & Sääksjärvi, sp. nov. Las especies, *S. alajuela* Khalaim & Broad, *S. heredia* Khalaim & Broad, *S. orosia* Khalaim & Broad, *S. propodeator* Khalaim & Broad, *S. puntarenasa* Khalaim & Broad y *S. sanjosea* Khalaim & Broad, conocidas únicamente de Costa Rica, son citadas por primera vez de Sudamérica. Asimismo se proporciona una clave parcial para la determinación de las especies del oeste de Amazonía.

Palabras clave: Perú, Ecuador, Región Neotropical, especies nuevas, nuevos registros, taxonomía, clave, fauna, parasitoides

Introduction

Stethantyx Townes, 1971 is a moderately large genus occurring almost entirely in the Neotropical region, where it is the most abundant and species-rich tersilochine genus. Only three species are known from the Nearctic region: the genotype *S. nearctica* Townes, 1971 occurring in U.S.A. and northern Mexico; *S. crassa* Horstmann, 2010 from

Material examined. Holotype female (ZMUT), Peru, Loreto reg., Iquitos Area, Allpahuayo, clayish soil, 17.XI–3.XII.1998, coll. I.E. Sääksjärvi *et al.*

Distribution. Peru (Loreto).

Acknowledgements

We are thankful to Gavin R. Broad (BMNH) and John Jennings (The University of Adelaide, Australia) for their important comments and corrections. This work was supported by the Russian Foundation for Basic Research (grants 10-04-00265 and 13-04-00026); Kone Foundation, Finland (grant for the research team of Ilari E. Sääksjärvi); and by project A/013484/07 from the Agencia Española de Cooperación Internacional para el Desarrollo (Ministerio de Asuntos Exteriores y de Cooperación, Spain). We thank the following institutions for help during the study: Universidad Nacional de la Amazonía Peruana, Peru; Instituto Nacional de Recursos Naturales, Peru; Instituto de Investigaciones de la Amazonía Peruana, Peru; Instituto Nacional de Investigación Agraria, Peru; Universidad Nacional Mayor de San Marcos, Peru; Universidad Nacional Agraria La Molina, Peru; the Natural History Museum of London, UK; Zoological Museum of the University of Turku, Finland; and the Smithsonian Institution, USA. We would also like to thank Terry Erwin for providing his Ecuadorean canopy specimens for study and Anu Veijalainen for preliminary sorting of this material into subfamilies, and Alexander Rodríguez Berriño and Mabel Alvarado from UNALM for collecting and sorting material belonging to the AECID project. The Ministry of Agriculture and the Ministry of Environment of Peru provided the collecting and export permits for the Peruvian samples. The Instituto de Investigaciones de la Amazonía Peruana provided vital support in the field studies conducted in the Loreto region.

References

- Blanchard, E.E. (1945) Dos nuevos Ichneumonidos, parásitos de *Listroderes* (Hym. Ichneum.). *Revista Sociedad Entomologica Argentina*, 12, 305–309.
- Clancy, D.W. (1969) A new parasite of the vegetable weevil in southern California. *Journal of Economic Entomology*, 62, 743–745.
- Gauld, I.D. (1984) An introduction to the Ichneumonidae of Australia. *Bulletin of the British Museum (Natural History) (Entomology)*, 895, 1–413.
- Graf, V. (1980) Ichneumofauna da regiao sudeste e sul do Brasil. II. Gênero *Stethantyx* (Tersilochinae, Ichneumonidae). *Dusenia*, 12 (4), 115–122.
- Horstmann, K. (2010) Revision of Nearctic Tersilochinae II. Genera *Allophrys* Förster, *Barycnemis* Förster, *Ctenophion* gen. nov., *Sathropterus* Förster, *Spinolochus* Horstmann and *Stethantyx* Townes (Hymenoptera, Ichneumonidae). *Spixiana*, 33, 73–109.
- Kerrich, G.J. (1961) A study of the Tersilochine parasites of vegetable weevils of the genus *Listroderes* (Hym., Ichneumonidae). *Eos*, 37, 497–503.
- Khalaim, A.I. (2011) Tersilochinae of South, Southeast and East Asia, excluding Mongolia and Japan (Hymenoptera: Ichneumonidae). *Zoosystematica Rossica*, 20, 96–148.
- Khalaim, A.I. & Broad, G.R. (2013) Tersilochinae (Hymenoptera: Ichneumonidae) of Costa Rica, part 2. Genera *Megalochus* gen. nov. and *Stethantyx* Townes. *Zootaxa*, 3693 (2), 221–266.
<http://dx.doi.org/10.11646/zootaxa.3693.2.8>
- Parker, H.L., Berry, P.A. & Silveira, A. (1950) Vegetable weevils and their natural enemies in Argentina and Uruguay. *United States Department of Agriculture, Technical Bulletin*, 1016, 1–28.
- Sääksjärvi, I.E., Haataja, S., Neuvonen, S., Gauld, I.D., Jussila, R., Salo, J. & Burgos, A.M. (2004) High local species richness of parasitic wasps (Hymenoptera: Ichneumonidae; Pimplinae and Rhyssinae) from the lowland rainforests of Peruvian Amazonia. *Ecological Entomology*, 29, 735–743.
<http://dx.doi.org/10.1111/j.0307-6946.2004.00656.x>
- Townes, H.K. (1969) The genera of Ichneumonidae, Part 1. *Memoirs of the American Entomological Institute*, 11, 1–300.
- Townes, H.K. (1971) The genera of Ichneumonidae, Part 4. *Memoirs of the American Entomological Institute*, 17, 1–372.
- Veijalainen, A., Sääksjärvi, I.E., Erwin, T., Gómez, I. & Longino J. (2013) Subfamily composition of Ichneumonidae (Hymenoptera) from western Amazonia: insights into diversity of tropical parasitoid wasps. *Insect Conservation and Diversity*, 6, 628–37.
<http://dx.doi.org/10.1111/j.1752-4598.2012.00185.x>
- Wilson, F. & Wearne, G.R. (1962) The introduction into Australia of parasites of *Listroderes obliquus* Klug. *Australian Journal of Agricultural Research*, 13, 249–257.
<http://dx.doi.org/10.1071/ar9620249>