

A new species of *Amazophrynellula* (Anura:Bufonidae) from the southwestern part of the Brazilian Guiana Shield

ROMMEL R. ROJAS^{1,2,7}, VINÍCIUS TADEU DE CARVALHO^{1,3,4}, MARCELO GORDO⁵, ROBSON W. ÁVILA⁶, IZENI PIRES FARIAS¹ & TOMAS HRBEK¹

¹Laboratório de Evolução e Genética Animal, Departamento de Biologia, Universidade Federal do Amazonas, Av. Gen. Rodrigo Octávio Jordão Ramos, 3000, CEP 69077-000, Manaus, AM, Brazil

²Programa de Pós-Graduação em Diversidade Biológica, Instituto de Ciências Biológicas, Universidade Federal do Amazonas, Av. Gen. Rodrigo Octávio Jordão Ramos, 3000, CEP 69077-000, Manaus, AM, Brazil

³Programa de Pós-Graduação em Biodiversidade e Biotecnologia, Instituto de Ciências Biológicas, Universidade Federal do Amazonas, Av. Gen. Rodrigo Octávio Jordão Ramos, 3000, CEP 69077-000, Manaus, AM, Brazil

⁴Instituto Nacional de Pesquisas da Amazônia, Coleção de Anfíbios e Répteis, Campus II, Av. André Araújo, 2936, Aleixo, CEP 69060-001, Manaus, AM, Brazil

⁵Departamento de Biologia, ICB, Universidade Federal do Amazonas, Av. Gen. Rodrigo Octávio Jordão Ramos, 3000, CEP 69077-000, Manaus, AM, Brazil

⁶Universidade Regional do Cariri, Centro de Ciências Biológicas e da Saúde, Departamento de Ciências Biológicas, Campus do Pimenta, Rua Cel. Antônio Luiz, 1161, Bairro do Pimenta, CEP 63105-100, Crato, CE, Brazil

⁷Corresponding author. E-mail: rommelrojaszamora@yahoo.es

Abstract

Amazophrynellula is a genus of the family Bufonidae, currently represented by three species. The type species of the genus, *Amazophrynellula minuta*, however, is a complex of species occurring throughout the Amazonian biome. This group remains problematic taxonomically; the difficulty lies principally in the lack of diagnostic characters in the original description of *A. minuta*, the lack of molecular data and refined taxonomic comparison of individuals through its wide distribution. We describe a new species of the genus *Amazophrynellula*, distributed in the southwestern part of the Guiana Shield of Brazil based on a series of morphological and molecular characters. The new species differs from others of the genus by presenting a slightly truncated triangular snout, ventral texture covered by many fine granules, white belly covered with black spots and by fifteen molecular autapomorphies in the 16S rDNA fragment. Uncorrected p-distances of a fragment of the 16S mitochondrial rDNA gene revealed high divergence among other *Amazophrynellula* species (9%–14%). Additionally we provide a new diagnosis of topotypic material of *Amazophrynellula minuta* including molecular data. Our results show the existence of a underestimated diversity in the genus *Amazophrynellula*.

Key words: Amphibian, Anura, Bufonidae, *Amazophrynellula*, Brazil

Resumo

Amazophrynellula é um gênero da família Bufonidae, atualmente representado por três espécies. *Amazophrynellula minuta* é considerada como um complexo de espécies que se distribui por todo o bioma amazônico. Este grupo permanece taxonomicamente problemático, sendo as principais dificuldades a ausência de caracteres diagnósticos na descrição original de *A. minuta*, a inexistência de dados moleculares e comparações taxonômicas detalhadas dos indivíduos, através de sua extensa distribuição. Descrevemos uma nova espécie do gênero *Amazophrynellula*, distribuída no sudeste da região do escudo da Guiana baseados em caracteres morfológicos e moleculares. A nova espécie difere de outras do gênero por apresentar um focinho triangular ligeiramente truncado, textura ventral coberto por muitos grânulos finos, barriga branca coberta por manchas pretas e quinze automorfias moleculares no fragmento do 16S rDNA. As distâncias não corrigidas (p-distâncias) no fragmento 16S rDNA revelaram uma alta divergência entre as outras espécies de *Amazophrynellula* (9%–14%). Adicionalmente providenciamos uma nova diagnose do material topotípico de *Amazophrynellula minuta*, incluindo dados moleculares. Nossos resultados mostram a existência de uma diversidade subestimada dentro do gênero *Amazophrynellula*.

Literature cited

- Ávila, R.W., Carvalho, V.T., Gordo, M., Ribeiro, K. & Morais, D. (2012) A new species of *Amazophrynellula* (Anura: Bufonidae) from southern Amazonia. *Zootaxa*, 3484, 65–74.
- Baum, D.A. & Donoghue, M.J. (1995) Choosing among alternative “phylogenetic” species concepts. *Systematic Botany*, 20, 560–573.
<http://dx.doi.org/10.2307/2419810>
- Bocxlaer, I., Van Loader, S.P., Roelants, K., Biju, S.D., Menegon, M. & Bossuyt, F. (2010) Gradual adaptation toward a range-expansion phenotype initiated the global radiation of toads. *Science*, 327, 679–682.
<http://dx.doi.org/10.1126/science.1181707>
- Cracraft, J. (1983) Species concepts and speciation analysis. *Current Ornithology*, 1, 159–187.
http://dx.doi.org/10.1007/978-1-4615-6781-3_6
- Cruz, C.A.G. & Fusinatto, L.A. (2008) A new species of *Dendrophryniscus*, Jiménez de la Espada, 1871 (Amphibia, Anura, Bufonidae) from the Atlantic Rain Forest of Rio Grande do Sul, Brazil. *South American Journal of Herpetology*, 3, 22–26.
[http://dx.doi.org/10.2994/1808-9798\(2008\)3\[22:ansodj\]2.0.co;2](http://dx.doi.org/10.2994/1808-9798(2008)3[22:ansodj]2.0.co;2)
- Darst, C.R. & Cannatella, D.C. (2004) Novel relationships among hyloid frogs inferred from 12S and 16S mitochondrial DNA sequences. *Molecular Phylogenetics and Evolution*, 31, 462–75.
<http://dx.doi.org/10.1016/j.ympev.2003.09.003>
- Davis, J.I. & Nixon, K.C. (1992) Populations, genetic variation, and the delimitation of phylogenetic species. *Systematic Biology*, 41, 421–435.
- de Queiroz, K. (2007) Species concepts and species delimitation. *Systematic Biology*, 56, 879–886.
- Duellman, W.E. & Mendelson, J.R. (1995) Amphibians and reptiles from northern Departamento Loreto, Peru: Taxonomy and biogeography. *University of Kansas Science Bulletin*, 55, 329–376.
- Duellman, W.E. (1978) The biology of an equatorial herpetofauna in Amazonian Ecuador. *Miscellaneous Publications, University of Kansas Museum of Natural History*, 65, 1–352.
- Felsenstein, J. (1981) Evolutionary trees from DNA sequences: a maximum likelihood approach. *Journal of Molecular Evolution*, 17, 396–376.
<http://dx.doi.org/10.1007/bf01734359>
- Felsenstein, J. (1981) Confidence limits on phylogenies: An approach using the bootstrap. *Evolution*, 39, 783–791.
<http://dx.doi.org/10.2307/2408678>
- Fouquet, A., Gilles, A., Vences, M., Marty, C., Blanc, M. & Gemmell, N.J. (2007) Underestimation of species richness in Neotropical frogs revealed by mtDNA analyses. *PLoS ONE*, 2, e1109.
<http://dx.doi.org/10.1371/journal.pone.0001109>
- Fouquet, A., Recoder, R., Teixeira Jr., M., Cassimiro, J., Amaro, R.C., Camacho, A., Damasceno, R., Carnaval, A.C., Moritz, C. & Rodrigues, M.T. (2012a) Molecular phylogeny and morphometric analyses reveal deep divergence between Amazonian and Atlantic Forest species of *Dendrophryniscus*. *Molecular Phylogenetics and Evolution*, 62, 823–838.
<http://dx.doi.org/10.1016/j.ympev.2011.11.023>
- Fouquet, A., Recoder, R., Teixeira Jr., M., Cassimiro, J., Amaro, R.C., Camacho, A., Damasceno, R., Carnaval, A.C., Moritz, C. & Rodrigues, M.T. (2012b) *Amazonella* Fouquet et al., 2012 (Anura: Bufonidae) junior homonym of *Amazonella* Lundblad, 1931 (Acari: Unionicolidae); proposed replacement by *Amazophrynellula* nom. nov. *Zootaxa*, 3244, 68.
- Frost, D.R. (2013) *Amphibian Species of the World: an Online Reference*. Version 5.4 (9 January 2013) American Museum of Natural History, New York, USA. Available from: <http://research.amnh.org/herpetology/amphibia/> (accessed 20 May 2013)
- Jobb, G. (2008) TREEFINDER version of March 2011, distributed by the author, Munich, Germany. Available from: <http://www.treefinder.de/> (accessed 27 May 2013)
- Hall, T.A. (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series*, 41, 95–98.
- Izecksohn, E. (1968) Nova espécie de *Dendrophryniscus* do Estado do Rio de Janeiro (Amphibia, Salientia). *Revista Brasileira Biologia*, 28, 357–362.
<http://dx.doi.org/10.1590/s0101-81751993000300006>
- Izecksohn, E. (1993) Nova espécie de *Dendrophryniscus* da região amazônica (Amphibia, Anura, Bufonidae). *Revista Brasileira de Zoologia*, 10, 407–412.
<http://dx.doi.org/10.1590/s0101-81751993000300006>
- Lima, A.P., Magnusson, W.E., Menin, M., Erdtmann, L.K., Rodrigues, D.J., Keller, C. & Hödl, W. (2006) *Guia de sapos da Reserva Adolpho Ducke, Amazônia Central—Guide to the frogs of Reserva Adolpho Ducke, Central Amazonian*. Áttema Design Editorial, Manaus, 168 pp.
- Magnusson, W.E. & Hero, J.H. (1991) Predation and the evolution of complex oviposition behaviour in Amazon rainforest frogs. *Oecologia*, 86, 310–318.
<http://dx.doi.org/10.1007/bf00317595>
- McDiarmid, R.W. (1971) Comparative morphology and evolution of frogs of the Neotropical genera *Atelopus*, *Dendrophryniscus*, *Melanophryniscus* and *Oreophrynellula*. *Bulletin of the Los Angeles County Museum of Natural History*, 12, 1–66.

- Melin, D.E. (1941) Contributions to the knowledge of the Amphibia of South America. *Göteborgs Kungl. Vetenskaps- och Vitterhets-samhälles. Handlingar. Serien B, Matematiska och Naturvetenskapliga Skrifter*, 1, 1–71.
- Palumbi, S.R. (1996) Nucleic acids II: the polymerase chain reaction. In: Hillis, D.M., Moritz, C. & Mable, K.B. (Eds.), *Molecular Systematics*. Sinauer & Associates Inc., Sunderland, Massachusetts, pp. 205–247.
- Pramuk, J.B. (2006) Phylogeny of South American *Bufo* (Anura: Bufonidae) inferred from combined evidence. *Zoological Journal of the Linnean Society*, 146, 407–452.
<http://dx.doi.org/10.1111/j.1096-3642.2006.00212.x>
- Pramuk, J.B., Robertson, B., Sites, J.W. & Noonan, B.P. (2008) Around the world in 10 million years: biogeography of the nearly cosmopolitan true toads (Anura: Bufonidae). *Global Ecology and Biogeography*, 17, 72–83.
<http://dx.doi.org/10.1111/j.1466-8238.2007.00348.x>
- Rach, J., DeSalle, R., Sarkar, I.N., Schierwater, B. & Hadrys, H. (2008) Character-based DNA barcoding allows discriminations of genera, species and populations in Odonata. *Proceedings of the Royal Society of London*, 275, 237–247.
<http://dx.doi.org/10.1098/rspb.2007.1290>
- Rodríguez, L.O. & Duellman, W.E. (1994) *Guide to the Frogs of the Iquitos Region. Amazonian Peru*. University of Kansas Natural History Museum, special Publications, 22, 80 pp.
- Sambrook, J., Fritsch, E.F. & Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual, second edition*. Cold Spring Harbor Laboratory Press, Cold Springs Harbor, NY, 1626 pp.
- Tamura, K., Dudley, J., Nei, M. & Kumar, S. (2007) MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Molecular Biology and Evolution*, 24, 1596–1599.
<http://dx.doi.org/10.1093/molbev/msm092>
- Thompson, J.D., Higgins, D.G. & Gibson, T.J. (1996) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position specific gap penalties and weight matrix choice. *Nucleic Acids Research*, 22, 4673–4680.
<http://dx.doi.org/10.1093/nar/22.22.4673>
- Vences, M., Teschke, M., van der Meijden, A., Chiari, Y. & Vieites, D. (2005) Comparative performance of the 16S rRNA gene in DNA barcoding of amphibians. *Frontiers in Zoology*, 2, 7–12.
- Zimmerman, B.L. & Rodrigues, M.T. (1990) Frogs, snakes, and lizards of INPA-WWF reserves near Manaus, Brazil. In: Gentry, A.H. (Eds.), *Four Neotropical Rainforests*. Yale University Press, Connecticut, USA, pp. 426–454.

APPENDIX 1. Specimens examined.

- Amazophrynellabokermanni*—INPA-H 31861, INPA-H 31864, INPA-H 31863, INPA-H 31862, INPA-H 31865, municipality of Juruti, Pará State, Brazil (50 km from type locality).
- Amazophrynellavote*—INPA-H 12256, 12331, 12255, 12342, 12343, 12366, 12267, Cachoeirinha, Madeira River, municipality of Manicoré, INPA-H 12326, Lago do Açaí, Aripuanã River, municipality of Novo Aripuanã, INPA-H 21558, Parque Estadual do Guariba, municipality of Manicoré, INPA-H 27412, 27417–27419, 27421–27423, 27425–27426, Parque Nacional Nascentes do Lago Jari, Tapauá municipality, Amazonas State, Brazil. UFMT-A 11138, Fazenda São Nicolau, municipality of Cotriguaçu, Mato Grosso State, Brazil.
- Amazophrynellaminuta*—INPA-H 32725, INPA-H 32723, INPA-H 32729, INPA-H32730, INPA-H32736, INPA-H32731 (females) and INPA-H 32724, INPA-H32728, INPA-H 32733, INPA-H 32735, INPA-H 32722, INPA-H 32738, INPA-H 32737, INPA-H 32739. INPA-H 32720, INPA-H 32732, INPA-H 32726, INPA-H 32730, INPA-H 32740, INPA-H 32734, INPA-H 32721 (males), Taracuá, right bank of the Uaupés River, Amazonas State, Brazil.

ONLINE SUPPLEMENT 1. Raw measurements of the 57 individuals of *Amazophrynellamanaos* sp. nov. examined in this study.